Cerebral Arterial Fenestrations

DANIEL L. COOKE¹, CHARLES E. STOUT¹, WARREN T. KIM¹, AKASH P. KANSAGRA¹, JOHN PAUL YU¹, AMY GU², NICHOLAS P. JEWELL², STEVEN W. HETTS¹, RANDALL T. HIGASHIDA¹, CHRISTOPHER F. DOWD¹, VAN V. HALBACH¹

¹ Department of Radiology & Biomedical Imaging, University of California; San Francisco, CA, USA

² University of California; Berkeley, CA, USA

Key words: fenestration, aneurysm, subarachnoid hemorrhage, stroke, anatomy

Summary

Arterial fenestrations are an anatomic variant with indeterminate significance. Given the controversy surrounding fenestrations we sought their prevalence within our practice along with their association with other cerebrovascular anomalies.

We retrospectively reviewed 10,927 patients undergoing digital subtraction angiography between 1992 and 2011. Dictated reports were searched for the terms “fenestration” or “fenestrated” with images reviewed for relevance, yielding 228 unique cases. A Medline database search from February 1964 to January 2013 generated 304 citations, 127 cases of which were selected for analysis.

Cerebral arterial fenestrations were identified in 228 patients (2.1%). At least one aneurysm was noted in 60.5% of patients, with an aneurysm arising from the fenestration in 19.6% of patients. Aneurysmal subarachnoid hemorrhage or non-aneurysmal subarachnoid hemorrhage were present in 60.1% and 15.8%, respectively. For the subset of patients with an aneurysm arising directly from a fenestration relative to those patients with an aneurysm not immediately associated with a fenestration, the prevalence of aneurysmal subarachnoid hemorrhage was 66.7% vs. 58.6% ($p = 0.58$). Fenestrations were more often within the posterior circulation (73.2%) than the anterior circulation (24.6%), though there was no difference in the prevalence of aneurysms within these groups (61.1% vs. 60.7%, $p = 1.0$).

Cerebral arterial fenestrations are an anatomic variant more often manifesting at the anterior communicating arterial complex and basilar artery and with no definite pathological relationship with aneurysms.

Introduction

A fenestration is an arterial anatomic variant where a segment of a single vessel divides into at least two channels, each comprising endothelial and muscular layers (they may share the adventitia) that coalesce to a single lumen along its more distal course. Varying series from autopsy, surgical dissection, digital subtraction angiography (DSA), and cross-sectional (computed tomography (CT) and magnetic resonance (MR)) imaging have suggested a widely varying prevalence of fenestrations ranging from 0.7 to 60%¹⁻⁷. This spacious interval largely reflects the small sample sizes, selection biases, and retrospective design of series, in addition to the sensitivity of the varying detection techniques. Advances in DSA, CT, and MR imaging have improved spatial resolution as well as the ability to manipulate anatomy in three-dimensional space with resultant higher prevalence rates of fenestrations noted in contemporary studies relative to their more dated technological equivalents².

The interest in fenestrations, aside from their intrinsic anatomic difference from conventional arteries, is their relationship to neurovascular pathology at large. Many studies have noted the presence of fenestrations in the setting of aneurysms, whether immediately adjacent or remote to one another. Others have noted their presence in the setting of arterial-venous malformations, ischemic stroke, and as well as other cerebrovascular anomalies, pathological or otherwise⁸. Series such as these suggest fenestrations may alter microvascular flow dynamics that may increase the risk of aneurysm formation in addition to their more generalized role

as markers for any pathological vascular manifestation. Others contend that fenestrations are not benign variants, but rather pathological and as such carry a risk of subarachnoid hemorrhage independent of any other identifiable source of bleeding⁹.

Given the controversy surrounding fenestrations, we sought to define their prevalence within our practice along with their varying anatomic locations, association with other cerebrovascular anomalies, and clinical presentation. We also reviewed the literature to note patterns within case series when examined collectively.

Methods

Under permission of the committee on human research (CHR 10-00936), we retrospectively reviewed our database of patients

(10,927) undergoing DSA between 1992 and 2011. Dictated reports were searched for the term “fenestration” or “fenestrated”. The initial query was reviewed and all duplicate medical records (18,437 examinations) removed leaving only unique cases. Three neurointerventional radiologists (DC, CS, and WK) reviewed the images from each of these cases to assess their relevance. Those considered pertinent then underwent more extensive review for demographic, clinical, imaging, and treatment variables.

A search of the Medline database <pubmed.gov> from February 1964 to January 2013 combining the terms “fenestration”, “artery or arterial”, and “brain or cerebral or intracranial” generated 304 citations. These were reviewed for relevance with 127 selected for analysis. Those considered pertinent were then reviewed for demographic, clinical, imaging, and treatment variables. The tabulated data were compared as proportions using the chi-squared test.

Table 1 Fenestration retrospective cohort analysis.

Location	Cases	Male (%)	Age (±SD)	Aneurysm	Aneurysm at site ^{1,3}	ICH ²	A-SAH ¹	NA-SAH ²
Total	228	93 (40.8%)	53 (17.0)	138 (60.5%)	27 (19.6%)	117 (51.3%)	83 (60.1%)	36 (15.8%)
Anterior	56 (24.6%)	21 (37.5%)	50 (18.2)	34 (60.7%)	11 (32.4%)	26 (46.4%)	19 (55.9%)	8 (14.3%)
Posterior	167 (73.2%)	71 (42.5%)	53 (16.3)	102 (61.1%)	15 (14.7%)	90 (53.9%)	64 (62.7%)	27 (16.2%)
Multiple	5 (2.2%)	1 (20.0%)	49 (25.0)	2 (40.0%)	1 (50.0%)	1 (20.0%)	0 (0%)	1 (20.0%)
Internal carotid	3 (1.3%)	1 (33.3%)	52 (48.6)	3 (100%)	0 (0%)	1 (33.3%)	1 (33.3)	0 (0%)
Anterior cerebral	43 (18.9%)	16 (37.2%)	51 (15.4)	26 (61.9%)	10 (38.5%)	21 (48.8%)	16 (61.5%)	6 (14.0%)
Middle cerebral	10 (4.4%)	4 (40.0%)	44 (19.0)	5 (50.0%)	1 (20.0%)	4 (40.0%)	2 (40.0%)	2 (20.0%)
Vertebral	45 (19.7%)	16 (35.6%)	53 (16.6)	24 (53.3%)	2 (8.3%)	23 (51.1%)	17 (70.8%)	6 (13.3%)
Basilar	120 (52.6%)	53 (44.2%)	54 (16.4)	77 (64.2%)	13 (16.9%)	65 (54.2%)	46 (59.7%)	20 (16.7%)
Posterior cerebral	1 (0.4%)	1 (100%)	41 (NA)	1 (100%)	0 (0%)	1 (100%)	1 (100%)	0 (0%)
PICA	1 (0.4%)	1 (100%)	62 (NA)	0 (0%)	0 (0%)	1 (100%)	0 (0%)	1 (100%)

¹ Denominator is number of aneurysms (at given location, if applicable)

² Denominator is total patients (at given location, if applicable)

³ For the subset of patients with an aneurysm arising directly from a fenestration (n = 27) relative to those patients with an aneurysm not immediately associated with a fenestration (n = 111), the prevalence of aneurysmal SAH was 66.7% vs. 58.6% (p = 0.58).

All analyses were performed using the software R and with a probability of significance of 0.05. To calculate 95% confidence intervals a t distribution based on sample size was employed.

Results

Review of 18,437 examinations, representing 10,927 patients yielded 228 unique fenestration cases, generating a prevalence of 2.1%. Table 1 details the general demographic and hemorrhagic details of the fenestration cohort in addition to their locations. The majority (59.2%) of these patients were female and had undergone evaluation within the sixth decade of life (53 ± 17 years). Many (60.5%) of these patients harbored at least one aneurysm at any location within the cerebral vasculature, though a smaller fraction (19.6%) manifested an aneurysm at the actual fenestration site. More than half of all patients (51.3%) presented clinically with intracranial hemorrhage, subarachnoid or otherwise, and of those patients with an aneurysm 60.1% had a history of aneurysmal rupture; 15.8% of patients had a history of a non-aneurysmal subarachnoid hemorrhage. Only five patients had multiple fenestrations.

As to the distribution within the intracranial circulation, nearly three quarters (73.2%) of fenestrations were located within the posterior circulation with the basilar artery being the most frequently (52.6%) affected vessel. Fenestrations were found at other locations, including the internal carotid (1.3%), middle and posterior cerebral (4.4% and 0.4%, respectively), and posterior-inferior cerebral (0.4%) arteries, though certainly more seldom than vertebral, anterior cerebral, and basilar arteries. Those fenestrations found within the anterior cerebral artery, defined as its primary and secondary segments and anterior communicating artery complex, most frequently (38.5%) had intimate aneurysmal association while the basilar artery less often (16.9%) showed such a relationship. For the subset of patients with an aneurysm arising directly from a fenestration ($n = 27$) relative to those patients with an aneurysm not immediately associated with a fenestration ($n = 111$), the prevalence of aneurysmal SAH was 66.7% vs. 58.6% ($p = 0.58$).

Collectively, the mean aneurysm size (Table 2) was 6.74 mm (5.98, 7.50) identical to those arising at the fenestration specifically (mean 6.74 mm 5.31, 8.17). Posterior circulation aneu-

rysms (5.52 mm 3.86, 7.18), in particular those from basilar artery (5.22 mm 3.71, 6.73), were smaller than those within the anterior circulation. As previously stated, fenestrations were more often within the posterior circulation than the anterior circulation (24.6%), though there was no difference in the prevalence of aneurysms within these groups (61.1% vs. 60.7%, $p = 1.0$).

Table 2 Aneurysm size mm [95% CI].

Mean overall	6.74 (n=121) [5.98, 7.50]
Mean A-SAH	6.58 (n=74) [5.77, 7.39]
Mean aneurysm at site	6.74 (n=27) [5.31, 8.17]
Location	
Mean of anterior	6.63 (n=52) [5.71, 7.56]
Mean of posterior	5.52 (n=25) [3.86, 7.18]
Mean of basilar	5.22 (n=18) [3.71, 6.73]

Discussion

The fenestration cohort herein described is the largest of its kind. The prevalence of fenestrations within our retrospective cohort was 2.1%. This is more than the 0.7% prevalence from a series using a comparable DSA technique, though fewer than other series using MR (2.8 – 3.0%), DSA (22.9 – 28%), and CT (3.5 – 12.9%) and significantly less than those reported from autopsy series^{3,4,6,7,9-12}.

To help put our results in an historical context we compiled a comparable number (204) of cases (Table 3) from the literature. These individual and serial reports span more than four decades and reflect the technological changes in kind. They also reflect reporting bias selecting for cases less frequently seen in practice as evidenced by the higher fractions relative to our cohort of fenestrations of the internal carotid (10.3% vs. 1.3%, p -value <0.001), and middle (11.3% vs. 4.4%, p -value 0.01) and posterior (2.0% vs. 0.4%, p -value 0.3) cerebral arteries, and lower fractions within the anterior cerebral (13.7% vs. 18.9% p -value 0.19), basilar (44.1% vs. 52.8%, p -value 0.1), and vertebral (11.8% vs. 19.7%, p -value 0.03) arteries. And just as such exotic locations are over represented so are the relative associations with other pathologies, namely hemorrhage and aneurysms. Within the meta-analysis nearly three quarters (72.5%) of patients had an aneurysm,

60.3% had an aneurysm arising at the site of the fenestration, and 72.1% manifested aneurysmal rupture.

Fenestrations and aneurysms

The relationship between fenestrations and aneurysms is controversial with many series, including our own, reporting a high prevalence of aneurysms in patients with fenestrations, and with other series noting no such relationship^{2,4,6}. Excluding our series, the prevalence of an aneurysm in the setting of a fenestration ranges from 2.5 to 17%^{2,6,10,12,13}. The distinction should be made regarding fenestrations and aneurysms as to whether the two are anatomically associated or remote relative one another². In the setting of fenestrations distant from aneurysms, a higher prevalence of aneurysms might suggest the fenestration represents a marker of more generalized vascular dysfunction, though

there is no evidence describing alterations in genetic expression unique to patients with fenestrations that might put them at such risk. Furthermore, though the number of patients was small⁵, we did not see a trend whereby patients with multiple fenestrations had more aneurysms, suggesting that more fenestrations are not an indicator of more vascular pathology.

The scenario of anatomically associated fenestrations and aneurysms, however, has been explored, particularly as to the changes in fenestrated arterial wall organization and alterations of flow around the fenestration and how both may be involved in aneurysm formation^{3,14-16}. Thinning or absence of the media has been noted at the proximal and distal margins of fenestrations, though similar histological arrangements are noted at other conventional anatomic arterial bifurcation locations. That saccular aneurysms tend to form at such branch points, fenestration-based or otherwise, is well-

Table 3 Fenestration meta-analysis.

Location	Cases	Male	Age (SD)	Aneurysm	Aneurysm at site	ICH	A-SAH	NA-SAH	p-value ^a
Total	204	103 (50.5%)	48 (16.8)	148 (72.5%)	123 (60.3%)	116 (56.9%)	106 (71.6%)	1 (0.5%)	
Anterior ⁴⁵⁻⁹⁶	72 (35.3%)	33 (45.8%)	52 (16.2)	45 (62.5%)	39 (86.7%)	33 (45.8%)	33 (73.3%)	0 (0%)	
Posterior ^{80,95,97-161}	122 (59.8%)	67 (54.9%)	46 (16.4)	94 (77.0%)	78 (83.0%)	75 (61.5%)	66 (70.2%)	1 (0.8%)	
Multiple ^{48,49,51,70,95,122,136,162,163}	9 (4.4%)	3 (33.3%)	37 (19.9)	7 (77.8%)	5 (71.4%)	7 (77.8%)	6 (85.7%)	0 (0%)	
Internal carotid ^{45,47,52,56,57,64,73,75,77-79,84-86,93,164-166}	21 (10.3%) ^a	6 (28.6%)	45 (13.9)	14 (66.7%)	13 (92.9%)	7 (33.3%)	7 (50.0%)	0 (0%)	< 0.001
Anterior cerebral ^{46,51,53,54,59,61,63,65,68-71,74,80,87,90,91,94,167,168}	28 (13.7%) ^a	15 (53.6%)	53 (14.5)	22 (78.6%)	19 (86.4%)	17 (60.7%)	18 (81.8%)	0 (0%)	0.19
Middle cerebral ^{50,55,60,62,67,72,76,81-83,92,96}	23 (11.3%) ^a	12 (52.2%)	56 (18.5)	9 (39.1%)	7 (77.8%)	9 (39.1%)	8 (88.9%)	0 (0%)	0.01
Vertebral ^{80,95,101,114,119,121,132,133,139,145,148,149,154,159,169}	24 (11.8%) ^a	13 (54.2%)	42 (18.6)	9 (37.5%)	3 (33.3%)	13 (54.2%)	8 (88.9%)	0 (0%)	0.03
Basilar ^{80,97-100,102,104-106,108-113,115-117,120,123-126,128-130,134,135,140,141,143,144,146,147,150-152,154,155,157,158,161,170,171}	90 (44.1%) ^a	48 (53.3%)	48 (15.3)	79 (87.8%)	72 (91.4%)	59 (65.6%)	56 (70.9%)	1 (1.1%)	0.1
Posterior cerebral ^{88,103,118,127,142,160}	4 (2.0%) ^a	3 (75.0%)	26 (10.8)	3 (75.0%)	3 (100%)	2 (50.0%)	2 (66.6%)	0 (0%)	0.30
PICA ^{107,137,138,153}	4 (2.0%) ^a	3 (75.0%)	56 (13.4)	3 (75.0%)	0 (0%)	1 (25.0%)	0 (0%)	0 (0%)	0.30

^a Chi-square test comparing retrospective cohort vs. meta cohort.

Figure 1 High magnification oblique 2D DSA (A) and 3D rotational angiographic (B) images of an internal carotid artery terminus fenestration. C) Lateral 2D DSA image of a cavernous internal carotid artery fenestration. Note the M1 middle cerebral segment aneurysm.

Figure 2 High magnification oblique 2D DSA (A) and 3D rotational angiographic (B) images of a middle cerebral artery fenestration. Note the anterior communicating artery aneurysm.

described, though there may be additional risks added by a fenestration such as changes in pressure and flow. Such hemodynamic alterations may not only predispose to aneurysm formation, but also to dissection and/or thromboembolic events^{3,17-26}. Ischemic injury may be related to an absence of anterograde filling of a more expansive distribution or to occlusion of small perforating arteries arising from a fenestrated segment. Arterial perforators are often occult on DSA, though need to be considered if occlusion of the limb of the fenestration is considered part of treatment^{27,28}. These small branches, best visualized on microsurgical dissection, are variable in their origins and may arise from either the smaller or larger of the fenestration limbs.

Fenestrations and location

We recognize the limitation of endowing equal meaning to all fenestrations. These variants may arise in multiple locations and can have vastly different apertures, limb lengths, and relationships with perforating vessels to which they might give rise. It may be that only a subset(s) of fenestrations carry greater risk of bleeding and/or aneurysmal formation. We noted fenestrations at multiple locations throughout the cerebral vasculature, though with more limited representation at the internal carotid (0.03%; Figure 1), middle cerebral (0.9%; Figure 2), posterior cerebral (0.01%; Figure 3), vertebral (0.4% (Figure 3) and posterior-inferior cerebellar (0.01%) arteries. Given such scarci-

Figure 3 A) High magnification lateral 2D DSA image of a cervical segment vertebral artery fenestration. B) High magnification Waters projection 2D DSA image of a left P1 posterior communicating artery fenestration. C,D) High magnification Waters 2D DSA and 3D rotational images of a basilar artery terminus fenestration.

ty, we focused our discussion on the two most frequently encountered fenestration locations: the anterior communicating arterial complex (Figure 4) and the basilar artery (Figure 5).

As with other morphological variations, arteries closer to the anatomic midline more frequently manifest fenestrations, namely the an-

terior communicating and basilar arteries^{3,5,8,16,24,29-32}. Fenestration of the anterior cerebral artery is rare with only 0.4% prevalence within our cohort and with other case series ranging from 0.1 to 7.2%^{11,19,27,30,33-35}, though if the anterior communicating artery is included the prevalence increases significantly rising up

to 40% by some accounts. The increased prevalence of fenestrations within the anterior communicating artery mirrors the relatively higher rate of fenestration within the basilar artery, both midline vascular structures, and supporting their etiology as one of abnormal embryonic arterial fusion^{2-4,8,11,19,33-39}. The anterior cerebral-anterior communicating artery complex was the most common site within our cohort to have an aneurysm arise directly from the fenestration (38.5%) with a similar fraction (27.6%) reported by others for aneurysms arising from A1 fenestrations³¹. The relatively increased aneurysm-fenestration co-location may in part be related to the manner in which the anterior communicating region is interrogated during 2D DSA. The standard anterior-posterior and lateral projections used in DSA are limited in their ability to resolve aberrant anatomy of the anterior communication region. In the setting of an aneurysm within this region, a finding easier to make on such routine AP and lateral projections, additional oblique projections (Figure 1) are performed to better delineate the aneurysmal architecture often facilitated by 3D rotational angiography^{11,33}. It is these tailored projections that will more often reveal such anatomic variants. For those fenestrations arising from the A1 segment special consideration must be paid to the presence of perforating branches supplying the basal ganglia and optic chiasm if vessel sacrifice is necessitated. Autopsy series have noted the predilection for perforators to arise from the more lateral aspects of the A1, though these vessels may arise from either or both limbs of a fenestration²⁷.

Fenestrations of the basilar artery are thought to arise from congenital non-fusion of the paired longitudinal neural arteries and are rare with only 1.1% prevalence within our cohort and from other case series ranging from 0.3 to 6.0%^{2-4,7,8,14,16,40-43}. Of all locations within our cohort, the basilar artery occurred most frequently (52.6%) with a similar relationship noted within the literature reviewed herein (44.1%). This finding is thought to be in part related to the relative ease of their detection using standard DSA projections. Juxtaposed against the difficulty in resolving the anterior communicating complex with routine 2D DSA, the basilar artery is nearly always larger in diameter and, more importantly, in a more orthogonal relationship with the image intensifier using either Townes or Waters projections

Figure 4 High magnification oblique 2D DSA (A) and 3D rotational angiographic (B) images of an anterior communicating artery fenestration.

(Figure 2). Fenestrations may occur anywhere along the length of the basilar artery, though they are most frequently noted at the proximal extent adjacent to the vertebral-basilar junction⁸. As such this location has been noted as a hotspot for aneurysm formation, with up to 70% of vertebral-basilar junction aneurysms having an associated fenestration, though such aneurysms remain rare overall with fewer than 80 reported in the literature^{3,28,44}.

There are limitations to this descriptive review imposed by its retrospective methodology. The source population, patients undergoing DSA (primarily 2D) at a large tertiary care medical center, certainly introduces bias and in turn limits any conclusion regarding the fenestration

Figure 5 A) Waters projection 2D DSA image of a proximal basilar artery fenestration. B) Townes projection 2D DSA image of a proximal basilar artery fenestration with anteriorly and posteriorly directed aneurysms arising from the fenestration. C,D) High magnification oblique 2D DSA images of the fenestration-related aneurysms.

tration prevalence within the general population. Furthermore, the use of document review to screen for cases, as opposed to prospectively reviewing all angiograms, ensures many fenestrations will have been missed. Given the controversy surrounding fenestrations, operators may have made note of them more frequently in the setting of hemorrhagic or aneurysmal pathology, while more permissive in documentation of cases where no such pathology existed. We certainly see such reporting bias within

the literature. Collectively, these case series (Table 3) do not reflect the natural history of fenestrations, but rather the attention to an incompletely understood phenomenon at extreme clinical spectra. No doubt some of that reporting bias is here as well and as such no conclusions of causation are made between fenestrations, aneurysms, and/or intracerebral hemorrhage.

Future studies on the topic would ideally be prospective in nature and from large, more

generalized populations (e.g. patients undergoing cross-sectional imaging in an emergency department setting). Such studies have been performed, though with cohorts of hundreds of patients leaving them relatively underpowered to answer questions concerning such rare phenomena. As it stands, declarations of causation are invalid, though we would be remiss to not stress the value in identifying and reporting fenestrations as part of clinical practice particularly when an aneurysm is in play as it may affect treatment. Neuroradiologists and neurointerventional surgeons are encouraged to note fenestrations in practice given the associ-

ations described herein and, if for no other reason, to help improve our overall understanding of neurovascular pathology.

Conclusions

Cerebral arterial fenestrations are a relatively rare anatomic variant and remain one of indeterminate and likely heterogeneous clinical significance. They more often manifest in the anterior communicating arterial complex and basilar artery and have no definite pathological relationship with aneurysms.

References

- 1 Dey M, Awad IA. Fenestration of supraclinoid internal carotid artery and associated aneurysm: embryogenesis, recognition, and management. *World Neurosurg.* 2011; 76 (6): 592.e1-5. doi: 10.1016/j.wneu.2011.04.019.
- 2 Bożek P, Pilch-Kowalczyk J, Kluczevska E, et al. Detection of cerebral artery fenestrations by computed tomography angiography. *Neurol Neurochir Pol.* 2012; 46 (3): 239-244. doi: 10.5114/ninp.2012.29132.
- 3 Arráez-Aybar LA, Villar-Martin A, Poyatos-Ruiperez C, et al. Prevalence of fenestrated basilar artery with magnetic resonance angiography: a transversal study. *Surg Radiol Anat.* 2013; 35 (6): 487-93. doi: 10.1007/s00276-012-1053-5.
- 4 Bayrak AH, Senturk S, Akay HO, et al. The frequency of intracranial arterial fenestrations: a study with 64-detector CT-angiography. *Eur J Radiol.* 2011; 77 (3): 392-396. doi: 10.1016/j.ejrad.2009.09.015.
- 5 Dimmick SJ, Faulder KC. Fenestrated anterior cerebral artery with associated arterial anomalies. Case reports and literature review. *Interv Neuroradiol.* 2008; 14 (4): 441-445.
- 6 Sun ZK, Li M, Li MH, et al. Fenestrations accompanied by intracranial aneurysms assessed with magnetic resonance angiography. *Neurol India.* 2012; 60 (1): 45-49. doi: 10.4103/0028-3886.93588.
- 7 Uchino A, Saito N, Okada Y, et al. Fenestrations of the intracranial vertebrobasilar system diagnosed by MR angiography. *Neuroradiology.* 2012; 54 (5): 445-450. doi: 10.1007/s00234-011-0903-x.
- 8 Vasović L, Trandafilović M, Jovanović I, et al. Human basilar artery abnormalities in the prenatal and postnatal period. *World Neurosurg.* 2013; 79 (3-4): 593.e15-23. doi: 10.1016/j.wneu.2012.06.033.
- 9 Hudák I, Lenzšér G, Lunenkova V, et al. Cerebral arterial fenestrations: a common phenomenon in unexplained subarachnoid haemorrhage. *Acta Neurochir (Wien).* 2013; 155 (2): 217-222. doi: 10.1007/s00701-012-1587-7.
- 10 Sanders WP, Sorek PA, Mehta BA. Fenestration of intracranial arteries with special attention to associated aneurysms and other anomalies. *Am J Neuroradiol.* 1993; 14 (3): 675-680.
- 11 van Rooij SB, van Rooij WJ, Sluzewski M, et al. Fenestrations of intracranial arteries detected with 3D rotational angiography. *Am J Neuroradiol.* 2009; 30 (7): 1347-1350. doi: 10.3174/ajnr.A1563.
- 12 Bharatha A, Aviv RI, White J, et al. Intracranial arterial fenestrations: frequency on CT angiography and association with other vascular lesions. *Surg Radiol Anat.* 2008; 30 (5): 397-401. doi: 10.1007/s00276-008-0340-7.
- 13 Uchino A, Saito N, Okada Y, et al. Fenestrations of the intracranial vertebrobasilar system diagnosed by MR angiography. *Neuroradiology.* 2012; 54 (5): 445-450. doi: 10.1007/s00234-011-0903-x.
- 14 Tsuei YS, Matsumoto Y, Ohta M, et al. Vertebrobasilar junction fenestration with dumbbell-shaped aneurysms formation: computational fluid dynamics analysis. *Surg Neurol.* 2009; 72 Suppl 2: S11-9. doi: 10.1016/j.surneu.2009.05.026.
- 15 Futami K, Yamashita J, Higashi S. Do cerebral aneurysms originate at the site of medial defects? Microscopic examinations of experimental aneurysms at the fenestration of the anterior cerebral artery in rats. *Surg Neurol.* 1998; 50 (2): 141-146. doi: 10.1016/S0090-3019(97)00154-7.
- 16 De Caro R, Serafini MT, Galli S, et al. Anatomy of segmental duplication in the human basilar artery. Possible site of aneurysm formation. *Clin Neuropathol.* 1995; 14 (6): 303-309.
- 17 Chen YY, Chang FC, Hu HH, et al. Fenestration of the supraclinoid internal carotid artery associated with aneurysm and ischemic stroke. *Surg Neurol.* 2007; 68 Suppl 1: S60-3; discussion S3. doi: 10.1016/j.surneu.2007.05.051.
- 18 Jeong SK, Kwak HS, Cho YI. Middle cerebral artery fenestration in patients with cerebral ischemia. *J Neurol Sci.* 2008; 275 (1-2): 181-184. doi: 10.1016/j.jns.2008.07.037.
- 19 San-Galli F, Leman C, Kien P, et al. Cerebral arterial fenestrations associated with intracranial saccular aneurysms. *Neurosurgery.* 1992; 30 (2): 279-283. doi: 10.1227/00006123-199202000-00026.
- 20 Ahn JY, Kim OJ, Joo YJ, et al. Fenestration of the internal carotid artery associated with arterial dissections. *J Clin Neurosci.* 2003; 10 (2): 257-260. doi: 10.1016/S0967-5868(02)00331-4.
- 21 Bernard TJ, Mull BR, Handler MH, et al. An 18-year-old man with fenestrated vertebral arteries, recurrent stroke and successful angiographic coiling. *J Neurol Sci.* 2007; 260 (1-2): 279-282. doi: 10.1016/j.jns.2007.04.032.
- 22 Hayashi T, Hirose Y, Sagoh M, et al. Spontaneous occlusion of ruptured vertebral artery dissection at the extradural fenestration associated with extradural ori-

- gin of the posterior inferior cerebellar artery--case report. *Neurol Med Chir (Tokyo)*. 2000; 40 (3): 164-168. doi: 10.2176/nmc.40.164.
- 23 Sim E, Vaccaro AR, Berzlanovich A, et al. Fenestration of the extracranial vertebral artery: review of the literature. *Spine (Phila Pa 1976)*. 2001; 26 (6): E139-142. doi: 10.1097/00007632-200103150-00007.
 - 24 Berry AD 3rd, Kepes JJ, Wetzel MD. Segmental duplication of the basilar artery with thrombosis. *Stroke*. 1988; 19 (2): 256-260. doi: 10.1161/01.STR.19.2.256.
 - 25 Kloska SP, Schlegel PM, Sträter R, et al. Causality of pediatric brainstem infarction and basilar artery fenestration? *Pediatr Neurol*. 2006; 35 (6): 436-438. doi: 10.1016/j.pediatrneurol.2006.06.013.
 - 26 Zhang QJ, Kobayashi S, Gibo H, et al. Vertebrobasilar junction fenestration associated with dissecting aneurysm of intracranial vertebral artery. *Stroke*. 1994; 25 (6): 1273-1275. doi: 10.1161/01.STR.25.6.1273.
 - 27 Aktüre E, Arat A, Niemann DB, et al. Bilateral A1 fenestrations: Report of two cases and literature review. *Surg Neurol Int*. 2012; 3: 43. doi: 10.4103/2152-7806.94928.
 - 28 Tanaka S, Tokimura H, Makiuchi T, et al. Clinical presentation and treatment of aneurysms associated with basilar artery fenestration. *J Clin Neurosci*. 2012; 19 (3): 394-401. doi: 10.1016/j.jocn.2011.04.043.
 - 29 Black SP, Ansbacher LE. Saccular aneurysm associated with segmental duplication of the basilar artery. A morphological study. *J Neurosurg*. 1984; 61 (6):1005-1008. doi: 10.3171/jns.1984.61.6.1005.
 - 30 Dey M, Awad IA. Fenestration of supraclinoid internal carotid artery and associated aneurysm: embryogenesis, recognition, and management. *World Neurosurg*. 2011; 76 (6): 592.e1-5. doi: 10.1016/j.wneu.2011.04.019.
 - 31 Friedlander RM, Oglivly CS. Aneurysmal subarachnoid hemorrhage in a patient with bilateral A1 fenestrations associated with an azygos anterior cerebral artery. Case report and literature review. *J Neurosurg*. 1996; 84 (4): 681-684. doi: 10.3171/jns.1996.84.4.0681.
 - 32 Ito J, Washiyama K, Kim CH, et al. Fenestration of the anterior cerebral artery. *Neuroradiology*. 1981; 21 (5): 277-280. doi: 10.1007/BF02100160.
 - 33 de Gast AN, van Rooij WJ, Sluzewski M, et al. Fenestrations of the anterior communicating artery: incidence on 3D angiography and relationship to aneurysms. *Am J Neuroradiol*. 2008; 29 (2): 296-298. doi: 10.3174/ajnr.A0807.
 - 34 de Oliveira JG, du Mesnil de Rochemont R, Beck J, et al. A rare anomaly of the anterior communicating artery complex hidden by a large broad-neck aneurysm and disclosed by three-dimensional rotational angiography. *Acta Neurochir (Wien)*. 2008; 150 (3): 279-284; discussion 84. doi: 10.1007/s00701-007-1401-0.
 - 35 Matsumura M, Nojiri K. Ruptured anterior communicating artery aneurysms associated with fenestration of the anterior cerebral artery. *Surg Neurol*. 1984; 22 (4): 371-376. doi: 10.1016/0090-3019(84)90142-3.
 - 36 Fujimura M, Seki H, Sugawara T, et al. Anomalous internal carotid artery-anterior cerebral artery anastomosis associated with fenestration and cerebral aneurysm. *Neurol Med Chir (Tokyo)*. 1996; 36 (4): 229-233. doi: 10.2176/nmc.36.229.
 - 37 Eustacchio S, Klein GE, Pendl G. Ruptured vertebrobasilar junction aneurysm associated with basilar artery fenestration. *Acta Neurochir (Wien)*. 1997; 139 (10): 923-927. doi: 10.1007/BF01411300.
 - 38 Tran-Dinh HD, Soo YS, Jayasinghe LS. Duplication of the vertebro-basilar system. *Australas Radiol*. 1991; 35 (3): 220-224. doi: 10.1111/j.1440-1673.1991.tb03012.x.
 - 39 Namiki J, Doumoto Y. Microsurgically critical anomaly of the anterior communicating artery complex during the pterional approach to a ruptured aneurysm: double fenestration of the proximal A2 segments. *Neurol Med Chir (Tokyo)*. 2003; 43 (6): 304-307. doi: 10.2176/nmc.43.304.
 - 40 Andrews BT, Brant-Zawadzki M, Wilson CB. Variant aneurysms of the fenestrated basilar artery. *Neurosurgery*. 1986; 18 (2): 204-207. doi: 10.1227/00006123-198602000-00017.
 - 41 Bentura JE, Figueiredo EG, de Monaco BA, et al. Vertebrobasilar artery junction aneurysm associated with fenestration. *Arq Neuropsiquiatr*. 2010; 68 (2): 312-314. doi: 10.1590/S0004-282X2010000200031.
 - 42 Dodevski A, Lazareska M, Tosovska-Lazarova D, et al. Basilar artery fenestration. *Folia Morphol (Warsz)*. 2011; 70 (2): 80-83.
 - 43 Tasker AD, Byrne JV. Basilar artery fenestration in association with aneurysms of the posterior cerebral circulation. *Neuroradiology*. 1997; 39 (3): 185-189. doi: 10.1007/s002340050389.
 - 44 Tanaka M, Kikuchi Y, Ouchi T. Neuroradiological analysis of 23 cases of basilar artery fenestration based on 2280 cases of MR angiographies. *Interv Neuroradiol*. 2006; 12 (Suppl 1): 39-44.
 - 45 Ahn JY, Kim OJ, Joo YJ, et al. Fenestration of the internal carotid artery associated with arterial dissections. *J Clin Neurosci*. 2003; 10 (2): 257-260. doi: 10.1016/S0967-5868(02)00331-4.
 - 46 Aktüre E, Arat A, Niemann DB, et al. Bilateral A1 fenestrations: Report of two cases and literature review. *Surg Neurol Int*. 2012; 3: 43. doi: 10.4103/2152-7806.94928.
 - 47 Banach MJ, Flamm ES. Supraclinoid internal carotid artery fenestration with an associated aneurysm. Case report. *J Neurosurg*. 1993; 79 (3): 438-441. doi: 10.3171/jns.1993.79.3.0438.
 - 48 Chen YY, Chang FC, Hu HH, et al. Fenestration of the supraclinoid internal carotid artery associated with aneurysm and ischemic stroke. *Surg Neurol*. 2007; 68 Suppl 1: S60-3; discussion S3.
 - 49 de Oliveira JG, du Mesnil de Rochemont R, et al. A rare anomaly of the anterior communicating artery complex hidden by a large broad-neck aneurysm and disclosed by three-dimensional rotational angiography. *Acta Neurochir (Wien)*. 2008; 150 (3): 2792-84; discussion 284. doi: 10.1007/s00701-007-1401-0.
 - 50 Deruty R, Pelissou-Guyotat I, Mottolese C, et al. Fenestration of the middle cerebral artery and aneurysm at the site of the fenestration. *Neurol Res*. 1992; 14 (5): 421-424.
 - 51 Dimmick SJ, Faulder KC. Fenestrated anterior cerebral artery with associated arterial anomalies. Case reports and literature review. *Interv Neuroradiol*. 2008; 14 (4): 441-445.
 - 52 Findlay JM, Chui M, Muller PJ. Fenestration of the supraclinoid internal carotid artery. *Can J Neurol Sci*. 1987; 14 (2): 159-161.
 - 53 Friedlander RM, Oglivly CS. Aneurysmal subarachnoid hemorrhage in a patient with bilateral A1 fenestrations associated with an azygos anterior cerebral artery. Case report and literature review. *J Neurosurg*. 1996; 84 (4): 681-684. doi: 10.3171/jns.1996.84.4.0681.
 - 54 Fujimura M, Seki H, Sugawara T, et al. Anomalous internal carotid artery-anterior cerebral artery anastomosis associated with fenestration and cerebral aneurysm. *Neurol Med Chir (Tokyo)*. 1996; 36 (4): 229-233. doi: 10.2176/nmc.36.229.
 - 55 Gailloud P, Albayram S, Fasel JH, et al. Angiographic and embryologic considerations in five cases of middle cerebral artery fenestration. *Am J Neuroradiol*. 2002; 23 (4): 585-587.
 - 56 Harnier S, Harzheim A, Limmroth V, et al. Duplication of the common carotid artery and the ipsilateral

- vertebral artery with a fenestration of the contralateral common carotid artery. *Neurol India*. 2008; 56 (4): 491-493. doi: 10.4103/0028-3886.44825.
- 57 Hattori T, Kobayashi H. Fenestration of the supraclinoid internal carotid artery associated with carotid bifurcation aneurysm. *Surg Neurol*. 1992; 37 (4): 284-288. doi: 10.1016/0090-3019(92)90154-F.
 - 58 Ichikawa T, Miyachi S, Izumi T, et al. Fenestration of a supraclinoid internal carotid artery associated with dual aneurysms: case report. *Neurosurgery*. 2011 Oct;69(4):E1005-8; discussion E1009. doi: 10.1227/NEU.0b013e318223b613.
 - 59 Ihara S, Uemura K, Tsukada A, et al. Aneurysm and fenestration of the azygos anterior cerebral artery--case report. *Neurol Med Chir (Tokyo)*. 2003; 43 (5): 246-249. doi: 10.2176/nmc.43.246.
 - 60 Ito J, Maeda H, Inoue K, et al. Fenestration of the middle cerebral artery. *Neuroradiology*. 1977; 13 (1): 37-39. doi: 10.1007/BF00335032.
 - 61 Ito J, Washiyama K, Kim CH, et al. Fenestration of the anterior cerebral artery. *Neuroradiology*. 1981; 21 (5): 277-280. doi: 10.1007/BF02100160.
 - 62 Jeong SK, Kwak HS, Cho YI. Middle cerebral artery fenestration in patients with cerebral ischemia. *J Neurol Sci*. 2008 Dec 15;275(1-2):181-4. doi: 10.1016/j.jns.2008.07.037.
 - 63 Kobayashi S, Yuge T, Sugita Y, et al. Azygos anterior cerebral artery aneurysm associated with fenestration of the anterior cerebral artery. *Kurume Med J*. 1986; 33 (3): 149-153. doi: 10.2739/ikumemedj.33.149.
 - 64 Koenigsberg RA, Zito JL, Patel M, et al. Fenestration of the internal carotid artery: a rare mass of the hypotympanum associated with persistence of the stapedial artery. *Am J Neuroradiol*. 1995; 16 (4 Suppl): 908-910.
 - 65 Koh JS, Kim EJ, Lee SH, et al. Ruptured aneurysm arising from the distal end of a proximal a(1) fenestration: case report and review of the literature. *J Korean Neurosurg Soc*. 2009; 45 (1): 43-45. doi: 10.3340/jkns.2009.45.1.43. doi: 10.3340/jkns.2009.45.1.43.
 - 66 Koh JS, Lee SH, Bang JS, et al. Three-dimensional angiographic demonstration of plexiform fenestrations of the proximal anterior cerebral artery associated with a ruptured aneurysm. *J Korean Neurosurg Soc*. 2008; 44 (5): 338-340. doi: 10.3340/jkns.2008.44.5.338.
 - 67 Lazar ML, Bland JE, North RR, et al. Middle cerebral artery fenestration. *Neurosurgery*. 1980; 6 (3): 297-300. doi: 10.1227/00006123-198003000-00014.
 - 68 Mantatzis M, Kizilkilic O, Albayram S, et al. Endovascular treatment of aneurysms associated with fenestrated A1 segment of anterior cerebral artery: report of two cases. *J Neuroimaging*. 2011; 21 (2): 165-169. doi: 10.1111/j.1552-6569.2009.00410.x.
 - 69 Matsumura M, Nojiri K. Ruptured anterior communicating artery aneurysms associated with fenestration of the anterior cerebral artery. *Surg Neurol*. 1984; 22 (4): 371-376. doi: 10.1016/0090-3019(84)90142-3.
 - 70 Minakawa T, Kawamata M, Hayano M, et al. Aneurysms associated with fenestrated anterior cerebral arteries. Report of four cases and review of the literature. *Surg Neurol*. 1985; 24 (3): 284-288. doi: 10.1016/0090-3019(85)90040-0.
 - 71 Mitsuhashi T, Sakamoto S, Kiura Y, et al. Endovascular coil embolization for ruptured kissing aneurysms associated with A1 fenestration. *Surg Neurol Int*. 2011;2:85. doi: 10.4103/2152-7806.82251.
 - 72 Nakamura H, Takada A, Hide T, et al. Fenestration of the middle cerebral artery associated with an aneurysm--case report. *Neurol Med Chir (Tokyo)*. 1994; 34 (8): 555-557. doi: 10.2176/nmc.34.555.
 - 73 Nakamura H, Yamada H, Nagao T, et al. Fenestration of the internal carotid artery associated with an ischemic attack--case report. *Neurol Med Chir (Tokyo)*. 1993; 33 (5): 306-308. doi: 10.2176/nmc.33.306.
 - 74 Namiki J, Doumoto Y. Microsurgically critical anomaly of the anterior communicating artery complex during the pterional approach to a ruptured aneurysm: double fenestration of the proximal A2 segments. *Neurol Med Chir (Tokyo)*. 2003; 43 (6): 304-307. doi: 10.2176/nmc.43.304.
 - 75 Ng PP, Steinfert B, Stoodley MA. Internal carotid artery fenestration with dual aneurysms. Case illustration. *J Neurosurg*. 2006; 104 (6): 979. doi: 10.3171/jns.2006.104.6.979.
 - 76 Nussbaum ES, Defillo A, Janjua TM, et al. Fenestration of the middle cerebral artery with an associated ruptured aneurysm. *J Clin Neurosci*. 2009; 16 (6): 845-847. doi: 10.1016/j.jocn.2008.08.039.
 - 77 Onoda K, Ono S, Tokunaga K, et al. Fenestration of the supraclinoid internal carotid artery with associated aneurysm. *Neurol Med Chir (Tokyo)*. 2008; 48 (3): 118-120. doi: 10.2176/nmc.48.118.
 - 78 Park SH, Lee CY. Supraclinoid internal carotid artery fenestration harboring an unruptured aneurysm and another remote ruptured aneurysm: case report and review of the literature. *J Cerebrovasc Endovasc Neurosurg*. 2012; 14 (4): 295-299. doi: 10.7461/jcen.2012.14.4.295.
 - 79 Plumb AA, Herwadkar A, Pickett G. Incidental finding of fenestration of the supraclinoid internal carotid artery with appearances on magnetic resonance angiography. *Surg Radiol Anat*. 2010; 32 (2): 165-169. doi: 10.1007/s00276-009-0555-2.
 - 80 San-Galli F, Leman C, Kien P, et al. Cerebral arterial fenestrations associated with intracranial saccular aneurysms. *Neurosurgery*. 1992; 30 (2): 279-283. doi: 10.1097/00006123-199202000-00026.
 - 81 Schmieder K, Hardenack M, Harders A. Proximal long fenestration associated with an internal carotid artery aneurysm. Case illustration. *J Neurosurg*. 1997; 86 (4): 733. doi: 10.3171/jns.1997.86.4.0733.
 - 82 Seo BS, Lee YS, Lee JH, et al. Mechanical thrombolysis using coil in acute occlusion of fenestrate m1 segment. *J Cerebrovasc Endovasc Neurosurg*. 2012; 14 (2): 108-112. doi: 10.7461/jcen.2012.14.2.108.
 - 83 Sim KB, Lee CS, Park JC, et al. Cerebral aneurysm in the long fenestration at the middle portion of m1 segment. *J Korean Neurosurg Soc*. 2010; 48 (5): 434-437. doi: 10.3340/jkns.2010.48.5.434.
 - 84 Stojanov D, Bosnjakovic P, Ristic S, et al. Multichannel fenestration of the petrous and cavernous segments of the internal carotid artery associated with two small carotid canals. *Neurol Sci*. 2013; 34 (2): 247-250. doi: 10.1007/s10072-012-0995-z.
 - 85 Takano S, Saitoh M, Miyasaka Y, et al. Fenestration of the intracranial internal carotid artery--case report. *Neurol Med Chir (Tokyo)*. 1991; 31 (11): 740-742. doi: 10.2176/nmc.31.740.
 - 86 Tourdias T, Berge J, Menegon P, et al. Fenestration of the internal carotid artery mimicking floating thrombus on CT and MR angiography. *Neurology*. 2011; 76 (21): 1846. doi: 10.1212/WNL.0b013e31821ccd28.
 - 87 Tran-Dinh HD, Dorsch NW, Soo YS. Ectasia and fenestration of the anterior cerebral artery associated with persistent trigeminal artery: case report. *Neurosurgery*. 1992; 31 (1): 125-128. doi: 10.1227/00006123-199207000-00019.
 - 88 Tripathi M, Goel V, Padma MV, et al. Fenestration of the posterior communicating artery. *Neurol India*. 2003; 51 (1): 75-76.
 - 89 Uchino A, Saito N, Okada Y, et al. Duplicate origin and fenestration of the middle cerebral artery on MR angiography. *Surg Radiol Anat*. 2012; 34 (5): 401-404. doi: 10.1007/s00276-012-0936-9.
 - 90 Wong GK, Wang K, Yu SC, et al. A rare anatomical

- variant: median anterior cerebral artery fenestration associated with an azygous infra-optic anterior cerebral artery. *J Clin Neurosci*. 2010; 17 (11): 1434-1436. doi: 10.1016/j.jocn.2010.02.016.
- 91 Yamada T, Inagawa T, Takeda T. Ruptured aneurysm at the anterior cerebral artery fenestration. Case report. *J Neurosurg*. 1982; 57 (6): 826-828. doi: 10.3171/jns.1982.57.6.0826.
 - 92 Yamaguchi S, Ito O, Suzuki S. Coil embolization of a ruptured aneurysm arising from a middle cerebral artery fenestration--case report. *Neurol Med Chir (Tokyo)*. 2010; 50 (3): 213-216. doi: 10.2176/nmc.50.213.
 - 93 Yock DH. Fenestration of the supraclinoid internal carotid artery with rupture of associated aneurysm. *Am J Neuroradiol*. 1984; 5 (5): 634-636.
 - 94 Yoshida M, Ezura M, Sasaki K, et al. Endovascular repair of ruptured aneurysm arising from fenestration of the horizontal segment of the anterior cerebral artery. *Neurol Med Chir (Tokyo)*. 2012; 52 (12): 924-927. doi: 10.2176/nmc.52.924.
 - 95 Uchino A, Kato A, Abe M, et al. Association of cerebral arteriovenous malformation with cerebral arterial fenestration. *Eur Radiol*. 2001; 11 (3): 493-496. doi: 10.1007/s003300000640.
 - 96 Ueda T, Goya T, Kinoshita K, et al. [Multiple anomalies of cerebral vessels. A case of multiple aneurysms associated with fenestration of the middle cerebral artery and persistent primitive trigeminal artery]. *No Shinkei Geka*. 1984; 12 (4): 531-536.
 - 97 Abe T, Fujita H, Higano S, et al. A saccular aneurysm arising from the fenestration of the basilar artery--a case report. *Kurume Med J*. 1987; 34 (4): 203-206. doi: 10.2739/kurumemedj.34.203.
 - 98 Albanese E, Russo A, Ulm AJ. Fenestrated vertebro-basilar junction aneurysm: diagnostic and therapeutic considerations. *J Neurosurg*. 2009; 110 (3): 525-529. doi: 10.3171/2008.9.JNS08170.
 - 99 Alurkar A, Karanam LS, Oak S. Endovascular treatment of ruptured saccular aneurysm from basilar artery fenestration. *Neurol India*. 2012; 60 (6): 682-683. doi: 10.4103/0028-3886.105227.
 - 100 Andrews BT, Brant-Zawadzki M, Wilson CB. Variant aneurysms of the fenestrated basilar artery. *Neurosurgery*. 1986; 18 (2): 204-207. doi: 10.1227/00006123-198602000-00017.
 - 101 Arai K, Endo S, Hirashima Y, et al. Posterior inferior cerebellar artery aneurysm associated with fenestration of the vertebral artery--case report. *Neurol Med Chir (Tokyo)*. 1989; 29 (1): 29-31. doi: 10.2176/nmc.29.29.
 - 102 Arráez-Aybar LA, Villar-Martin A, Poyatos-Ruiperez C, et al. Prevalence of fenestrated basilar artery with magnetic resonance angiography: a transversal study. *Surg Radiol Anat*. 2013; 35 (6): 487-493. doi: 10.1007/s00276-012-1053-5.
 - 103 Baba S, Fukuda Y, Mizota S, et al. Fusiform aneurysm associated with fenestration of the posterior communicating artery. *Neurol Med Chir (Tokyo)*. 2010; 50 (7): 568-570. doi: 10.2176/nmc.50.568.
 - 104 Bentura JE, Figueiredo EG, de Monaco BA, et al. Vertebrobasilar artery junction aneurysm associated with fenestration. *Arq Neuropsiquiatr*. 2010; 68 (2): 312-314. doi: 10.1590/S0004-282X2010000200031.
 - 105 Berry AD 3rd, Kepes JJ, Wetzel MD. Segmental duplication of the basilar artery with thrombosis. *Stroke*. 1988; 19 (2): 256-260. doi: 10.1161/01.STR.19.2.256.
 - 106 Black SP, Ansbacher LE. Saccular aneurysm associated with segmental duplication of the basilar artery. A morphological study. *J Neurosurg*. 1984; 61 (6): 1005-1008. doi: 10.3171/jns.1984.61.6.1005.
 - 107 Cho YD, Han MH, Lee JY. Double origin of the posterior inferior cerebellar artery with juxta-proximal fenestration of caudal component. *Surg Radiol Anat*. 2011; 33 (3): 271-273. doi: 10.1007/s00276-010-0747-9.
 - 108 Eustacchio S, Klein GE, Pendl G. Ruptured vertebro-basilar junction aneurysm associated with basilar artery fenestration. *Acta Neurochir (Wien)*. 1997; 139 (10): 923-927. doi: 10.1007/BF01411300.
 - 109 Ezaki Y, Kazekawa K, Tsutsumi K, et al. A vertebro-basilar junction aneurysm associated with fenestration treated by intra-aneurysmal embolization. *Acta Neurochir (Wien)*. 2003; 145 (9): 807-808; discussion 808-809. doi: 10.1007/s00701-003-0070-x.
 - 110 Fabrikant H. Basilar artery fenestration with an associated aneurysm: case report. *Australas Radiol*. 1988; 32 (1): 127-130. doi: 10.1111/j.1440-1673.1988.tb02704.x.
 - 111 Foroni LH, Figueiredo EG, Teixeira MJ, et al. Saccular aneurysms at middle basilar trunk fenestration. *Arq Neuropsiquiatr*. 2010; 68 (2): 309-311. doi: 10.1590/S0004-282X2010000200030.
 - 112 Fujimoto K, Kawai S, Yonezawa T, et al. Basilar trunk aneurysms with associated fenestration treated by using Guglielmi detachable coils: two cases reports. *J Stroke Cerebrovasc Dis*. 2007; 16 (2): 84-87. doi: 10.1016/j.jstrokecerebrovasdis.2006.10.004.
 - 113 Fujimura M, Sugawara T, Higuchi H, et al. A ruptured aneurysm at the distal end of the basilar artery fenestration associated with multiple fenestrations of the vertebrobasilar system: case report. *Surg Neurol*. 1997; 47 (5): 469-472. doi: 10.1016/S0090-3019(96)00460-0.
 - 114 Gard AP, Kebriaei MA, Thorell WE. Review of vertebral artery fenestration and novel extracranial fenestration. *Clin Neurol Neurosurg*. 2013; 115 (7): 1107-1109. doi: 10.1016/j.clineuro.2012.08.023.
 - 115 Graves VB, Strother CM, Weir B, et al. Vertebrobasilar junction aneurysms associated with fenestration: treatment with Guglielmi detachable coils. *Am J Neuroradiol*. 1996; 17 (1): 35-40.
 - 116 Greenberg E, Katz JM, Janardhan V, et al. Treatment of a giant vertebrobasilar artery aneurysm using stent grafts. Case report. *J Neurosurg*. 2007; 107 (1): 165-168. doi: 10.3171/JNS-07/07/0165.
 - 117 Gruber TJ, Ogilvy CS, Hauck EF, et al. Endovascular treatment of a large aneurysm arising from a basilar trunk fenestration using the waffle-cone technique. *Neurosurgery*. 2010; 67 (3 Suppl Operative): ons140-4; discussion ons144. doi: 10.1227/01.NEU.0000382977.55504.6C.
 - 118 Hacıin-Bey L, Muszynski CA, Varelak PN. Saccular aneurysm associated with posterior cerebral artery fenestration manifesting as a subarachnoid hemorrhage in a child. *Am J Neuroradiol*. 2002; 23 (8): 1291-1294.
 - 119 Hasegawa T, Ito H, Hwang WZ, et al. Single extracranial-intracranial duplication of the vertebral artery. *Surg Neurol*. 1986; 25 (4): 369-372. doi: 10.1016/0090-3019(86)90212-0.
 - 120 Hattori T, Inoue S, Sakai N. Fenestration of the basilar artery associated with persistent primitive trigeminal artery. *Neurol Med Chir (Tokyo)*. 1997; 37 (11): 841-843. doi: 10.2176/nmc.37.841.
 - 121 Hayashi T, Hirose Y, Sagoh M, et al. Spontaneous occlusion of ruptured vertebral artery dissection at the extradural fenestration associated with extradural origin of the posterior inferior cerebellar artery--case report. *Neurol Med Chir (Tokyo)*. 2000; 40 (3): 164-168. doi: 10.2176/nmc.40.164.
 - 122 Hemmati M, Kim KS. A ruptured aneurysm at the basilar artery fenestration. *Radiology*. 1979; 130 (1): 174.
 - 123 Hoshimaru M, Hashimoto N, Kikuchi H, et al. Aneurysm of the fenestrated basilar artery: report of two cases. *Surg Neurol*. 1992; 37 (5): 406-409. doi: 10.1016/0090-3019(92)90013-D.

- 124 Iihara K, Murao K, Yamada N, et al. Growth potential and response to multimodality treatment of partially thrombosed large or giant aneurysms in the posterior circulation. *Neurosurgery*. 2008; 63 (5): 832-842; discussion 842-844. doi: 10.1227/01.NEU.0000313625.15571.1B.
- 125 Im SH, Kwon BJ, Jung C, et al. Coil embolization of "kissing aneurysms" associated with distal basilar artery fenestration. *Clin Neurol Neurosurg*. 2007; 109 (2): 210-213. doi: 10.1016/j.clineuro.2006.10.001.
- 126 Islak C, Kocer N, Kantarci F, et al. Endovascular management of basilar artery aneurysms associated with fenestrations. *Am J Neuroradiol*. 2002; 23 (6): 958-964.
- 127 Iwashita T, Tanaka Y, Hongo K, et al. Aneurysm originating from the fenestration of the posterior cerebral artery: case report. *Neurosurgery*. 2002; 50 (4): 881-884; discussion 884. doi: 10.1097/00006123-200204000-00040.
- 128 Juszkat R, Nowak S, Moskal J, et al. Endovascular treatment of basilar artery aneurysms associated with distal fenestration. A case report. *Interv Neuroradiol*. 2009; 15 (1): 109-111.
- 129 Kai Y, Hamada J, Morioka M, et al. Endovascular treatment of ruptured aneurysms associated with fenestrated basilar artery. Two case reports. *Neurol Med Chir (Tokyo)*. 2006; 46 (5): 244-247. doi: 10.2176/nmc.46.244.
- 130 Kanematsu M, Satoh K, Nakajima N, et al. Ruptured aneurysm arising from a basilar artery fenestration and associated with a persistent primitive hypoglossal artery. Case report and review of the literature. *J Neurosurg*. 2004; 101 (3): 532-535. doi: 10.3171/jns.2004.101.3.0532.
- 131 Kloska SP, Schlegel PM, Sträter R, et al. Causality of pediatric brainstem infarction and basilar artery fenestration? *Pediatr Neurol*. 2006; 35 (6): 436-438. doi: 10.1016/j.pediatrneurol.2006.06.013.
- 132 Komiya M, Nakajima H, Nishikawa M, et al. Treatment of a saccular aneurysm at the fenestration of the intracranial vertebral artery with Guglielmi detachable coils. *Acta Neurochir (Wien)*. 1999; 141 (10): 1125-1127. doi: 10.1007/s007010050495.
- 133 Kowada M, Yamaguchi K, Takahashi H. Fenestration of the vertebral artery with a review of 23 cases in Japan. *Radiology*. 1972; 103 (2): 343-346.
- 134 Krings T, Baccin CE, Alvarez H, et al. Segmental unfused basilar artery with kissing aneurysms: report of three cases and literature review. *Acta Neurochir (Wien)*. 2007; 149 (6): 567-574; discussion 574. doi: 10.1007/s00701-007-1118-0.
- 135 Kubo M, Haccin-Bey L, Varelas PN, et al. Ruptured saccular aneurysm of distal vertebral artery fenestration managed with Guglielmi detachable coils and intraventricular tissue plasminogen activator. *Surg Neurol*. 2005; 63 (3): 244-248; discussion 248. doi: 10.1016/j.surneu.2004.02.038.
- 136 Kumar S, Justin EM, Mishra NK. Fenestrated posterior inferior cerebellar artery with concomitant vertebro-basilar junction fenestration and vertebral artery aneurysm. *Clin Neuroradiol*. 2012; 22 (3): 235-237. doi: 10.1007/s00062-011-0070-9.
- 137 Lee SH, Koh JS, Ryu CW, et al. Fenestration of the double origin of the posterior inferior cerebellar artery associated with a contralateral vertebral artery dissection. *Cerebellum*. 2009; 8 (3): 382-384. doi: 10.1007/s12311-009-0108-6.
- 138 Lesley WS. Fenestration of the posterior inferior cerebellar artery. *Cerebellum*. 2008; 7 (3): 240-241. doi: 10.1007/s12311-008-0002-7.
- 139 Miyazaki S, Kamata K, Yamaura A. Multiple aneurysms of the vertebrobasilar system associated with fenestration of the vertebral artery. *Surg Neurol*. 1981; 15 (3): 192-195. doi: 10.1016/0090-3019(81)90140-3.
- 140 Morelli N, Gori S, Mancuso M, et al. Basilar artery fenestration in a woman with basilar-type migraine. *Neurol Sci*. 2007; 28 (2): 116-117. doi: 10.1007/s10072-007-0800-6.
- 141 Nagashima H, Okudera H, Orz Y, et al. Endovascular treatment of basilar trunk aneurysm associated with fenestration of the basilar artery. *Neurosurg Rev*. 1999; 22 (4): 219-221. doi: 10.1007/s101430050020.
- 142 Pasaoglu L, Hatipoglu HG, Vural M, et al. Persistent primitive hypoglossal artery and fenestration of posterior cerebral artery: CT and MR angiography. *Neuroradiology (Astur)*. 2009; 20 (6): 563-566; discussion 566. doi: 10.4321/S1130-14732009000600007.
- 143 Picard L, Roy D, Bracard S, et al. Aneurysm associated with a fenestrated basilar artery: report of two cases treated by endovascular detachable balloon embolization. *Am J Neuroradiol*. 1993; 14 (3): 591-594.
- 144 Prabhakaran S, Lopes DK. The "OK" sign: an unusual angiographic appearance of basilar artery fenestration. *Arch Neurol*. 2008; 65 (6): 836. doi: 10.1001/archneur.65.6.836.
- 145 Rieger P, Huber G. Fenestration and duplicate origin of the left vertebral artery in angiography. Report of three cases. *Neuroradiology*. 1983; 25 (1): 45-50. doi: 10.1007/BF00327480.
- 146 Saatci I, Cekirge HS, Karcaaltincaba M, et al. Endovascular treatment of kissing aneurysms at the fenestrated basilar artery. Case report with literature review. *Surg Neurol*. 2002; 58 (1): 54-58; discussion 58. doi: 10.1016/S0090-3019(02)00748-6.
- 147 Scherer A, Siebler M, Aulich A. Virtual arterial endoscopy as a diagnostic aid in a patient with basilar artery fenestration and thromboembolic pontine infarct. *Am J Neuroradiol*. 2002; 23 (7): 1237-1239.
- 148 Shigemori M, Shirahama M, Tokutomi T, et al. Bilateral fenestration of the vertebral artery in a case with cerebral aneurysm. *Kurume Med J*. 1981; 28 (1): 79-82. doi: 10.2739/kurumemedj.28.79.
- 149 Sim E, Vaccaro AR, Berzlanovich A, et al. Fenestration of the extracranial vertebral artery: review of the literature. *Spine (Phila Pa 1976)*. 2001; 26 (6): E139-142.
- 150 Sugita S, Shigemori M, Watanabe M, et al. A case of multiple cerebral aneurysms associated with fenestration of the basilar artery. *Kurume Med J*. 1986; 33 (3): 139-141. doi: 10.2739/kurumemedj.33.139.
- 151 Tanaka S, Tokimura H, Makiuchi T, et al. Clinical presentation and treatment of aneurysms associated with basilar artery fenestration. *J Clin Neurosci*. 2012; 19 (3): 394-401. doi: 10.1016/j.jocn.2011.04.043.
- 152 Tasker AD, Byrne JV. Basilar artery fenestration in association with aneurysms of the posterior cerebral circulation. *Neuroradiology*. 1997; 39 (3): 185-189. doi: 10.1007/s002340050389.
- 153 Theodosopoulos PV, Lawton MT. Fenestration of the posteroinferior cerebellar artery: case report. *Neurosurgery*. 2000; 47 (2): 463-465. doi: 10.1097/00006123-200008000-00041.
- 154 Tran-Dinh HD, Soo YS, Jayasinghe LS. Duplication of the vertebro-basilar system. *Australas Radiol*. 1991; 35 (3): 220-224. doi: 10.1111/j.1440-1673.1991.tb03012.x.
- 155 Tsuei YS, Matsumoto Y, Ohta M, et al. Vertebrobasilar junction fenestration with dumbbell-shaped aneurysms formation: computational fluid dynamics analysis. *Surg Neurol*. 2009; 72 Suppl 2: S11-19. doi: 10.1016/j.surneu.2009.05.026.
- 156 Uchino A, Sawada A, Takase Y, et al. Extreme fenestration of the right vertebral artery: magnetic resonance angiographic demonstration. *Eur Radiol*. 2002; 12 Suppl 3: S32-34.
- 157 Yim NY, Ha HI, Park JH, et al. Agenesis of bilateral

- internal carotid artery associated with basilar artery fenestration mimicking intra-arterial thrombus: a case report. *Vasc Endovascular Surg.* 2010; 44 (1): 69-74. doi: 10.1177/1538574409347393.
- 158 Yoon SM, Chun YI, Kwon Y, et al. Vertebrobasilar junction aneurysms associated with fenestration: experience of five cases treated with Guglielmi detachable coils. *Surg Neurol.* 2004; 61 (3): 248-254. doi: 10.1016/S0090-3019(03)00485-3.
 - 159 Yoshimoto H, Maeda H, Aoyama H, et al. Enlargement of cerebellar arteriovenous malformation associated with fenestration of the vertebral artery--case report. *Neurol Med Chir (Tokyo).* 1992; 32 (8): 585-588. doi: 10.2176/nmc.32.585.
 - 160 Zanini MA, Pereira VM, Jory M, et al. Giant fusiform aneurysm arising from fenestrated posterior cerebral artery and basilar tip variation: case report. *Neurosurgery.* 2009; 64 (3): E564-565; discussion E565. doi: 10.1227/01.NEU.0000338431.70709.81.
 - 161 Zhang QJ, Kobayashi S, Gibo H, et al. Vertebrobasilar junction fenestration associated with dissecting aneurysm of intracranial vertebral artery. *Stroke.* 1994; 25 (6): 1273-1275. doi: 10.1161/01.STR.25.6.1273.
 - 162 Katsuta T, Matsubara T, Fujii K. Fenestration of the supraclinoid internal carotid artery. *Neuroradiology.* 1993; 35 (6): 461. doi: 10.1007/BF00602832.
 - 163 Uchino A, Sawada A, Takase Y, et al. Extreme fenestration of the basilar artery associated with cleft palate, nasopharyngeal mature teratoma, and hypophyseal duplication. *Eur Radiol.* 2002; 12 (8): 2087-2090. doi: 10.1007/s00330-001-1194-0.
 - 164 Uchino A, Saito N, Kurita H, et al. Persistent trigeminal artery arising from the arterial ring/fenestration of the cavernous segment of the internal carotid artery. *Surg Radiol Anat.* 2012; 34 (7): 651-654. doi: 10.1007/s00276-011-0927-2.
 - 165 Dey M, Awad IA. Fenestration of supraclinoid internal carotid artery and associated aneurysm: embryogenesis, recognition, and management. *World Neurosurg.* 2011; 76 (6): 592.e1-5. doi: 10.1016/j.wneu.2011.04.019.
 - 166 Ichikawa T, Miyachi S, Izumi T, et al. Fenestration of a supraclinoid internal carotid artery associated with dual aneurysms: case report. *Neurosurgery.* 2011; 69 (4): E1005-8; discussion E1009. doi: 10.1227/NEU.0b013e318223b613.
 - 167 Kachhara R, Nair S, Gupta AK. Fenestration of the proximal anterior cerebral artery (A1) with aneurysm manifesting as subarachnoid hemorrhage--case report. *Neurol Med Chir (Tokyo).* 1998; 38 (7): 409-412. doi: 10.2176/nmc.38.409.
 - 168 Kobari M, Ishihara N, Yunoki K, et al. Triplication of the middle cerebral artery associated with fenestration of the anterior cerebral artery. *Keio J Med.* 1988; 37 (4): 429-433. doi: 10.2302/kjm.37.429.
 - 169 Bernard TJ, Mull BR, Handler MH, et al. An 18-year-old man with fenestrated vertebral arteries, recurrent stroke and successful angiographic coiling. *J Neurol Sci.* 2007; 260 (1-2): 279-282. doi: 10.1016/j.jns.2007.04.032.
 - 170 Kan P, Abila AA, Dumont TM, et al. Double-barrel stent-assisted coiling of a basilar artery fenestration aneurysm. *J Neuroimaging.* 2013; 23 (3): 496-499. doi: 10.1111/j.1552-6569.2012.00720.x.
 - 171 Tanaka M, Kikuchi Y, Ouchi T. Neuroradiological Analysis of 23 Cases of Basilar Artery Fenestration Based on 2280 Cases of MR Angiographies. *Interv Neuroradiol.* 2006; 12 (Suppl 1): 39-44.

Daniel L. Cooke, MD
 Assistant Professor in-Residence
 Neurointerventional Radiology
 Dept. Radiology & Biomedical Imaging
 University of California San Francisco
 505 Parnassus Ave. Long 352
 Box 0628
 San Francisco, CA 94143-0628, USA
 Tel.: 415-353-1863
 Fax: 415-353-8606
 E-mail: daniel.cooke@ucsf.edu