Published in final edited form as:

Br J Sports Med. 2015 January ; 49(2): 118–122. doi:10.1136/bjsports-2013-092536.

High knee abduction moments are common risk factors for patellofemoral pain (PFP) and anterior cruciate ligament (ACL) injury in girls: Is PFP itself a predictor for subsequent ACL injury?

Gregory D Myer^{1,2,3,4}, Kevin R Ford^{1,2,5}, Stephanie L Di Stasi³, Kim D Barber Foss¹, Lyle J Micheli^{4,6,7}, and Timothy E Hewett^{1,2,3,4,8}

¹Division of Sports Medicine, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio, USA

²Departments of Pediatrics and Orthopaedic Surgery, University of Cincinnati, Cincinnati, Ohio, USA

³The Sports Health and Performance Institute, OSU Sports Medicine, Ohio State University Medical Center, Columbus, Ohio, USA

⁴The Micheli Center for Sports Injury Prevention, Waltham, Massachusetts, USA

⁵Department of Physical Therapy, School of Health Sciences, High Point University, High Point, North Carolina, USA

⁶Division of Sports Medicine, Department of Orthopaedics, Boston Children's Hospital, Boston, Massachusetts, USA

⁷Harvard Medical School, Boston, Massachusetts, USA

⁸Departments of Physiology and Cell Biology, Family Medicine and of Orthopaedic Surgery and Biomedical Engineering, The Ohio State University, Columbus, Ohio, USA

Abstract

Background—Identifying risk factors for knee pain and anterior cruciate ligament (ACL) injury can be an important step in the injury prevention cycle.

Correspondence to Dr Gregory D Myer, Division of Sports Medicine, Cincinnati Children's Hospital Medical Center, 3333 Burnet Avenue; MLC 10001, Cincinnati, OH 45229, USA; greg.myer@cchmc.org.

Contributors GDM contributed to the conception and design, acquisition of data, analysis and interpretation of data, drafting of the manuscript, obtaining funding and supervision. KRF contributed to conception and design, critical revision of the manuscript for important intellectual content, analysis and interpretation of data, and administrative, technical or material support. KDBF contributed to acquisition of data, drafting of the manuscript, and administrative, technical or material support. SLDS contributed to the analysis and interpretation of data, critical revision of the manuscript for important intellectual content, and administrative, technical or material support. TEH contributed to the conception and design, critical revision of the manuscript for important intellectual content, obtaining funding and supervision.

Competing interests None.

Ethics approval Cincinnati Children's Hospital Institutional Review Board.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Objective—We evaluated two unique prospective cohorts with similar populations and methodologies to compare the incidence rates and risk factors associated with patellofemoral pain (PFP) and ACL injury.

Methods—The ‘PFP cohort’ consisted of 240 middle and high school female athletes. They were evaluated by a physician and underwent anthropometric assessment, strength testing and three-dimensional landing biomechanical analyses prior to their basketball season. 145 of these athletes met inclusion for surveillance of incident (new) PFP by certified athletic trainers during their competitive season. The ‘ACL cohort’ included 205 high school female volleyball, soccer and basketball athletes who underwent the same anthropometric, strength and biomechanical assessment prior to their competitive season and were subsequently followed up for incidence of ACL injury. A one-way analysis of variance was used to evaluate potential group (incident PFP vs ACL injured) differences in anthropometrics, strength and landing biomechanics. Knee abduction moment (KAM) cut-scores that provided the maximal sensitivity and specificity for prediction of PFP or ACL injury risk were also compared between the cohorts.

Results—KAM during landing above 15.4 Nm was associated with a 6.8% risk to develop PFP compared to a 2.9% risk if below the PFP risk threshold in our sample. Likewise, a KAM above 25.3 Nm was associated with a 6.8% risk for subsequent ACL injury compared to a 0.4% risk if below the established ACL risk threshold. The ACL-injured athletes initiated landing with a greater knee abduction angle and a reduced hamstrings-to-quadriceps strength ratio relative to the incident PFP group. Also, when comparing across cohorts, the athletes who suffered ACL injury also had lower hamstring/quadriceps ratio than the players in the PFP sample ($p < 0.05$).

Conclusions—In adolescent girls aged 13.3 years, >15 Nm of knee abduction load during landing is associated with greater likelihood of developing PFP. Also, in girls aged 16.1 years who land with >25 Nm of knee abduction load during landing are at increased risk for both PFP and ACL injury.

INTRODUCTION

Patellofemoral pain (PFP) is the most common disorder of the knee, with its greatest incidence in young, physically active girls.¹⁻⁵ Early identification of risk factors related to PFP may be critical for the prevention of PFP in young and highly active female athletes.

In young girls, injuries to the anterior cruciate ligament (ACL) result in the greatest total time lost from sport and recreational participation.⁶ Adolescent girls are affected with ACL injury 2–10 times more often than their male counterparts.⁷⁻⁹

Abnormal neuromuscular control strategies and anthropometric differences in patients with symptomatic PFP and following ACL injury are observed in both sexes (figure 1A). Several retrospective sex-specific investigations sought to identify risk factors for knee pain and injury in female populations (figure 1B).¹⁰⁻¹⁹ However, these retrospective study designs limit the potential to determine if the differences in lower extremity power and static alignments found in male and female patients with symptomatic PFP also contribute to the genesis of PFP.

Prospective studies of the risk factors predictive of the development of debilitating PFP and ACL injury in girls are lacking (figure 1C). Increased external knee abduction moment (KAM) during a drop vertical jump (DVJ) task is associated with increased risk for primary ACL rupture²⁰ and the development of PFP.¹⁹ There is substantial ACL-related research focused on retrospective analysis²¹⁻⁴⁴ and mixed populations;²²⁴⁵⁻⁶¹ this limits the potential to determine sex-specific risk factors that predispose young girls to ACL injury (figure 1).

The purpose of this study was to evaluate two prospective datasets with similar populations and methodologies with injury surveillance methods focused on incidence rates and risk factors for increased risk of PFP or ACL injury. The similar methods and populations provide an opportunity to compare incidence rates and risk factors. We hypothesised that the prospective measure of external KAM would be increased in magnitude in those athletes who sustained an ACL injury compared to those who sustained incident PFP.

METHODS

The current study evaluated and compared two different prospective datasets with similar populations and methodologies with injury surveillance methods focused on incidence rates and risk factors for increased risk of PFP or ACL injury.

PFP incident cohort

Two hundred-and-forty adolescent female athletes were recruited from a county public school district in Kentucky, USA, with five middle schools and three high schools. Participants were evaluated by a physician for prevalence of PFP (n=39) and other exclusionary (N=49) criteria prior to the surveillance period. PFP was operationally defined as retropatellar or peripatellar pain around the patellofemoral joint. At screening, participants were also evaluated for anthropometrics, lower extremity strength and landing biomechanics (DVJ) prior to their basketball season. Participants (N=152; 4 excluded at postseason follow-up) were monitored by certified athletic trainers for PFP during their competitive season. Fourteen participants developed incident PFP during the competitive season.¹⁹

ACL injury cohort

A similar prospective investigation in our laboratory also evaluated anthropometrics, lower extremity strength and landing biomechanics of 205 young female athletes preseason and monitored for ACL injury during their competitive season. Nine participants sustained an ACL rupture during their competitive season.²⁰ Therefore, the outcome of interest was the development of pathology in these two cohorts and the involved limb was utilised for between-group comparisons. The cumulative incidence rate for new PFP was 2.2-fold greater (95% CI 1.1 to 4.3) relative to ACL injury when normalised per 100 athlete seasons (PFP 9.7 vs ACL 4.4) in the two compared datasets.

The Institutional Review Board approved the data collection procedures and consent forms. Parental consent and athlete assent were obtained before data collection. Participants were tested prior to the start of their competitive season. The average age, height and mass for the

combined cohorts was 15.0 (95% CI 14.7 to 15.2 years), 162.6 (95% CI 161.8 to 163.4 cm) and 57.2 (95% CI 56.1 to 58.2 kg), respectively (table 1).

Landing biomechanics

Three-dimensional (3D) hip, knee and ankle kinematic and kinetic data were quantified for the contact phase of three DVJ tasks. Following instrumentation with retro-reflective markers, a static trial was conducted in which the participant was instructed to stand still with foot placement standardised to the laboratory coordinate system. This static measurement was used as each participant's neutral (zero) alignment; subsequent kinematic measures were referenced in relation to this position. The DVJ involved the participant standing on top of a box (31 cm high) with her feet positioned 35 cm apart. Each athlete was instructed to drop directly down off the box and immediately perform a maximum effort vertical jump, raising both arms while jumping for a basketball that was suspended overhead at a height corresponding to her predetermined maximum vertical jump height.⁶²

Data analysis

For each participant, biomechanical variables were calculated during the DVJ manoeuvre and the mean of three trials was utilised for analyses. Specifically, lower limb joint data were generated via the 3D coordinates of externally mounted skin markers.^{62,63} Net external KAM was described in this paper and has previously demonstrated high reliability for outcome measurement in our laboratory.⁶³

Athlete surveillance

The athletes were monitored on a weekly basis for athletic exposures (AEs) for either incident ACL injury or PFP by a certified athletic trainer. AEs were defined as one game or practice session. The AEs of participants with preseason PFP, or other participants who were excluded, were not included in the compilation of total AEs. These data were compiled to calculate in-season injury incidence for each of the compared studies.⁶⁴

Statistical analyses

Statistical analyses were conducted in SPSS (SPSS, V.21.0, Chicago, Illinois, USA). A one-way analysis of variance was used to evaluate potential group (PFP vs ACL) differences in anthropometrics, strength and landing biomechanics. Logistic regression analyses were used to determine threshold KAM cut-off scores that provided the maximal sensitivity and specificity for prediction of PFP and ACL injury risk. The predictive accuracy of the multivariable regression model was quantified with the use of the C statistic, which measures the area under the receiver operating characteristic curve. Incidence rates were expressed per 100 athlete competitive seasons. Statistical significance was established a priori at $p < 0.05$.

RESULTS

Maximal sensitivity and specificity for prediction of PFP increased in athletes who demonstrated a peak external KAM > 15.4 Nm; ACL injury risk increased with peak landing KAM > 25.3 Nm. Cut-scores relative to ACL injury and PFP incidence were compared

between the two evaluated investigations (figure 2). KAM during landing above 15.4 Nm was associated with a 6.8% risk to develop PFP compared to a 2.9% risk if below the PFP risk threshold in our sample. Likewise, a KAM above 25.3 Nm was associated with a 6.8% risk for subsequent ACL injury compared to a 0.4% risk if below the established ACL risk threshold. The study participants who developed PFP demonstrated reduced initial contact knee abduction angle relative to the ACL-injured cohort. Specifically, the incident PFP group initiated landing with 1.4° (95% CI -0.5° to 3.2°) compared to ACL injured who impacted the ground with 5.0° (3.4° to 6.5°; $p<0.05$) of knee abduction angle.

Hamstring flexibility and vertical jump height were not different between the ACL injured and incident PFP groups ($p>0.05$). The hamstrings-to-quadriceps (HQ) strength ratio was lower in the ACL-injured group than the PFP group. The ACL-injured group demonstrated a 51% HQ strength ratio (42–59%) and the PFP group demonstrated a 68% HQ strength ratio (57–79%; $p<0.05$). The ACL-injured group demonstrated higher quadriceps strength force production compared to the PFP group ($p<0.05$), but did not show a similar adaptive increase in hamstrings strength ($p>0.05$). As noted in table 1, the overall study cohorts were not entirely homogenous in sampling. However, there were group differences in demographic and anthropometric measures between the subsample of incident PFP and ACL-injured cohorts. The ACL-injured group was 15.8 years old compared to the incident PFP group who were 12.7 years of age ($p<0.05$). Likewise, the ACL-injured cohort was 5.2% taller than the PFP group ($p<0.05$); however, there were no group differences in participant mass ($p>0.05$).

DISCUSSION

The purpose of this study was to evaluate two prospective datasets with similar populations to compare the incidence rates and risk factors associated with increased risk of PFP and ACL injury. We hypothesised that the external KAM would be increased in magnitude in those athletes who had an ACL injury compared to those who sustained incident PFP. The results of this study found that girls who demonstrated >15 Nm of knee abduction during landing may be at increased risk for the development of PFP and those who demonstrate >25 Nm of knee abduction during landing may be at increased risk for both PFP and ACL injury.¹⁹²⁰⁶⁵ The increased incidence of PFP compared to that of ACL injury may be associated with the reduced threshold of knee abduction associated with increased risk of PFP. In other words, increased prevalence of PFP may be related to the chronic mechanisms of aberrant knee loading that do not reach the magnitude required for ACL injury during early maturation years. However, as young girls increase in height (specifically tibia length) and mass, small magnitude knee abduction loads naturally increase and can exceed ACL injury risk thresholds in the later adolescent years.⁶⁶⁶⁷

Differences in anthropometric and strength measures between athletes with PFP and those with ACL injury may also help explain their unique risk profiles. The ACL-injured group was both older and taller than the PFP group, indicating that PFP may be more likely to occur earlier in the maturation cycle. HQ strength measures were also different between groups. While the ACL-injured group demonstrated higher quadriceps strength than the PFP group, their HQ ratio was lower than that of their counterparts. Previous work has

highlighted HQ ratios less than 60% as a risk factor for ACL injury,⁶⁸ and this may have implications for the jump-landing strategies of these athletes. Girls land with greater net knee extensor compared to net hip extensor moments.⁶⁹ In addition, female athletes tend to increase quadriceps activation with increased plyometric intensity without a balanced increase in hamstrings activation.⁷⁰ Conversely, during changes through maturation, boys exhibit an increase in net hip extensor torque compared to net knee extensor torque during a landing task.⁶⁹ These opposing strategies may, in part, explain the divergence in relative injury risk of girls and boys during the maturation process. Adequate co-contraction of the knee flexors may balance contraction of the quadriceps, compress the joint and potentially limit high knee abduction torques during landing.

Aberrant neuromuscular strategies appear to manifest as reduced frontal plane control of the knee in young female athletes.⁷¹ Reduced lower extremity frontal plane control, combined with a reduction in posterior chain strength, is exacerbated as young girls grow both in height and mass with maturation.⁶⁷⁷⁰⁷² Therefore, focused training on the posterior chain at early maturational stages may be warranted in girls that exhibit high risk (eg, increased KAM) for ACL or PFP.⁷³⁻⁷⁸

The high recurrence rate of PFP and second ACL injury combined with the poor long-term prognosis indicate that prevention strategies are the best approach to reduce the morbidity associated with PFP developed during youth. Work in our laboratory has successfully employed neuromuscular training interventions to correct the abnormal knee biomechanics that manifest during maturational development in young girls.¹⁹⁷³⁷⁴⁷⁶ Preseason exercise interventions targeted at the neuromuscular deficits known to increase knee injury risk may be warranted for maturing, adolescent girls. Specifically, adolescent female athletes who land with >15 Nm of knee abduction, may benefit from only preseason training, while those girls who land with >25 Nm of knee abduction may benefit from increased treatment dosage gained from both preseason and in-season neuromuscular training protocols aimed to reduce knee abduction and PFP/ACL injury incidence.⁶⁵⁷³⁷⁹ Particularly, training protocols that focus on improved neuromuscular control, with technical feedback provided in the maturing years, appear most effective in the reduction of risk factors and knee injury incidence in female athletes.⁷⁶⁸⁰⁻⁸² In addition, in young, adolescent (middle-school aged) girls in whom new PFP incidence appears to peak, increased knee abduction during landing appears to be an early indicator of increased PFP risk and is highly related to increased ACL injury during later adolescence.¹⁹

An optimal window of opportunity may be present for the initiation of integrative neuromuscular training based on measures of somatic maturity (growth). From these data, it appears most beneficial to initiate integrative training programmes during preadolescence *prior* to the period of peak height velocity, when youth are growing the fastest. Children with earlier growth may particularly benefit from earlier participation in integrative neuromuscular training.⁸³⁸⁴ While growth helps define periods optimal for gaining motor competence with integrative training, it is ultimately the child's psychosocial status (ability to follow coaching instructions, adhere to safety rules and handle the attention demands of a training programme) that is paramount in the decision process regarding his or her participation in a structured training programme.⁸³⁸⁴ In a recent longitudinal study, it was

noted that pubertal girls have an increase in abnormal landing mechanics over time.⁷² Contributing risk factors for knee injury were significantly greater across consecutive years in young postpubertal female athletes compared to males. Importantly, integrative neuromuscular training programmes successfully reduce these abnormal biomechanics⁷³⁷⁴⁷⁶⁸⁵ and appear to decrease PFP and ACL injury rates in female athletes.⁸⁶⁸⁷ Therefore, preadolescence may be the optimal time to institute programmes aimed to reduce neuromuscular deficits (increased knee abduction) that accelerate during maturation and lead to increased musculoskeletal injury risk.¹⁹²⁰⁶⁵

The point prevalence of active symptomatic PFP at the beginning of the season in a cohort of adolescent female athletes followed by our research group was 16.3 per 100 participants (age 13.4 years; range 9.6–18.2). The cumulative incidence risk and rate for the development of new unilateral PFP over the subsequent competitive season was 9.7 per 100 participants and 1.1 per 1000 athlete-exposures, respectively.¹⁹ These incidence data are nearly identical to the 10% reported in a study of older student athletes (age 18.6 years; range 17–21).⁴ However, if the participants identified with preseason PFP were included in the analyses, our monitored sample population demonstrated a prevalence of 22.0 per 100 participants. Cumulatively, these data provide strong indications of the need for preseason screening in younger female athletes to identify and minimise the effect of predisposing risk factors of PFP and future ACL injury as young girls mature.¹⁹ Longitudinal data indicate knee pain and injury influences an immediate increase in BMI z-score and body fat percentage in the year of reported incidence.⁸⁸ Future research should focus on determination of the underlying pathomechanics of PFP that increase its prevalence in young growing girls to development of sex-specific and age-appropriate interventions. Based on the current report it may be speculated that effective interventions implemented at an early age could be beneficial to prevent PFP prevalence and potentially reduce future risk of ACL injury.

The current report raises the important issue of the use of exercise interventions for prevention. Prior prospective biomechanical reports indicate that exercise interventions may reduce the KAM risk factor common to PFP and ACL injury risk reported in this study.⁷³⁷⁶⁸⁵ A more clear understanding of whether reductions in KAM translate into subsequent reductions in injuries such as PFP and ACL will need to be tested in a randomised controlled trial setting.

The clinical interventions that have reduced lower limb injuries appear to be most effective when implemented in younger athletes⁸¹ and must provide adequate dosage of prescribed exercise.⁸⁹ Relative to exercise selection, plyometric, strengthening and proximal neuromuscular control exercises appear to reduce ACL injury incidence to a greater extent than programmes without these exercise components.⁹⁰ Likewise, it is also likely that proper supervision and feedback of exercises are the common thread throughout the effective training protocols.⁸¹ Qualified instruction is integral to ensure that young athletes are performing exercises properly and not reinforcing bad movement mechanics such as high KAM strategies which may put them at risk for PFP and ACL injury.⁷⁶⁹¹⁹² Clinical tools that can identify athletes with increased KAM for targeted neuromuscular training might further optimise the efficacy and efficiency of exercise interventions focused to reduce PFP

and ACL injury.⁹²⁻⁹⁵ While the evidence is accumulating, further research in children and adolescents is warranted to further determine the potential to prevent PFP and ACL injury risk with interventions targeting underlying KAM mechanics.

LIMITATIONS

We evaluated two prospective datasets to compare the risk factors associated with increased risk of PFP and ACL injury. Future studies should prospectively screen and identify, within the same population, the predictive value of knee abduction to identify risk of both PFP and ACL rupture. Specifically, future studies with homogeneity in anthropometrics and demographics that are longitudinally monitored for PFP and ACL injury incidence are needed to validate our speculations of the current report.

The generalisability of this study is limited to girls performing a DVJ with similar methodology. Lastly, ACL injury and PFP likely have multifactorial aetiology, with unmeasured factors influencing outcome that were not compared in the current investigation. Injury data demonstrate that many physical and psychological parameters affect injury rates. There are several possible contributing and confounding variables that were not controlled for in the study design, which included school, team, age/grade, aggressiveness, foot pronation, quadriceps angle, femoral notch width, reliable menstrual status reporting and blood hormone levels (figure 1). However, neuromuscular parameters appear to be a major determinant of risk for knee injury. Future investigations with larger sample sizes should aim to develop more robust prediction models which include other potential contributing parameters (eg, sport, training error, hormonal measures and potentially psychological parameters) to further elucidate risk factors for PFP or ACL injury.

CONCLUSIONS

The current data indicate that there is minimal overlap between incident ACL injured and incident PFP in the magnitude of the KAM values as their risk factor for injury. It appears that this kinetic parameter measured during landing is, in itself, a risk factor common to both conditions, but lower values are able to predispose young girls to PFP risk and higher values predispose to both PFP and ACL injury. Specifically, young girls who demonstrate >15 Nm of knee abduction during landing may be at increased risk for the development of PFP and is likely reflective of their reduced mass and height (ie, lighter and shorter stature).

Likewise, the limited number of younger female athletes who demonstrate >25 Nm of knee abduction during landing suggests that the increase in mass and height increases frontal plane knee loads. We speculate that this increased frontal load likely propagates beyond just increased risk for the development of PFP to the more severe ACL injury in maturing young girls. Young girls with PFP may possess risk factors that put them at future risk for ACL injury. This may be due to increased mass and height during their adolescent years. Bearing in mind the limitations of our data (above), we contend that preventive exercise interventions for girls with a history of PFP during their growing years may benefit from a targeted neuromuscular training programme to reduce KAM landing mechanics. Similarly,

adolescent girls who land with >25 Nm of knee abduction may benefit from such a programme.

What are the new findings?

- ▶ Knee abduction moment during landing above 15.4 Nm is associated with a 6.8% risk to develop patellofemoral pain (PFP) compared to a 2.9% risk if below the PFP risk threshold in our sample.
- ▶ Knee abduction moment above 25.3 Nm is associated with a 6.8% risk for subsequent anterior cruciate ligament (ACL) injury compared to a 0.4% risk if below the established ACL risk threshold.
- ▶ Young girls with PFP may have risk factors that put them at future risk for ACL injury as they mature during their adolescent years.

How might it impact on clinical practice in the near future?

- ▶ Preseason exercise interventions focused on plyometric, strengthening and proximal neuromuscular control exercises may be warranted for girls with a history of patellofemoral pain (PFP) during their growing years.
- ▶ Adolescent girls with a history of PFP and/or who land with >25 Nm of knee abduction may benefit from increased treatment dosage gained from both preseason and in-season neuromuscular training protocols to lower risk of PFP and/or ACL injury.

Acknowledgements

The authors would like to thank Boone County Kentucky, School District, especially School Superintendent Randy Poe, for participation in this study. They would also like to thank Mike Blevins, Ed Massey, Dr Brian Blavatt and the athletes of Boone County public school district for their participation in this study.

Funding This work was supported by the National Institutes of Health/NIAMS Grants #R01-AR049735, #R01-AR05563 and #R01-AR056259.

REFERENCES

1. Heintjes, E.; Berger, M.; Bierma-Zeinsträ, S., et al. Exercise therapy for patellofemoral pain syndrome. John Wiley & Sons, Ltd; Hobken, NJ: 2005.
2. Loudon JK, Gajewski B, Goist-Foley HL, et al. The effectiveness of exercise in treating patellofemoral-pain syndrome. J Sport Rehab. 2004; 13:323–42.
3. Natri A, Kannus P, Jarvinen M. Which factors predict the long-term outcome in chronic patellofemoral pain syndrome? A 7-yr prospective follow-up study. Med Sci Sports Exerc. 1998; 30:1572–7. [PubMed: 9813868]
4. Witvrouw E, Lysens R, Bellemans J, et al. Intrinsic risk factors for the development of anterior knee pain in an athletic population. A two-year prospective study. Am J Sports Med. 2000; 28:480–9. [PubMed: 10921638]
5. Kannus P, Aho H, Jarvinen M, et al. Computerized recording of visits to an outpatient sports clinic. Am J Sports Med. 1987; 15:79–85. [PubMed: 3812865]

6. Alentorn-Geli E, Myer GD, Silvers HJ, et al. Prevention of non-contact anterior cruciate ligament injuries in soccer players. Part 1: Mechanisms of injury and underlying risk factors. *Knee Surg Sports Traumatol Arthrosc.* 2009; 17:705–29. [PubMed: 19452139]
7. Malone TR, Hardaker WT, Garrett WE, et al. Relationship of gender to anterior cruciate ligament injuries in intercollegiate basketball players. *J South Orthop Assoc.* 1993; 2:36–9.
8. Arendt E, Dick R. Knee injury patterns among men and women in collegiate basketball and soccer. NCAA data and review of literature. *Am J Sports Med.* 1995; 23:694–701. [PubMed: 8600737]
9. Hewett TE, Lindenfeld TN, Riccobene JV, et al. The effect of neuromuscular training on the incidence of knee injury in female athletes. A prospective study. *Am J Sports Med.* 1999; 27:699–706. [PubMed: 10569353]
10. Boling M, Padua D, Marshall S, et al. Gender differences in the incidence and prevalence of patellofemoral pain syndrome. *Scand J Med Sci Sports.* 2010; 20:725–30. [PubMed: 19765240]
11. Powers CM. Patellar kinematics, part II: the influence of the depth of the trochlear groove in subjects with and without patellofemoral pain. *Phys Ther.* 2000; 80:965–78. [PubMed: 11002432]
12. Kaya D, Citaker S, Kerimoglu U, et al. Women with patellofemoral pain syndrome have quadriceps femoris volume and strength deficiency. *Knee Surg Sports Traumatol Arthrosc.* 2011; 19:242–7. [PubMed: 20953760]
13. Long-Rossi F, Salsich GB. Pain and hip lateral rotator muscle strength contribute to functional status in females with patellofemoral pain. *Physiother Res Int.* 2010; 15:57–64. [PubMed: 19790109]
14. Prins MR, van der Wurff P. Females with patellofemoral pain syndrome have weak hip muscles: a systematic review. *Aust J Physiother.* 2009; 55:9–15. [PubMed: 19226237]
15. Magalhaes E, Fukuda TY, Sacramento SN, et al. A comparison of hip strength between sedentary females with and without patellofemoral pain syndrome. *J Orthop Sports Phys Ther.* 2010; 40:641–7. [PubMed: 20508327]
16. Willson JD, Kernozek TW, Arndt RL, et al. Gluteal muscle activation during running in females with and without patellofemoral pain syndrome. *Clin Biomech.* 2011; 26:735–40.
17. Powers CM, Maffucci R, Hampton S. Rearfoot posture in subjects with patellofemoral pain. *J Orthop Sports Phys Ther.* 1995; 22:155–60. [PubMed: 8535473]
18. Ireland ML, Willson JD, Ballantyne BT, et al. Hip strength in females with and without patellofemoral pain. *J Orthop Sports Phys Ther.* 2003; 33:671–6. [PubMed: 14669962]
19. Myer GD, Ford KR, Barber Foss KD, et al. The incidence and potential pathomechanics of patellofemoral pain in female athletes. *Clin Biomech.* 2010; 25:700–7.
20. Hewett TE, Myer GD, Ford KR, et al. Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: a prospective study. *Am J Sports Med.* 2005; 33:492–501. [PubMed: 15722287]
21. Adachi N, Nawata K, Maeta M, et al. Relationship of the menstrual cycle phase to anterior cruciate ligament injuries in teenaged female athletes. *Arch Orthop Trauma Surg.* 2008; 128:473–8. [PubMed: 17909824]
22. Bisson LJ, Gurske-DePerio J. Axial and sagittal knee geometry as a risk factor for noncontact anterior cruciate ligament tear: a case-control study. *Arthroscopy.* 2010; 26:901–6. [PubMed: 20620789]
23. Brophy R, Silvers HJ, Gonzales T, et al. Gender influences: the role of leg dominance in ACL injury among soccer players. *Br J Sports Med.* 2010; 44:694–7. [PubMed: 20542974]
24. Chappell JD, Creighton RA, Giuliani C, et al. Kinematics and electromyography of landing preparation in vertical stop-jump: risks for noncontact anterior cruciate ligament injury. *Am J Sports Med.* 2007; 35:235–41. [PubMed: 17092926]
25. Dienst M, Schneider G, Altmeyer K, et al. Correlation of intercondylar notch cross sections to the ACL size: a high resolution MR tomographic in vivo analysis. *Arch Orthop Trauma Surg.* 2007; 127:253–60. [PubMed: 16807752]
26. Durall CJ, Kernozek TW, Kersten M, et al. Associations between single-leg postural control and drop-landing mechanics in healthy women. *J Sport Rehabil.* 2011; 20:406–18. [PubMed: 22012495]

27. Huston LJ, Wojtys EM. Neuromuscular performance characteristics in elite female athletes. *Am J Sports Med.* 1996; 24:427–36. [PubMed: 8827300]
28. Malinzak RA, Colby SM, Kirkendall DT, et al. A comparison of knee joint motion patterns between men and women in selected athletic tasks. *Clin Biomech.* 2001; 16:438–45.
29. Zeller BL, McCrory JL, Kibler WB, et al. Differences in kinematics and electromyographic activity between men and women during the single-legged squat. *Am J Sports Med.* 2003; 31:449–56. [PubMed: 12750142]
30. Chappell JD, Yu B, Kirkendall DT, et al. A comparison of knee kinetics between male and female recreational athletes in stop-jump tasks. *Am J Sports Med.* 2002; 30:261–7. [PubMed: 11912098]
31. Salci Y, Kentel BB, Heycan C, et al. Comparison of landing maneuvers between male and female college volleyball players. *Clin Biomech.* 2004; 19:622–8.
32. James CR, Sizer PS, Starch DW, et al. Gender differences among sagittal plane knee kinematic and ground reaction force characteristics during a rapid sprint and cut maneuver. *Res Q Exerc Sport.* 2004; 75:31–8. [PubMed: 15532359]
33. Khan MS, Seon JK, Song EK. Risk factors for anterior cruciate ligament injury: assessment of tibial plateau anatomic variables on conventional MRI using a new combined method. *Int Orthop.* 2011; 35:1251–6. [PubMed: 21340682]
34. Kramer LC, Denegar CR, Buckley WE, et al. Factors associated with anterior cruciate ligament injury: history in female athletes. *J Sports Med Phys Fitness.* 2007; 47:446–54. [PubMed: 18091686]
35. Mendiguchia J, Ford KR, Quatman CE, et al. Sex differences in proximal control of the knee joint. *Sports Med.* 2011; 41:541–57. [PubMed: 21688868]
36. Olsen OE, Myklebust G, Engebretsen L, et al. Relationship between floor type and risk of ACL injury in team handball. *Scand J Med Sci Sports.* 2003; 13:299–304. [PubMed: 14507295]
37. Posthumus M, September AV, Keegan M, et al. Genetic risk factors for anterior cruciate ligament ruptures: COL1A1 gene variant. *Br J Sports Med.* 2009; 43:352–6. [PubMed: 19193663]
38. Posthumus M, September AV, O’Cuinneagain D, et al. The COL5A1 gene is associated with increased risk of anterior cruciate ligament ruptures in female participants. *Am J Sports Med.* 2009; 37:2234–40. [PubMed: 19654427]
39. Posthumus M, September AV, O’Cuinneagain D, et al. The association between the COL12A1 gene and anterior cruciate ligament ruptures. *Br J Sports Med.* 2010; 44:1160–5. [PubMed: 19443461]
40. Ruedl G, Ploner P, Linortner I, et al. Are oral contraceptive use and menstrual cycle phase related to anterior cruciate ligament injury risk in female recreational skiers? *Knee Surg Sports Traumatol Arthrosc.* 2009; 17:1065–9. [PubMed: 19333573]
41. Todd MS, Lalliss S, Garcia E, et al. The relationship between posterior tibial slope and anterior cruciate ligament injuries. *Am J Sports Med.* 2010; 38:63–7. [PubMed: 19737987]
42. Slauterbeck JR, Fuzie SF, Smith MP, et al. The menstrual cycle, sex hormones, and anterior cruciate ligament injury. *J Athl Train.* 2002; 37:275–8. [PubMed: 12937585]
43. Beynnon BD, Johnson RJ, Braun S, et al. The relationship between menstrual cycle phase and anterior cruciate ligament injury: a case-control study of recreational alpine skiers. *Am J Sports Med.* 2006; 34:757–64. [PubMed: 16436538]
44. Posthumus M, Collins M, van der Merwe L, et al. Matrix metalloproteinase genes on chromosome 11q22 and the risk of anterior cruciate ligament (ACL) rupture. *Scand J Med Sci Sports.* 2012; 22:523–33. [PubMed: 21410539]
45. Chaudhari AM, Zelman EA, Flanigan DC, et al. Anterior cruciate ligament-injured subjects have smaller anterior cruciate ligaments than matched controls: a magnetic resonance imaging study. *Am J Sports Med.* 2009; 37:1282–7. [PubMed: 19307330]
46. Domzalski M, Grzelak P, Gabos P. Risk factors for anterior cruciate ligament injury in skeletally immature patients: analysis of intercondylar notch width using magnetic resonance imaging. *Int Orthop.* 2010; 34:703–7. [PubMed: 20333378]
47. Evans KN, Kilcoyne KG, Dickens JF, et al. Predisposing risk factors for non-contact ACL injuries in military subjects. *Knee Surg Sports Traumatol Arthrosc.* 2012; 20:1554–9. [PubMed: 22080351]

48. Fong CM, Blackburn JT, Norcross MF, et al. Ankle-dorsiflexion range of motion and landing biomechanics. *J Athl Train.* 2011; 46:5–10. [PubMed: 21214345]
49. Dowling AV, Corazza S, Chaudhari AM, et al. Shoe-surface friction influences movement strategies during a sidestep cutting task: implications for anterior cruciate ligament injury risk. *Am J Sports Med.* 2010; 38:478–85. [PubMed: 20194954]
50. Hoteya K, Kato Y, Motojima S, et al. Association between intercondylar notch narrowing and bilateral anterior cruciate ligament injuries in athletes. *Arch Orthop Trauma Surg.* 2011; 131:371–6. [PubMed: 21246376]
51. Uhorchak JM, Scoville CR, Williams GN, et al. Risk factors associated with noncontact injury of the anterior cruciate ligament: a prospective four-year evaluation of 859 West Point cadets. *Am J Sports Med.* 2003; 31:831–42. [PubMed: 14623646]
52. Charlton WP, St John TA, Ciccotti MG, et al. Differences in femoral notch anatomy between men and women: a magnetic resonance imaging study. *Am J Sports Med.* 2002; 30:329–33. [PubMed: 12016071]
53. Wojtys EM, Huston LJ, Taylor PD, et al. Neuromuscular adaptations in isokinetic, isotonic, and agility training programs. *Am J Sports Med.* 1996; 24:187–92. [PubMed: 8775118]
54. Malila S, Yuktanandana P, Saowaprut S, et al. Association between matrix metalloproteinase-3 polymorphism and anterior cruciate ligament ruptures. *Genet Mol Res.* 2011; 10:4158–65. [PubMed: 22057989]
55. Posthumus M, Collins M, September AV, et al. The intrinsic risk factors for ACL ruptures: an evidence-based review. *Phys Sportsmed.* 2011; 39:62–73. [PubMed: 21378488]
56. Simon RA, Everhart JS, Nagaraja HN, et al. A case-control study of anterior cruciate ligament volume, tibial plateau slopes and intercondylar notch dimensions in ACL-injured knees. *J Biomech.* 2010; 43:1702–7. [PubMed: 20385387]
57. Sonnery-Cottet B, Archbold P, Cucurulo T, et al. The influence of the tibial slope and the size of the intercondylar notch on rupture of the anterior cruciate ligament. *J Bone Joint Surg Br.* 2011; 93:1475–8. [PubMed: 22058297]
58. Vyas S, van Eck CF, Vyas N, et al. Increased medial tibial slope in teenage pediatric population with open physes and anterior cruciate ligament injuries. *Knee Surg Sports Traumatol Arthrosc.* 2011; 19:372–7. [PubMed: 20676607]
59. Myklebust G, Maehlum S, Holm I, et al. A prospective cohort study of anterior cruciate ligament injuries in elite Norwegian team handball. *Scand J Med Sci Sports.* 1998; 8:149–53. [PubMed: 9659675]
60. Harner CD, Paulos LE, Greenwald AE, et al. Detailed analysis of patients with bilateral anterior cruciate ligament injuries. *Am J Sports Med.* 1994; 22:37–43. [PubMed: 8129108]
61. Flynn RK, Pedersen CL, Birmingham TB, et al. The familial predisposition toward tearing the anterior cruciate ligament: a case control study. *Am J Sports Med.* 2005; 33:23–8. [PubMed: 15610995]
62. Ford KR, Myer GD, Hewett TE. Valgus knee motion during landing in high school female and male basketball players. *Med Sci Sports Exerc.* 2003; 35:1745–50. [PubMed: 14523314]
63. Ford KR, Myer GD, Hewett TE. Reliability of landing 3D motion analysis: implications for longitudinal analyses. *Med Sci Sports Exerc.* 2007; 39:2021–8. [PubMed: 17986911]
64. Knowles SB, Marshall SW, Guskiewicz KM. Issues in estimating risks and rates in sports injury research. *J Athl Train.* 2006; 41:207–15. [PubMed: 16791309]
65. Myer GD, Ford KR, Hewett TE. Quantification of knee load associated with increased risk for specific knee injury incidence. *J Athl Train* 2011;46:S–30.
66. Myer GD, Ford KR, Khoury J, et al. Biomechanics laboratory-based prediction algorithm to identify female athletes with high knee loads that increase risk of ACL injury. *Br J Sports Med.* 2011; 45:245–52. [PubMed: 20558526]
67. Quatman-Yates CC, Myer GD, Ford KR, et al. A longitudinal evaluation of maturational effects on lower extremity strength in female adolescent athletes. *Pediatr Phys Ther.* 2013; 25:271–6. [PubMed: 23619561]

68. Myer GD, Ford KR, Barber Foss KD, et al. The relationship of hamstrings and quadriceps strength to anterior cruciate ligament injury in female athletes. *Clin J Sport Med.* 2009; 19:3–8. [PubMed: 19124976]
69. Ford KR, Myer GD, Hewett TE. Longitudinal effects of maturation on lower extremity joint stiffness in adolescent athletes. *Am J Sports Med.* 2010; 38:1829–37. [PubMed: 20522830]
70. Ford KR, Myer GD, Schmitt LC, et al. Preferential quadriceps activation in female athletes with incremental increases in landing intensity. *J Appl Biomech.* 2011; 27:215–22. [PubMed: 21844610]
71. Hewett TE, Myer GD, Ford KR. Decrease in neuromuscular control about the knee with maturation in female athletes. *J Bone Joint Surg Am.* 2004; 86-A:1601–8. [PubMed: 15292405]
72. Ford KR, Shapiro R, Myer GD, et al. Longitudinal sex differences during landing in knee abduction in young athletes. *Med Sci Sports Exerc.* 2010; 42:1923–31. [PubMed: 20305577]
73. Myer GD, Ford KR, Palumbo JP, et al. Neuromuscular training improves performance and lower-extremity biomechanics in female athletes. *J Strength Cond Res.* 2005; 19:51–60. [PubMed: 15705045]
74. Myer GD, Ford KR, McLean SG, et al. The effects of plyometric versus dynamic stabilization and balance training on lower extremity biomechanics. *Am J Sports Med.* 2006; 34:445–55. [PubMed: 16282579]
75. Myer GD, Ford KR, Brent JL, et al. The effects of plyometric versus dynamic balance training on power, balance and landing force in female athletes. *J Strength Cond Res.* 2006; 20:345–53. [PubMed: 16686562]
76. Myer GD, Ford KR, Brent JL, et al. Differential neuromuscular training effects on ACL injury risk factors in “high-risk” versus “low-risk” athletes. *BMC Musculoskelet Disord.* 2007; 8:39. [PubMed: 17488502]
77. Myer GD, Chu DA, Brent JL, et al. 2008; 27:425–48. ix.
78. Myer GD, Brent JL, Ford KR, et al. A pilot study to determine the effect of trunk and hip focused neuromuscular training on hip and knee isokinetic strength. *Br J Sports Med.* 2008; 42:614–19. [PubMed: 18308886]
79. Klugman MF, Brent JL, Myer GD, et al. Does an in-season only neuromuscular training protocol reduce deficits quantified by the tuck jump assessment? *Clin Sports Med.* 2011; 30:825–40. [PubMed: 22018323]
80. Myer GD, Stroube BW, DiCesare CA, et al. Augmented feedback supports skill transfer and reduces high-risk injury landing mechanics: a double-blind, randomized controlled laboratory study. *Am J Sports Med.* 2013; 41:669–77. [PubMed: 23371471]
81. Myer GD, Sugimoto D, Thomas S, et al. The influence of age on the effectiveness of neuromuscular training to reduce anterior cruciate ligament injury in female athletes: a meta-analysis. *Am J Sports Med.* 2013; 41:203–15. [PubMed: 23048042]
82. Stroube BW, Myer GD, Brent JL, et al. Effects of task-specific augmented feedback on deficit modification during performance of the tuck-jump exercise. *J Sport Rehabil.* 2013; 22:7–18. [PubMed: 23238301]
83. Myer GD, Faigenbaum AD, Ford KR, et al. When to initiate integrative neuromuscular training to reduce sports-related injuries and enhance health in youth? *Curr Sports Med Rep.* 2011; 10:157–66.
84. Myer GD, Faigenbaum AD, Chu DA, et al. Integrative training for children and adolescents: techniques and practices for reducing sports-related injuries and enhancing athletic performance. *Phys Sportsmed.* 2011; 39:74–84. [PubMed: 21378489]
85. Hewett TE, Stroupe AL, Nance TA, et al. Plyometric training in female athletes. Decreased impact forces and increased hamstring torques. *Am J Sports Med.* 1996; 24:765–73. [PubMed: 8947398]
86. Sugimoto D, Myer GD, Bush HM, et al. Effect of compliance with neuromuscular training on ACL injury risk reduction in female athletes: a meta-analysis. *J Athl Train.* 2012; 47:714–23. [PubMed: 23182020]
87. LaBella CR, Huxford MR, Smith TL, et al. Preseason neuromuscular exercise program reduces sports-related knee pain in female adolescent athletes. *Clin Pediatr (Phila).* 2009; 48:327–30. [PubMed: 18832542]

88. Myer GD, Faigenbaum AD, Foss KB, et al. Injury initiates unfavourable weight gain and obesity markers in youth. *Br J Sports Med.* 2013 epub doi: [published Online First: Epub Date].
89. Sugimoto D, Myer GD, Barber Foss KD, et al. Dosage effects of neuromuscular training intervention to reduce anterior cruciate ligament injuries in female athletes: meta- and sub-group analyses. *Sports Med.* 2013 In Press. doi:. [published Online First: Epub Date].
90. Sugimoto D, Myer GD, Hewett TE. Sports specific effectiveness of neuromuscular training on anterior cruciate ligament injury risk reduction in female athletes. *Br J Sports Med.* 2014 Under Review.
91. Myer GD, Brent JL, Ford KR, et al. Real-time assessment and neuromuscular training feedback techniques to prevent ACL injury in female athletes. *Strength Cond J.* 2011; 33:21–35. [PubMed: 21643474]
92. Myer GD, Ford KR, Hewett TE. New method to identify athletes at high risk of ACL injury using clinic-based measurements and freeware computer analysis. *Br J Sports Med.* 2011; 45:238–44. [PubMed: 21081640]
93. Myer GD, Ford KR, Brent JL, et al. An integrated approach to change the outcome part II: targeted neuromuscular training techniques to reduce identified ACL injury risk factors. *J Strength Cond Res.* 2012; 26:2272–92. [PubMed: 22580980]
94. Myer GD, Ford KR, Brent JL, et al. An integrated approach to change the outcome part I: neuromuscular screening methods to identify high ACL injury risk athletes. *J Strength Cond Res.* 2012; 26:2265–71. [PubMed: 22580976]
95. Myer GD, Ford KR, Barber Foss KD, et al. A predictive model to estimate knee abduction moment: Implications for development of clinically applicable patellofemoral pain screening tool in female athletes. *J Athl Train.* 2014 In Press.
96. Schmitz RJ, Ficklin TK, Shimokochi Y, et al. Varus/valgus and internal/external torsional knee joint stiffness differs between sexes. *Am J Sports Med.* 2008; 36:1380–8. [PubMed: 18535251]
97. Myer GD, Ford KR, Hewett TE. The effects of gender on quadriceps muscle activation strategies during a maneuver that mimics a high ACL injury risk position. *J Electromyogr Kinesiol.* 2005; 15:181–9. [PubMed: 15664147]
98. Hewett TE, Myer GD. The mechanistic connection between the trunk, hip, knee, and anterior cruciate ligament injury. *Exerc Sport Sci Rev.* 2011; 39:161–6. [PubMed: 21799427]
99. Zebis MK, Andersen LL, Bencke J, et al. Identification of athletes at future risk of anterior cruciate ligament ruptures by neuromuscular screening. *Am J Sports Med.* 2009; 37:1967–73. [PubMed: 19574475]

Figure 1.

Graphical illustration of reported risk factors for ACL injury and PFP incidence. (A) $ACL_{224547-4952-5456-586061} PFP^{19}$; (B) $ACL_{21-445596} PFP^{10-18}$ and (C) $ACL_{3442515962666897-99} PFP^{19}$. ACL, anterior cruciate ligament; BMI, body mass index; Ext, external; PFP, patellofemoral pain. ITB, iliotibial band; COF, center of force; ROM, range of motion; PF, patellofemoral; ER, external rotation; VMO, vastus medialis oblique; GRF, ground reaction force; VL, vastus lateralis; EMG, electromyography; ST, sartorius.

Figure 2.

Relative cut-off score for increased ACL and PFP injury risk. The highlighted red patella (grey knee) indicates the participants who developed PFP following screening. The red knee represents those participants who went on to ACL injury following screening. ACL, anterior cruciate ligament; CTRL, control; PFP, patellofemoral pain.

Table 1

Cohort descriptive demographics and anthropometrics

Mean	95% CI for mean	
	Lower bound	Upper bound
Height (cm)		
ACL	164.4	163.6
PFP	160.1	158.7
<i>Total</i>	162.6	161.8
Mass (kg)		
ACL	59.2	58.1
PFP	54.3	52.3
<i>Total</i>	57.2	56.1
Age (years)		
ACL	16.1	15.9
PFP	13.3	13.1
<i>Total</i>	15.0	14.7

ACL, anterior cruciate ligament; PFP, patellofemoral pain.