CORRESPONDENCE

Post-Traumatic Stress Disorder—a Diagnostic and Therapeutic Challenge

by PD Dr. med. Dipl.-Biol. Ulrich Frommberger, Dr. phil. Dipl.-Psych. Jörg Angenendt, and Prof. Dr. med. Mathias Berger in issue 5/2014

Incomplete Description

The overall highly successful review article is incomplete with regard to one issue that is crucial for all those who are not experts in psychotrauma. The incompletely listed criteria for relevant trauma (A criterion) may create the impression in non-specialist readers that only "... life-threatening or potentially life-threatening external events and those associated with serious injury, ..." allow a diagnosis of PTSD according to DSM-IV.

This would mean that post-traumatic stress disorder (PTSD) according to DSM-IV cannot be diagnosed in a patient who has been raped but where the rape took place without intention to kill and did not result in serious physical injury, or in a child who experienced a serious physical attack on its mother, even if all other criteria are met.

Obviously the diagnosis in both cases is PTSD, because the complete text in the DSM-IV says:

- The person has been exposed to a traumatic event in which both of the following have been present:
 - "The person experienced, witnessed, or was confronted with an event or events that involved actual or threatened death or serious injury, or a threat to the physical integrity of self or others (2)."
 - "[...] the person's response involved intense fear, helplessness, or horror. Note: In children, this may be expressed instead by disorganized or agitated behavior."

DOI: 10.3238/arztebl.2014.0488a

Dr. med. Roland Ebner

Kinder- und Jugendpsychiatrie am Klinikum Deggendorf
Außenstelle des BKH Landshut
Perlasberger Straße 41, Deggendorf
ebner@degnet.de

REFERENCES

1. American Psychiatric Association: Diagnostic and statistical manual of mental disorders. DSM-IV-TR. 4th edition, Text Revision. Washington DC; American Psychiatric Association 2000.
2. Frommberger U, Angenendt J, Berger M: Post-traumatic stress disorder—a diagnostic and therapeutic challenge. Dtsch Arztebl Int 2014; 111: 59–65.

Conflict of interest statement

The author declares that no conflict of interest exists.

Critical Additions Required

In Stuttgart, Germany, hospital physicians and out-patient care providers have joined in an initiative that aims to improve healthcare for persons who have been through trauma, because the sequelae of trauma are still being overlooked in primary healthcare and are not treated in a manner that is sensitive to trauma. The article is welcome but requires some additions:

- In your article, you warn against possible inflation of the term trauma, but you abbreviate this to "life-threatening or potentially life-threatening external events and those associated with serious injury ...". In the German S3 guideline on post-traumatic stress disorder, the underlying trauma is described more comprehensively as "... reaction following one or more traumatic events, for example the experience of physical or sexual violence, even in childhood ..., rape, violent attacks against self ..." (followed by a list of traumatic situations) "that may be experienced by oneself but also by others. In many cases this results in feelings of helplessness and ... a shock to the perception of self and of the world." It is not only the threat of death and serious physical injury, but also sexualized violence and witnessing such traumatic experiences that can lead to PTSD. DSM-V even includes reports of such experiences gained first-hand as the trauma that triggered PTSD. Unfortunately, neglect and psychological violence in childhood are not included.
- The article does not clarify that SSRIs can be helpful in treating the associated depression, but only in combination with trauma-specific psychotherapy.
- Psychodynamic-imaginative trauma therapy has proved to be successful, especially in complex post-traumatic disorders. Criteria setting out which therapy is most effective for which constellation of symptoms should be developed for the purposes of assessing therapeutic effectiveness.

DOI: 10.3238/arztebl.2014.0488b

Dr. Gabriele Weigel

Specialist in neurology and psychiatry, Stuttgart
dr.g.weigel@web.de

REFERENCES

1. Guido Flatten (eds.): Posttraumatische Belastungsstörung. Leitlinie und Quellentext. Stuttgart: Schattauer Verlag, 2013.
2. American Psychiatric Association: www.DSM5.org (last accessed on 23 May 2014)
3. Luise Reddemann: Psychodynamisch imaginative Traumtherapie PITT – das Manual. 7th edition. Stuttgart, Klett-Cotta 2012.
4. Frommberger U, Angenendt J, Berger M: Post-traumatic stress disorder—a diagnostic and therapeutic challenge. Dtsch Arztebl Int 2014; 111: 59–65.

Conflict of interest statement

The author declares that no conflict of interest exists.

Drugs Can Only Affect Symptoms

The comment relating to the risk of “overdiagnosis” with all its sequelae should be stressed, especially in view of certain figures—for example, it is claimed that 60% of the US population have experienced at least one traumatic event (defined as life-threatening).

It is inevitable that a review article will remain somewhat superficial, but in this case, important associations were not mentioned. The diagnostic criteria and the described therapeutic approaches are valid in that form only in the setting of “simple trauma” (type 1) and, even then, only for persons whose psychological stability is otherwise good. However, we often deal with persons from very problematic family backgrounds. It is especially in this setting that violent and sexual attacks are not altogether rare, which will result in serious psychological disorders, especially in young persons. This group is likely to include many of the persons with comorbid personality disorders such as and in particular borderline personality disorder, addiction disorders, affective disorders, somatoform disorders, but also psychoses. In this setting, the post-traumatic symptoms are difficult to distinguish from any comorbid psychopathologies. Accordingly, different therapeutic strategies need to be employed, which lie in the domain of psychodynamic approaches to therapy and usually necessitate long-term, comprehensive courses of treatment (1). Treatment that is narrowed to trauma-specific subjects would be contraindicated in this setting and might even be massively harmful, although the attribution of symptoms to a single cause and the handy therapeutic approach may hold great appeal for patients, treating healthcare professionals, and the public—including statutory health insurance funds and politicians.

Medications can only affect symptoms; they cannot cure PTSD.

DOI: 10.3238/arztebl.2014.0489a

Dr. med. Peter Brdiczka
Specialist in psychiatry and psychotherapy
Porta Westfalica
mail@praxis-brdiczka.de

REFERENCES

1. Sachse U: Traumazentrierte Psychotherapie. Stuttgart: Schattauer 2004; 115–37.
2. Frommberger U, Angenendt J, Berger M: Post-traumatic stress disorder—a diagnostic and therapeutic challenge. Dtsch Arztebl Int 2014; 111: 59–65.

Conflict of interest statement

The author declares that no conflict of interest exists.

Large Group of Patients Cannot Be Found

Part of the challenge in diagnosing and treating PTSD is based on the unsatisfactory differentiation of section F43 in the ICD-10 (Reaction to severe stress ...), as this includes only three diagnoses: acute stress reaction (F43.0), PTSD (F43.1), and adjustment disorders (F43.2).

A large group of patients cannot be found in section F43 of the ICD-10 and will therefore be grouped under one of these three diagnoses, which are incorrect for such patients—namely, those with “cumulative traumatic stress disorder” (CTSD) (1). The traumatic event has not reached its conclusion but continues in a perpetrator–victim constellation that is usually characterized by psychological violence. Such patients are continually exposed to the perpetrators and cannot remove themselves from the psychological terror. CTSD includes the subdiagnoses “mobbing (adult bullying) syndrome” (2), “stalking syndrome,” “domestic violence syndrome,” etc.

Patients with CTSD either do not meet all the criteria for PTSD or exceed the time limit for the duration of acute stress reactions, or they are even accused by their therapists, with a diagnosis of “adjustment disorder,” of not adjusting to the psychoterror of mobbing, stalking, or domestic violence. In contradiction to the misdiagnosis of an adjustment disorder, CTSD has a traumatic component and does not have any discrepancy between the (low-level) trigger and the (high degree of) suffering. Non-compliance and non-responsiveness to therapy are pre-programmed for patients with CTSD if they are misdiagnosed or even accused of being victims because of an unsatisfactory range of diagnoses and therapeutic options. How are therapies meant to work if patients are continually exposed to the trauma of psychological violence, which therapists cannot stop and can even exaggerate?

Including CTSD in the ICD-10 would document the prevalence rates in section F43.- and thus emphasize the urgency of intervention by the legislators.

DOI: 10.3238/arztebl.2014.0489b

Dr. med. Argeo Bämayer
Specialist in psychiatry and psychotherapy
Coburg
drbaemayer@ungwana.de

REFERENCES

1. Fischer G, Riedesser P: Lehrbuch der Psychotraumatologie. München, Basel: Reinhard-Verlag 1998; 332.
2. Bämayer A: Das Mobbingssyndrom, Diagnostik, Therapie und Begutachtung im Kontext zur in Deutschland ubiquitär praktizierten psychischen Gewalt. Europäischer Universitätsverlag 2012; 136–9.
3. Bämayer A: Kumulative traumatische Belastungsstörung, Verworrene Diagnostik bei Patienten mit Mobbing- und Stalkingsyndrom, Neurotransmitter 2013; 24: 43–5.
4. Frommberger U, Angenendt J, Berger M: Post-traumatic stress disorder—a diagnostic and therapeutic challenge. Dtsch Arztebl Int 2014; 111: 59–65.

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply

Dr. Bämayer points out that the diagnostic criteria for post-traumatic disorders in the ICD-10 reflect some forms of stress to an unsatisfactory degree. It should be borne in mind that these categories include a multitude

of traumatizations and stresses of different grades of severity, but this does not mean these are incorrect because they do not spell out each individual one. It is unfortunate that some patients interpret the term “adjustment disorder” as an accusation. In the ICD-10, the term “adjustment disorder” describes a “reaction,” and several stressors may be present; it is by no means intended to be an accusation. Rather, it characterizes the mental overload that challenges the individual’s personal coping strategies throughout the stressful situation. The potential cumulative effects of stressors is taken into account in the DSM-5 in the trauma A criterion for PTSD as well as for the adjustment disorders.

Dr. Ebner points out the incomplete list of DSM criteria for PTSD in the article. As the complete presentation of the DSM alone would have gone far beyond the scope of the article, we summarized the criteria in key words. The term “potentially life threatening” was intended to include the concrete situations laid out by Dr. Ebner. We did not mention explicitly that this experience can also include witnessing a traumatic event, with all the psychological sequelae afterwards.

Dr. Weigel also points out that our summary of the examples for traumatizations could have been far more detailed, as in the German S3 guideline, for example.

She limits the treatment with SSRIs to depression, which often accompanies PTSD. In doing so she does not take into consideration that the effectiveness of the SSRIs does not solely extend to depression but to all three symptoms clusters of DSM-IV-PTSD. Pharmacotherapy using the SSRIs sertraline or paroxetine alone is effective in PTSD. In spite of this, from a clinical perspective, the S3 guideline from the Association of Scientific Medical Societies in Germany (AWMF) regards psychotherapy as the first-line treatment.

The Cochrane analysis of the effectiveness of combination treatment for psychotherapy with pharmacotherapy in PTSD did not confirm the definite superiority of combination treatment versus monotherapy with psychotherapy or pharmacotherapy. This contradicts the clinical course of action, in which combination treatment is often used in cases of severe disorder and for the clinical reasons given.

The problem with psychodynamic therapies lies in the fact that their effectiveness has not been the subject of much scientific study. A single study from 1989, with a small number of cases, was conducted as a randomized controlled trial. And the only study of the effectiveness of inpatient psychodynamic imaginative trauma therapy (PITT) showed only small effect sizes. This is not to say that these approaches are ineffective. They merely have not been subject to enough scientific study; what is urgently required is for this situation to improve and change, so that more studies are undertaken, in much the way that approaches such as cognitive behavioral therapy, EMDR, and other therapeutic methods are being evaluated.

Unfortunately, robust data are lacking to answer the last mentioned and clinically important question regarding the criteria for differential indications—that is,

which therapy should be used in which constellation of symptoms.

Dr. Brdiczka puts PTSD into context with type 1 traumata and regards the concept as too narrow. How, then, can it be explained that using this concept and the trauma-focused therapies derived from it achieved a substantial reduction in symptoms and clinical improvement within a short time even in those patients whose treatment Dr. Brdiczka suspects of being “contraindicated” and potentially “massively harmful”? Consideration of some of the mentioned problems did, after all, result in an additional symptom cluster being introduced to the DSM-V and in the recommendation to introduce “complex PTSD” in the ICD-11. Although this makes perfect sense for many clinicians, it is still scientifically controversial.

Another question relates to the term “cure.” In the studies we referred to, this term was not the object of study, and neither was the definition of a cure from trauma. A substantial reduction of symptoms was the primary objective in the studies investigating psychotherapy as well as those investigating pharmacotherapy. In the therapies found to be most successful, both categories are achieved to a relevant degree in about two-thirds of cases. Many chronically ill patients continue to have symptoms. Once some important therapeutic objectives have been achieved, the trauma loses its horror and can be better integrated into the individual’s biography; handling everyday life in an appropriate manner becomes possible again.

Classification systems are always an expression of the insights and influences of their time.

DSM-V as well as the changes planned for ICD-11 characterize the continual developments in the subject of psychotraumatology, but they will not be able to clarify all the controversial points of view expressed in the correspondence.

In conclusion, each topic mentioned in our article could have been presented in much more detail. We provided an extensive list of references and relevant literature for those who want to explore the subject in greater detail.

DOI: 10.3238/arztebl.2014.0489c

PD Dr. med. Dipl.-Biol. Ulrich Frommberger
Offenburg
ulrich.frommberger@medclin.de

REFERENCES

1. Frommberger U, Angenendt J, Berger M: Post-traumatic stress disorder—a diagnostic and therapeutic challenge. *Dtsch Arztebl Int* 2014; 111: 59–65.

Conflict of interest statement

Dr. Frommberger has received reimbursement of conference fees and travel expenses from Servier, Astra-Zeneca, Lilly, Janssen-Cilag, GlaxoSmithKline, and Pfizer. He has received lecture fees from Astra-Zeneca and Servier. He is a supervisor and assistant professor at the Freiburg Behavioral Therapy Training Institute.