Selective and Superselective Angiography of Pediatric Moyamoya Disease Angioarchitecture: the Anterior Circulation

GERASIMOS BALTSAVIAS¹, ANTON VALAVANIS¹, VENKO FILIPCE¹, NADIA KHAN²

¹ Department of Neuroradiology, University Hospital Zurich; Zurich, Switzerland

² Moyamoya Centre, University Children's Hospital Zurich; Zurich, Switzerland

Key words: Moyamoya paediatric disease, moyamoya network, moyamoya vessels, superselective angiography

Summary

The angioarchitecture of the so-called moyamoya vessels in children has not been explicitly analyzed. We aimed to investigate the precise anatomy of the vascular anastomotic networks in patients with childhood moyamoya disease.

Six children diagnosed with moyamoya disease for the first time underwent an angiographic investigation with selective and superselective injections. We recorded the arterial branches feeding the moyamoya anastomotic networks, their connections and the recipient vessels.

Depending on the level of the steno-occlusive lesion, the feeding vessels included the medial striate arteries, the perforators of the choroidal segment of the carotid, the uncus artery, the medial and lateral branches of the intraventricular segment of the anterior choroidal artery, perforators of the communicating segment, the superior hypophyseal arteries, the prechiasmatic branches of the ophthalmic artery, the ethmoidal arteries and the dural branches of the cavernous carotid. Through connections, which are described, the recipient vessels were the lateral striate arteries and the middle cerebral, the medial striate arteries and the anterior cerebral, medullary arteries around the ventricular system, anterior temporal branches of the middle cerebral, orbitofrontal and frontopolar branches of the anterior cerebral, as well as other cortical branches of the anterior and middle cerebral territories.

The use of high quality selective and superselective angiography enabled us to clearly demonstrate for the first time aspects of the microangiographic anatomy of the moyamoya anastomotic network previously only vaguely or incompletely described.

Introduction

The angiographic characteristics of moyamoya disease described to date¹⁻⁷ were exclusively based on the images obtained during a selective ICA or vertebral artery (VA) contrast injection. Among the factors influencing the information that can be extracted from an angiographic investigation is the selectivity of the injection. It is a fact that the exact relations, connections and territory of the arterial branches which participate in the complex collateral network mostly at the base of the brain cannot be reliably analysed when multiple overprojections inevitably occur in an ICA or VA injection.

In this study, the high-quality selective and magnified angiographic images were in some cases supplemented by several superselective injections so that the angioarchitecture of the collateral circulation of those cases was more clearly demonstrated.

Many authors agree that basal moyamoya vessels are dilated perforating vessels or branches of the ICA, which ordinarily supply brain structures, anastomosing to distal cortical branches through a collateral network in re-

sponse to the increasing hypoperfusion⁸⁻¹¹. However, other descriptions refer to new anastomotic vessels¹² and true neoangiogenesis around the circle of Willis¹³⁻¹⁴.

Materials and Methods

During 2012, 19 children diagnosed with moyamoya disease or syndrome, were hospitalized in the Moyamoya Centre, Children's University Hospital in Zurich. All children underwent digital subtraction angiography (DSA) at our department. Five patients had a follow-up angiogram only after a revascularization operation with initial angiograms having been performed in referring institutes, whereas in 14 cases the angiogram performed at our department was the first diagnostic one. Eight of these patients were considered moyamoya syndrome and were not included in this study, keeping the studied population homogeneous. The DSA imaging of six Caucasian patients (male:female = 3:3), aged 2-12 years, with newly diagnosed moyamoya disease, were retrospectively analysed.

The present report only describes the angiographic features of the anterior circulation. The posterior communicating artery (Pcom) supplied either through the ICA or the basilar ar-

tery (BA) corresponds to the caudal division of the ICA^{15,16} and was considered part of the posterior circulation, which will be analyzed in a separate publication.

All six patients underwent six-vessel selective angiography. In three patients, superselective microcatheterizations (Elite 1.5F Stryker Neurovascular, Fremont, CA, USA) were performed in the anterior circulation by the first author (GB). Informed consent was obtained in all cases. The motivation-indication for superselective injections was diagnostic uncertainty about stenosis or occlusion at the area of ICA bifurcation and degree of involvement of the MCA and ACA, as well as the extent of collateralization from moyamoya vessels. Depending on the stage of angiopathy, distal arterial filling and extent of the deep and dural collaterals, optimal planning of the number, location and type of revascularisation procedures in all the affected arterial territories could be undertaken. The microcatheterizations were performed with the same angiographic setup, through the angiographic catheter (5Fr "Val" catheter, Cook Medical Inc., Bloomington, IN, USA). The microcatheter (Spinnaker Elite 1.5 Fr Target Therapeutics/ Boston Scientific, Fremont, California, USA) used was exclusively flow-directed with the microguidewire used only for proximal support. In this way, the distal supraclinoid ICA

Table 1 Suzuki grades and steno-occlusive changes. P1, P2, P3: segments of the posterior cerebral artery.

Pt	Suzuki grade		Age	M/F	Right hemisphere anterior circulation	Right hemisphere posterior circulation	Left hemisphere anterior circulation	Left hemisphere posterior circulation
	R	L						
A	5	4	11	M	ICA occlusion at ophthalmic segment	Severe stenosis of P1 segment	Severe stenosis at communicating segment	Slight stenosis of P2 segment
B	3	3	7	F	ICA occlusion at choroidal segment	No steno-occlusion	Severe stenosis at choroidal, A1, M1 segments	No steno-occlusion
C	3	3	2	F	Stenosis at ICA bifurcation	Mild stenosis of P3 segment	Severe stenosis at ICA bifurcation	Occlusion of P2 segment
D	2	2	12	F	Severe stenosis at ICA bifurcation	No steno-occlusion	Severe stenosis at ICA bifurcation, M1 occlusion	No steno-occlusion
E	3	3	2	M	Severe stenosis at ICA bifurcation, M1 occlusion	No steno-occlusion	Severe stenosis at ICA bifurcation, A1 occlusion	No steno-occlusion
F	2	2	3	M	Severe stenosis at ICA bifurcation	No steno-occlusion	Severe stenosis at ICA bifurcation	Occlusion of P2 segment

Figure 1 Patient F. Anteroposterior view of a superselective injection of a distal right highly stenotic ICA. Distal to the tip of the microcatheter both M1 (arrows) and A1 (thin arrows) are opacified. None of the striate perforators normally arising from M1 are anterogradely opacified. An early origin of the orbitofrontal branch of the ACA from the A1 is noticed (arrowhead).

A

B

Figure 2 Patient F. A) Anteroposterior view of a superselective injection of a medial striate artery (arrow) arising just proximal to the major stenotic site of the ICA bifurcation. An opacified intrastriatal network of vessels fills retrogradely a lateral striate artery which exits (short arrow) at the MCA trunk (thin arrows left) distal to the level of the major stenosis as well as to a small medial striate branch of the ACA exiting at the level of the mid A1 (arrowhead) and opacifying the distal ACA (thin arrows right). B) The same injection in lateral view. The tip of the microcatheter and the injected medial striate artery (arrow) opacifying the striatum and through retrograde filling of a lateral striate and a more medial striate artery, supplying the distal ACA (arrowhead) and MCA (short arrow).

and/or the anterior choroidal artery were spontaneously catheterized without wire manipulations. Slow withdrawal of the microcatheter taking advantage of the pulse-induced movements of its tip, enabled more proximal catheteriza-

tions. In none of the superselective injections did the microcatheter have a wedged position. No complications occurred.

Particular attention was paid to identification of the individual arterial branches, which

Figure 3 Patient E. A) Anteroposterior view of a superselective injection of the anterior choroidal artery (long arrows) with the origin of the rostral uncus artery just distal to the tip of the microcatheter with demonstration of its characteristic meandric collateral connection (small arrowheads) with the anterior temporal branch (large arrowhead) of the MCA (short arrows). B) Lateral view of the same injection, where the exact site of origin of the uncus artery (small arrowhead) from the anterior choroidal (long arrows) is shown, as well as the anterior temporal branch (large arrowhead) and the distal MCA (short arrows).

Figure 4 A) Patient E. Another example of an uncus artery (short arrow) with the characteristic meandric (in anteroposterior view) course of its connection with the temporal branch (arrowhead) of the MCA (thin arrows) in a superselective injection of a left anterior choroidal artery (long arrow). B) Patient D. Anteroposterior view of an ICA injection where the meandric laterally directed connection (small thin arrows) of the uncus artery to the temporal branch of the MCA (arrowheads) can be distinguished. Anterior choroidal artery (long thin arrows).

made up the so-called basal “moyamoya anastomotic network”, as well as their connections and direction of blood flow. Included here are the regularly visible branches as well as arterial

branches usually not visible in a normal angiogram which may both appear dilated. When part of the entire course and potential anastomoses of an arterial branch were not clearly

Figure 5 Patient F. Anteroposterior (A) and lateral (B) views of a superselective superior hypophyseal artery (long arrow) injection. Superomedial course of the artery and anastomoses (arrowheads) with diencephalic perforators of distal A1 and reconstruction of both ACA (short arrows).

Figure 6 Patient C. Lateral view of an ICA injection with clear opacification of a dilated proximal prechiasmatic ophthalmic branch (arrow) following a superomedial course and probably connected to other chiasmatic and hypothalamic perforators.

Figure 7 Patient E. Lateral view of a superselective injection of the ophthalmic artery (long arrow) which through the ethmoidal arteries retrogradely supply the orbitofrontal branches and the A2 segment (arrowheads) as well as the polar branches (short arrow – through the artery of the falx) of the ACA.

seen, then it was recorded as “not identified”. The description of the supraclinoid ICA segments based on anatomical-surgical landmarks¹⁷ rather than embryologic considerations, was

deemed more convenient for the purpose of this study.

For this retrospective study no Institutional Review Board approval was necessary.

Figure 8 Patient B. Anteroposterior (A) and lateral (B) views of a superselective injection of a left anterior choroidal artery (long thick arrow) with supply through its medial and lateral branches (long thin arrows) of medial striate arteries (short thin arrows) exiting to the M1 (thick arrowhead) and A1 (thick short arrow). A very tortuous arterial branch (thin arrowheads) running laterally (anteroposterior view) and retrogradely opacified through the lateral intraventricular branch of the anterior choroidal is also exiting into the MCA at the M1-M2 level.

Figure 9 Patient E. A) Anteroposterior view of a right anterior choroidal (long arrow) injection. Its lateral intraventricular branch (long thick arrow) retrogradely supplies (better visible in this view) transinsular fine branches exiting into the distal M2 segment (thin short arrows) as well as very tortuous branches exiting into the M1-M2 segment (arrowheads) of the MCA (long thin arrows). Short arrow: exit of a striate artery into the M1 segment retrogradely opacified through the same lateral intraventricular branch. Thick arrowhead: choroid plexus. B) Lateral view of the same injection. Long arrow: tip of the microcatheter in the anterior choroidal artery. Thick short arrow: its lateral intraventricular branch. Thick arrowheads: choroid plexus. Thick long arrow: connection (better visible in this view – overprojected in the anteroposterior view) of the lateral branch with the distal segment of a lateral striate (thin arrowheads). The very tortuous branches exiting (arrowheads) into the M1-M2 segment of the MCA (long thin arrows) and the striate artery (short arrows) exiting into the M1 segment (washed out).

Figure 10 Patient B. Lateral view of a superselective left anterior choroidal (short thin arrow) injection – late arterial phase. Its lateral intraventricular branch (short thick arrow) mainly supplies the atrial and temporal dilated medullary arteries (lower oval). The more medial branch (thin long arrow) in its course supplies the distal segment of a striate artery (long thick arrow) which opacifies the dilated medullary arteries of the posterior body of the lateral ventricle (upper oval) and then more anteriorly is connected with the distal segment of a medial striate artery opacifying the vascular network of the head of the caudate (circle) and retrogradely the proximal medial striate artery (thin arrowhead) which exits to the distal A1 (lower thick arrowhead) MCA (upper thick arrowhead).

Results

According to Susuki's classification², one hemisphere was at stage 5, one at stage 4, six hemispheres were at stage 3 and four hemispheres at stage 2. A more detailed description of the stenocclusive changes is shown in Table 1.

Table 2 presents the moyamoya anastomotic networks in a coherent way. The identified arterial branches participating in the networks are presented from distal to proximal carotid segments with their connections as well as the patients (hemispheres) in which these vessels appeared.

The superselective injection of a stenotic but still not occluded distal ICA showed that neither the M1 segment of the MCA nor the A1 segment of the anterior cerebral artery (ACA) supplied any of their perforating branches (Figure 1).

Perforators arising proximal to the steno-occlusive site through collateral connections retrogradely supplied the perforators of the distal M1 segment and then the distal cortical territories. More precisely, a medial striate artery arising from the distal choroidal segment just proximal to the most stenosed ICA site was injected and demonstrated the intrastriate collateral network retrogradely supplying the lateral striate branches and then the distal MCA (Figure 2).

Our study identified a rostral uncus branch originating from the proximal anterior choroidal artery. It was a significant collateral to the

MCA at the M1 level through its anterior temporal branches. This collateral connection to the anterior temporal branches of the MCA had a characteristic meandric tortuosity and a lateral course in the anteroposterior projection (Figure 3) and in retrospect can be relatively easily recognised even in non-superselective injections (Figure 4).

In a superselective injection of a dilated superior hypophyseal artery, the bilateral A2 segments of the ACA were opacified through an anastomotic network at the level of the lamina terminalis which retrogradely irrigated distal diencephalic perforators of the A1, possibly the commissural and/or preoptic arteries¹⁸ (Figure 5).

The distal ACA, like the orbital and frontopolar branches, were often supplied from the posterior and anterior ethmoidal arteries and artery of the falx (Figure 7).

In one case the superselective injection of the anterior choroidal artery retrogradely opacified the striate perforators of the M1 and a medial striate perforator of the A1 supplied through the medial branch of its intraventricular segment. The lateral branch was connected to a remarkably tortuous artery running laterally parallel to the insula and opacified the MCA at the proximal M2 segment (Figure 8).

In the contralateral hemisphere of the same patient, the lateral striate arteries as well as a similarly tortuous arterial branch running lateral to the lateral striates were retrogradely

opacified emptying into the MCA, supplied by the lateral branch of the intraventricular segment of the anterior choroidal artery (Figure 9).

The retrograde filling of the dilated medullary network at the area of the temporal horn, atrium and occipital horn as well as at the posterior part of the body of the lateral ventricles was supplied through branches of the intraventricular segment of the anterior choroidal artery itself. The medullary arteries located in the more anterior lateral ventricles were supplied by the distal segment of the lateral striate arteries fed either by medial striate branches or Pcom perforators or the intraventricular branches of the anterior choroidal artery (Figure 10).

Discussion

Since the first descriptions and classification^{2,19} of the moyamoya disease, many aspects related to its angiographic features have been described^{1,4,6,7,20}. However, an accurate and detailed description of the vessels constituting the so-called “moyamoya vessels”, their connections and direction of flow using high quality selective and superselective angiographic injections has never been done before. In fact, these vessels are often presented as chaotic in the literature¹⁴.

The original 1968 publication of Nishimoto et al. mentioned the moyamoya anastomotic network playing the role of collateral supply due to the stenosis of the ICA only as a hypothesis. In 1969, Suzuki et al., after studying 11 paediatric cases of moyamoya disease who underwent cerebral angiography, supposed that the “abnormal rete vasculosum” represented a new type of collateral circulation, but they did not analyse it further. In 1972, Handa et al. studied the angiographic features of 16 children with moyamoya and reported their autopsy findings confirming that the vascular network was constituted by the perforating branches of ICA, MCA, ACA, PCA, Pcom and choroidal arteries. They pointed out that angiographically the relation of the vessels in the moyamoya anastomotic network was difficult to determine because of the heavy stain. However, based on five angiographic investigations, they noticed that the cortical branches of the MCA and ACA distal to the stenosed or occluded ICA bifurcation were reconstructed by the moyamoya anastomotic network. In 1974, Crouzet et al. described some characteristics of the moyamoya network more

specifically and defined three types of moyamoya collaterals: those supplying retrogradely the lateral striates and the M1 segment of the MCA (type 1); those supplying retrogradely insular perforators and the M2 segment of the MCA (type 2); and those supplying retrogradely the cortical convexial perforating branches of the M3 and M4 segments of the MCA. The authors failed to define which vessels of the network specifically fed each of those collateral routes although the concept of afferent vessels-moyamoya network-efferent vessels was well applied in their analysis of the ethmoidal collaterals. In 1980, Takahashi's angiographic study of seven patients (three of them children) with moyamoya disease attempted a more detailed description of the vessels which constitute the moyamoya anastomotic network and its interconnections. He identified the vessels comprising the network as dilated perforating arteries similarly to previous authors and referred to numerous anastomoses among them. He reported that all moyamoya networks anastomosed with the insular segments of the sylvian MCA through the claustrum and the insular cortex. He also reported that the anastomotic vessels usually arose from the lateral lenticulostriate arteries and that the direction of flow was usually from the network to the sylvian vessels but it was reversed when the MCA was “not completely occluded”. In 1988, Satoh et al. described the angiographic findings of 34 newly diagnosed paediatric cases with moyamoya disease and distinguished the feeding branches of the moyamoya anastomotic network derived from the ICA group and PCA group. The feeding branches of the ICA group were identified as the “supraclinoid ICA, the Pcom, the anterior choroidal artery and the medial and lateral lenticulostriate arteries”. This statement together with most of the descriptions in the above-mentioned reports summarize and actually perpetuate an ambiguity regarding the exact feeding vessels, the recipient vessels and the direction of flow. Additionally, they fail to identify the individual feeding branches and their interconnections.

The arterial vessels which participate in the known basal moyamoya anastomotic network in childhood moyamoya disease are dilated but otherwise normal proximal twigs of the ICA and its branches, which provide collateral supply to arterial territories distal to the steno-occlusive sites of ICA termination and proximal A1 and M1 segments of the ACA and MCA.

Table 2 Vessels constituting the moyamoya anastomotic networks.

Vessel of origin	Feeding vessel	Course	Recipient vessel(s)	Hemispheres
Choroidal segment ICA	Medial striate artery	Endostriatal network	Lateral striate artery to M2	F
	Medial striate artery	Not identified	Not identified	A,E,E
	Perforators of the choroidal segment	Not identified	Not identified	A,B,B,C,C,D,F,F
Anterior choroidal	Uncal artery	Anterolateral to the temporal pole	Ant. temporal branch of MCA	B,B,D,D,E,E,F
	Medial intraventricular branches	Subependymal network	Medullary arteries	B,E
	Medial intraventricular branches	Subependymal network	Medial striate artery to A1	B
	Lateral intraventricular branches	Subependymal network	Medullary arteries and/or distal convexial cortical arteries	B, D,E,C,C,F
	Lateral intraventricular branches	Subependymal network	Lateral striate a. and claustral (?) artery to M2	D,E,E
Communicating segment ICA	Diencephalic perforator	Probably subependymal to the lateral aspect of striatum	Medullary arteries at the body of lateral ventricle	C
	Not identified	Not identified	Not identified	E
Ophthalmic segment ICA	Superior hypophyseal artery	Circum-infudibular plexus	Diencephalic perforators to A1	F
	Not identified	Not identified	Not identified	B,C,D,D,E,E,F
Ophthalmic art.	Prechiasmal arteries	Not identified	Not identified	B,C,C
	Post ethmoidal artery, ant ethmoidal artery, artery of the falx	Duro-cortical subarachnoid connections	Fronto-orbital and frontopolar aa. of ACA	B,B,C,D,D,E,E,F,F
Extradural ICA	ILT and/or MHT	Sup. orbital fissure	Ophthalmic a.	A
	Tentorial artery	Tentorial edge to duro-cortical connections	Medial occipital lobe cortical a./or not identified	B,B,C
	MHT-inf Hypophyseal artery	Circum-infudibular plexus	Diencephalic perforators	A
	MHT-Inf Hypophyseal artery	Circum-infudibular plexus	Not identified	B,B
ECA	MMA, occipital artery -mastoid branch		Parietal and Temporal cortical branches	A,B,B,C,D

Display of a patient's hemisphere twice (e.g. B,B) signifies appearance of the described vessels in both hemispheres.
MHT: meningohypophyseal trunk, MMA: middle meningeal artery, ILT: inferolateral trunk.

These vessels will vary according to the progression of the disease and the level of major affection of the vasculature. The example of the uncal artery is noteworthy. The rostral uncal artery is usually the most proximal lateral branch of the anterior choroidal artery. It can also arise separately from the choroidal segment of the ICA or can share a common origin with the anterior choroidal. It may supply the rostral uncus, part of the amygdala, the pyriform cortex or the anterior parahippocampal gyrus. It anastomoses with PCA-Pcom branches and early branches of the MCA²¹. To our knowledge, the participation and role of this artery in supplying the MCA through temporal leptomeningeal anastomoses has never been reported. Additionally, it demonstrates that the basal moyamoya vessels are not composed exclusively of perforators but also of classic leptomeningeal vessels and network similar to the retrosplenial one.

Superselective atraumatic microcatheterization is a valuable diagnostic tool established and routinely used in our department since 1986. Its diagnostic yield and safety for children and adults has been proved for almost every aspect of cerebrovascular pathology²²⁻²⁹. The role of superselective angiography for the evaluation of newly formed collaterals after revascularization procedures in children with moyamoya, as well as its safety despite catheterization of the vasospasm-prone middle meningeal and temporal arteries or of the small-calibre anterior choroidal artery for an endovascular approach to distal aneurysms, was also reported by other groups³⁰⁻³⁴. In moyamoya disease, superselective injections can offer more reliable information on the angioarchitecture and dynamic vascular collaterals where many small-sized vessels are usually overprojected. For surgical planning, the superselective injections in our patients provided not only additional information complementary to the clinical and haemodynamic evaluation but helped highlight the need for multiple revascularisation procedures i.e. revascularisation not only for the MCA territory but also for the ACA and PCA territories.

Takahashi's report that the anastomotic vessels usually arose from the lateral lenticulostriate arteries was not confirmed by our findings. In our study, the lateral lenticulostriate arteries were in all cases recipient vessels retrogradely opacified subsequently supplying the distal M1 or M2 segments of the MCA which is proximally occluded or stenosed. Therefore, the anastomotic vessels arose from more proximal arteri-

al branches, including 1) the medial striate branches; 2) other perforators of the choroidal segment of the ICA just proximal to the major stenotic site; 3) anterior choroidal branches; 4) Pcom branches; 5) more proximal ICA segments; 6) the ophthalmic artery. Based on our findings, Takahashi's description of flow direction and its reversal from the MCA to the network vessels when the MCA was "not completely occluded" is confusing. Reversal of flow from the MCA to the network in cases of high-grade stenosis but incomplete occlusion of the MCA would mean actually anterograde flow to the lateral striate arteries. This was not observed in our series and it is rather unlikely to occur since such an anterograde flow to the lateral striates (and then apparently to medullary arteries) would correspond to a by-pass of a steno-occlusion located distal to their origin, whereas the primary site of steno-occlusion in moyamoya paediatric disease is typically proximal to their origin.

Our finding of reversed flow in the lateral striates coming from anastomotic connections with proximal perforating branches does not mean that a stenotic ICA bifurcation always induces reversal of flow in the striate perforators, regardless the stage of the disease. Some of the medial perforators, depending on their sites of origin and on the availability of other collateral channels may have anterograde flow even in highly stenotic, pre-occlusive stages. An antero-grad flow pattern is unlikely to exist in the lateral striates which arise distal to the site of major and progressive stenosis in the course of the paediatric disease. In adult moyamoya or other syndromic phenotypes where steno-occlusion of the MCA follows more atypical patterns, antero-grad flow may be observed³⁵.

Our study confirmed the retrograde supply of the lateral striates and the M1 or M2 (depending on the site of their origin) segment of the MCA reported by Crouzet et al. and Takahashi, and showed that the lateral striates were actually fed by intraventricular branches of the anterior choroidal artery, which was not previously demonstrated.

The question whether moyamoya vessels are dilated pre-existing vessels or newly formed connections could not be convincingly addressed in this study. The remarkably tortuous arteries supplied by the anterior choroidal artery and running parallel to the insula and exiting into the proximal M2 segment, was a considerable finding. In case these arteries are pre-ex-

isting vessels, based on published anatomic descriptions they could correspond to the claustral arteries³⁶. To our knowledge, such vessels were not previously demonstrated by angiography.

The retrograde supply of insular perforators was also demonstrated, but fed by lateral recurrent intraventricular branches of the anterior choroidal artery. Although the possibility of a connection through intraventricular branches of the anterior choroidal is supported by our findings, this certainly does not verify the universal connection between lateral striates and insular perforators in all anastomotic networks as Takahashi hypothesized. Moreover, no other vessels supplying the insular segments of the sylvian MCA through the claustrum and the insular cortex were identified in this study.

Additionally, Takahashi pointed out that most often the lenticulostriates anastomosed with medullary arteries retrogradely supplying the cortical vessels. In our cases this was demonstrated only for medullary arteries connected with the body of the lateral ventricles. The remainder of the medullary arteries, connected to the posterior body of the lateral ventricles, the atrium and the posterior and inferior horn, were supplied retrogradely through intraventricular branches of the anterior choroidal artery. The role of the anterior choroidal artery in the supply of the striate arteries, M1 and M2 segments and the medullary arteries as demonstrated in this study has not been previously reported.

Although it appears in Table 2 that in most of our cases the multiple fine vessels arising from the ophthalmic segment of the ICA and their connections were not identified, we assumed that most if not all of those contributed to the hypothalamic network identified in the superselective injection of the superior hypophyseal artery. In a few cases the prechiasmal branches of the ophthalmic artery were dilated, apparently due to their participation in the collateral network. Most probably, although not demonstrated, their collateral contribution to

the vasculature distal to the steno-occlusive sites would follow a pathway through connections with other chiasmatic and hypothalamic branches (Figure 6).

Several factors may limit our study. In particular, only half of our cases included superselective injections. The level of each superselective injection varied, the stages of the disease varied from patient to patient, and potential anatomic variations are always possible.

Conclusions

Previous descriptions of a chaotic “network of abnormal vessels” create a vaguely defined picture of the anastomotic connections in moyamoya disease angioarchitecture. High quality selective and superselective angiography of the anterior territory revealed previously unreported connections within the moyamoya anastomotic network and helped in clarifying the anatomy of a rather organised network which has distinct feeding branches, characteristic course and anatomically expected recipient vessels.

Integrating these findings with those in the posterior circulation in moyamoya paediatric disease will facilitate more comprehensive conclusions on the architecture of collateral networks in moyamoya patients and this may also contribute to a new staging of the disease with clinical and therapeutic relevance.

Moreover these findings have possible implications for response of the normal cerebral vasculature to ischaemic conditions.

Acknowledgments

We wish to thank Professor Scott W. Atlas and Professor Val M. Runge for their valuable comments on the manuscript and knowledgeable suggestions.

References

- 1 Satoh S, Shibuya H, Matsushima Y, et al. Analysis of the angiographic findings in cases of childhood moyamoya disease. *Neuroradiology*. 1988; 30 (2): 111-119. doi: 10.1007/BF00395611.
- 2 Suzuki J, Takaku A. Cerebrovascular “moyamoya” disease. Disease showing abnormal net-like vessels in base of brain. *Arch Neurol*. 1969; 20 (3): 288-299. doi: 10.1001/archneur.1969.00480090076012.
- 3 Hasuo K, Tamura S, Kudo S, et al. Moyamoya disease: use of digital subtraction angiography in its diagnosis. *Radiology*. 1985; 157 (1): 107-111.
- 4 Takahashi M. Magnification angiography in moyamoya disease: new observations on collateral vessels. *Radiology*. 1980; 136 (2): 379-386.
- 5 Nishimoto A, Takeuchi S. Abnormal cerebrovascular network related to the internal carotid arteries. *J Neurosurg*. 1968; 29 (3): 255-260. doi: 10.3171/jns.1968.29.3.0255.

- 6 Crouzet G, Aguetaz G, Pellat J, et al. Collateral circulation in Moyamoya. *J Neuroradiol.* 1974; 1: 87-101.
- 7 Handa J, Handa, H. Progressive cerebral arterial occlusive disease: analysis of 27 cases. *Neuroradiology.* 1972; 3 (3): 119-133. doi: 10.1007/BF00341494.
- 8 Suzuki J, Kodama N. Moyamoya disease--a review. *Stroke.* 1983; 14 (1): 104-109. doi: 10.1161/01.STR.14.1.104.
- 9 Takanashi J. Moyamoya disease in children. *Brain Dev.* 2011; 33 (3): 229-34. doi: 10.1016/j.braindev.2010.09.003.
- 10 Takebayashi S, Matsuo K, Kaneko M. Ultrastructural studies of cerebral arteries and collateral vessels in moyamoya disease. *Stroke.* 1984; 15 (4): 728-732. doi: 10.1161/01.STR.15.4.728.
- 11 Scott RM, Smith JL, Robertson RL, et al. Long-term outcome in children with moyamoya syndrome after cranial revascularization by pial synangiosis. *J Neurosurg.* 2004; 100 (2 Suppl Pediatrics): 142-149.
- 12 Smith ER. Moyamoya arteriopathy. *Curr Treat Options Neurol.* 2012; 14 (6): 549-556. doi: 10.1007/s11940-012-0195-4.
- 13 Currie S, Raghavan A, Batty R, et al. Childhood moyamoya disease and moyamoya syndrome: a pictorial review. *Pediatr Neurol.* 2011; 44 (6): 401-413. doi: 10.1016/j.pediatrneurol.2011.02.007.
- 14 Kassner A, Zhu XP, Li KL, et al. Neoangiogenesis in association with moyamoya syndrome shown by estimation of relative recirculation based on dynamic contrast-enhanced MR images. *Am J Neuroradiol.* 2003; 24 (5): 810-818.
- 15 De Vriese B. Sur la signification morphologique des artères cérébrales. *Arch Biol.* 1904/5; (21): 357-457.
- 16 Lasjaunias P, Berenstein A., terBrugge KG. Intradural arteries. Clinical vascular anatomy and variations. In: *Surgical Neuroangiography. Volume 1.* Berlin, Germany: Springer-Verlag; 2001. p. 479-629. doi: 10.1007/978-3-662-10172-8.
- 17 Gibo H, Lenkey C, Rhoton AL Jr. Microsurgical anatomy of the supraclinoid portion of the internal carotid artery. *J Neurosurg.* 1981; 55 (4): 560-574. doi: 10.3171/jns.1981.55.4.0560.
- 18 Marinković SV, Milisavljević MM, Marinković ZD. Microanatomy and possible clinical significance of anastomoses among hypothalamic arteries. *Stroke.* 1989; 20 (10): 1341-1352. doi: 10.1161/01.STR.20.10.1341.
- 19 Takeuchi K. Occlusive diseases of the carotid artery: Especially on their surgical treatment. *Shinkei Shimpō.* 1961; 5: 511-543.
- 20 Rosengren K. Moya-moya vessels. Collateral arteries of the basal ganglia. Malignant occlusion of the anterior cerebral arteries. *Acta Radiol Diagn (Stockh).* 1974; 15 (2): 145-151.
- 21 Marinković SV, Milisavljević MM, Vucković VD. Experimental and clinical studies: microvascular anatomy of the uncus and the parahippocampal gyrus. *Neurosurgery.* 1991; 29 (6): 805-814. doi: 10.1227/00006123-199112000-00001.
- 22 Wehrli M, Lieberherr U, Valavanis A. Superselective embolization for intractable epistaxis: experiences with 19 patients. *Clin Otolaryngol Allied Sci.* 1988; 13 (6): 415-420. doi: http://dx.doi.org/10.1111/j.1365-2273.1988.tb00314.x.
- 23 Valavanis A. The role of angiography in the evaluation of cerebral vascular malformations. *Neuroimaging Clin N Am.* 1996; 6 (3): 679-704.
- 24 Khan N, Hajek M, Antonini A, et al. Cerebral metabolic changes (18F-FDG PET) during selective anterior temporal lobe amobarbital test. *Eur Neurol.* 1997; 38 (4): 268-275. doi: 10.1159/000113393.
- 25 Wieser HG, Muller S, Schiess R, et al. The anterior and posterior selective temporal lobe amobarbital tests: angiographic, clinical, electroencephalographic, PET, SPECT findings, and memory performance. *Brain Cogn.* 1997; 33 (1): 71-97. doi: 10.1006/brcg.1997.0885.
- 26 Hajek M, Valavanis A, Yonekawa Y, et al. Selective amobarbital test for the determination of language function in patients with epilepsy with frontal and posterior temporal brain lesions. *Epilepsia.* 1998; 39 (4): 389-398. doi: 10.1111/j.1528-1157.1998.tb01391.x.
- 27 Valavanis A, Yasargil MG. The endovascular treatment of brain arteriovenous malformations. *Adv Tech Stand Neurosurg.* 1998; 24: 131-214. doi: 10.1007/978-3-7091-6504-1_4.
- 28 Tanaka M, Valavanis A. Role of superselective angiography in the detection and endovascular treatment of ruptured occult arteriovenous malformations. *Interv Neuroradiol.* 2001; 7 (4): 303-311.
- 29 Tanaka M, Imhof HG, Schucknecht B, et al. Correlation between the efferent venous drainage of the tumor and peritumoral edema in intracranial meningiomas: superselective angiographic analysis of 25 cases. *J Neurosurg.* 2006; 104 (3): 382-388. doi: 10.3171/jns.2006.104.3.382.
- 30 Kashiwagi S, Kato S, Yasuhara S, et al. Use of a split dura for revascularization of ischemic hemispheres in moyamoya disease. *J Neurosurg.* 1996; 85 (3): 380-383. doi: 10.3171/jns.1996.85.3.0380.
- 31 Kashiwagi S, Kato S, Yamashita K, et al. Revascularization with split duro-encephalo-synangiosis in the pediatric moyamoya disease--surgical result and clinical outcome. *Clin Neurol Neurosurg.* 1997; 99 (Suppl 2): S115-117. doi: 10.1016/S0303-8467(97)00069-3.
- 32 King JA, Armstrong D, Vachhrajani S, et al. Relative contributions of the middle meningeal artery and superficial temporal artery in revascularization surgery for moyamoya syndrome in children: the results of superselective angiography. *J Neurosurg Pediatr.* 2010; 5 (2): 184-189. doi: 10.3171/2009.9.PEDS0932.
- 33 Yang S, Yu JL, Wang HL, et al. Endovascular embolization of distal anterior choroidal artery aneurysms associated with moyamoya disease. A report of two cases and a literature review. *Interv Neuroradiol.* 2010; 16 (4): 433-441.
- 34 Yu JL, Wang HL, Xu K, et al. Endovascular treatment of intracranial aneurysms associated with moyamoya disease or moyamoya syndrome. *Interv Neuroradiol.* 2010; 16 (3): 240-248.
- 35 Cho HJ, Song DB, Choi HY, et al. Lenticulostriate-medullary artery anastomoses in moyamoya disease. *Neurology.* 2007; 68 (15): E21. doi: 10.1212/01.wnl.0000259084.38361.f2.
- 36 Shellshear J. The basal arteries of the forebrain and their functional significance. *J Anat.* 1920, 55 (Pt 1): 27-35.

Gerasimos Baltasvias, MD
 Department of Neuroradiology
 University Hospital Zurich
 Frauenklinikstrasse 10
 8091, Zurich, Switzerland
 Tel: +41442558657
 Fax: +41442554504
 E-mail: Gerasimos.Baltasvias@usz.ch