Published in final edited form as:

Ann Epidemiol. 2014 August ; 24(8): 612–619. doi:10.1016/j.annepidem.2014.05.014.

Associations of Allostatic Load with Sleep Apnea, Insomnia, Short Sleep Duration, and Other Sleep Disturbances: Findings from the National Health and Nutrition Examination Survey 2005-2008

Xiaoli Chen, MD, PhD, MPH^{*,1}, Susan Redline, MD, MPH², Alexandra E. Shields, PhD³, David R. Williams, PhD, MPH⁴, and Michelle A. Williams, PhD¹

¹Department of Epidemiology, Harvard School of Public Health, Boston, MA 02115, USA

²Brigham and Women's Hospital and Beth Israel Deaconess Medical Center, Department of Medicine, 221 Longwood Ave., Room 225, Boston, MA 02115, USA

³Department of Medicine, Harvard Medical School and Harvard/MGH Center for Genomics, Vulnerable Populations and Health Disparities, Mongan Institute for Health Policy, Massachusetts General Hospital, Boston, MA, USA

⁴Department of Social and Behavioral Sciences, Harvard School of Public Health, Boston, MA 02115, USA

Abstract

Purpose—To examine whether allostatic load (AL), a measure of cumulative physiologic dysregulation across biological systems, was associated with sleep apnea, insomnia, and other sleep disturbances.

Methods—Data from the National Health and Nutrition Examination Survey 2005-2008 were used. AL was measured using nine biomarkers representing cardiovascular, inflammatory, and metabolic system functioning. A total of 7,726 US adults aged 18 years and older were included in this study.

Results—The prevalence of high AL (AL score ≥ 3) was the highest among African Americans (25.2%), followed by Hispanic Americans (21.0%), Whites (18.8%), and other racial/ethnic group (16.5%). After adjustment for sociodemographic and lifestyle factors and depression status, high AL was significantly associated with sleep apnea (odds ratio (OR)=1.49, 95% confidence interval (CI): 1.14-1.95), snoring (1.65, 95% CI: 1.41-1.93), snoring/stop breathing (OR=1.62, 95% CI: 1.25-2.10), prolonged sleep latency (OR=1.33, 95% CI: 1.13-1.56), short sleep duration (<6 hours)

© 2014 Elsevier Inc. All rights reserved.

***Corresponding author:** Xiaoli Chen, MD, PhD, MPH Department of Epidemiology Harvard School of Public Health 677 Huntington Ave, Kresge 500 Boston, MA 02115 Phone: 617-432-0067 Fax: 617 566-7805 xchen@hsph.harvard.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

DECLARATION OF INTEREST: The authors report no conflicts of interest

(OR=1.29, 95% CI: 1.04-1.61), and diagnosed sleep disorder (OR=1.85, 95% CI: 1.53-2.24).

There was no clear evidence that observed associations varied by sociodemographic characteristics or depression status.

Conclusions—This study suggests significant associations of high AL with sleep apnea, sleep apnea symptoms, insomnia component, short sleep duration, and diagnosed sleep disorder among US adults.

Keywords

adult; allostatic load; insomnia; national survey; short sleep duration; sleep apnea; sleep disorder; sleep disturbance

INTRODUCTION

Quality sleep is fundamental for health and wellness. Increasing epidemiologic studies have linked sleep characteristics such as short sleep duration and poor sleep to obesity [1], hypertension [2], diabetes [3], cardiovascular disease [4, 5], mortality [6], and decreased health-related quality of life [7]. Additionally, sleep research and health disparities has been prioritized by the Institute of Medicine and Healthy People 2020 [8, 9]. Allostatic load (AL) is a multisystem construct theorized to quantify stress-induced biological risk, defined as the cumulative dysregulation of biological systems with prolonged or poorly regulated responses to internal and external stressors [10]. Several studies including one recent systematic review have shown that elevated AL is associated with cardiovascular diseases [11, 12], chronic fatigue [13], pain [14], declines in health and cognition [15], and mortality [16, 17]. Differences in AL may reflect the accumulation of physiological changes induced by differences in exposure to chronic stress, and thus might provide a mechanistic link for understanding the differential burden of life stressors and health disparities [18]. AL has been proposed as a possible mechanism contributing to health disparities observed in racial/ethnic minority groups [19]. Recent research has demonstrated that African Americans exhibit higher levels of AL than other racial/ethnic groups [20-22].

However, only limited research has been conducted on possible associations between AL and sleep disturbances (e.g., poor sleep quality). Some researchers have proposed that poor sleep might act as a neurobiological and physiologic stressor that could impair neurophysiological functions, and lead to allostatic changes throughout the body by increasing proinflammatory cytokines, oxidative stress, and evening cortisol and insulin concentrations [23, 24]. Other researchers consider sleep as a component of AL [21, 25]. Recurrent or chronic stress attributable from environmental and social factors can alter physiological functioning and behavior (e.g., sleep) which may lead to increased AL in a vicious cycle [26]. There might be bi-directional associations between AL and sleep disturbances. Prolonged sleep deprivation, poor sleep quality, and sleep apnea-associated stresses may contribute to AL and in turn high AL might contribute to sleep disturbances. To our knowledge, no research findings to date have been reported assessing any associations between AL and sleep characteristics, and certainly none of previous studies were based on national survey data.

Using nationally representative data, we examined whether AL served as a predictor of insufficient sleep or sleep disturbances. We conducted stratified analyses to evaluate whether the associations between AL and sleep disturbances varied by sex, race/ethnicity, and country of birth. We hypothesized that AL would be significantly associated with sleep apnea, insomnia, short sleep duration, and other sleep disturbances. We also hypothesized that such associations would be stronger for African and Hispanic Americans.

METHODS

Dataset and study population

Data were obtained from continuous biennial cycles of the National Health and Nutrition Examination Survey (NHANES) during 2005-2008. The NHANES is a stratified multistage probability survey conducted by the Department of Health and Human Services in the non-institutionalized population and administered by the National Center for Health Statistics. The NHANES incorporates a series of cross-sectional surveys providing health and nutrition data on a nationally representative sample. Using a computer-assisted personal interview system, trained interviewers interviewed participants in their homes to collect sociodemographic data and sleep-related information. Questionnaires that included assessment of sleep habits, sleep quality, and sleep disorders were administered by interviewers in the homes of participants aged ≥16 years. Participants were also asked to visit the NHANES Mobile Examination Center, where they completed additional questionnaires, underwent physical examinations, and provided a blood sample for laboratory measurements. Measured weight and height were used to calculate body mass index (BMI). All survey information is confidential and approved by the National Center for Health Statistics Institutional Review Board. Detailed information on the study design and data collection are described on the NHANES website <http://www.cdc.gov/nchs/nhanes.htm> and a number of NHANES papers have been published elsewhere [28, 29].

In this study, participants aged ≥18 years with complete sleep data were included. Participants were excluded from this study if they were <18 years of age (1,397 participants with sleep information aged 16-17, due to our interest in an adult population ≥18 years of age), pregnant (428 participants, due to associated changes in sleep physiology), or for whom incomplete data on AL existed (1,475 participants). A total of 3,330 individuals were included as our analytic sample in this study.

Sleep characteristics

Sleep characteristics, which were classified on the basis of participant self-reports, included sleep apnea; sleep apnea symptoms such as habitual snoring, snorting, or stop breathing; insomnia; short sleep duration; and any sleep disorder diagnosed by a physician or other health professional. Sleep apnea was defined based on an affirmative answer to the following question: "Have you ever been told by a doctor or other health professional that you have a sleep disorder: sleep apnea?"

Regarding sleep apnea symptoms of habitual snoring and snorting/stop breathing, the following two questions were asked: 1) "In the past 12 months, how often did you snore

while you were sleeping?” (snoring), and 2) “In the past 12 months, how often did you snort, gasp, or stop breathing while you were asleep?” (snorting or stop breathing). Participants who answered “frequently (5 or more nights per week)” were considered as having “habitual snoring” and “snorting/stop breathing”, respectively, whereas those with response “never”, “rarely (1-2 nights/week)”, or “occasionally (3-4 nights/week)” were considered as having no snoring or snorting/stop breathing, respectively.

Insomnia symptoms were based on the following standard questions: 1) “trouble falling asleep” (prolonged sleep latency), 2) “waking up during the night and had trouble getting back to sleep” (frequent nocturnal awakenings), 3) “waking up too early in the morning and unable to get back to sleep” (early morning awakening), and 4) “feeling unrested during the day, no matter how many hours of sleep had” (un-refreshed sleep). Responses to each insomnia symptom were collapsed as follows: occurring 2-4 times per month (considered a negligible symptom), 5-15 times per month (some level of insomnia, or mild or moderate insomnia), and >15 times/month (severe insomnia).

The study also collected data on functional impairments related to daytime sleepiness, including difficulties carrying out specific regular daily activities over the last month: 1) “concentrating on the things”, 2) “remembering things”, 3) “getting things done because too sleepy or tired to drive or take public transportation”, 4) “performing employed or volunteer work or attending school”, 5) “working on a hobby, for example, sewing, collecting, gardening”, and 6) “taking care of financial affairs and doing paperwork”. In this study, insomnia (yes/no) was defined by using the National Heart, Lung, and Blood Institute Working Group definition as one of the four insomnia symptoms plus at least one self-reported daytime functional impairment due to lack of sleep [30]. This approach has been used in other NHANES research [29].

Sleep duration was identified based on participants’ responses to the question: “How much sleep do you usually get at night on weekdays or workdays?” Previous studies have defined short habitual sleep using two cut-points: <7 and <6 hours per week-night [29, 31]. We found similar results using both cut-points. In this study, we reported findings based on the cut-point: sleep <6 hours/night (yes/no). AL was not related to long sleep duration (≥ 9 hours/night) in this study (data not shown).

Diagnosed sleep disorder was based on the question: “Have you ever been told by a doctor or other health professional that you have a sleep disorder?” Those with the response “yes” were considered as having a diagnosed sleep disorder.

Allostatic load (AL)

AL levels were measured using nine biomarkers representing cardiovascular, inflammatory, and metabolic system functioning [32]. The nine biomarkers and their corresponding cutoffs were indicated as: (1) systolic blood pressure ≥140mm Hg, (2) diastolic blood pressure ≥90mm Hg, (3) heart rate ≥90beats/minute, (4) total cholesterol level ≥240mg/dL, (5) high-density lipoprotein cholesterol<40mg/dL, (6) BMI ≥30kg/m², (7) glycosylated hemoglobin ≥6.4%, (8) C-reactive protein ≥0.3mg/dL, and (9) albumin<3.8 g/dL. Each cutoff was coded as a dichotomous variable (1, if the respondent had indicated the condition; 0, if

otherwise). The AL score was defined as the sum of the indicators for the nine components, and was then converted into a dichotomous variable, with high AL defined as the AL score ≥ 3 . The same cutoff values and measures have been used with the NHANES dataset in previous studies [32-34].

Covariates

Sociodemographic characteristics—Participants' sociodemographic characteristics included sex, age, race/ethnicity, marital status, education, poverty income ratio (PIR), and country of birth. Participants were grouped into three age categories: young (18-39 years), middle-aged (40-59 years), and old group (≥ 60 years). Based on self-reported information, participants were categorized as Whites, African Americans, Hispanic Americans, and other racial/ethnic group. Education levels were grouped into 3 categories: '<high school' (<12 years of education), 'high school', and '>high school' (≥ 12 years of education). PIR divides family income by the poverty threshold, taking into account family size. As such PIR represents adjusted income, which is more informative about available resources than income alone [14]. PIR was categorized as three levels: low income (PIR<1.0, below the poverty threshold), middle (PIR: 1-2), and high income (PIR ≥ 3). The question "In what country were you born?" was used to categorize respondents as 'US-born' or 'foreign-born'.

Lifestyle factors—Participants were asked the following questions about their participation in sport(s), exercise, or any other recreational activities: 1) "Over the past 30 days, did you do any vigorous activities for at least 10 minutes that caused heavy sweating or large increases in breathing or heart rate?"; 2) "Over the past 30 days, did you do any moderate activities for at least 10 minutes that cause only light sweating or a slight to moderate increase in breathing or heart rate?". Participants were grouped into two categories: without physical activity participation (individuals did not participate in any moderate or vigorous recreational activities) and with physical activity participation (individuals participated in moderate and/or vigorous recreational activities). Alcohol consumption and cigarette smoking were also asked and included as covariates in this study (yes vs. no).

Statistical analysis

The primary exposure variable was AL, whereas the outcome variables were sleep apnea, sleep apnea symptoms of snoring and snorting/ stop breathing, insomnia, short sleep duration, any sleep disorder diagnosed by a physician or other health professional. AL was examined both as a continuous and a dichotomous variable (high AL, defined as the AL score ≥ 3 ; low AL, defined as the AL score < 3). We used chi-square tests to examine any differences in the distribution of elevated AL across sociodemographic, lifestyle, and sleep characteristics. Multivariable logistic regression models were fit to examine the associations between AL and sleep parameters with adjustment for sociodemographic and lifestyle factors. Odds ratios (ORs) and 95% confidence intervals (95% CIs) were calculated. We further conducted stratified analyses to test whether the AL-sleep associations varied by sex, race/ethnicity, and country of birth. Interaction terms were included and tested in separate models. Our sensitivity analysis showed that the sample included in the study was not significantly different from the entire NHANES population.

All statistical analyses were conducted in SAS (version 9.3; SAS Institute, Inc., Cary, North Carolina). Survey-related commands including sampling design variables and sampling weights were applied to account for the complex multistage sampling design. Statistical significance was set at two-tailed $P < 0.05$.

RESULTS

Prevalence of high AL and sleep characteristics

Approximately 21.1% of U.S. adults had high AL. There was no significant sex difference in the prevalence of high AL (Table 1). The prevalence of high AL was higher among participants 60 years of age or older than in the youngest age group (18-39 years), was higher among widowed, separated, or divorced individuals than among those married or living with a partner, and was higher for US-born than for foreign-born individuals. Participants with less than high school of education levels were more likely to have high AL than those with high school or more than high school of education levels. We did not find significant differences in the prevalence of high AL across PIR categories. The prevalence of high AL was the highest among African Americans (26.3%), followed by Hispanic Americans (20.3%), Whites (17.7%), and other racial/ethnic group (13.8%). Those with high education levels and recreational activity participation had a lower prevalence of high AL than their counterparts.

As shown in Table 2, the prevalence of high AL was higher among adults with diagnosed sleep apnea, snoring, snorting/stop breathing, insomnia, short sleep duration (<6 hours), diagnosed sleep disorder than those without sleep disturbances.

Figure 1a shows that the mean AL score was high for African Americans, followed by Hispanic Americans, and Whites. Individuals in other racial/ethnic group had the lowest AL score. The mean AL score was the highest among African American women, while the mean AL score was the lowest in other racial/ethnic women (Figure 1b). The elevated mean AL score in African Americans was solely due to African American women.

Table 3 shows that African Americans had a higher prevalence of insomnia and short sleep duration compared with other racial/ethnic groups (both $P < 0.05$). There were no significant differences in the distributions of sleep apnea, sleep apnea symptoms of snoring and snorting, and diagnosed sleep disorder across racial/ethnic groups.

Associations between high AL and sleep characteristics: mean differences

As shown in Figure 2, adults with sleep apnea, sleep apnea symptoms of snoring and snorting/stop breathing, insomnia, short sleep duration, and diagnosed sleep disorder had higher mean AL scores than those without these sleep disturbances (all $P < 0.001$).

Associations between high AL and sleep characteristics: logistic regression analysis

Table 4 shows the results of logistic regression models for the associations between AL and sleep characteristics. High AL was associated with sleep apnea (OR=1.92, 95% CI: 1.40-2.63), insomnia (OR=1.70, 95% CI: 1.16-2.47), and short sleep duration (OR=1.35, 95% CI: 1.00-1.82), controlling for age, sex, race/ethnicity, marital status, education, PIR,

country of birth, physical activity participation, alcohol consumption, and cigarette smoking. Individuals with a high AL were about two times as likely to have snoring (OR=2.20, 95% CI: 1.79-2.69), snorting/stop breathing (OR=2.16, 95% CI: 1.46-3.21), and diagnosed sleep disorder (OR=2.26, 95% CI: 1.66-3.08). High AL was related to increased risk for prolonged sleep latency (one of the insomnia components), the adjusted OR was 1.42 (95% CI: 1.08-1.88). High AL was not associated with other components of insomnia, including frequent nocturnal awakenings, early morning awakening, or un-refreshed sleep.

Stratified analysis

As shown in Table 5, the associations of high AL with sleep apnea, snoring, and diagnosed sleep disorder were statistically significant and similar for both men and women. Although the association between high AL and insomnia was significant for women (OR=1.90; 95% CI: 1.10-3.44) but not for men (OR=1.43; 95% CI: 0.83-2.46), the interaction term was not significant. The associations of high AL with sleep apnea, snorting, insomnia, short sleep duration, and diagnosed sleep disorder were statistically significant and stronger for Whites but not for African Americans, Hispanic Americans, or other racial/ethnic group. However, interaction terms were not statistically significant. There were significant associations between high AL and habitual snoring across racial/ethnic groups. The associations of high AL with sleep disturbances did not vary by country of birth (data not shown in tables).

DISCUSSION

Minority racial/ethnic and low socioeconomic groups tend to have a higher prevalence of sleep disturbances and are more likely to be exposed to stress or high AL, a measure of physiological instability across biological systems from cumulative or repeated adaptation to stressors [17]. Using nationally representative data, we found a high prevalence of AL (21.1%) and racial/ethnic differences in AL among U.S. adults. High AL was associated with sleep apnea, sleep apnea symptoms (snoring, snorting/stop breathing), insomnia, short sleep duration, and diagnosed sleep disorder. These associations were independent of sociodemographic and lifestyle factors. Short sleep duration was most prevalent in African Americans. To our knowledge, this is the first study to examine the associations between AL and sleep characteristics based on national survey data.

Increasing research has indicated the relationships between AL and sociodemographic, lifestyle, and genetic characteristics [17, 20-22]. Several studies have investigated the racial/ethnic differences in AL levels [20-22]. A study of 129 US adults found that African Americans had higher AL scores than Whites, and more African Americans had high AL (67.9%) than did Whites (48.9%) [21]. The NHANES 1999-2004 data showed that African American women had the highest AL scores relative to other racial/ethnic groups [22]. Our study, based on the NHANES 2005-2008, found similar results, which expand upon previous research by evaluating the AL-sleep association using the national survey data. The NHANES III (1988-1994) data demonstrated that lower individual socioeconomic status was associated with higher AL, and this association existed for all major racial/ethnic groups [35]. In this study, we found that being African Americans or Hispanic Americans, having less education, being older, widowed, separated, or divorced, and alcohol consumption were

associated with high AL, while recreational physical activity participation was related to low AL. These results were consistent with previous findings [17, 36]. Our findings add to the growing literature regarding AL and its risk factors.

Several studies have suggested that country of birth may explain some health disparities in the United States [29, 37-39]. Seicean et al reported that Mexican-born US immigrants had more favorable sleep patterns than the general US population [29]. In a population-based sample of adults living in Texas City, foreign-born Mexicans were the least likely group to score in the higher AL categories [20]. US-born Mexican Americans had higher AL scores than foreign-born Mexicans, and acculturation measures did not account for the difference. We found that the prevalence of high AL was higher for US-born than for foreign-born individuals, supporting the healthy immigrant hypothesis [20].

Although there has been an increasing number of studies on the association between AL and health outcomes [14, 40], ours is the first study investigating the association between AL and sleep disturbances based on national survey data. The NHANES 1999-2004 data showed that greater AL was associated with elevated prevalence of pain in US adults [14]. The NHANES 1999-2004 data also indicated that US adults with high AL were more likely to have periodontitis than their counterparts with low AL [40]. In our study, we found that US adults with greater biological “wear and tear” as measured by AL were more likely to have insufficient sleep as measured by short sleep duration as well as diagnosed sleep apnea or sleep apnea symptoms, or other sleep disorders. Furthermore, these relationships were independent of sociodemographic and lifestyle factors. Our findings indicate that greater biological “wear and tear” is independently associated with insufficient sleep and sleep disturbances, thereby suggesting their importance in enhancing physiological functioning among US adults. Contrary to our hypothesis, we did not find that the association between AL and sleep was stronger among African Americans or Hispanic Americans. This might be due to self-reported sleep information used in our study. African Americans and Hispanic Americans might have underestimated their sleep problems. One recent study, for example, revealed that African and Hispanic respondents were relatively optimistic in their ratings than Whites [41]. Studies that deploy objective measures of sleep behavior and sleep quality may overcome limitations associated with reliance on self-reported sleep measures.

Our study has several strengths. First, it is based on nationally representative and recent NHANES data collected between 2005 and 2008, with balanced gender representation and geographic diversity, and likely to be free of the referral biases that may occur from studies of sleep clinic-based samples [29]. This allows greater generalizability than previous studies. Second, it has a large sample size, which uses highly structured protocols and allows us to conduct stratified analyses to examine whether the associations between AL and sleep parameters differ by sociodemographics characteristics. Third, we conducted a robust set of statistical analyses by applying multivariable logistic regression and stratified analyses to examine the associations between AL and sleep disturbances.

This study has limitations. A major limitation is its cross-sectional study design, thus we are unable to detect whether AL leads to sleep disturbances or vice versa, or neither may be true. Our findings could be explained by an underlying mechanism that affects both AL and

sleep disturbances. Future research should seek an answer to this foundational question. Another limitation, as noted above, is that sleep characteristics were assessed by self-reported questionnaire, which is subject to measurement error and/or report bias. In addition, there is considerable heterogeneity in the biomarkers selected to assess AL [17, 20, 21, 36, 40] and in the cut-points for defining high AL [14, 16, 20-22, 42, 43]. A recent review of 58 published papers on AL indicated that interpretations and comparisons across studies was challenging due to different AL biomarker measurement [17]. In our study, we applied nine biomarkers representing cardiovascular, inflammatory, and metabolic system functioning to evaluate the AL levels, as in prior research with the NHANES dataset [32-34].

In summary, results from this cross-sectional study with a nationally representative sample suggest that high AL is associated with sleep apnea, sleep apnea symptoms, insomnia, short sleep duration, and diagnosed sleep disorder. These data support a potentially pervasive influence of sleep-related stresses on physiological functioning associated with cardio-metabolic disease and mortality. Individuals with sleep problems have a high burden of physical and psychological health conditions. Future research should focus on refining the definition of AL, or the development of a consensus opinion. Further prospective longitudinal studies, with objective measures of sleep duration and quality, are needed to examine the possible bi-directionality and causal associations between AL and sleep characteristics. Priorities for future research also include examining a broad range of antecedent allostatic challenges, and collecting reliable measures of multisystem dysregulation explicitly designed to assess AL at multiple time points in population-based prospective studies [17].

Acknowledgments

This study was supported by National Institutes of Health (NIH) and National Institute on Minority Health and Health Disparities (NIMHD) grants (T37-MD001449) and NIH/NCRR/NCATS (8UL1TR000170). We thank Dr. Bizu Gelaye for his technical assistance.

Abbreviations

AL	allostatic load
BMI	body mass index
CI	confidence interval
NHANES	National Health and Nutrition Examination Survey
OR	odds ratio
PIR	poverty income ratio

REFERENCES

1. Chen X, Beydoun MA, Wang Y. Is sleep duration associated with childhood obesity? A systematic review and meta-analysis. *Obesity* (Silver Spring). 2008; 16:265–74. [PubMed: 18239632]
2. Gangwisch JE, Heymsfield SB, Boden-Albala B, et al. Short sleep duration as a risk factor for hypertension: analyses of the first National Health and Nutrition Examination Survey. *Hypertension*. 2006; 47:833–9. [PubMed: 16585410]

3. Yaggi HK, Araujo AB, McKinlay JB. Sleep duration as a risk factor for the development of type 2 diabetes. *Diabetes Care*. 2006; 29:657–61. [PubMed: 16505522]
4. Currie A, Cappuccio FP. Sleep in children and adolescents: a worrying scenario: can we understand the sleep deprivation-obesity epidemic? *Nutr Metab Cardiovasc Dis*. 2007; 17:230–2. [PubMed: 17321728]
5. Redline S, Foody J. Sleep disturbances: time to join the top 10 potentially modifiable cardiovascular risk factors? *Circulation*. 2011; 124:2049–51. [PubMed: 22064955]
6. Kripke DF, Garfinkel L, Wingard DL, et al. Mortality associated with sleep duration and insomnia. *Arch Gen Psychiatry*. 2002; 59:131–6. [PubMed: 11825133]
7. Silva GE, An MW, Goodwin JL, et al. Longitudinal evaluation of sleep-disordered breathing and sleep symptoms with change in quality of life: the Sleep Heart Health Study (SHHS). *Sleep*. 2009; 32:1049–57. [PubMed: 19725256]
8. Institute of Medicine, Committee on Sleep Medicine and Research. Sleep disorders and sleep deprivation: An unmet public health problem. National Academy of Sciences Press; Washington, DC: 2006.
9. US Department of Health and Human Services. Office of Disease Prevention and Health Promotion. [August 8, 2013] Healthy People 2020. Nov. 2010 Available at: <http://www.healthypeople.gov/2020/about/disparitiesAbout.aspx>.
10. McEwen BS, Seeman T. Protective and damaging effects of mediators of stress. Elaborating and testing the concepts of allostasis and allostatic load. *Ann N Y Acad Sci*. 1999; 896:30–47. [PubMed: 10681886]
11. Mattei J, Demissie S, Falcon LM, et al. Allostatic load is associated with chronic conditions in the Boston Puerto Rican Health Study. *Soc Sci Med*. 2010; 70:1988–96. [PubMed: 20381934]
12. Sabbah W, Watt RG, Sheiham A, et al. Effects of allostatic load on the social gradient in ischaemic heart disease and periodontal disease: evidence from the Third National Health and Nutrition Examination Survey. *J Epidemiol Community Health*. 2008; 62:415–20. [PubMed: 18413454]
13. Goertzel BN, Pennachin C, de Souza Coelho L, et al. Allostatic load is associated with symptoms in chronic fatigue syndrome patients. *Pharmacogenomics*. 2006; 7:485–94. [PubMed: 16610958]
14. Slade GD, Sanders AE, By K. Role of allostatic load in sociodemographic patterns of pain prevalence in the U.S. population. *J Pain*. 2012; 13:666–75. [PubMed: 22677453]
15. Juster RP, McEwen BS, Lupien SJ. Allostatic load biomarkers of chronic stress and impact on health and cognition. *Neurosci Biobehav Rev*. 2010; 35:2–16. [PubMed: 19822172]
16. Borrell LN, Dallo FJ, Nguyen N. Racial/ethnic disparities in all-cause mortality in U.S. adults: the effect of allostatic load. *Public Health Rep*. 2010; 125:810–6. [PubMed: 21121226]
17. Beckie TM. A systematic review of allostatic load, health, and health disparities. *Biol Res Nurs*. 2012; 14:311–46. [PubMed: 23007870]
18. Szanton SL, Gill JM, Allen JK. Allostatic load: a mechanism of socioeconomic health disparities? *Biol Res Nurs*. 2005; 7:7–15. [PubMed: 15919999]
19. Carlson ED, Chamberlain RM. Allostatic load and health disparities: a theoretical orientation. *Res Nurs Health*. 2005; 28:306–15. [PubMed: 16028266]
20. Peek MK, Cutchin MP, Salinas JJ, et al. Allostatic load among non-Hispanic Whites, non-Hispanic Blacks, and people of Mexican origin: effects of ethnicity, nativity, and acculturation. *Am J Public Health*. 2010; 100:940–6. [PubMed: 19834005]
21. Deuster PA, Kim-Dorner SJ, Remaley AT, et al. Allostatic load and health status of African Americans and whites. *Am J Health Behav*. 2011; 35:641–53. [PubMed: 22251756]
22. Chyu L, Upchurch DM. Racial and ethnic patterns of allostatic load among adult women in the United States: findings from the National Health and Nutrition Examination Survey 1999-2004. *J Womens Health (Larchmt)*. 2011; 20:575–83. [PubMed: 21428732]
23. Tremblay A, Chaput JP. Obesity: the allostatic load of weight loss dieting. *Physiol Behav*. 2012; 106:16–21. [PubMed: 21627975]
24. McEwen BS. Sleep deprivation as a neurobiologic and physiologic stressor: Allostasis and allostatic load. *Metabolism*. 2006; 55:S20–3. [PubMed: 16979422]

25. Hamilton NA, Affleck G, Tennen H, et al. Fibromyalgia: the role of sleep in affect and in negative event reactivity and recovery. *Health Psychol.* 2008; 27:490–7. [PubMed: 18643007]
26. Borsook D, Maleki N, Becerra L, et al. Understanding migraine through the lens of maladaptive stress responses: a model disease of allostatic load. *Neuron.* 2012; 73:219–34. [PubMed: 22284178]
27. Patel SR, White DP, Malhotra A, et al. Continuous positive airway pressure therapy for treating sleepiness in a diverse population with obstructive sleep apnea: results of a meta-analysis. *Arch Intern Med.* 2003; 163:565–71. [PubMed: 12622603]
28. Ogden CL, Carroll MD, Kit BK, et al. Prevalence of obesity and trends in body mass index among US children and adolescents, 1999–2010. *JAMA.* 2012; 307:483–90. [PubMed: 22253364]
29. Seicean S, Neuhauser D, Strohl K, et al. An exploration of differences in sleep characteristics between Mexico-born US immigrants and other Americans to address the Hispanic Paradox. *Sleep.* 2011; 34:1021–31. [PubMed: 21804664]
30. Wolfe WS, Campbell CC, Frongillo EA Jr. et al. Overweight schoolchildren in New York State: prevalence and characteristics. *Am J Public Health.* 1994; 84:807–13. [PubMed: 8179053]
31. Hagen EW, Mirer AG, Palta M, et al. The sleep-time cost of parenting: sleep duration and sleepiness among employed parents in the wisconsin sleep cohort study. *Am J Epidemiol.* 2013; 177:394–401. [PubMed: 23378502]
32. Parente V, Hale L, Palermo T. Association between breast cancer and allostatic load by race: National Health and Nutrition Examination Survey 1999–2008. *Psycho-oncology.* 2013; 22:621–8. [PubMed: 22290849]
33. Crimmins EM, Kim JK, Seeman TE. Poverty and biological risk: the earlier “aging” of the poor. *J Gerontol A Biol Sci Med Sci.* 2009; 64:286–92. [PubMed: 19196637]
34. Crimmins EM, Johnston M, Hayward M, et al. Age differences in allostatic load: an index of physiological dysregulation. *Exp Gerontol.* 2003; 38:731–4. [PubMed: 12855278]
35. Seeman T, Merkin SS, Crimmins E, et al. Education, income and ethnic differences in cumulative biological risk profiles in a national sample of US adults: NHANES III (1988–1994). *Soc Sci Med.* 2008; 66:72–87. [PubMed: 17920177]
36. Bird CE, Seeman T, Escarce JJ, et al. Neighbourhood socioeconomic status and biological ‘wear and tear’ in a nationally representative sample of US adults. *J Epidemiol Community Health.* 2010; 64:860–5. [PubMed: 19759056]
37. Montez JK, Eschbach K. Country of birth and language are uniquely associated with intakes of fat, fiber, and fruits and vegetables among Mexican-American women in the United States. *J Am Diet Assoc.* 2008; 108:473–80. [PubMed: 18313430]
38. Eldeirawi KM, Persky VW. Associations of acculturation and country of birth with asthma and wheezing in Mexican American youths. *J Asthma.* 2006; 43:279–86. [PubMed: 16809241]
39. Sundquist J, Winkleby M. Country of birth, acculturation status and abdominal obesity in a national sample of Mexican-American women and men. *Int J Epidemiol.* 2000; 29:470–7. [PubMed: 10869319]
40. Borrell LN, Crawford ND. Social disparities in periodontitis among US adults: the effect of allostatic load. *J Epidemiol Community Health.* 2011; 65:144–9. [PubMed: 19996354]
41. Dowd JB, Todd M. Does self-reported health bias the measurement of health inequalities in U.S. adults? Evidence using anchoring vignettes from the Health and Retirement Study. *J Gerontol B Psychol Sci Soc Sci.* 2011; 66:478–89. [PubMed: 21666144]
42. Merkin SS, Basurto-Davila R, Karlamangla A, et al. Neighborhoods and cumulative biological risk profiles by race/ethnicity in a national sample of U.S. adults: NHANES III. *Ann Epidemiol.* 2009; 19:194–201. [PubMed: 19217002]
43. Mattei J, Noel SE, Tucker KL. A meat, processed meat, and French fries dietary pattern is associated with high allostatic load in Puerto Rican older adults. *J Am Diet Assoc.* 2011; 111:1498–506. [PubMed: 21963016]

a) Both men and women

b) For men and women separately

Figure 1.

Race/ethnicity distributions and mean allostatic load scores among 3,330 US adults in the National Health and Nutrition Examination Survey 2005-2008: means and 95% confidence limits of means

Figure 2.

Sleep disturbances and mean allostatic load scores among 3,330 US adults in the National Health and Nutrition Examination Survey 2005-2008: Means and 95% confidence limits of means

There were significant differences in the allostatic load scores between those with and without sleep problems (all P values <0.001). Tests were conducted using the 'surveyreg' commands.

Table 1

Sociodemographic and lifestyle characteristics of 3,330 US adults in the National Health and Nutrition Examination Survey 2005-2008, according to allostatic load status

Characteristic	N	Low AL ^a (n =2,627) % (SE)	High AL ^a (n =703) % (SE)	P value ^b
Sex				
Men	1,749	82.4 (1.0)	17.6 (1.0)	0.260
Women	1,581	80.6 (1.4)	19.4 (1.4)	
Age (years)				
18-39	1,273	87.1 (1.3)	12.9 (1.3)	<0.001
40-59	1,115	78.1 (1.6)	21.9 (1.6)	
60	942	77.3 (1.8)	22.7 (1.8)	
Race/ethnicity				
White	1,593	82.3 (1.1)	17.7 (1.1)	<0.001
African American	712	73.6 (1.9)	26.3 (1.9)	
Hispanic American	635	79.7 (2.0)	20.3 (2.0)	
Other	390	86.1 (2.2)	13.8 (2.2)	
Marital status				
Married/living with a partner	2,121	82.3 (1.0)	17.7 (1.0)	<0.001
Widowed/separated/divorced	557	71.6 (2.4)	28.4 (2.4)	
Single	652	86.7 (1.7)	13.3 (1.7)	
Education level				
< High school	892	74.4 (1.0)	25.6 (1.6)	<0.001
High school	861	78.4 (1.6)	21.6 (1.6)	
>High school	1,577	85.5 (1.1)	14.9 (1.1)	
Poverty income ratio				
< 1	757	79.8 (2.2)	20.2 (2.2)	0.086
1-2	1,286	79.3 (1.4)	20.7 (1.4)	
3	1,287	83.4 (1.4)	16.6 (1.4)	
Country of birth				
US-born	2,618	80.7 (1.0)	19.3 (1.0)	<0.001
Foreign-born	712	86.9 (1.1)	13.1 (1.1)	
Physical activity participation ^c				
No	1,462	74.1 (1.4)	25.9 (1.4)	<0.001
Yes	1,868	86.0 (1.1)	14.0 (1.1)	
Alcohol consumption				
No	2,002	79.5 (1.3)	20.5 (1.3)	0.002
Yes	1,328	84.0 (1.0)	16.0 (1.0)	
Cigarette smoking				
No	2,637	81.8 (1.0)	18.2 (1.0)	0.557
Yes	693	80.5 (2.1)	19.5 (2.1)	

Abbreviations: AL, Allostatic load; SE, standard error.

^a Allostatic load was measured using the following nine components and their corresponding cutoffs: (1) systolic blood pressure ≥ 140 mm Hg, (2) diastolic blood pressure ≥ 90 mm Hg, (3) heart rate ≥ 90 beats/minute, (4) total cholesterol level ≥ 240 mg/dL, (5) high-density lipoprotein cholesterol < 40 mg/dL, (6) body mass index ≥ 30 kg/m², (7) glycosylated hemoglobin $\geq 6.4\%$, (8) C-reactive protein ≥ 0.3 mg/dL, and (9) albumin < 3.8 g/dL. Each cutoff was coded as a dichotomous variable, and the AL score was defined as the sum of the indicators for the nine components. High AL was defined as the AL score ≥ 3 , while low AL was defined as the AL score < 3 .

^b Chi-square test was conducted.

^c Insomnia (yes/no) was defined by using the National Heart, Lung, and Blood Institute Working Group definition as one of the four insomnia symptoms plus at least one self-reported daytime functional impairment due to lack of sleep.

Table 2

Sleep characteristics of 3,330 US adults in the National Health and Nutrition Examination Survey 2005-2008, according to allostatic load status

Characteristic	N	Low AL ^a (n =2,627) % (SE)	High AL ^a (n =703) % (SE)	P value ^b
Sleep apnea				
No	2,895	83.4 (0.9)	16.6 (0.9)	<0.001
Yes	435	69.3 (2.6)	30.6 (2.6)	
Sleep apnea symptoms				
Snoring				
No	2,232	86.0 (1.0)	14.0 (1.0)	<0.001
Yes	1,098	72.3 (1.2)	27.7 (1.4)	
Snorting/stop breathing				
No	3,120	82.8 (0.9)	17.2 (0.9)	<0.001
Yes	210	64.7 (3.9)	35.2 (3.9)	
Insomnia ^c				
No	3,109	82.1 (0.9)	17.9 (0.9)	0.002
Yes	221	72.8 (3.6)	27.2 (3.6)	
Components of insomnia				
Prolonged sleep latency				
No	2,763	82.8 (0.9)	17.2 (0.9)	<0.001
Yes	567	75.5 (1.9)	24.5 (1.9)	
Frequent nocturnal awakenings				
No	2,667	81.9 (1.0)	18.1 (1.0)	0.267
Yes	663	80.0 (1.6)	20.0 (1.6)	
Early morning awakening				
No	2,753	82.1 (0.9)	17.9 (0.9)	0.041
Yes	575	78.9 (1.6)	21.1 (1.6)	
Un-refreshed sleep				
No	2,449	82.0 (0.9)	17.9 (0.9)	0.283
Yes	881	80.3 (1.6)	19.7 (1.7)	
Short sleep duration				
6 hours	2,793	82.6 (1.0)	17.4 (1.0)	0.001
< 6 hours	537	75.3 (2.0)	24.7 (2.0)	
Diagnosed sleep disorder				
No	3,076	82.8 (0.9)	17.2 (0.9)	<0.001
Yes	254	65.5 (2.9)	34.5 (2.9)	

Abbreviations: AL, Allostatic load; SE, standard error.

^a Allostatic load was measured using the following nine components and their corresponding cutoffs: (1) systolic blood pressure ≥140mm Hg, (2) diastolic blood pressure ≥90mm Hg, (3) heart rate ≥90beats/minute, (4) total cholesterol level ≥240mg/dL, (5) high-density lipoprotein cholesterol<40mg/dL (6) body mass index ≥30kg/m², (7) glycosylated hemoglobin ≥6.4%, (8) C-reactive protein ≥0.3mg/dL, and (9) albumin<3.8g/dL. Each cutoff was coded as a dichotomous variable, and the AL score was defined as the sum of the indicators for the nine components. High AL was defined as the AL score ≥3, while low AL was defined as the AL score<3.

^b Chi-square test was conducted.

^c Insomnia (yes/no) was defined by using the National Heart, Lung, and Blood Institute Working Group definition as one of the four insomnia symptoms plus at least one self-reported daytime functional impairment due to lack of sleep.

Table 3

Distributions of sleep disturbances in the National Health and Nutrition Examination Survey 2005-2008, according to race/ethnicity

	White % (SE)	African American % (SE)	Hispanic American % (SE)	Other % (SE)	P value ^a
Total sample size	1,593	712	635	390	
Sleep apnea					
No	86.2 (1.2)	85.8 (2.0)	90.2 (1.5)	87.1 (1.9)	0.364
Yes	13.8 (1.2)	14.2 (2.0)	9.8 (1.5)	12.9 (1.9)	
Sleep apnea symptoms					
Snoring					
No	67.2 (1.6)	68.7 (1.8)	66.3 (1.7)	66.9 (3.4)	0.45
Yes	32.8 (1.6)	31.3 (1.8)	33.7 (1.7)	33.1 (3.4)	
Snoring/stop breathing					
No	92.9 (0.9)	93.8 (0.9)	94.7 (1.3)	93.7 (1.4)	0.605
Yes	7.1 (0.9)	6.2 (0.9)	5.3 (1.3)	6.3 (1.4)	
Insomnia ^b					
No	94.4 (0.6)	90.5 (1.4)	95.1 (1.0)	91.9 (1.6)	0.013
Yes	5.6 (0.6)	9.5 (1.4)	4.9 (1.0)	8.1 (1.6)	
Short sleep duration					
6 hours	87.4 (0.8)	72.1 (1.4)	90.0 (1.5)	82.6 (2.4)	<0.001
< 6 hours	12.6 (0.8)	27.9 (1.4)	10.0 (1.5)	17.4 (2.4)	
Diagnosed sleep disorder					
No	92.5 (0.7)	91.0 (1.2)	95.9 (1.0)	92.2 (1.7)	0.109
Yes	7.5 (0.7)	9.0 (1.2)	4.1 (1.0)	7.8 (1.7)	

Abbreviations: SE, standard error.

^a Rao-Scott Chi-square test was conducted.

^b Insomnia (yes/no) was defined by using the National Heart, Lung, and Blood Institute Working Group definition as one of the four insomnia symptoms plus at least one self-reported daytime functional impairment due to lack of sleep.

Table 4
Multivariate logistic regression models: associations of allostatic load with sleep disturbances among 3,330 US adults in the National Health and Nutrition Examination Survey 2005-2008^a

Outcome variable	Unadjusted	P value	Adjusted ^b	P value
	OR (95% CI)		(95% CI)	
Sleep apnea	2.23 (1.64-3.01)	<0.001	1.92 (1.40-2.63)	<0.001
Sleep apnea symptoms				
Snoring	2.37 (1.94-2.89)	<0.001	2.20 (1.79-2.69)	<0.001
Snorting/stop breathing	2.62 (1.78-3.84)	<0.001	2.16 (1.46-3.21)	<0.001
Insomnia ^d	1.71 (1.21-2.41)	0.002	1.70 (1.16-2.47)	0.006
Insomnia component				
Prolonged sleep latency	1.57 (1.26-1.94)	<0.001	1.42 (1.08-1.88)	0.013
Frequent nocturnal awakenings	1.13 (0.91-1.41)	0.269	0.99 (0.77-1.27)	0.944
Early morning awakening	1.23 (1.01-1.49)	0.041	1.03 (0.83-1.28)	0.793
Un-refreshed sleep	1.12 (0.91-1.38)	0.281	1.13 (0.88-1.45)	0.347
Short sleep duration (< 6 hours)	1.56 (1.19-2.06)	0.002	1.35 (1.00-1.82)	0.048
Diagnosed sleep disorder	2.54 (1.95-3.32)	<0.001	2.26 (1.66-3.08)	<0.001

Abbreviations: OR, odds ratio; 95% CI, 95% confidence interval.

^c Insomnia (yes/no) was defined by using the National Heart, Lung, and Blood Institute Working Group definition as one of the four insomnia symptoms plus at least one self-reported daytime functional impairment due to lack of sleep.

^a Allostatic load was measured using the following nine components and their corresponding cutoffs: (1) systolic blood pressure 140mm Hg, (2) diastolic blood pressure 90mm Hg, (3) heart rate 90beats/minute, (4) total cholesterol level 240mg/dL, (5) high-density lipoprotein cholesterol<40mg/dL (6) body mass index 30kg/m², (7) glycosylated hemoglobin 6.4%, (8) C-reactive protein 0.3mg/dL, and (9) albumin<3.8g/dL. Each cutoff was coded as a dichotomous variable, and the AL score was defined as the sum of the indicators for the nine components. High AL was defined as a total of AL score 3, while low AL was defined as the AL score<3.

^b The following variables were adjusted for in the models: age, sex, marital status, education, household income, race/ethnicity, country of birth, alcohol consumption, cigarette smoking, and regular physical activity participation.

Table 5

Stratified analyses for the associations between high allostatic load and sleep disturbances among 3,330 US adults in the National Health and Nutrition Examination Survey 2005-2008, by sex and race/ethnicity^a

	Sleep apnea		Snoring		Snoring/stop breathing		Insomnia ^b		Short sleep duration <6 hours		Sleep disorder	
	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)
Sex												
Men	1.96	(1.35-2.84)***	2.00	(1.44-2.77)***	2.23	(1.44-3.45)***	1.43	(0.83-2.46)	1.30	(1.00-1.75)*	2.30	(1.54-3.53)***
Women	1.78	(1.09-2.89)*	2.24	(1.55-3.23)***	1.96	(0.98-3.90)	1.90	(1.10-3.44)*	1.31	(0.78-2.19)	2.30	(1.53-3.35)***
<i>P for interaction</i>	0.931		0.363		0.890		0.455		0.832		0.973	
Race/ethnicity												
White	2.01	(1.39-2.91)***	1.97	(1.50-2.60)***	2.44	(1.50-3.98)***	1.90	(1.15-2.99)*	1.40	(1.01-2.05)*	2.60	(1.68-4.03)***
AA	1.80	(0.90-3.60)	2.44	(1.69-3.54)***	1.28	(0.52-3.15)	1.36	(0.71-2.62)	1.03	(0.69-1.53)	1.15	(0.57-2.33)
HA	1.43	(0.59-3.44)	3.29	(2.09-5.19)***	1.75	(0.57-5.35)	1.09	(0.49-2.44)	0.85	(0.47-1.57)	1.91	(0.59-6.16)
Other	1.43	(0.45-4.54)	2.85	(1.25-6.50)*	1.51	(0.30-7.20)	1.81	(0.56-5.83)	1.90	(0.78-4.67)	2.73	(0.73-10.2)
<i>P for interaction</i>	0.498		0.118		0.413		0.891		0.181		0.331	

Abbreviations: OR, odds ratio; 95% CI, 95% confidence interval; AA, African American; HA, Hispanic American.

**P<0.01

^a Allostatic load was measured using the following nine components and their corresponding cutoffs: (1) systolic blood pressure ≥140mm Hg, (2) diastolic blood pressure ≥90mm Hg, (3) heart rate ≥90beats/minute, (4) total cholesterol level ≥240mg/dL, (5) high-density lipoprotein cholesterol<40mg/dL (6) body mass index ≥30kg/m², (7) glycosylated hemoglobin ≥6.4%, (8) C-reactive protein ≥0.3mg/dL, and (9) albumin<3.8g/dL. Each cutoff was coded as a dichotomous variable, and the AL score was defined as the sum of the indicators for the nine components. High AL was defined as score ≥3, while low AL was defined as AL score<3. The following variables were adjusted for in the models: age, sex, marital status, education, household income, country of birth, alcohol consumption, cigarette smoking, and regular physical activity participation.

^b Insomnia (yes/no) was defined by using the National Heart, Lung, and Blood Institute Working Group definition as one of the four insomnia symptoms plus at least one self-reported daytime functional impairment due to lack of sleep.

* P<0.05

*** P<0.001