Published in final edited form as:

J Mol Biol. 2002 August 9; 321(2): 379–380. doi:10.1016/S0022-2836(02)00614-9.

Note to the Paper by Guasch *et al.* (2002) Detailed Architecture of a DNA Translocating Machine: The High-resolution Structure of the Bacteriophage ϕ 29 Connector Particle

Alicia Guasch¹, Joan Pous¹, Borja Ibarra², F. Xavier Gomis-Rüth¹, José Maria Valpuesta², Natalia Sousa², José L. Carrascosa², Miquel Coll¹, Alan A. Simpson³, Yizhi Tao³, Petr G. Leiman³, Mohammed O. Badasso⁴, Yongning He³, Paul J. Jardine⁴, Norman H. Olson³, Marc C. Morais³, Shelley Grimes⁴, Dwight L. Anderson⁴, Timothy S. Baker³, and Michael G. Rossmann³

Michael G. Rossmann: mgr@indiana.bio.purdue.edu

¹Institut de Biologia Molecular de Barcelona, CSIC, Jordi Girona 18-26, E-08034 Barcelona, Spain

²Centro Nacional de Biotecnología, CSIC, Universidad Autónoma de Madrid, Cantoblanco, E-28049 Madrid, Spain

³Department of Biological Sciences, Purdue University, West Lafayette, IN 47907-1392, USA

⁴Departments of Microbiology and Oral Science, 18-246 Moos Tower, University of Minnesota, Minneapolis, MN 55455, USA

The recent paper by Guasch *et al.*¹ describing the structure of the ϕ 29 bacteriophage head–tail connector at 2.1 Å resolution (PDB code 1h5w), makes reference to the original 3.2 Å resolution structure (PDB code 1fou) published by Simpson *et al.*² and used by Guasch *et al.* to solve the 1h5w structure by molecular replacement. The Guasch paper states that there are marked differences between the 1h5w and 1fou structures in the chain tracing and the side-chain positioning in the wide domain. As these and a number of other statements in the Guasch paper may have led to the impression that these differences are larger than they really are, we give here Tables summarizing the similarities and differences (Tables 1 and 2).

Another paper by Simpson *et al.*,³ which was accidentally omitted from the text of the Guasch *et al.* paper, reports two additional structures of the ϕ 29 connector: at 3.2 Å resolution and at 2.9 Å resolution (PDB codes 1jnb and 1ijg, respectively). The 2.9 Å resolution structure shows no significant differences from the structure subsequently reported by Guasch *et al.* (Figure 1; Tables 1 and 2).

References

1. Guasch A, Pous J, Ibarra B, Gomis-Rüth FX, Valpuesta JM, Sousa N, et al. Detailed architecture of a DNA translocating machine: the high-resolution structure of the bacteriophage ϕ 29 connector particle. *J. Mol. Biol.* 2002; 315:663–676. [PubMed: 11812138]
2. Simpson AA, Tao Y, Leiman PG, Badasso MO, He Y, Jardine PJ, et al. Structure of the bacteriophage ϕ 29 DNA packaging motor. *Nature.* 2000; 408:745–750. [PubMed: 11130079]
3. Simpson AA, Leiman PG, Tao Y, He Y, Badasso MO, Jardine PJ, et al. Structure determination of the head–tail connector of bacteriophage ϕ 29. *Acta Crystallog. sect. D.* 2001; 57:1260–1269.

Figure 1.
Stereo diagram showing the superposition of the C α back-bone of monomer A in the 2.1 Å resolution structure 1h5w (green) and the 2.9 Å resolution 1ijg structure (red).

Table 1

Statistical comparison of the $\phi 29$ connector wide domains in different structures

	RMS deviation		Maximum deviation		Number of C α atoms deviating more than 2 Å
	C α atoms	All atoms	C α atoms	All atoms	
1h5w <i>versus</i> 1fou	1.1	2.2	4.9	10.6	15
1h5w <i>versus</i> 1jnb	1.2	2.0	5.2	11.1	10
1h5w <i>versus</i> 1jig	0.8	1.6	6.4	11.4	5

The wide domain is defined as residues 38–128 and 247–284. All distances are in Å and correspond to comparisons of the A monomers in all structures.

Table 2

Superposition of the entire monomers in the four reported structures

	RMS deviation		Maximum deviation		Number of C α atoms deviating more than 2 Å
	C α atoms	All atoms	C α atoms	All atoms	
1h5w <i>versus</i> 1fou	1.1	1.8	4.9	10.6	20
1h5w <i>versus</i> 1jnb	1.1	1.8	5.1	11.3	19
1h5w <i>versus</i> 1jig	0.9	1.3	7.0	11.6	9 ^a
1jig <i>versus</i> 1jnb	1.1	1.9	6.0	11.8	18
Chain A <i>versus</i> chain B in 1jig	0.9	1.5	7.5	13.8	6
Chain A <i>versus</i> chain B in 1h5w	0.5	0.8	4.4	6.5	2

The number of the aligned C α atoms is 251. All distances are in Å.

^aThe number of differences greater than 2 Å varies from 1 to 9 depending on which of the 12 monomers are being compared.