Published in final edited form as:

Depress Anxiety. 2014 October ; 31(10): 834–842. doi:10.1002/da.22284.

Amygdala Response to Negative Stimuli Predicts PTSD Symptom Onset following a Terrorist Attack

Katie A. McLaughlin, Ph.D.¹, Daniel S. Busso, M.A.², Andrea Duys, B.A.¹, Jennifer Greif Green, Ph.D.³, Sonia Alves, B.A.², Marcus Way, B.A.⁴, and Margaret A. Sheridan, Ph.D.⁴

¹University of Washington

²Harvard Graduate School of Education

³Boston University School of Education

⁴Boston Children's Hospital, Harvard Medical School

Abstract

OBJECTIVE—Individuals with post-traumatic stress disorder (PTSD) exhibit heightened amygdala reactivity and atypical activation patterns in the medial prefrontal cortex (mPFC) in response to negative emotional information. It is unknown whether these aspects of neural function are risk factors for PTSD or consequences of either trauma exposure or onset of the disorder. We had a unique opportunity to investigate this issue following the terrorist attacks at the 2013 Boston Marathon and the ensuing manhunt and shelter in place order. We examined associations of neural function measured prior to the attack with PTSD symptom onset related to these events.

METHODS—A sample of 15 adolescents (mean age=16.5 years) who previously participated in a neuroimaging study completed a survey assessing posttraumatic symptoms related to the terrorist attack. We examined blood oxygen-level dependent (BOLD) response to viewing and actively down-regulating emotional responses to negative stimuli in regions previously associated with PTSD, including the amygdala, hippocampus, and mPFC, as prospective predictors of posttraumatic symptom onset.

RESULTS—Increased BOLD signal to negative emotional stimuli in the left amygdala was strongly associated with posttraumatic symptoms following the attack. Reduced bilateral hippocampal activation during effortful attempts to down-regulate emotional responses to negative stimuli was also associated with greater posttraumatic symptoms. Associations of amygdala reactivity with posttraumatic symptoms were robust to controls for pre-existing depression, anxiety, and PTSD symptoms and prior exposure to violence.

CONCLUSIONS—Amygdala reactivity to negative emotional information might represent a neurobiological marker of vulnerability to traumatic stress and, potentially, a risk factor for PTSD.

The corresponding author is Katie A. McLaughlin. Contact information is as follows: Katie McLaughlin, Ph.D., Assistant Professor of Psychology, University of Washington, Box 351525, Seattle, WA 98195, Phone: 206-616-7863, mclaughk@uw.edu.

The authors declare that, except for income received from their primary employers, no financial support or compensation has been received from any individual or corporate entity over the past three years for research or professional service and there are no personal financial holdings that could be perceived as constituting a potential conflict of interest.

Keywords

amygdala; hippocampus; trauma; terrorism; post-traumatic stress disorder; stress

Most people will experience a traumatic event at some point in their lives. Population-based data indicate that at least two-thirds of U.S. adults and youths will be exposed to a lifetime traumatic event. [1–4] Although trauma exposure is pervasive, only a minority of individuals develop post-traumatic stress disorder (PTSD). [1–4] The conditional risk of PTSD is less than 50% even for severe events. [5; 6] Despite extensive efforts to identify risk factors that confer vulnerability to PTSD following trauma exposure, meta-analyses suggest that psychosocial and environmental risk factors explain only about 20% of the variance in risk for PTSD among trauma-exposed individuals. [7; 8]

Recent work has attempted to identify neurobiological markers associated with PTSD vulnerability, such as autonomic nervous system and hypothalamic-pituitary-adrenal (HPA) axis activity in acutely traumatized individuals. However, these markers have not reliably predicted PTSD across studies. [9–11] In contrast, reduced hippocampal volume has been consistently observed among individuals with PTSD. [12] Small hippocampal volume has been observed among veterans with PTSD and their monozygotic twins discordant for trauma exposure, suggesting that reduced hippocampal volume could be a vulnerability marker for PTSD. [13] Additionally, numerous studies have observed differences in neural function among individuals with PTSD, including heightened amygdala activation, reduced activity in the ventromedial prefrontal cortex (vmPFC) and rostral anterior cingulate cortex (rACC) in response to emotional or threatening cues, and elevated activity in the dorsal anterior cingulate cortex (dACC) during fear conditioning, extinction learning recall, and response selection. [14–19] The extent to which these disruptions in neural function represent vulnerability markers for the disorder or consequences of trauma exposure or PTSD onset is unknown. However, atypical medial PFC function has been identified as a potential familial risk factor for PTSD, indicating that some of these differences might increase vulnerability to PTSD. [20; 21]

Identifying neurobiological vulnerability markers for PTSD *prior* to trauma exposure is challenging, in part because most traumatic events do not occur at random. A variety of individual-level characteristics predict trauma exposure, including socio-demographic factors, prior trauma, and psychopathology. [2; 4; 22–24] As such, it is difficult to disentangle, even in longitudinal studies, whether neurobiological indicators are markers of vulnerability to trauma-related psychopathology or simply predict differential risk of trauma exposure. One strategy for overcoming these challenges is to study traumatic events that are unrelated to pre-existing characteristics, such as natural disasters and terrorist attacks. However, neurobiological markers are not generally available in trauma-exposed individuals prior to these kinds of unpredictable and low-probability traumatic events. The only existing evidence regarding premorbid neural function and vulnerability to psychopathology following traumatic stressors comes from an innovative study in which neuroimaging data were collected on a sample of new recruits to the Israeli Defense Forces. Amygdala reactivity prior to combat exposure during anticipation of outcomes following a risky choice

and presentation of military-themed content predicted severity of PTSD symptoms following deployment. [25; 26] However, these findings could reflect, a) a marker of PTSD risk; b) a predictor of greater combat exposure, which has been linked to individual-level characteristics in military samples; [27; 28] or c) the effects of prior psychopathology or trauma exposure, factors strongly linked to PTSD [4; 22; 29] that were not examined in these studies.

In the current study, we examined whether aspects of neural function predicted onset of posttraumatic symptoms in adolescents following the terrorist attack at the 2013 Boston Marathon. The attack killed three spectators and injured hundreds of bystanders. Four days after the attack, a manhunt for the perpetrators resulted in an unprecedented shelter in place order that required residents of Boston and surrounding communities to remain indoors. The public transportation system, educational institutions, local government offices, and most businesses were closed. Although direct exposure to the attack was limited to spectators at the finish line of the marathon, hundreds of thousands of Boston residents watched the manhunt unfold live on television while the shelter in place order was in effect (about 12 hours). This kind of indirect exposure to terrorist attacks and their sequelae has been shown to precipitate PTSD symptoms in children and adults living in close proximity to the attack. [30–33] Here, we examine if neural function assessed in the year prior to the terrorist attack predicts posttraumatic symptom onset following the attack. These data provide a unique opportunity to examine pre-existing neural markers of risk for posttraumatic symptoms following a terrorist attack, and the only study, to our knowledge, to examine such predictors of response to an unpredictable traumatic event.

Materials and Methods

Sample

Following the Boston Marathon terrorist attack, we sent an online survey to adolescents who participated in studies in our lab in the two years prior to the event. Requests for parental permission to re-contact their children and adolescent surveys for those who provided consent were sent beginning one month following the attack and were open for a two-week period. Data for the current report were drawn specifically from an ongoing fMRI study. A total of 40 adolescents completed fMRI scans prior to the terrorist attack (time since scan=2.0–59.6 weeks). We obtained parental permission and received completed surveys from 15 of these participants (37.5% of the original sample). Participation in the survey was unrelated to pre-existing internalizing symptoms or child maltreatment, but adolescents with high community violence exposure were less likely to respond ($t=4.0$, $p<.001$). Participants ranged in age from 14.1 to 19.1 years at the time of scan ($M=16.49$ years) and from 14.8 to 19.9 years at the time of the survey ($M=17.25$ years). All participants reported exposure to media coverage of the bombings on the day of the marathon and the shelter in place order (with the exception of two participants who skipped the latter survey item). See Table 1 for sample socio-demographic and pre-attack characteristics and degree of exposure to the shelter in place order.

fMRI Task

Participants engaged in a well-established event-related task designed to assess neural markers of emotional reactivity and emotion regulation [34] that has previously been used in dozens of investigations, [35] including with children and adolescents. [36] Task design and construction of contrasts for analysis were based on the substantial prior literature on this task. Specifically, participants viewed neutral and negative images drawn from the International Affective Picture System (IAPS). [37] Prior to each image, participants were shown an instructional cue to either “look” or “decrease.” During *look* trials, participants were instructed to allow their emotions to unfold naturally, and to not engage in active strategies to modify their emotional response. During *decrease* trials, participants engaged in specific cognitive reappraisal strategies to try to reduce their emotional response.

All participants completed a training session prior to the MRI where they received detailed instructions about how to respond to each cue, observed examples completed out loud by a research assistant, and practiced with sample images not included in the task. On decrease trials, participants were instructed to think about the image in a way that made it psychologically more distant (e.g., imagine the scene as far away, that the situation did not involve them, that the people in the image were actors, etc.). These strategies have been used in previous studies with this task. [34; 36]

Stimuli were presented in 4 runs lasting 9 minutes each. The average valence and arousal of images and the number of faces within each image were equivalent for look and decrease trials. The stimuli were matched on valence and arousal during *look* and *decrease* trials. The instructional cue appeared for 2 seconds, the emotional stimulus appeared for 6–10 seconds, the rating screen appeared for 4 seconds, and the inter-trial interval (ITI) lasted from 1–6.5 seconds. Because the stimulus and ITI were jittered, each part of the task was modeled independently.

Image Acquisition

Scanning was performed on a 3T Siemens Trio scanner at the Harvard Center for Brain Science using a 32-channel head coil. Anatomical scans (T1-weighted multi-echo MPRAGE volumes) were acquired for co-registration with fMRI (TR=2530ms, TE=1640–7040ms, flip angle=7°, FOV=220mm², 176 slices, in-plane voxel size=1mm³). To reduce motion-related artifacts a navigator echo was used prior to scan acquisition, which compares slices to this echo online and permits up to 20% of slices be re-acquired.

Blood oxygenation level dependent (BOLD) signal during functional runs was acquired using a gradient-echo T2*-weighted EPI sequence. Thirty-two 3mm thick slices were acquired parallel to the AC-PC line (TR=2000ms, TE=30ms, flip angle=90degrees, bandwidth=2300, echo spacing=0.5, FOV=256×256, matrix size=64×64). Prior to each scan, four images were acquired and discarded to allow longitudinal magnetization to reach equilibrium. An online prospective motion correction algorithm (PACE) was used to reduce the effect of motion artifacts.

Image Processing

T1-weighted scans were processed using FreeSurfer version 5.0. [38–42] Automatic image segmentation was used to identify sub-cortical gray matter structures. Gray/white matter and gray matter/CSF boundaries were constructed using spatial intensity gradients across tissue classes. Following reconstruction, the cortex was parcellated based on the structure of gyri and sulci. [40; 43] The results were inspected and manually edited to optimize accurate placement of gray/white and gray/CSF borders based on shifts in the image intensity gradient. [38; 41] FreeSurfer morphometric procedures have demonstrated good test-retest reliability across scanner manufacturers and field strengths, [44; 45] been validated against manual measurement [46; 47] and histological analysis, [48] and used in children. [49–51]

Pre-processing and statistical analysis of fMRI data was performed in Nipype (<http://nipy.sourceforge.net/nipype/>), a platform that implements analysis tools from multiple software packages using the Python programming language. [52] fMRI pre-processing included spatial realignment, slice-time correction, and spatial smoothing (6mm FWHM), implemented in FSL. [53] Data were inspected for artifacts using custom artifact detection software (ART). [54] Volumes with motion >3 mm or >3 SD change in signal intensity were excluded from analysis, and 6 rigid-body motion regressors were included in person-level models. Person- and group-level models were estimated in FSL. A component-based anatomical noise correction method [55] was used to reduce noise associated with physiological fluctuations, including cardiac pulsations and respiratory-induced modulations of the magnetic field. Following estimation of the person-level models, the resulting contrast images were normalized into standard anatomical space, and anatomical co-registration of the functional data with each participant's T1-weighted image was performed. Normalization was implemented in Advanced Normalization Tools (ANTs) software. [56] These registration and normalization procedures are superior to standard techniques found in other software packages, particularly for children. [57]

Posttraumatic Symptoms

Posttraumatic symptoms *specifically related to the attack* were assessed using a brief version of the Impact of Events Scale-Revised (IES-R), [58] which has sound psychometric properties and discriminates between individuals with and without PTSD. [59; 60] The IES-6 is an abbreviated form of the IES-R, and is a widely used screener for PTSD symptoms. Respondents rated the frequency of hyperarousal (e.g. “I had trouble concentrating”), intrusive thoughts (e.g. “I thought about it when I didn’t want to”) and avoidance (e.g. “I tried not to think about it”) experienced since the bombings, on a 5-point Likert scale ranging from 0 (“almost never”) to 4 (“almost always”). Previous studies have shown that the IES-6 explains most of the variance of the IES-R[61], and the measure demonstrated good internal consistency in our sample ($\alpha=.89$).

Pre-Attack Survey Measures

As part of the study that took place prior to the terrorist attack, we collected information on internalizing symptoms, child maltreatment, and community violence exposure. Depressive symptoms were assessed with the Children’s Depression Inventory (CDI)[62], anxiety symptoms were assessed with the Multidimensional Anxiety Scale for Children (MASC),

[63; 64], and PTSD symptoms were assessed with the Youth Self Report Form. [65] Child maltreatment was assessed with the Childhood Trauma Questionnaire (CTQ), [66] a 28-item scale that assesses the frequency of maltreatment exposure during childhood and adolescence. [66; 67] We summed items from the physical, sexual, and emotional abuse sub-scales. Community violence exposure was assessed using the Screen for Adolescent Violence Exposure (SAVE)[68], a 32-item measure assessing violence exposure in school, home, and neighborhood contexts. We standardized the CTQ and SAVE scores and summed them to create an index of prior violence exposure.

Statistical Analysis

fMRI Analysis—To identify task related activity, regressors were created for each phase of the task: instructional cue, stimulus, and rating periods separately for look and decrease trials for neutral and negative stimuli. Using FSL, a general linear model (GLM) was constructed to estimate the association between variation in BOLD signal and task demands across time for each subject, prior to normalization. Using this GLM, individual-level estimates of BOLD activity were identified and submitted to group-level random effects models that contrasted activity across conditions. Here, we focus on neural activity to the emotional stimulus. We report a) emotional reactivity to negative stimuli (look negative > look neutral trials); and b) emotion regulation (decrease > look trials for negative stimuli), which are the standard contrasts used in studies of this task (see [35] for a meta-analysis). In the whole brain analysis we corrected for multiple comparisons using cluster-level correction in FSL, which estimates cluster-level significance using Gaussian Random Field Theory; our primary threshold was $p < .001$, as recommended [69].

We examined task-related activity in five regions-of-interest (ROIs), selected based on prior evidence for atypical patterns of neural activity to emotional stimuli in individuals with PTSD[70]. ROIs were included if there was significant task-related activation in those regions in the whole-brain analysis. Potential ROIs were the amygdala, hippocampus, dACC, rACC, and vmPFC. We constructed structural ROIs in each participant's native space using in FreeSurfer (Figure 1). We extracted the average estimate of neural activity within the entire ROI for each participant. For ROIs that met our *a priori* and task-active criteria, we examined associations with PTSD symptoms following the terrorist attack. Gender, age, and time since scan were included as covariates in all analysis. Finally, we conducted sensitivity analysis to determine whether our results were explained by pre-existing internalizing symptoms (depression, anxiety, or PTSD) or exposure to violence by re-estimating our final models after controlling for these factors.

Results

Neural Activity and Posttraumatic Symptoms

Emotional Reactivity—In whole-brain cluster-level corrected analysis, the amygdala and vmPFC exhibited significantly greater activation during viewing of negative emotional stimuli relative to neutral stimuli. ROI analyses revealed that left amygdala activation, $\beta = .72$, $p = .013$, to negative emotional stimuli was positively associated with posttraumatic symptoms related to the terrorist attack (Figure 2). No association was observed with right

amygdala activation, $\beta = .11$, $p = .78$, or vmPFC activation in either hemisphere, $\beta = .21-.41$, $p = .46-.15$.

Emotion Regulation—In whole-brain analysis, the hippocampus and vmPFC were recruited during regulation trials relative to trials involving simple viewing of negative stimuli. ROI analyses revealed that hippocampal activation during emotion regulation was negatively related to posttraumatic symptoms, such that lower hippocampal activation in both the right, $\beta = -.78$, $p = .046$, and left hemisphere, $\beta = -.67$, $p = .030$, was associated with higher symptoms (Figure 3).

Sensitivity Analysis

After controlling for pre-existing symptoms, the association between amygdala reactivity and posttraumatic symptoms remained significant in the left amygdala, $\beta = .98$, $p = .007$, and became marginally significant in the right amygdala, $\beta = 1.05$, $p = .070$. Hippocampal activation during emotion regulation remained marginally associated with posttraumatic symptoms in the left, $\beta = -.69$, $p = .077$, but not right, $\beta = -.71$, $p = .20$, hemisphere.

Activation in the left amygdala, $\beta = .70$, $p = .019$, during emotional reactivity trials also remained a significant predictor of posttraumatic symptoms after controlling for prior violence exposure. Neural activity during emotion regulation trials remained significantly associated with posttraumatic symptoms in the left hippocampus, $\beta = -.69$, $p = .025$, and marginally in the right hippocampus, $\beta = -.77$, $p = .053$, after controlling for prior violence.

When both internalizing symptoms and prior violence exposure were controlled, amygdala activation was a significant predictor of posttraumatic symptoms in both the left, $\beta = .97$, $p = .009$, and right, $\beta = 1.27$, $p = .027$, hemisphere. Hippocampal activation was no longer associated with posttraumatic symptoms, $\beta = -.66-.67$, $p = .14-.30$.

Discussion

Trauma exposure is pervasive, yet only a minority of individuals develops PTSD following a traumatic event. Identifying factors associated with PTSD risk is important not only for informing models of disorder etiology but also for improving efforts to target early interventions at trauma-exposed individuals most likely to develop PTSD. Here, we provide novel evidence that amygdala reactivity to negative emotional stimuli, measured prior to a terrorist attack, are associated with subsequent onset of posttraumatic symptoms. These findings suggest that amygdala reactivity to negative emotional information might represent a useful marker of PTSD risk.

Heightened amygdala activity in response to threatening or negative emotional stimuli has been observed consistently in individuals with PTSD. [15; 17; 18; 70] However, these studies have been unable to disentangle whether amygdala reactivity is a risk factor for PTSD or a consequence of trauma exposure or PTSD. Our findings suggest that heightened amygdala activity is associated with vulnerability to trauma-related psychopathology. Pre-deployment amygdala recruitment predicts PTSD symptoms following combat exposure. [25; 26] We extend this previous work in several ways. First, we measured posttraumatic

symptoms in response to a terrorist attack—an event for which exposure is unpredictable and independent of individual-level characteristics. In military samples, combat exposure is likely to be predicted by genetics and personality characteristics; [27; 28] even where service is mandatory, personality factors might shape degree of trauma exposure during combat, meaning that associations between pre-deployment measures of neural function and PTSD symptoms could reflect associations between neural function and degree of combat exposure. Second, our findings were robust to controls for pre-existing internalizing symptoms and prior violence exposure, factors not accounted for in previous military studies. Third, our sample was comprised of adolescents, indicating that individual differences in amygdala reactivity are associated with posttraumatic symptoms during the developmental period of highest risk for trauma exposure. [2] Finally, the task we used to assess amygdala response was designed to assess emotional reactivity and regulation in an ecologically valid manner, using realistic stimuli that have parallels to emotional situations encountered in daily life. Together with previous evidence from military samples, [25; 26] our findings suggest that amygdala reactivity represents a vulnerability marker for PTSD.

Reduced hippocampal activation during active efforts to reduce negative emotional reactions through cognitive reappraisal predicted higher posttraumatic symptoms related to the attack, although our sensitivity analysis suggests that this finding was explained by pre-existing internalizing symptoms. Theoretical models of neural function in PTSD argue that disruptions in hippocampal function are a central feature of the disorder. [71; 72] Deficits in contextual discrimination and in generalization of extinction learning to novel contexts might contribute to poor contextualization of fear cues and deficits in identifying safety cues in PTSD. [71; 72] However, the precise nature of hippocampal deficits in PTSD remains to be elucidated. Previous studies have examined hippocampal function largely within the context of long-term memory tasks, with some reporting reduced activation during encoding and retrieval, [73] and others documenting elevated activation. [74] Our findings suggest that hippocampal response to emotional stimuli might be important to consider in future neuroimaging studies of PTSD.

We found no evidence for an association between medial PFC (mPFC) activation during emotional reactivity or regulation trials and posttraumatic symptoms. A variety of deficits in functioning of mPFC regions have been observed in PTSD, including reduced vmPFC activation and elevated dACC activation during fear extinction recall, [14–19] some of which appear to be familial risk factors for PTSD. [20; 21] This finding should be interpreted with caution given the small size of our sample and the fact that our task was not designed to assess mPFC function. In addition, because we utilized structural ROIs, the lack of association could reflect the relatively larger size of cortical than sub-cortical ROIs.

Study findings should be interpreted in light of several key limitations. Most notably, the number of participants in the study was small. This limitation is particularly important to consider in interpreting null findings. Second, the response rate to the survey was low. This was due largely to parental non-response to our email requesting permission to contact their children, and was likely influenced by the short period of time (2 weeks) in which we actively collected survey responses. Our analysis of selection bias with regard to survey completion suggests that pre-existing internalizing symptoms and child maltreatment did not

influence the likelihood of responding, but that adolescents with high exposure to community violence were less likely to participate. Third, we were unable to collect follow-up neuroimaging data, which would have allowed us to determine whether exposure to the attacks was associated with changes in neural function. Finally, we assessed posttraumatic symptoms in relation to an event of low severity with regard to objective life threat that no longer qualifies as a Criterion A traumatic event in DSM-5 [75]. However, considerable variation in posttraumatic symptoms associated with the event was observed, consistent with prior evidence indicating that terrorist attacks can trigger PTSD symptoms even in individuals who were not directly exposed. [30–33] Moreover, the unique nature of the terrorist attacks at the Boston Marathon and the ensuing manhunt likely resulted in appraisals of perceived threat for youths who were not present at the bombings but were required to stay indoors during the shelter in place order. [76] Nonetheless, future research is needed to replicate the patterns observed here in response to other types of traumatic events.

Conclusion

Heightened amygdala reactivity to negative emotional information is associated with future onset of posttraumatic symptoms following a terrorist attack, independent of prior internalizing symptoms—including symptoms of PTSD—and violence exposure. These findings suggest that elevated amygdala reactivity to negative emotional information could represent a neurobiological marker of vulnerability to traumatic stress and, potentially, a risk factor for PTSD.

Acknowledgments

This research was supported by grants from the National Institutes of Health (K01-MH092526, and K01-MH092555). These funders provided support for all data collection and analysis.

References

1. Kessler RC, Sonnega A, Bromet E, et al. Posttraumatic stress disorder in the National Comorbidity Survey. *Archives of General Psychiatry*. 1995; 52:1048–1060. [PubMed: 7492257]
2. Breslau N, Kessler RC, Chilcoat HD, et al. Trauma and posttraumatic stress disorder in the community: The 1996 Detroit Area Survey of Trauma. *Archives of General Psychiatry*. 1998; 55:626–632. [PubMed: 9672053]
3. Roberts AL, Gilman SE, Breslau J, et al. Race/ethnic differences in exposure to traumatic events, development of post-traumatic stress disorder, and treatment-seeking for post-traumatic stress disorder in the United States. *Psychological Medicine*. 2011; 41:71–83. [PubMed: 20346193]
4. McLaughlin KA, Koenen KC, Hill E, et al. Trauma exposure and posttraumatic stress disorder in a US national sample of adolescents. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2013; 52:815–830. [PubMed: 23880492]
5. Dohrenwend BP, Turner JB, Turse NA, et al. The psychological risks of Vietnam for U.S. veterans: A revisit with new data and methods. *Science*. 2006; 313:979–982. [PubMed: 16917066]
6. Widom CS. Posttraumatic stress disorder in abused and neglected children grown up. *American Journal of Psychiatry*. 1999; 156:1223–1229. [PubMed: 10450264]
7. Ozer EJ, Best SR, Lipsey TL, Weiss DS. Predictors of posttraumatic stress disorder and symptoms in adults: A meta-analysis. *Psychological Bulletin*. 2003; 129:52–73. [PubMed: 12555794]
8. Brewin CR, Andrews B, Valentine JD. Meta-analysis of risk factors for posttraumatic stress disorder in trauma-exposed adults. *Journal of Consulting and Clinical Psychology*. 2000; 68:748–766. [PubMed: 11068961]

9. Bryant RA, Creamer M, O'Donnell M, et al. A multisite study of initial respiration rate and heart rate as predictors of posttraumatic stress disorder. *Journal of Clinical Psychiatry*. 2008; 69:1694–1701. [PubMed: 19014750]
10. Shalev AY, Videlock EJ, Peleg T, et al. Stress hormones and post-traumatic stress disorder in civilian trauma victims: a longitudinal study. Part I: HPA axis responses. *International Journal of Neuropsychopharmacology*. 2008; 11:365–372. [PubMed: 17971262]
11. Videlock EJ, Peleg T, Segman R, et al. Stress hormones and post-traumatic stress disorder in civilian trauma victims: a longitudinal study. Part II: the adrenergic response. *International Journal of Neuropsychopharmacology*. 2008; 11:373–380. [PubMed: 17971259]
12. Kitayama N, Vaccarino V, Kutner M, et al. Magnetic resonance imaging (MRI) measurement of hippocampal volume in posttraumatic stress disorder: A meta-analysis. *Journal of Affective Disorders*. 2005; 88:79–86. [PubMed: 16033700]
13. Gilbertson MW, Shenton ME, Ciszewski A, et al. Smaller hippocampal volume predicts pathologic vulnerability to psychological trauma. *Nature Neuroscience*. 2002; 5:1242–1247.
14. Pitman RK, Rasmusson AM, Koenen KC, et al. Biological studies of post-traumatic stress disorder. *Nature Reviews Neuroscience*. 2012; 13:769–787.
15. Etkin A, Wager TD. Functional neuroimaging of anxiety: A meta-analysis of emotional processing in PTSD, social anxiety disorder, and specific phobia. *American Journal of Psychiatry*. 2007; 164:1476–1488. [PubMed: 17898336]
16. Milad MR, Pitman RK, Ellis CB, et al. Neurobiological basis of failure to recall extinction memory in posttraumatic stress disorder. *Biological Psychiatry*. 2009; 66:1075–1082. [PubMed: 19748076]
17. Shin LM, Wright CL, Cannistraro PA, et al. A functional magnetic resonance imaging study of amygdala and medial prefrontal cortex responses to overtly presented fearful faces in posttraumatic stress disorder. *Archives of General Psychiatry*. 2005; 62:273–281. [PubMed: 15753240]
18. Rauch SL, Whalen PJ, Shin LM, et al. Exaggerated amygdala response to masked facial stimuli in posttraumatic stress disorder: a functional MRI study. *Biological Psychiatry*. 2000; 47:769–776. [PubMed: 10812035]
19. Shin LM, Whalen PJ, Pitman RK, et al. An fMRI study of anterior cingulate function in posttraumatic stress disorder. *Biological Psychiatry*. 2001; 50:932–942. [PubMed: 11750889]
20. Shin LM, Lasko NB, Macklin ML, et al. Resting metabolic activity in the cingulate cortex and vulnerability to posttraumatic stress disorder. *Archives of General Psychiatry*. 2009; 66:1099–1107. [PubMed: 19805700]
21. Shin LM, Bush G, Milad MR, et al. Exaggerated activation of dorsal anterior cingulate cortex during cognitive interference: A monozygotic twin study of posttraumatic stress disorder. *American Journal of Psychiatry*. 2011; 168:979–985. [PubMed: 21724666]
22. Koenen KC, Moffit TE, Caspi A, et al. The developmental mental-disorder histories of adults with posttraumatic stress disorder: A prospective longitudinal birth cohort study. *Journal of Abnormal Psychology*. 2008; 117:460–466. [PubMed: 18489223]
23. Koenen KC, Moffit TE, Poulin R, et al. Early childhood factors associated with the development of post-traumatic stress disorder: results from a longitudinal birth cohort. *Psychological Medicine*. 2007; 37:181–192. [PubMed: 17052377]
24. Breslau N, Wilcox HC, Storr CL, et al. Trauma exposure and posttraumatic stress disorder: A study of youths in urban America. *Journal of Urban Health: Bulletin of the New York Academy of Medicine*. 2004; 81:530–544. [PubMed: 15466836]
25. Admon R, Lubin G, Rosenblatt JD, et al. Imbalanced neural responsivity to risk and reward indicates stress vulnerability in humans. *Cerebral Cortex*. 2013; 23:28–35. [PubMed: 22291028]
26. Admon R, Lubin G, Stern O, et al. Human vulnerability to stress depends on amygdala's predisposition and hippocampal plasticity. *Proceedings of the National Academy of Sciences*. 2009; 106:14120–14125.
27. Koenen KC, Harley R, Lyons M, et al. A twin registry study of familial and individual risk factors for trauma exposure and posttraumatic stress disorder. *Journal of Nervous and Mental Disease*. 2002; 190:209–218. [PubMed: 11960081]

28. Lyons MJ, Goldberg J, Eisen SA, et al. Do genes influence exposure to trauma? A twin study of combat. *American Journal of Medical Genetics*. 1993; 48:22–27. [PubMed: 8357033]
29. Storr CL, Ialongo N, Anthony JC, Breslau N. Childhood antecedents of exposure to traumatic events and posttraumatic stress disorder. *American Journal of Psychiatry*. 2007; 164:119–125. [PubMed: 17202553]
30. Comer JS, Kendall PC. Terrorism: The psychological impact on youth. *Clinical Psychology: Science and Practice*. 2007; 14:179–212.
31. Galea S, Vlahov D, Resnick H, et al. Trends of probable post-traumatic stress disorder in New York City after the September 11 terrorist attacks. *American Journal of Epidemiology*. 2003; 158:514–524. [PubMed: 12965877]
32. Ahern J, Galea S, Resnick H, et al. Television images and psychological symptoms after the September 11 terrorist attacks. *Psychiatry*. 2002; 65:289–300. [PubMed: 12530330]
33. Otto MW, Henin A, Hirschfeld-Becker DR, et al. Posttraumatic stress disorder symptoms following media exposure to tragic events: Impact of 9/11 on children at risk for anxiety disorders. *Journal of Anxiety Disorders*. 2007; 21:888–902. [PubMed: 17276653]
34. Ochsner KN, Ray RD, Cooper JC, et al. For better or worse: neural systems supporting the cognitive down- and up-regulation of negative emotion. *Neuroimage*. 2004; 23:483–499. [PubMed: 15488398]
35. Buhle J, Silvers JA, Wager TD, et al. Cognitive Reappraisal of Emotion: A Meta-Analysis of Human Neuroimaging Studies. *Cerebral Cortex*. in press.
36. McRae K, Gross JJ, Weber J, et al. The development of emotion regulation: an fMRI study of cognitive reappraisal in children, adolescents, and young adults. *SCAN*. 2012; 7:11–22. [PubMed: 22228751]
37. Lang, PJ.; Bradley, MM.; Cuthbert, BN. International Affective Picture System (IAPS): Affective ratings of pictures and instruction manual. Gainesville, FL: University of Florida; 2008.
38. Fischl B, Dale AM. Measuring the thickness of the human cerebral cortex from magnetic resonance images. *Proceedings of the National Academy of Sciences*. 2000; 97:11050–11055.
39. Fischl B, Salat DH, Busa E, et al. Whole brain segmentation: automated labeling of neuroanatomical structures in the human brain. *Neuron*. 2002; 33:341–355. [PubMed: 11832223]
40. Fischl B, van der Kouwe A, Destrieux C, et al. Automatically parcellating the human cerebral cortex. *Cerebral Cortex*. 2004; 14:11–22. [PubMed: 14654453]
41. Dale AM, Fischl B, Sereno MI. Cortical surface-based analysis. I. Segmentation and surface reconstruction. *Neuroimage*. 1999; 9:179–194. [PubMed: 9931268]
42. Fischl B, Sereno MI, Dale AM. Cortical surface-based analysis. II. Inflation, flattening, and a surface-based coordinate system. *Neuroimage*. 1999; 9:195–207. [PubMed: 9931269]
43. Desikan RS, Segonne F, Fischl B, et al. An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *Neuroimage*. 2006; 31:968–980. [PubMed: 16530430]
44. Han X, Jovicich J, Salat D, et al. Reliability of MRI-derived measurements of human cerebral cortical thickness: the effects of field strength, scanner upgrade and manufacturer. *Neuroimage*. 2006; 32:180–194. [PubMed: 16651008]
45. Reuter M, Schmansky NJ, Rosas HD, Fischl B. Within-subject template estimation for unbiased longitudinal image analysis. *Neuroimage*. 2012; 61:1402–1418. [PubMed: 22430496]
46. Kuperberg GR, Broome MR, McGuire PK, et al. Regionally localized thinning of the cerebral cortex in schizophrenia. *Archives of General Psychiatry*. 2003; 60:878–888. [PubMed: 12963669]
47. Salat DH, Buckner RL, Snyder AZ, et al. Thinning of the cerebral cortex in aging. *Cerebral Cortex*. 2004; 14:721–730. [PubMed: 15054051]
48. Rosas HD, Liu AK, Hersch S, et al. Regional and progressive thinning of the cortical ribbon in Huntington's disease. *Neurology*. 2002; 58:695–701. [PubMed: 11889230]
49. Sheridan MA, Fox NA, Zeanah CH, et al. Variation in neural development as a result of exposure to institutionalization early in childhood. *Proceedings of the National Academy of Sciences*. 2012; 109(32):12927–12932.

50. Tottenham N, Hare T, Quinn BT, et al. Prolonged institutional rearing is associated with atypically larger amygdala volume and difficulties in emotion regulation. *Developmental Science*. 2010; 13:46–61. [PubMed: 20121862]
51. Ostby Y. Heterogeneity in subcortical brain development: A structural magnetic resonance imaging study of brain maturation from 8 to 30 years. *Journal of Neuroscience*. 2009; 29:11772–11782. [PubMed: 19776264]
52. Gorgolewski K, Burns CD, Madison C, et al. Nipype: a flexible, lightweight and extensible neuroimaging data processing framework in Python. *Frontiers in Neuroinformatics*. 2011; 5:Article 13.
53. Jenkinson M, Beckmann CF, Behrens TE, et al. FSL. *Neuroimage*. 2012; 62:782–790. [PubMed: 21979382]
54. Mazaika, PK.; Whitfield-Gabrieli, S.; Reiss, AL.; Glover, G. *Human Brain Mapping*. Chicago: 2007. Artifact repair of fMRI data from high motion clinical subjects.
55. Behzadi Y, Restom K, Liao J, Liu TT. A component based noise correction method (CompCor) for BOLD and perfusion based fMRI. *Neuroimage*. 2007; 37:90–101. [PubMed: 17560126]
56. Klein A, Andersson J, Ardekani BA, et al. Evaluation of 14 nonlinear deformation algorithms applied to human brain MRI registration. *Neuroimage*. 2009; 46:786–802. [PubMed: 19195496]
57. Ghosh SS, Kakunoori S, Augustinack J, et al. Evaluating the validity of volume-based and surface-based brain image registration for developmental cognitive neuroscience studies in children 4 to 11 years of age. *Neuroimage*. 2010; 53:85–93. [PubMed: 20621657]
58. Weiss, DS.; Marmar, CR. The impact of event scale-revised. In: Wilson, JP.; Keane, TM., editors. *Assessing psychological trauma and PTSD*. New York, NY: Guilford; 2004. p. 168-189.
59. Sundin EC, Horowitz MJ. Impact of Event Scale: psychometric properties. *British Journal of Psychiatry*. 2002; 180:205–209. [PubMed: 11872511]
60. Creamer M, Bell R, Failla S. Psychometric properties of the Impact of Events Scale-Revised. *Behaviour Research and Therapy*. 2003; 41:1489–1496. [PubMed: 14705607]
61. Thoresen S, Tambs K, Hussain A, et al. Brief measure of posttraumatic stress reactions: Impact of Event Scale-6. *Social Psychiatry and Psychiatric Epidemiology*. 2010; 45(3):405–412. [PubMed: 19479171]
62. Kovacs, M. *Children's Depression Inventory manual*. North Tonawanda, NY: Multi-Health Systems; 1992.
63. March JS, Parker JDA, Sullivan K, et al. The Multidimensional Anxiety Scale for Children (MASC): Factor structure, reliability, and validity. *Journal of the American Academy of Child & Adolescent Psychiatry*. 1997; 36:554–565. [PubMed: 9100431]
64. Muris P, Merckelbach H, Ollendick T, et al. Three traditional and three new childhood anxiety questionnaires: their reliability and validity in a normal adolescent sample. *Behaviour Research and Therapy*. 2002; 40:753–772. [PubMed: 12074371]
65. Achenbach, TM. *Integrative guide for the 1991 CBCL/4-18, YSR and TRF Profiles*. Burlington, VT: Department of Psychiatry, University of Vermont; 1991.
66. Bernstein DP, Ahluvalia T, Pogge D, Handelsman L. Validity of the Childhood Trauma Questionnaire in an adolescent psychiatric population. *Journal of the American Academy of Child & Adolescent Psychiatry*. 1997; 36:340–348. [PubMed: 9055514]
67. Bernstein DP, Fink L, Handelsman L, et al. Initial reliability and validity of a new retrospective measure of child abuse and neglect. *American Journal of Psychiatry*. 1994; 151:1132–1136. [PubMed: 8037246]
68. Hastings TL, Kelley ML. Development and validation of the Screen for Adolescent Violence Exposure (SAVE). *Journal of Abnormal Child Psychology*. 1997; 25:511–520. [PubMed: 9468111]
69. Woo C-W, Krishnan A, Wager TD. Cluster-extent based thresholding in fMRI analyses: Pitfalls and recommendations. *Neuroimage*. 2014; 91:412–419. [PubMed: 24412399]
70. Shin LM, Rauch SL, Pitman RK. Amygdala, medial prefrontal cortex, and hippocampal function in PTSD. *Annals of the New York Academy of Sciences*. 2006; 1071:67–79. [PubMed: 16891563]

71. Rauch SL, Shin LM, Phelps EA. Neurocircuitry models of posttraumatic stress disorder and extinction: Human neuroimaging research-past, present, and future. *Biological Psychiatry*. 2006; 60:376–382. [PubMed: 16919525]
72. Liberzon I, Sripada CS. The functional neuroanatomy of PTSD: a critical review. *Progress in Brain Research*. 2008; 167:151–169. [PubMed: 18037013]
73. Bremner JD, Vythilingam M, Vermetten E, et al. MRI and PET study of deficits in hippocampal structure and function in women with childhood sexual abuse and posttraumatic stress disorder. *American Journal of Psychiatry*. 2003; 160:924–932. [PubMed: 12727697]
74. Werner NS, Meindl T, Engel RR, et al. Hippocampal function during associative learning in patients with posttraumatic stress disorder. *Journal of Psychiatric Research*. 2009; 43:309–318. [PubMed: 18490028]
75. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 5. Washington, DC: American Psychiatric Press; 2013. (DSM-5)
76. Marshall RD, Bryant RA, Amsel L, et al. The psychology of ongoing threat: Relative risk appraisal, the September 11 attacks, and terrorism-related fears. *American Psychologist*. 2007; 62:304–316. [PubMed: 17516775]

Figure 1.

Structural regions of interest (ROIs) for cortical and subcortical regions.

Three regions within medial prefrontal cortex (mPFC) were examined. These included dorsal anterior cingulate cortex (light purple; caudal_acc), rostral anterior cingulate cortex (dark purple; rostral_acc) and ventromedial prefrontal cortex (pink; medial orbitofrontal and frontal pole). All regions of interest are defined anatomically based on the individual's own anatomy using FreeSurfer and shown on the expanded cortical surface for viewing purposes; cortical regions were defined using the 2005 segmentation atlas. Within the subcortex two regions were identified: the amygdala (blue) and hippocampus (yellow). All regions shown here are from individual representative subject's aparc (cortex) and aseg (subcortex) segmentation. Activation during contrasts of interest was extracted from within the entire structure for each ROI.

Cluster-level corrected activation for look negative > look neutral contrast. Amygdala activation in the left but not right hemisphere was significantly associated with PTSD symptoms.

Figure 2.

Amygdala reactivity to negative vs. neutral images and PTSD symptoms related to the Boston Marathon terrorist attack.

Left amygdala activation for the contrast of look negative > look neutral. Model controls for age, gender, time since scan, and symptoms of depression and anxiety prior to the attack; R^2 represents contribution of amygdala activity over and above these covariates.

Cluster-level corrected activation for decrease > look negative contrast. Hippocampal activation in both hemispheres was significantly associated with PTSD symptoms.

Figure 3.

Hippocampal activation during effortful emotion regulation vs. viewing negative images and PTSD symptoms related to the Boston Marathon terrorist attack.

Left hippocampal activation for the contrast of decrease > look negative. Model controls for age, gender, time since scan, and symptoms of depression and anxiety prior to the attack; R^2 represents contribution of hippocampal activity over and above these covariates.

Table 1

Distribution of socio-demographic factors, exposure to media coverage and the shelter in place order, prior trauma exposure, and pre-existing psychopathology

	%	(n)
Female	66.7	10
Race/Ethnicity		
White	46.7	7
Black	26.7	4
Latino	13.3	2
Other/Biracial	13.3	2
Exposure to Media Coverage [/]	100.0	13
Exposure to Shelter in Place [/]	76.9	10
	M	(SD)
Prior Maltreatment (CTQ)	23.53	(10.64)
Prior Community Violence (SAVE)	44.93	(7.91)
Pre-existing Depressive Symptoms (CDI)	3.08	(1.83)
Pre-existing Anxiety Symptoms (MASC)	25.58	(17.05)
Pre-existing PTSD Symptoms (YSR)	55.29	(7.35)

[/]Two participants skipped the survey questions about exposure to the shelter in place order and media coverage during that day. Proportions are reported based on the sample of participants who responded to these questions.

Abbreviations: CTQ, Childhood Trauma Questionnaire; SAVE, Screen for Adolescent Violence Exposure; CDI, Children's Depression Inventory; MASC, Multidimensional Anxiety Scale for Children; YSR, Youth Self-Report Form.