Nutritional State of Children in the Democratic People's Republic of Korea (DPRK): Based on the DPRK Final Report of the National Nutrition Survey 2012

Jeong Eun Kim

Department of Pediatrics, Shincheon Union Hospital, Siheung, Korea

A nationwide nutrition survey began when the Democratic People's Republic of Korea (DPRK) requested assistance from international relief organizations in 1997 due to flooding in 1995-1996, followed by the worsening food shortage peak in 1997. According to reported data for the 15 years since the active intervention and assistance from international societies, the malnutrition and mortality rates of children in the DPRK have improved. However, the prevalence of the stunting reported in the latest 2012 report is 27.9%, which is still a moderate – severe level, and worrisome in terms of international standards. In particular, the prevalence in Ryangangdo, which is regarded as the worst region in the DPRK, is 39.6%, which is a very high level of stunting. To alleviate such regional deviation will be a major task for future assistance. In addition, one cannot emphasize too highly the importance of early nutritional assistance for pregnant women and infants, considering that the recovery from stunted growth is low after two years of age, and the aftereffects would continue for the rest of their lives.

Key Words: Nutrition status, Child, Democratic People's Republic of Korea, Nutrition surveys

INTRODUCTION

The Democratic People's Republic of Korea (DPRK) has conducted a Nutritional Examination Survey every two years since 1998, together with the aid of the United Nations Children's Fund (UNICEF) and World Food Program. There are two survey types of nutrition assessment report in the DPRK: Multiple Indicator Cluster Surveys (MICS) and National Nutritional Surveys (NS). The MICS is a survey sponsored by

UNICEF to evaluate the results of the achievements and action plans of each nation for 10 years, and includes a mid-term evaluation. It has been active since the UN Convention on the Rights of the Child in 1990 to promote the well-being and welfare of children. The NS has been performed to provide and monitor basic data for program plans and executions conducted to rebuild infrastructure and food assistance at the initial phase of humanitarian assistance for the DPRK.

It was conducted without the assistance of inter-

Received : September 13, 2014, Accepted : September 21, 2014

Corresponding author: Jeong Eun Kim, Department of Pediatrics, Shincheon Union Hospital, 57, Bokji-ro, Siheung 429-810, Korea.
Tel: +82-31-310-6357, Fax: +82-31-310-6400, E-mail: galmuae1@naver.com

Copyright © 2014 by The Korean Society of Pediatric Gastroenterology, Hepatology and Nutrition

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

national organizations independently in both 2000 and 2006. However, the data published in 2006 is not cited much, due to its low credibility. Since then, after a dormant period, it was resumed in 2009 in the name of the MICS, while the NS published in March 2012 was both the next publication and the latest survey report. This survey report was conducted September 17 to October 17, 2012, and it included 10 regions in the DPRK (including 40 clusters in each region) with 8,040 children at 0-59 months, and 7,649 mothers 15-49 years old. For the survey indicators, nutritional status, anemia, Mid-Upper Arm Circumference, multiple micronutrient supplements for pregnant women, dietary habits, and food diversity were selected as subjects and were divided into children and pregnant women. The main results are shown in Table 1 [1].

NUTRITIONAL STATUS OF CHILDREN UNDER FIVE YEARS OLD

Since a low height is a symptom of long-term malnutrition, height-for-age was set as an indicator of

stunting, while weight-for-height was set as an indicator of wasting, since weight can be reduced because of a short-term food intake reduction. "Underweight" refers to weight-for-age, which was

Fig 1. Trends of prevalence by malnutrition type. Data source: Excerpt from the reports of multiple indicator cluster surveys (MICS) [2]/nutrition surveys (NS) [3] in 1998, MICS in 2000 [4], NS in 2002 [5], NS in 2004 [6], MICS in 2009 [7] and NS in 2012 [1].

Table 1. Main Results of National Nutrition Survey 2012, Democratic People's Republic of Korea, October 2012 [1]

Indicator	Age group	Prevalence	95% confidence interval
Children			
Global chronic malnutrition (stunting)	0-59 months	27.9%	26.3-29.4
Severe chronic malnutrition (severe stunting)	0-59 months	7.2%	6.4-7.9
Global acute malnutrition (wasting)	0-59 months	4.0%	3.5-4.5
Severe acute malnutrition (severe wasting)	0-59 months	0.6%	0.4-0.7
Underweight	0-59 months	15.2%	14.2-16.3
Low MUAC (<125 mm)	0-59 months	4.1%	3.6-4.6
Anaemia	0-59 months	28.7%	26.5-30.9
Early initiation of breastfeeding	0-23.9 months	28.1	26.2-30.0
Exclusive breastfeeding	0-5.9 months	68.9	65.1-72.8
Timely introduction of solid, semi-solid and soft food	6-8 months	66.0	61.4-70.5
Mean number of food groups consumed (maximum 7 food groups)	6-23.9 months	2.5	2.4-2.6
Minimum dietary diversity (consumption of 4 out of 7 food groups)	6-23.9 months	26.5	24.0-28.9
Women			
Low MUAC (<225 mm)	15-49 years	23.2%	21.9-24.5
(<210 mm)	15-49 years	5.3%	4.7-5.9
Anaemia	15-49 years	31.2%	29.6-32.9
Multi-micronutrient supplementation for at least 6 months during pregnancy	15-49 years	26.9%	24.6-29.1
Mean number of food groups consumed (maximum 9 food groups)	15-49 years	3.6	3.5-3.6
Minimum Dietary Diversity (consumption of at least 4 out of 9 food groups)	15-49 years	49.6	47.7-51.4

MUAC: Mid-Upper Arm Circumference

set as an indicator of wasting and stunting.

The 2012 survey showed improved results overall compared to 2009 (Fig. 1) [1-7]. However, regional deviation became more severe. For example, Ryangangdo, where stunting prevalence was the highest, showed 43% in 2004 to 39.6% in 2012, which is a decrease of 3.4%. Meanwhile, Pyongyang, where stunting prevalence was the lowest, showed 24% in

Fig. 2. Prevalence of global ($H/A < -2$ Z-scores), moderate ($H/A \geq -3$ Z-scores), and severe ($H/A < -3$ Z-scores) chronic malnutrition based on height of age Z-score per province in children aged 0-59 months. Democratic People's Republic of Korea, October 2012 (95% confidence interval) [1]. H/A: height/age.

2004 and 15.6% in 2012, revealing an 8.4% decrease. This result shows that it is urgently necessary to assist Ryangangdo, Hamgyongnamdo, and Jagangdo, which were chronically vulnerable districts in which stunting rates exceed 9-10%. Fig. 2 shows the regional prevalence of stunting in 2012 [1].

Fig. 3 shows the prevalence of stunting in 2012 by age, while Table 2 summarizes the trends by age since 1998 [1-7]. It is noted that the recovery rate of stunting stagnated after two years old, despite the

Fig. 3. Global, moderate & severe chronic malnutrition prevalence by age groups in months. Democratic People's Republic of Korea, October 2012 (95% confidence interval) [1].

Table 2. Prevalence of Nutrition Types and Trends by Age [1-7]

Indicator	Stunting Height-for-age $< -2SD$						Wasting Weight-for-height $< -2SD$						Underweight Weight-for-age $< -2SD$					
	1998	2000	2002	2004	2009	2012	1998	2000	2002	2004	2009	2012	1998	2000	2002	2004	2009	2012
Survey year	1998	2000	2002	2004	2009	2012	1998	2000	2002	2004	2009	2012	1998	2000	2002	2004	2009	2012
Age (m)																		
0-5	-	21.9	17.3	19.4	12.5	3.7	-	7.8	5.3	6.0	1.8	4.3				14.1	5.6	
6-11	14.5	31.9	22.7		23.6	8.4	17.6	10.4	6.7		4.8	4.9						17.4
12-23	48.4	50.2	41.6	27.1	23.9	27.0	30.9	11.9	11.9	8.7	5.7	4.3				20.5	16.2	
24-35	62.2	47.5	45.6	34.7	32.1	36.8	20.5	9.7	8.4	7.3	6.2	3.9				27.3	21.7	
36-47	75.1	58.6	47.6	46.7	39.5	33.3	13.4	10.2	7.4	7.5	5.0	3.6				26.3	21.7	
48-59	77.5	60.3	47.5	50.8	46.5	35.4	8.9	11.9	6.3	6.2	5.2	3.3				26.6	21.3	
60-84	74.8	-	44.2	46.8			7.8	-	5.6	5.3						22.5		
Average (m)	62.3	45.2	39.2	37.0	32.4	27.9	15.6	10.4	8.1	7.0	5.2	4	60.6	27.9	21.2	23.4	18.8	15.2

Children under six months old were removed from the survey in 1998.

In 2004, the survey's scope was reduced by combining 0-11 months and 60-72 months.

SD: standard deviation.

overall reduction of prevalence. The prevalence of stunting in 24-35 months old children showed an increasing trend from 32.1% in 2009 to 36.8%. A period of about 1,000 days from pre-pregnancy to 24 months of age that can prevent children from developing stunted growth is called the window of opportunity, and it indicates that the early intervention of nutritional assistance for pregnant women and infants is essential from the prenatal period to 24 months of age in which neurodevelopment is completed. Malnutrition in this period can result in

Fig. 4. Causes of under-five deaths in Democratic People's Republic of Korea count down to 2015, report 2010 (source: World Health Organization, 2010). HIV: human immunodeficiency virus, AIDS: acquired immune deficiency syndrome.

Fig. 5. Child Mortality Trend in Democratic People's Republic of Korea 1990-2011. IMR: infant mortality rate.

irrevocable aftereffects, not only for physical growth, but also for developmental and cognitive statuses at later stages of life.

DEATH OF CHILDREN UNDER FIVE

Cause of death (World Health Organization, 2010)

Around 35% of the cause of children's death around the world is related to malnutrition (Lanset Maternal and Child Undernutrition series 2008). In light of this finding, 4,000 children under five who have died early every year in the DPRK were in a state of under-nutrition before they died (Fig. 4) [8].

Children mortality

Children mortality under five has decreased. However, 11,406 children (31 children daily) still die before they reach their fifth birthday according to the State of the World's Children (SOWC) report in 2010. Approximately 6,000 children (16 children daily) die before they reach 28 days of age. Furthermore, approximately 289 women (five per week) die due to childbirth and related complications (Fig. 5) [8].

CHILDREN ANEMIA AND BREASTFEEDING

The incidence of child anemia in the DPRK was 28.7%, which is very high. This high incidence is likely associated with insufficient breastfeeding and supplemented food that contains low quality nutrients, and insufficient supplemented food intake. A low breastfeeding rate, maternal anemia, and malnutrition can also be attributed to main causes of high children anemia. Wasting children showed a high rate of anemia compared to children in a normal group, so the prevention of anemia can be an important measure that can prevent wasting.

CONCLUSION

The latest 2012 report for nutrition status of the DPRK showed the malnutrition and mortality rates

of children in the DPRK have improved. However, the prevalence of the stunting reported is still a moderate – severe level and regional deviation became more severe. The 1,000 days between pregnancy and age two are window of opportunity for appropriate growth & development. So early nutritional assistance have high importance.

ACKNOWLEDGEMENTS

This paper, written by the author, was published in “Annual Report of Healthcare in the DPRK in 2013” by Medical Aid for Children Non-Profit Organization and excerpted with the permission of the organization.

REFERENCES

1. UNICEF. Democratic People’s Republic of Korea final report of the National Nutrition Survey 2012. Pyongyang: UNICEF; 2012 [cited 2013 Mar]. Available from: http://www.unicef.org/eapro/DPRK_National_Nutrition_Survey_2012.pdf.
2. UNICEF Pyongyang. Report on the Multiple Indicator Cluster Survey in the DPRK. Pyongyang: UNICEF; 1998.
3. EU, UNICEF, WFP. Nutrition Survey of the DPRK. Pyongyang: UNICEF; 1998 [cited 1998 Nov]. Available from: http://www.pwdigby.co.uk/pdf/Report_on_the_DPRK_Nutrition_Assessment_1998.pdf.
4. DPRK Central Bureau of Statistics. Report of the Second Multiple Indicator Cluster Survey 2000, DPRK. Pyongyang: DPRK Central Bureau of Statistics; 2000 [cited 2000 Oct]. Available from: <http://www.childinfo.org/files/dprk.pdf>.
5. DPRK Central Bureau of Statistics. Report of the DPRK Nutrition Assessment. Pyongyang: UNICEF; 2002 [cited 2003 Feb]. Available from: http://www.unicef.org/dprk/nutrition_assessment.pdf.
6. DPRK Central Bureau of Statistics. DPRK 2004 Nutrition Assessment Report of Survey Results. Pyongyang: DPRK Central Bureau of Statistics, Institute of Child Nutrition; 2004 [cited 2005 Feb]. Available from: http://www.unicef.org/dprk/dprk_national_nutrition_assessment_2004_final_report_07_03_05.pdf.
7. UNICEF CBS. DPRK Multiple Indicator Cluster Survey 2009. Pyongyang: DPRK Central Bureau of Statistics; 2009 [cited 2010 Dec]. Available from: http://planipolis.iiep.unesco.org/upload/Korea%20DPR/Korea_DPR_MICS_2009.pdf.
8. UNICEF. DPRK at a glance 2013 (Feb). Pyongyang: UNICEF; 2013 [cited 2013 Feb]. Available from: http://www.unicef.org/dprk/DPRK_at_a_glance_April_2013.pdf.