Published in final edited form as:

J Cardiovasc Electrophysiol. 2014 November ; 25(11): 1257–1258. doi:10.1111/jce.12532.

Cross talk between renal and cardiac autonomic nerves: Is this how renal denervation works?

Wei-Chung Tsai, M.D.^{1,2} and Peng-Sheng Chen, M.D.¹

¹The Krannert Institute of Cardiology and Division of Cardiology, Department of Medicine, Indiana University School of Medicine, Indianapolis, IN

²Division of Cardiology, Department of Internal Medicine, Kaohsiung Medical University Hospital, Kaohsiung Medical University, Kaohsiung, Taiwan

Keywords

stellate ganglion; renal nerve; sympathetic system; renal artery ablation; autonomic nervous system

Editorial Comment

Surgical renal sympathetic denervation (RSD) has been used to control resistant hypertension.¹ With the development of catheter based RSD, this old technique has been resuscitated for the treatment of hypertension. While initial clinical studies showed effective hypertension control,² a recent sham-controlled randomized clinical trial failed to confirm its antihypertensive efficacy.³ These negative studies notwithstanding, RSD has been actively pursued as a tool to control various types of cardiac arrhythmias.^{4, 5} The mechanisms by which RSD helps arrhythmia control remain poorly understood. In the present issue of the *Journal*, Huang et al⁶ reported in anesthetized dogs that left renal nerve stimulation facilitates ischemia induced ventricular arrhythmia by increasing the function of the left stellate ganglion (LSG). LSG ablation attenuated these arrhythmias. This report suggests that there is a direct interaction between the left renal sympathetic nerve and the LSG. Is it possible that the cross talk between these two nerve structures underlies the antiarrhythmic mechanisms of RSD?

Left Stellate Ganglion and Cardiac Arrhythmia

LSG nerve activity is a direct trigger of cardiac arrhythmias.^{7–9} Left cardiac sympathetic denervation (LCSD) is effective in managing patients with long-QT syndrome and catecholaminergic polymorphic ventricular tachycardia.^{10–12} In addition to genetic arrhythmias, both LCSD and bilateral cardiac sympathetic denervation may be effective in controlling ventricular tachycardia (VT) storm or refractory VT.¹³ These studies indicate that LSG is an important arrhythmogenic structure.

Correspondence: Peng-Sheng Chen, 1800 N. Capitol Ave, Suite E475, Indianapolis, IN, 46202-1228, Telephone number: (317) 274-0909, Fax: (317) 962-0588, chenpp@iu.edu.

Dr. Chen has equity interest in Arrhythmotech, LLC. Medtronic, St Jude and Cyberonics Inc. donated research equipment to Dr. Chen's laboratory.

RSD and Cardiac Sympathetic Nerve Activity

Various neuromodulation methods have been tested for cardiac arrhythmia control.¹⁴ Among them, low-level tragus stimulation¹⁵ reduces ganglionated plexi neural activity while low level vagal nerve stimulation inhibits both ganglionated plexi and stellate ganglion nerve activities.^{16, 17} Stimulating the cardiac sympathetic nerves may cause reflex renal sympathetic nerve activity in dogs.¹⁸ Renal nerve stimulation may significantly alter the renal and cardiovascular function and contribute to the initiation, development and maintenance of hypertension.¹⁹ Li et al²⁰ demonstrated that RSD reduces BP by partially inhibiting sympathetic drive and systemic sympathetic outflow in hypertensive canine models. In obese spontaneously hypertensive rats, renal sympathetic activation increases while RSD decreases BP, renal injury and cardiac remodeling.²¹ In SYMPPLICITY-HTN1 trial, RSD reduced renal noradrenaline spillover 47% in patients with refractory hypertension.² Muscle sympathetic nerve activity in patients with resistant hypertension can be substantially and rapidly reduced by RSD.²² Huang et al⁶ extended these observations by demonstrating that left renal nerve stimulation facilitates ischemia induced ventricular arrhythmia by increasing the function of the left stellate ganglion (LSG). They also demonstrated increased nerve growth factor (NGF) in the LSG after left renal nerve stimulation. While the authors did not perform immunohistochemical studies of the heart, a previous study²³ showed that NGF infusion to the LSG can cause cardiac nerve sprouting and sympathetic hyperinnervation. An implication of the present study is that hyperactive renal sympathetic nerves may induce LSG remodeling and increase sympathetic tone, which facilitates the development of arrhythmias.^{23, 24} Therefore, ablation of either the renal sympathetic nerve or the LSG may be effective in cardiac arrhythmia control. Consistent with this hypothesis, we²⁵ have recently found that in ambulatory dogs that RSD is associated with reduced LSG nerve activity and atrial tachycardia episodes.

In summary, LSG plays an important role in cardiac arrhythmogenesis, and that modulating the stellate ganglion nerve activity by RSD may be effective in arrhythmia control. Huang et al⁶ provided important information on the interaction between renal sympathetic nerve and LSG. It is therefore possible that the RSD achieves arrhythmia control through the reduction of the stellate ganglion activity and cardiac nerve sprouting. More studies in ambulatory animals and in humans are needed to test the latter hypotheses.

Acknowledgments

Funding source: NIH Grants P01 HL78931, R01 HL71140, R41 HL124741, a Medtronic-Zipes Endowment and the Indiana University Health-Indiana University School of Medicine Strategic Research Initiative.

References

1. Page IH, Heuer GJ. The effect of renal denervation on the level of arterial blood pressure and renal function in essential hypertension. *J Clin Invest.* 1935; 14:27–30. [PubMed: 16694275]
2. Krum H, Schlaich M, Whitbourn R, Sobotka PA, Sadowski J, Bartus K, Kapelak B, Walton A, Sievert H, Thambar S, Abraham WT, Esler M. Catheter-based renal sympathetic denervation for resistant hypertension: a multicentre safety and proof-of-principle cohort study. *Lancet.* 2009; 373:1275–1281. [PubMed: 19332353]
3. Bhatt DL, Kandzari DE, O'Neill WW, D'Agostino R, Flack JM, Katzen BT, Leon MB, Liu M, Mauri L, Negoita M, Cohen SA, Oparil S, Rocha-Singh K, Townsend RR, Bakris GL. Investigators

- SH. A controlled trial of renal denervation for resistant hypertension. *N Engl J Med*. 2014; 370:1393–1401. [PubMed: 24678939]
4. Pokushalov E, Romanov A, Katritsis DG, Artyomenko S, Bayramova S, Losik D, Baranova V, Karaskov A, Steinberg JS. Renal Denervation for Improving Outcomes of Catheter Ablation in Patients With Atrial Fibrillation and Hypertension: Early Experience. *Heart Rhythm*. 2014
 5. Remo BF, Preminger M, Bradfield J, Mittal S, Boyle N, Gupta A, Shivkumar K, Steinberg JS, Dickfeld T. Safety and efficacy of renal denervation as a novel treatment of ventricular tachycardia storm in patients with cardiomyopathy. *Heart Rhythm*. 2014; 11:541–546. [PubMed: 24389229]
 6. Huang B, Yu L, Scherlag BJ, Wang S, He BO, Yang K, Liao KAI, Lu Z, He W, Zhang L, Po SS, Jiang H. Left Renal Nerves Stimulation Facilitates Ischemia-induced Ventricular Arrhythmia by Increasing Nerve Activity of Left Stellate Ganglion. *J Cardiovasc Electrophysiol*. 2014 (editor to fill in).
 7. Zhou S, Jung BC, Tan AY, Trang VQ, Gholmieh G, Han SW, Lin SF, Fishbein MC, Chen PS, Chen LS. Spontaneous stellate ganglion nerve activity and ventricular arrhythmia in a canine model of sudden death. *Heart Rhythm*. 2008; 5:131–139. [PubMed: 18055272]
 8. Ogawa M, Zhou S, Tan AY, Song J, Gholmieh G, Fishbein MCLH, Siegel RJ, Karagueuzian HS, Chen LS, Lin SF, Chen PS. Left stellate ganglion and vagal nerve activity and cardiac arrhythmias in ambulatory dogs with pacing-induced congestive heart failure. *J Am Coll Cardiol*. 2007; 50:335–343. [PubMed: 17659201]
 9. Tan AY, Zhou S, Ogawa M, Song J, Chu M, Li H, Fishbein MC, Lin SF, Chen LS, Chen PS. Neural mechanisms of paroxysmal atrial fibrillation and paroxysmal atrial tachycardia in ambulatory canines. *Circulation*. 2008; 118:916–925. [PubMed: 18697820]
 10. Wilde AA, Bhuiyan ZA, Crotti L, Facchini M, De Ferrari GM, Paul T, Ferrandi C, Koolbergen DR, Odero A, Schwartz PJ. Left cardiac sympathetic denervation for catecholaminergic polymorphic ventricular tachycardia. *N Engl J Med*. 2008; 358:2024–2029. [PubMed: 18463378]
 11. Schwartz PJ, Priori SG, Cerrone M, Spazzolini C, Odero A, Napolitano C, Bloise R, De Ferrari GM, Klersy C, Moss AJ, Zareba W, Robinson JL, Hall WJ, Brink PA, Toivonen L, Epstein AE, Li C, Hu D. Left cardiac sympathetic denervation in the management of high-risk patients affected by the long-QT syndrome. *Circulation*. 2004; 109:1826–1833. [PubMed: 15051644]
 12. Moss AJ, McDonald J. Unilateral cervicothoracic sympathetic ganglionectomy for the treatment of long QT interval syndrome. *N Engl J Med*. 1971; 285:903–904. [PubMed: 5112730]
 13. Vaseghi M, Gima J, Kanaan C, Ajijola OA, Marmureanu A, Mahajan A, Shivkumar K. Cardiac sympathetic denervation in patients with refractory ventricular arrhythmias or electrical storm: Intermediate and long-term follow-up. *Heart Rhythm*. 2014; 11:360–366. [PubMed: 24291775]
 14. Chen PS, Chen LS, Fishbein MC, Lin SF, Nattel S. Role of the autonomic nervous system in atrial fibrillation: pathophysiology and therapy. *Circ Res*. 2014; 114:1500–1515. [PubMed: 24763467]
 15. Yu L, Scherlag BJ, Li S, Fan Y, Dyer J, Male S, Varma V, Sha Y, Stavrakis S, Po SS. Low-level transcutaneous electrical stimulation of the auricular branch of the vagus nerve: a noninvasive approach to treat the initial phase of atrial fibrillation. *Heart Rhythm*. 2013; 10:428–435. [PubMed: 23183191]
 16. Yu L, Scherlag BJ, Li S, Sheng X, Lu Z, Nakagawa H, Zhang Y, Jackman WM, Lazzara R, Jiang H, Po SS. Low-level vagosympathetic nerve stimulation inhibits atrial fibrillation inducibility: direct evidence by neural recordings from intrinsic cardiac ganglia. *J Cardiovasc Electrophysiol*. 2010; 22:455–463. [PubMed: 20946225]
 17. Shen MJ, Shinohara T, Park HW, Frick K, Ice DS, Choi EK, Han S, Maruyama M, Sharma R, Shen C, Fishbein MC, Chen LS, Lopshire JC, Zipes DP, Lin SF, Chen PS. Continuous low-level vagus nerve stimulation reduces stellate ganglion nerve activity and paroxysmal atrial tachyarrhythmias in ambulatory canines. *Circulation*. 2011; 123:2204–2212. [PubMed: 21555706]
 18. Ma R, Zucker IH, Wang W. Central gain of the cardiac sympathetic afferent reflex in dogs with heart failure. *Am J Physiol*. 1997; 273:H2664–2671. [PubMed: 9435602]
 19. DiBona GF, Kopp UC. Neural control of renal function. *Physiol Rev*. 1997; 77:75–197. [PubMed: 9016301]

20. Li H, Yu H, Zeng C, Fang Y, He D, Zhang X, Wen C, Yang C. Renal denervation using catheter-based radiofrequency ablation with temperature control: renovascular safety profile and underlying mechanisms in a hypertensive canine model. *Clin Exp Hypertens*. 2014; 0:1–5.
21. Linz D, Hohl M, Schütze J, Mahfoud F, Speer T, Linz B, Hübschle T, Juretschke H-P, Dechend R, Geisel J, Rütten H, Böhm M. Progression of Kidney Injury and Cardiac Remodeling in Obese Spontaneously Hypertensive Rats: The Role of Renal Sympathetic Innervation. *Am J Hypertens*. 2014
22. Hering D, Lambert EA, Marusic P, Walton AS, Krum H, Lambert GW, Esler MD, Schlaich MP. Substantial reduction in single sympathetic nerve firing after renal denervation in patients with resistant hypertension. *Hypertension*. 2013; 61:457–464. [PubMed: 23172929]
23. Cao JM, Chen LS, KenKnight BH, Ohara T, Lee MH, Tsai J, Lai WW, Karagueuzian HS, Wolf PL, Fishbein MC, Chen PS. Nerve sprouting and sudden cardiac death. *Circ Res*. 2000; 86:816–821. [PubMed: 10764417]
24. Miyauchi Y, Zhou S, Miyauchi M, Omichi C, Okuyama Y, Hamabe A, Hayashi H, Mandel WJ, Fishbein MC, Chen LS, Chen PS, Karagueuzian HS. Induction of atrial sympathetic nerve sprouting and increased vulnerability to atrial fibrillation by chronic left ventricular myocardial infarction. *Circulation*. 2001; 104:II-77.
25. Tsai W-C, Chan Y-H, Chinda K, Lai W-T, Lin S-F, Chen P-S. Renal sympathetic denervation decreases the incidence of atrial tachycardia and improves baroreflex sensitivity in ambulatory dogs. *HeartRhythm*. 2014:S236. (abstract).