CORRESPONDENCE

Case Management Program for Patients With Chronic Heart Failure: Effectiveness in Terms of Mortality, Hospital Admissions and Costs

by Dipl.-Oek./Medizin (FH) Verena Hendricks M.Sc.,
Dipl.-Oek./Medizin (FH) Simone Schmidt, Prof. Dr. med. Achim Vogt,
Dr. med. Detlef Gysan, Dipl.-Oek./Medizin (FH) Volker Latz,
Dr. med. Ines Schwang, Prof. Dr. med. Reinhard Griebenow,
Prof. Dr. med. Dipl.-Kfm. (FH) Rainer Riedel in issue 15/2014

Cross-Sector Case Management

The authors demonstrate that a case management program (CMP) for patients with heart failure can result in a significant decrease in hospital admissions and readmissions as well as hospital costs. Here improvements in the outpatient sector are crucial (guideline-based pharmacotherapy, early specialist integration, regular follow-ups) (1).

Disease management programs (DMPs) are systematic, centrally organized, evidence-based treatment programs for people living with chronic illness (2). In contrast, case management models (CMs) focus on achieving the best possible treatment outcome for individual patients (2). Consequently, this paper is about a DMP, not about a CM.

Given the “close cooperation between the various service providers” (1), the hospital’s role in the reduction of the readmission rates remains elusive. The German SHI Health Care Delivery Structure Act of 2011 aims at achieving compulsory improvements of the hospitals’ discharge management. This is important for surgical fields, such as vascular surgery. For example, multimorbid patients with diabetes mellitus and peripheral artery occlusive disease (PAOD) receiving bypass grafts to avoid major amputations have readmission rates after discharge from hospital of 30% after seven weeks (3). Cross-sector case management can significantly reduce the readmission rate among these patients as well as the rate of case consolidations, without extending the length of the hospital stay (3). Here the improvement of inter-sector communication and collaboration is vital to coordinate an outpatient care network before the patient is discharged from the hospital. Consequently, the activities of the hospital providing the care are critical, while the study by Hendricks (1) assumes that the outpatient sector is responsible for the main effect.

Our CM has resulted in cost savings for hospitals. We do not know whether it creates additional costs for the outpatient sector. In the above study (1), it is also conceivable that the lowered hospital costs may be contrasted by increased outpatient costs, for example, as the result of more frequent visits to the doctor.

DOI: 10.3238/arztebl.2014.0680a

REFERENCES

- Hendricks V, Schmidt S, Vogt A, Gysan D, et al.: Case management program for patients with chronic heart failure—effectiveness in terms of mortality, hospital admissions and costs. *Dtsch Arztebl Int* 2014; 111: 264–70.
- Case Management Society of America: „Definition of Case Management“. www.cmsa.org/Home/CMSA/WhatIsACaseManager/tabid/224/Default.aspx (last accessed on 22 July 2014)

- Rümenapf G, Geiger S, Schneider B, et al.: Readmissions of patients with diabetes and foot ulcers after infra-popliteal bypass surgery: attacking the problem by an integrated case management model. *Vasa (Eur J Vasc Med)* 2013; 42: 56–67.

Prof. Dr. med. Gerhard Rümenapf

Diakonissen-Stiftungs-Krankenhaus, Speyer
gerhard.ruemenapf@diakonissen.de

Dr. rer. nat. Johannes Georg Boettrich

Dr. rer. nat. Norbert Nagel
Melsungen

Dr. med. Stephan Morbach

Marienkrankenhaus Soest gGmbH, Soest

Conflict of interest statement

Prof. Rümenapf has received lecture fees from B. Braun Melsungen AG.

Dr. Boettrich and Dr. Nagel are employees of B. Braun Melsungen AG.

Dr. Morbach has received consultancy fees (Advisory Board) from Novo Nordisk, Urgo Pharma and Olympus.

Methodological Shortcomings

This investigation is not a field study as the data were not collected to answer a research question.

In studies aimed at comparing the efficacy of different treatments, the intervention and control groups must be comparable (1). If in the intervention group a ventricular dysfunction and cardiologist contact are required, while this is not the case in the control group, it casts doubt on the comparability of the two groups. In this study, propensity score matching was used in an attempt to establish comparability. When using this method, all variables should be included which have an impact on the event or the allocation to a certain group, if possible, not only age, sex and NYHA stage. For this, stepwise regression is not suitable (1). Important parameters related to the severity of the disease, such as the number of hospital admissions prior to study commencement, were neither considered nor reported at all.

By excluding patients who participated in the program for less than 6 months, an immeasurable time bias is introduced (2). Patients who die during this period are not considered in the intervention group, while they are in the control group. CorBene was developed in 2005; thus it can be assumed that many patients were only registered in the program in the following years. By starting the follow-up on July 1, 2005, regardless of whether or not the patient was already enrolled in the program at that time, further adds to this bias.

The method used to identify heart failure-related physician contacts remains unclear, especially since in an outpatient environment diagnoses are billed on a case basis. Two primary care physician and cardiologist contacts annually appear far too few, considering that each person insured has an average of 18 physician contacts per year (10 of these visits to primary care physicians/internists) and these numbers can increase in the 70+ age group to more than 30 contacts annually (3).

Thus we have arrived at the conclusion that the study does not constitute an appropriate basis for political decisions due to its methodological shortcomings.

DOI: 10.3238/arztebl.2014.0680b

REFERENCES

1. Johnson ML, Crown W, Martin BC, Dormuth CR, Siebert U: Good research practices for comparative effectiveness research: analytic methods to improve causal inference from nonrandomized studies of treatment effects using secondary data sources: the ISPOR Good Research Practices for Retrospective Database Analysis Task Force Report-Part III. *Value Health* 2009; 12: 1062–73.
2. Suissa S: Immeasurable time bias in observational studies of drug effects on mortality. *Am J Epidemiol* 2008; 168: 329–35.
3. Grobe TG, Döring H, Schwartz FW: Barmer GEK Arztreport. Sankt Augustin: Asgard-Verlag 2010.
4. Hendricks V, Schmidt S, Vogt A, et al.: Case management program for patients with chronic heart failure—effectiveness in terms of mortality, hospital admissions and costs. *Dtsch Arztebl Int* 2014; 111: 264–70.

PD Dr. PH Falk Hoffmann

Universität Bremen, ZeS, Abteilung Gesundheitsökonomie,
Gesundheitspolitik und Versorgungsforschung, Bremen
hoffmann@zes.uni-bremen.de

Dr. rer. medic. Tania Schink

Leibniz-Institut für Präventionsforschung und Epidemiologie – BIPS GmbH,
Abteilung Klinische Epidemiologie, Bremen

Conflict of interest statement

The authors declare that no conflict of interest exists.

In Reply:

The research colleagues Hoffmann and Schink have highlighted potential limitations in outpatient healthcare research. This, in principle, is to be welcomed. Unfortunately, there are limitations to the healthcare research-based approach to patients and treatment outcomes in routine care. Nobody thought about establishing a platform prior to the introduction of a disease-specific healthcare program that would allow to generate high-quality scientific evidence for the evaluation of costs and health benefits. One of the reasons for this is that under everyday conditions it is rather difficult to get a primary care physician or a specialist interested in collecting relevant patient and treatment data for evaluation and research purposes. And it is even more difficult to collect matching data of a control group for comparison. Therefore, we used propensity score matching for our study to be able to make use of routine data of the statutory health insurer involved (1).

Due to the limited amount of data available, the approach chosen for our study has qualitative limitations with regard to the theoretical scientific requirements for data quality in healthcare research. As the insurer was not in the position to provide additional data on comorbidities or sociodemographic data for our study, only the variables “age, sex and severity of the condition (NYHA stage)” were used for the matching. The authors are fundamentally aware of the limitations of this approach with regard to group heterogeneity (selection bias) and have discussed these in their paper. The question, however, arises to what extent we have to be prepared in the current discussion about health technology assessment (HTA)-based data and their limitations to use the available routine data as a base for early forward-looking statements. This approach may help us to further improve the quality of patient care, especially in the light of limited resources.

From their perspective, the readers may be concerned about a potential immeasurable time bias – however, such considerations would not meet the authors’ requirements. According to the available data from the statutory health insurer, no patients enrolled in the healthcare program died during the first six months. Furthermore, there is the concern regarding the “reported physician contacts”; here an evaluation method has to define a standard which can be used to integrate, for example, physician contacts as a study criterion. In this study, the criterion “diagnosis-related physician contacts” was defined.

Any study incorporating an analysis of statutory health insurance routine data, which were primarily collected for billing purposes, has certain limitations (2–4). Especially with regard to the limitations in the use of HTA as well as prospective randomized controlled trials (RCTs) for the evaluation of data derived from routine care of multimorbid patients, there is a pressing need to evaluate and publish healthcare research data (5). It would be a mistake to not make use of the data collected so far and the insights derived from these on how to optimize the current healthcare situation because of methodological limitations, which can unfortunately be found in most health economic evaluations of the treatment programs for multimorbid patients established to date. Rather, the results from healthcare routine studies should stimulate healthcare decision makers to generate valid and reliable data that can be used to create highest quality scientific evidence and to take the HTA perspective into account prior to the implementation of healthcare programs.

DOI: 10.3238/arztebl.2014.0681

REFERENCES

1. Hendricks V, Schmidt S, Vogt A, et al.: Case management program for patients with chronic heart failure—effectiveness in terms of mortality, hospital admissions and costs. *Dtsch Arztebl Int* 2014; 111: 264–70.
2. Stock S, Drabik A, Büscher G, et al.: German diabetes management programs improve quality of care and curb costs. *Health Aff (Millwood)* 2010; 29: 2197–205.
3. Miksch A, Laux G, Ose D, et al.: Is there a survival benefit within a German primary care-based disease management program? *Am J Manag Care* 2010; 16: 49–54.
4. Lindner R, Ahrens S, Köppel D, Heilmann T, Verheyen F: The benefit and efficiency of the disease management program for type 2 diabetes. *Dtsch Arztebl Int* 2011; 108: 155–62.
5. Gartlehner G, West SL, Mansfield AJ, et al.: Clinical heterogeneity in systematic reviews and health technology assessments: synthesis of guidance documents and the literature. *Int J Technol Assess Health Care* 2012; 28: 36–43.

On behalf of the authors

Prof. Dr. med. Dipl.-Kfm. (FH) Rainer Riedel

Verena Hendricks, M.Sc.

Institut für Medizin-Ökonomie & Medizin

Versorgungsforschung Rheinische Fachhochschule Köln gGmbH

University of Applied Sciences, Köln

riedel@rfh-koeln.de

Conflict of interest statement

The data collection and analysis expenses of the Institute for Health Economics and Healthcare Research (Institut für Medizin-Ökonomie & Medizinische Versorgungsforschung) at the University of Applied Sciences, Cologne, which was entrusted with the scientific support of the CorBene Program, were reimbursed by the participating company health insurers.

The authors declare that no conflict of interest exists.