In brief

Survey finds longer waits for cancer treatment: Cancer patients now wait two weeks longer for treatment than they did six years ago, in three out of four NHS centres. This is a finding of an audit by the Royal College of Radiologists of all radiotherapy centres in the United Kingdom. The report attributes the rise to an increase in the number of patients being referred for treatment. It will be published next month on the college's website: www.rcr.ac.uk

Dutch institute warns of polio threat: The Dutch Institute of Public Health and the Environment is warning doctors to be alert to the threat of poliomyelitis among African population groups who may have had contact with the current outbreak in northern Nigeria. The institute has urged Dutch health services with large West African populations to maintain a 95% level of childhood vaccinations (www.rivm.nl)

Hookah pipes to focus in antismoking campaign: Israel's health ministry and the Israel Cancer Association are launching a publicity campaign aimed at discouraging youngsters from smoking hookahs (tobacco water pipes), traditionally used by elderly Arab men but now "fashionable" among Jewish teenagers. A survey showed some 40% to 57% of Jewish pupils from sixth to 10th grades had tried a water pipe, and many used one regularly.

Norway launches smoking ban: Norway is the second country in the world to have introduced a smoking ban in all indoor places. The health minister, Dagfinn Hoybraten, said that the blanket ban was designed to protect people working in bars and restaurants.

Cancer Bank is launched: The Wales Cancer Bank was launched this week by the Welsh Assembly health minister, Jane Hutt. Patients with possible or confirmed cancer will be asked for their permission for a portion of their tissue and a blood sample to be stored for future scientific studies.

Thyroid cancer has increased 12-fold in women since Chernobyl

Roger Dobson *Abergavenny*

Rates of thyroid cancer among women in Belarus have increased up to 12-fold since the explosion at the Chernobyl nuclear power plant, near Kiev, Ukraine, in April 1986.

Among young women aged under 14 in higher risk areas of the republic the rates have increased almost 30-fold since the disaster, new research has shown.

"This study documents dramatic increases in the incidence of thyroid cancer among both higher exposure and lower exposure areas within the republic of Belarus and among all age groups studied," says an "advance access" report of the study published online on 27 May in the *International Journal of Epidemiology* (www.ije.oupjournals.org (doi: 10.1093/ije/dyh201)).

It adds: "The magni-

tude of increases observed is remarkable given the relatively limited time interval since Chernobyl and argues for continued surveillance in Belarus as well as other affected areas."

The explosion at Chernobyl resulted in the release of substantial amounts of radioactive materials over western regions of what was then the Soviet

Union, including radioisotopes of iodine, caesium, strontium, and plutonium. The most significant contamination was in Belarus and Ukraine, as well as the western region of the Russian Federation.

The report says that although previous studies of the incidence of thyroid cancer in Belarus have shown an increase since the Chernobyl explosion, the size of the increase is not well quantified.

The authors, from the Roswell Park Cancer Institute, New York, and the Institute of Oncology, Minsk, used data from the Belarus national cancer registry and trends in average annual incidence of thyroid cancer.

The data show big increases since 1986. Between 1980 and 1986, for example, there were 0.15 cases per 100 000 diagnosed in girls aged under 14 in high risk areas. By the period 1997-2001, the rate had gone up to 43.84. Among males of the same age, the rate went up from 0.08 to 18.81. □

Early localised prostate cancer may accelerate after 15 years

Janice Hopkins Tanne *New York*

Watchful waiting is not a good idea for some patients with early localised prostate cancer because local progression and aggressive metastatic disease may develop after 15 years. This finding comes from a 20 year study—one of the largest and longest yet—from a Swedish group and published in *JAMA* (2004;291:2713-9).

"We had not anticipated this increased risk after 15 years," the lead author, Dr Jan-Erik Johansson of Örebro University Hospital, told the *BMJ*.

The study followed 223 patients with early stage, localised prostate cancer. They received no initial treatment if the cancer was localised to the prostate as

judged by digital rectal examination and there were no distant metastases. If the cancer progressed to symptomatic disease, patients were treated with oestrogens or orchidectomy. Local progression without generalised disease may cause substantial suffering, the authors wrote. About 70 patients received hormonal treatment, Dr Johansson told the *BMJ*.

During the first 15 years, most cancers followed an indolent course: 25 (11%) of patients died of prostate cancer. Ten more died during follow up.

Follow up from 15 years to 20 years showed a "substantial decrease in cumulative progression-free survival (from 45% to 36%), survival without metastases (from 76.9% to 51.2%), and prostate-cancer-specific survival (from 78.7% to 54.4%)." Mortality from prostate cancer increased from 15 per 1000 (95% confidence interval 10 to 21 per 1000) person years during the first 15 years to 44 per 1000 (22 to 88 per 1000) person years beyond 15 years of follow up ($P=0.01$).

Dr Johansson and colleagues followed the group for a mean of 21 years, beginning in 1977. About half the patients were diagnosed as a result of cancer being found during resection for supposedly benign prostatic enlargement. The mean age at diagnosis was 72 (range 41 to 91) years.

All 223 patients were followed until death or September 2001; none were lost to follow up. Patients who agreed had a new fine needle biopsy every other year during the first six years of follow up.

During the study period, 89 patients (40%) had progression of disease, with 39 of them (17% of the entire cohort) developing generalised disease; 203 patients (91%) died—35 (16%) of prostate cancer. Patients younger than 70 and those with poorly differentiated cancer were more likely to die of cancer than older patients.

The unexpected finding was that disease and death from prostate cancer accelerated after 15 years, regardless of the stage at which the tumours were first diagnosed. □