party considered that optimal care would require new models of service provision, championed by clinicians and working in partnership with patients. It suggested that joint approaches should be developed at local level, including primary care trusts and local hospitals, to improve coordination of patient care.

Setting up "joint clinical directorates"—clinical directorates that spanned primary and secondary care—should be considered as a way of facilitating the development of the infrastructure and clinical governance requirements for services across different sectors.

Professional training should also be reviewed and ways of sharing information should also be developed. The report recommended that primary care trusts and acute trusts should develop local shared information pathways, including electronic transfer of pathology results.

Susan Mayor *London*

Clinicians, Services and Commissioning in Chronic Disease Management in the NHS can be obtained by phoning the Royal College of Physicians (tel 020 7935 1174) and asking for the publications office.

West Nile virus infections are occurring early in North America

The United States is seeing West Nile virus infections earlier than usual. The Centers for Disease Control and Prevention (CDC) had reported two cases in humans—one in Arizona and one in New Mexico—and avian, animal, or mosquito infections in 17 states by 1 June. Last year more than 260 people died from the disease in the United States.

ProMedmail.org, a service of the International Society for Infectious Diseases, said West Nile virus was now widespread throughout the United States and in nearby regions in Canada and Mexico. It would probably move south into Mexico.

Arizona and New Mexico have a border with Mexico, and by 27 May the Mexican authori-

ties had reported 18 seropositive birds and 287 seropositive horses. Health Canada had reported eight dead birds infected with West Nile virus in Ontario by 4 June.

The mosquito season in Arizona is usually August and September. ProMedmail reported that Dr Jonathan Weisbuch, head of the Maricopa County Department of Public Health, which includes the city of Phoenix, said "This is much earlier than we expected. It's going to be a longer season. We could see several hundred cases."

Janice Hopkins Tanne *New York*

Doctors demand national screening for chlamydia

Public health doctors have derided the slow progress towards a national screening programme for chlamydia as "disgraceful."

The BMA's annual conference of public health medicine called on the government to roll out screening across the United Kingdom as a matter of urgency.

Dr Howard Barnes, who proposed the motion, said the absence of a national programme had led to a high volume of uncoordinated testing.

"In a district near me in Yorkshire 14 000 people have been tested. That is out of a population of 200 000," he said.

Chlamydia is the United Kingdom's most common sexually transmitted infection. Nearly 70 000 cases of chlamydia were reported by clinics in 2001, up 10% on the previous 12 months. The infection, which is often asymptomatic in women, can cause pelvic inflammatory disease and infertility.

Dr Barnes attacked the government for setting up a series of pilot groups but failing to take the plunge and construct a formal screening programme. The initial sites are in Portsmouth and the Wirral, followed by two more waves of pilots in subsequent years. The schemes primarily target women aged 16 to 24, but men are also offered tests. "With screening we would see a reduction in pelvic inflammatory disease and infertility," he said.

Rebecca Coombes *London*

Smoking habits of ethnic groups lead to higher risk of heart disease

Lynn Eaton *London*

Smoking, one of the many increased risk factors for coronary heart disease, is much higher among Black Caribbean, Bangladeshi, and Irish men than in the general population for England, according to the British Heart Foundation's latest statistics on coronary heart disease. And Bangladeshi men are at the highest risk of all because of their smoking habit.

The proportion of Irish women smoking is also higher than the general female population, putting them at greater risk too.

The ethnic differences in smoking habits are of particular concern to the foundation, said Qaim Zaidi, its ethnic strategy coordinator.

"In particular, South Asians living in the UK have a significantly higher premature death rate from coronary heart disease than the rest of the population—and the gap is widening. The death rate is 46% higher than the average for men and 51% higher for women, and it is not falling as fast as it is in the rest of the population," he said.

"One of the main problems to overcome is the high numbers of smokers in the Bangladeshi community—42% of Bangladeshi men smoke, as opposed to 27% in the general male population," he added.

The report looks at a number of factors that affect the risk of

heart disease, including diet, physical activity, alcohol consumption, blood pressure, blood cholesterol, and diabetes.

"Bangladeshi and Pakistani communities also eat the fewest fruits and vegetables, and South Asians are less likely to participate in physical activity," said Mr Zaidi. "Consequently, they are more likely than the general population to have central obesity... placing an extra strain on the heart."

"The BHF [British Heart Foundation] is working hard to tackle this critical issue and has made progress in educating South Asian communities about lifestyle and heart disease in recent years," he said, adding that the charity had invested millions into medical research to benefit ethnic groups. "However, the message is not sinking in quickly enough," he said.

The report says the overall number of coronary heart deaths is down from nearly 121 000 in 2001 to just under 117 500 in 2002. But because fewer people are dying of the disease, many more are living with it. An estimated 2.68 million people currently live with the disease, 30 000 higher than last year. □

The British Heart Foundation's coronary heart disease statistics for 2004 are available on www.heartstats.org