Published in final edited form as:

Curr Opin Crit Care. 2014 December ; 20(6): 656–661. doi:10.1097/MCC.0000000000000147.

Economic implications of end-of-life care in the ICU

Nita Khandelwal, MD, MSc¹ and J. Randall Curtis, MD, MPH²

¹Department of Anesthesiology and Pain Medicine, Harborview Medical Center, University of Washington, Seattle, WA

²Division of Pulmonary and Critical Care Medicine, Harborview Medical Center, University of Washington, Seattle, WA

Abstract

Purpose of review—Advance care planning and palliative care interventions can improve the quality of end-of-life care by reducing unwanted high intensity care at the end of life. This may have important economic implications and may reduce financial burden of patients' families. We review the literature to examine the impact advance care planning and palliative care have on ICU utilization, specifically ICU admissions and ICU LOS, to provide insight into ways to reduce costs and financial burden of care while simultaneously improving quality of care.

Recent findings—We identified 3 studies assessing the impact of palliative care consultation on ICU admissions for patients with life-limiting illness; all 3 demonstrate reduced ICU admissions for patients receiving palliative care consultation. Among 16 studies evaluating ICU LOS as an outcome, 5 report no change and 11 report decrease in LOS for patients receiving advance care planning or palliative care. These studies are heterogeneous in design and target population; however, a trend towards reduced ICU utilization exists.

Summary—Advance care planning and palliative care can reduce ICU utilization at the end of life. The degree to which reducing ICU utilization decreases emotional and financial burden of end-of-life care for patients and families is unknown.

Keywords

economics; critical illness; palliative care; end-of-life care

Introduction

In the United States, 1 in 5 adults die during a hospitalization that includes a stay in the ICU and 25% of our healthcare costs are spent on the 6% of people who die each year (1-3). Our aging population, coupled with advances in management of acute and chronic illness, has led to a steadily increasing number of critical care beds and increasing costs associated with critical care (1, 4, 5). In 2005, in the United States, critical care costs were estimated to be \$82 billion, accounting for 13% of inpatient hospital costs (4, 5). The U.S. spends more

Corresponding author: Nita Khandelwal, MD, MSc, Harborview Medical Center, 325 9th Avenue, Box 359724, Seattle, WA 98104, Phone: 206.744.4371, Fax: 206.744.8090, khandel@uw.edu.

Conflicts of interest: The authors have no conflicts of interest to declare.

hospital resources on critical care medicine than any other country, as evidenced by its having among the highest ratios of ICU-bed-to-population (20 ICU beds per 100K) and ICU-to-hospital-bed (9 per 100 hospital beds) in the world (4).

According to Medicare claims data, ICU use in the last 30 days of life increased between 2000 and 2009 despite public opinion surveys reporting that most patients would prefer to die at home if diagnosed with a terminal illness (6, 7). This apparent inconsistency raises concern about the use of potentially unwanted intensive care at the end of life. Additionally, costly interventions to support critically ill patients may be ineffective and can cause significant discomfort. For example, endotracheal tubes, intravascular lines, feeding tubes and restraints may reduce mobility, ability to communicate, and autonomy and may cause pain (8). Importantly, this care at the end of life may not be consistent with patient preferences and values, and may place unnecessary emotional, physical, and financial burden on dying patients and their family members.

Interventions including early advance care planning, time-limited trials (9), and palliative care consultation, seek to ensure that care at the end of life remains patient-centered and is respectful of individual preferences and values (10). There is evidence to suggest that ensuring care for patients with a high risk of death remains patient-centered can lead to a reduction in intensity of care near the end of life; for example, there is emerging evidence that advance care planning early during an acute care hospitalization can reduce ICU admissions (11-13) and that time-limited trials and proactive early palliative care consultation in the ICU can reduce the LOS and intensity of treatment for patients who die in the ICU *(14).

The rationale for improving palliative care for patients with, or at risk for, critical illness can be considered in three areas: first, improving quality of care and thereby improving patient and family outcomes; second, reducing the overall costs of care by reducing unwanted high intensity care at the end of life; and three, reducing the financial burden of end-of-life care on patients' families by reducing unwanted intensity of care. We believe the first rationale above must be the primary rationale and that the latter two are only relevant if they are achieved through improving quality and ensuring that patients receive the care they would choose if truly informed about all their options. In this review, we will explore the financial implication of the latter two rationales. We will first review the current literature to better understand the impact advance care planning, primary palliative care (by ICU clinicians), and specialty palliative care consultation (by palliative care or ethics consultants) have on ICU utilization, specifically ICU admissions and ICU LOS, as this may have important economic implications and provide insight into ways to reduce costs and the financial burden of care while simultaneously improving the quality of care at the end of life and quality of dying. We will then explore the potential financial burden of high intensity care at the end of life on patients' families and the ethical implications of considering this burden.

In the Institute of Medicine report on *Approaching Death: Improving Care at the End of Life*, various strategies to reduce costs at the end of life were put forth, including: broader use of hospice care, consumer choice strategies, futility guidelines, and expanded use of advanced directives and advance care planning (15). Advance care planning and palliative

care interventions may have the potential to improve the quality of end-of-life care and quality of dying while simultaneously reducing costs by decreasing intensive therapy at the end of life to match care with the goals and preferences of informed patients. We will review the current literature evaluating the effect of these interventions on ICU utilization.

Studies evaluating ICU admissions

Few studies in the existing literature have reported on the impact of palliative care consultation and advance care planning on ICU admissions. We found 3 trials, all assessing palliative care consultation, that reported on the effect of palliative care consultation on ICU admissions. All demonstrated a significant reduction in patients who received palliative care consultation for patients with life-limiting illness when compared to similar patients who did not receive palliative care consultation. Among these 3 trials, one was a randomized trial (11) and 2 were retrospective observational studies (12, 13). The randomized trial by Gade et al included adult patients hospitalized with at least one life-limiting diagnosis and whose attending physician indicated they “would not be surprised if the patient died within 1 year” while the Penrod 2006 study included decedents only. Both of these studies reported similar relative risk reductions in ICU admission (50%, 51%, respectively). The 2nd observational study conducted by Penrod and colleagues included any veteran admitted to one of five Veterans Affairs hospitals with at least one advanced disease (12). The relative risk reduction for this group was 11%, suggesting that the degree of impact from palliative care consultation is heavily influenced by the selection of the patient population. Studies selecting those at highest risk of death showed a larger relative risk reduction.

Studies reporting on ICU LOS

We found 16 studies evaluating ICU LOS as an outcome for various types of advance care planning and palliative care interventions. Settings and targets for interventions were diverse, as were methods and types of trials. Five of these trials reported no change in LOS in patients receiving advance care planning or a palliative care intervention compared to usual care; 11 trials reported a reduction in LOS.

Studies reporting no change in LOS

Five trials reported no change in ICU LOS in patients receiving advance care planning or a palliative care intervention compared to usual care. Project ENABLE II, conducted by Bakitas et al, was designed to improve palliative care for patients with advanced cancer in the outpatient setting. This study randomized patients to a multi-component palliative care intervention consisting of 4 weekly educational sessions; ICU LOS did not differ when compared to the control group receiving usual care (16). Andereck et al investigated whether the proactive intervention of a clinical ethicist in patients with prolonged ICU lengths of stay (> 5 days) in a mixed medical/surgical ICU setting reduced ICU LOS (17). Patients in both the intervention and control arms had the same number of ICU days. In a multi-center cluster RCT conducted by Curtis et al, a quality improvement intervention targeted at hospitals and clinicians to integrate palliative care in the ICU also did not result in a significant decrease in LOS (18). Shelton et al evaluated the effect of adding a full-time family support coordinator to a surgical ICU team in a pre-post study design; no differences

in LOS were observed in the pre versus post-implementation period (19). Lastly, Daly et al enrolled patients from 5 different ICUs and evaluated the effectiveness of an intensive communication strategy involving proactive multidisciplinary family conferences using a pre-post design; there were no significant differences in ICU LOS (20).

Studies reporting a reduction in LOS

Eleven studies reported a reduction in ICU LOS for patients who received an advance care planning or palliative care intervention (either primary palliative care by ICU clinicians or specialty consultation inclusive of palliative care and ethics consultation). One of these studies involved patient-centered interventions taking place outside of the ICU setting, enrolling medical inpatients age 80 and older for an advance care planning intervention (21). In this randomized trial, Detering and colleagues assessed the impact of advance care planning on end-of-life care in elderly hospitalized patients (21). They randomized eligible patients admitted under internal medicine, cardiology or respiratory medicine in a large university hospital. Upon request from the authors, we were able to obtain ICU LOS data as this outcome was not reported in the original manuscript; mean LOS was 10.9 days in the control arm and 5.3 days in the intervention arm (21).

The remaining 10 studies evaluated interventions that took place in the ICU setting. Two separate randomized trials conducted by Schneiderman et al (22, 23) examined the effect of routine ethics consultation, enrolling patients in whom value-related treatment conflicts arose. Both of these studies of routine ethics consultation in the ICU (one single-center study and one multicenter study) demonstrated significant reductions in ICU LOS for decedents in the intervention group as compared to usual care. Neither trial found a difference in ICU LOS for patients who survived to hospital discharge.

Eight of the studies reporting a decrease in ICU LOS associated with a palliative care intervention were non-randomized studies using historical or concurrent controls. Ahrens et al evaluated the impact of a communication team consisting of a physician and clinical nurse specialist; patients in the intervention group had shorter LOS compared to the control group receiving usual care (24). Campbell and Guzman's study of routine palliative care consults for patients with global cerebral ischemia after CPR using historical controls also found a decrease in LOS. In this same study, patients with multi-system organ failure did not spend a significantly longer time in the ICU when compared to historical controls receiving usual care; however, when assessed from the onset of multiple-system organ failure, there was a reduction in days in the ICU prior to death (25). The following year, Campbell and Guzman published another study of patients with advanced dementia using historical controls and found that proactive case finding facilitated by an inpatient palliative care service led to a significant reduction in ICU LOS (26). Using non-randomized controls, Dowdy et al reported a reduction in LOS when the ethics service intervened proactively after patients received >96 hours of continuous mechanical ventilation (27). In a single-center study Curtis et al evaluated the impact of a quality improvement intervention targeted to ICU personnel to improve palliative care in the ICU; median ICU LOS was shorter in the post-implementation period (28). This difference was not observed in the follow-up multicenter cluster RCT (18). Mosenthal et al evaluated the impact of integrating a structured palliative

care intervention, consisting of assessment of patient prognosis and preferences, an interdisciplinary family meeting, and family bereavement support, into standard care in a trauma ICU. Among decedents, median LOS also decreased in the post-implementation period (29).

Lilly et al evaluated the impact of a multidisciplinary family meeting held within 72 hours of admission to the medical ICU; they found a reduction in median LOS in the post-implementation phase when compared to the baseline period (30). However, the multicenter study using this same intervention lead by Daly did not show reduction in ICU LOS (20). Norton and colleagues evaluated the impact of proactive palliative care consultation for patients admitted to the medical ICU and identified to be at high risk of death. This study, using a pre-post design, found a significant reduction in ICU LOS when compared to usual care (31).

Summary of studies reporting on ICU utilization

All 3 studies reporting on ICU admissions involved an intervention targeted directly at the patient or provider, rather than on a system level. All 3 demonstrated a reduction in ICU admissions in patients receiving a palliative care consultation.

While the results for studies reporting on ICU LOS were less consistent, a trend towards a reduction in the LOS for patients receiving an advance care planning or palliative care intervention seems apparent. Additionally, interventions that were targeted directly at providers and patients at the highest risk of death appeared to be more effective than interventions targeted at the ICU or hospital level.

Financial burden of high intensity care at the end of life

Chronic and terminal illnesses have serious financial consequences for patients and their families (15). These financial consequences stem from both out-of-pocket medical expenses and from reduced patient or family income that results from decreased working hours or job loss because of illness or demands of caring for an ill family member (15). Seriously ill patients and their families have reported several end-of-life priorities, one of which is having health care costs covered to avoid placing a financial burden on loved ones (32, 33). Despite this finding that minimizing the financial burden of end-of-life care is a top priority for dying patients, little is known about the costs of dying, including what aspects are most concerning to which types of patients, how informed patients are about the costs of high intensity care at the end of life and, importantly, if there are ways to improve the quality of dying by addressing and reducing this financial burden. Recent research has demonstrated that insurance status is an important mediator of the relationship between ICU costs and family-rated quality of dying, suggesting that financing of care influences how families perceive quality of dying and satisfaction with care (34). Specifically, this study found that for patients who were uninsured or under-insured, high costs of care at the end of life were associated with higher ratings of quality of dying while there was no association for patients with Medicare or private insurance *(34).

Marshall and colleagues reported that spending in the last year of life for decedents represents a substantial portion of liquid wealth (35). While the majority of end-of-life care costs are financed by public programs, especially Medicare, there is strong evidence that out-of-pocket expenditures are growing over time (15, 36). Using data from the Health and Retirement Study, Marshall et al. found that out-of-pocket expenditures near the end of life were large relative to the decedent's median non-housing wealth (35). The degree to which hospital bills and reducing ICU LOS and ICU admissions can impact this financial burden is unknown and worthy of future investigation. Importantly, the primary goal must be to improve the quality of care and the quality of dying while providing care that is consistent with patient preferences and values. In addition, it is important to acknowledge and recognize that the financial burden for patients and families and the potential to reduce their costs may not align with the financial burdens and incentives of either the hospital or healthcare system.

Consideration of the financial burden on family members raises important issues of ethics and justice. Incorporating financial costs into decisions made about end-of-life care raises concern in situations where conflicts of interest between surrogate decision makers and patients or physicians may exist. Ideally, patients with terminal or chronic illness admitted to the ICU have had previous discussions about values, goals, and preferences for end-of-life care early in the course of their disease so that well documented goals of care exist to guide surrogate decision makers and physicians. Often times, however, this is not the case. Good communication and shared decision making may help to reveal situations in which potential conflicts of interest may exist. In a study investigating conflicts between physicians' practices and patients' wishes, Asch and colleagues concluded that physicians do not reflexively limit or continue life-sustaining treatments based solely on patients' or surrogates' requests. These requests are one of many factors, including assessments of prognosis, and perceptions of other ethical and legal guidelines, that help to guide decision making (37). Additionally, one of the goals of good communication and family engagement is to try and ascertain patients' thoughts on quality of life values before any surrogate decisions are made on the patients' behalf (38). This helps to evaluate the appropriateness of decisions in a larger context.

There are also important issues of justice that must be considered when considering the role of financial burden in decision-making about intensity of care at the end of life. The intensity of care and the quality of care that seriously ill patients and their families receive should not be based on ability to pay or ability to weather financial burden, rather it should be based on patient values, goals, and preferences for what constitutes the best quality of end-of-life care and quality of dying. Discussing costs of care for critically ill and dying patients is a sensitive and controversial topic. However, patients have the right to be informed of all of their options and for patients who consider the financial burden of care a top priority, they have the right to be informed about the financial consequences of care as well. Importantly, the concept of "rationing" care is not what is at play here. Rationing of care is defined as denying a potentially beneficial treatment to a patient on the grounds of scarcity (39, 40). Advance care planning and palliative care interventions seek to individualize care and put decision-making in the hands of informed patients. The goal here is to reduce unwanted intensive care at the end of life, not limit it for patients who desire it.

Conclusions

Studies investigating the effect of advance care planning and palliative care consultation on ICU utilization suggest that these interventions have the potential to reduce intensity of care at the end of life by reducing ICU admissions and decreasing ICU LOS. While the economic implications of this are unknown, providing care that is consistent with patient values and preferences can improve the quality of end-of-life care and the quality of dying. Future research is needed to better understand whether or not advance care planning and palliative care consultation have the ability to improve care while simultaneously decreasing overall costs. In addition, such interventions may also provide an opportunity to reduce the financial burden on the family members of these patients. Although reduction in financial burden should not be the primary rationale for palliative care interventions, it may be a useful secondary benefit provided that patients are receiving the care they would choose if fully informed and that the implementation of palliative care improves quality of care and improves patient and family outcomes. As we transition to the Accountable Care Organization environment, interventions that have the ability to improve quality of care delivered and reduce health care costs at the same time will be of special interest.

Acknowledgments

N.K. is currently receiving funding from the National Institute of General Medical Sciences (5T32GM086270).

References

1. Curtis JR, Engelberg RA, Bensink ME, et al. End-of-life care in the intensive care unit: can we simultaneously increase quality and reduce costs? *American journal of respiratory and critical care medicine*. 2012; 186(7):587–592. [PubMed: 22859524]
2. Riley GF, Lubitz JD. Long-term trends in Medicare payments in the last year of life. *Health services research*. 2010; 45(2):565–576. [PubMed: 20148984]
3. Seshamani M, Gray AM. A longitudinal study of the effects of age and time to death on hospital costs. *Journal of health economics*. 2004; 23(2):217–235. [PubMed: 15019753]
4. Pastores SM, Dakwar J, Halpern NA. Costs of critical care medicine. *Critical care clinics*. 2012; 28(1):1–10. v. [PubMed: 22123095]
5. Halpern NA, Pastores SM. Critical care medicine in the United States 2000–2005: an analysis of bed numbers, occupancy rates, payer mix, and costs. *Critical care medicine*. 2010; 38(1):65–71. [PubMed: 19730257]
6. Teno JM, Gozalo PL, Bynum JP, et al. Change in end-of-life care for Medicare beneficiaries: site of death, place of care, and health care transitions in 2000, 2005, and 2009. *JAMA : the journal of the American Medical Association*. 2013; 309(5):470–477.
7. G, G, Jr. Spiritual beliefs and the dying process: a report on a national survey. Conducted for the Nathan Cummings Foundation and the Fetzer Institute; 1997.
8. Lilly CM, De Meo DL, Sonna LA, et al. An intensive communication intervention for the critically ill. *The American journal of medicine*. 2000; 109(6):469–475. [PubMed: 11042236]
9. Quill TE, Holloway R. Time-limited trials near the end of life. *JAMA : the journal of the American Medical Association*. 2011; 306(13):1483–1484.
10. Scheunemann LP, McDevitt M, Carson SS, et al. Randomized, controlled trials of interventions to improve communication in intensive care: a systematic review. *Chest*. 2011; 139(3):543–554. [PubMed: 21106660]
11. Gade G, Venohr I, Conner D, et al. Impact of an inpatient palliative care team: a randomized control trial. *Journal of palliative medicine*. 2008; 11(2):180–190. [PubMed: 18333732]

12. Penrod JD, Deb P, Dellenbaugh C, et al. Hospital-based palliative care consultation: effects on hospital cost. *Journal of palliative medicine*. 2010; 13(8):973–979. [PubMed: 20642361]
13. Penrod JD, Deb P, Luhrs C, et al. Cost and utilization outcomes of patients receiving hospital-based palliative care consultation. *Journal of palliative medicine*. 2006; 9(4):855–860. [PubMed: 16910799]
- 14*. Aslakson R, Cheng J, Vollenweider D, et al. Evidence-based palliative care in the intensive care unit: a systematic review of interventions. *Journal of palliative medicine*. 2014; 17(2):219–235. This systematic review highlights the impact of consultative and integrative palliative care interventions on ICU and hospital LOS, satisfaction and mortality. [PubMed: 24517300]
15. Institute of Medicine. *Approaching Death: Improving Care at the End of Life*. Washington DC: The National Academic Press; 1997.
16. Bakitas M, Lyons KD, Hegel MT, et al. Effects of a palliative care intervention on clinical outcomes in patients with advanced cancer: the Project ENABLE II randomized controlled trial. *JAMA : the journal of the American Medical Association*. 2009; 302(7):741–749.
17. Andereck WS, M J, Schneiderman LJ, et al. Seeking to Reduce Nonbeneficial Treatment in the ICU: An Exploratory Trial of Proactive Ethics Consultation. *Critical Care Medicine*. 2014; 42(4)
18. Curtis JR, Nielsen EL, Treece PD, et al. Effect of a quality-improvement intervention on end-of-life care in the intensive care unit: a randomized trial. *American journal of respiratory and critical care medicine*. 2011; 183(3):348–355. [PubMed: 20833820]
19. Shelton W, Moore CD, Socaris S, et al. The effect of a family support intervention on family satisfaction, length-of-stay, and cost of care in the intensive care unit. *Critical care medicine*. 2010; 38(5):1315–1320. [PubMed: 20228678]
20. Daly BJ, Douglas SL, O'Toole E, et al. Effectiveness trial of an intensive communication structure for families of long-stay ICU patients. *Chest*. 2010:1340–1348. [PubMed: 20576734]
21. Detering KM, Hancock AD, Reade MC, et al. The impact of advance care planning on end of life care in elderly patients: randomised controlled trial. *BMJ*. 2010; 340:c1345. [PubMed: 20332506]
22. Schneiderman LJ, Gilmer T, Teetzel HD. Impact of ethics consultations in the intensive care setting: a randomized, controlled trial. *Critical care medicine*. 2000; 28(12):3920–3924. [PubMed: 11153636]
23. Schneiderman LJ, Gilmer T, Teetzel HD, et al. Effect of ethics consultations on nonbeneficial life-sustaining treatments in the intensive care setting: a randomized controlled trial. *JAMA : the journal of the American Medical Association*. 2003; 290(9):1166–1172.
24. Ahrens T, Yancey V, Kollef M. Improving family communications at the end of life: implications for length of stay in the intensive care unit and resource use. *American journal of critical care : an official publication, American Association of Critical-Care Nurses*. 2003; 12(4):317–323. discussion 324.
25. Campbell ML, Guzman JA. Impact of a proactive approach to improve end-of-life care in a medical ICU. *Chest*. 2003; 123(1):266–271. [PubMed: 12527629]
26. Campbell ML, Guzman JA. A proactive approach to improve end-of-life care in a medical intensive care unit for patients with terminal dementia. *Critical care medicine*. 2004; 32(9):1839–1843. [PubMed: 15343010]
27. Dowdy MD, Robertson C, Bander JA. A study of proactive ethics consultation for critically and terminally ill patients with extended lengths of stay. *Critical care medicine*. 1998; 26(2):252–259. [PubMed: 9468161]
28. Curtis JR, Treece PD, Nielsen EL, et al. Integrating palliative and critical care: evaluation of a quality-improvement intervention. *American journal of respiratory and critical care medicine*. 2008; 178(3):269–275. [PubMed: 18480429]
29. Mosenthal AC, Murphy PA, Barker LK, et al. Changing the culture around end-of-life care in the trauma intensive care unit. *The Journal of trauma*. 2008; 64(6):1587–1593. [PubMed: 18545128]
30. Lilly CM, Sonna LA, Haley KJ, et al. Intensive communication: four-year follow-up from a clinical practice study. *Critical care medicine*. 2003; 31(5 Suppl):S394–399. [PubMed: 12771590]
31. Norton SA, Hogan LA, Holloway RG, et al. Proactive palliative care in the medical intensive care unit: effects on length of stay for selected high-risk patients. *Critical care medicine*. 2007; 35(6):1530–1535. [PubMed: 17452930]

32. Downey L, Engelberg RA, Curtis JR, et al. Shared priorities for the end-of-life period. *J Pain Symptom Manage*. 2009; 37(2):175–188. [PubMed: 18722084]
33. Steinhauser KE, Christakis NA, Clipp EC, et al. Factors considered important at the end of life by patients, family, physicians, and other care providers. *JAMA : the journal of the American Medical Association*. 2000; 284(19):2476–2482.
- 34*. Khandelwal N, Engelberg RA, Benkeser D. End-of-Life expenditure in the ICU and perceived quality of dying. *Chest*. 2014 In press. This article highlights the important role of insurance status as a mediator in the relationship between costs of ICU care and family-assessed quality of dying.
35. Marshall, S.; McGarry, KM.; Skinner, JS. NBER Working Paper Series: The risk of out-of-pocket health care expenditure at end of life. National Bureau of Economic Research; 2010. Working Paper 16170
36. Paez KA, Zhao L, Hwang W. Rising out-of-pocket spending for chronic conditions: a ten-year trend. *Health Aff (Millwood)*. 2009; 28(1):15–25. [PubMed: 19124848]
37. Asch DA, Hansen-Flaschen J, Lanken PN. Decisions to limit or continue life-sustaining treatment by critical care physicians in the United States: conflicts between physicians' practices and patients' wishes. *American journal of respiratory and critical care medicine*. 1995; 151(2 Pt 1): 288–292. [PubMed: 7842181]
38. Henig NR, Faul JL, Raffin TA. Biomedical ethics and the withdrawal of advanced life support. *Annual review of medicine*. 2001; 52:79–92.
39. Truog RD, Brock DW, Cook DJ, et al. Rationing in the intensive care unit. *Critical care medicine*. 2006; 34(4):958–963. quiz 971. [PubMed: 16484912]
40. Scheunemann LP, White DB. The ethics and reality of rationing in medicine. *Chest*. 2011; 140(6): 1625–1632. [PubMed: 22147821]

Abbreviations

ICU	Intensive care unit
LOS	Length of stay

Key points

- Advance care planning and palliative care interventions may have the ability to improve the quality of dying while simultaneously reducing overall costs.
- Evidence suggests that advance care planning and palliative care interventions may reduce intensity of care at the end of life by reducing ICU LOS and ICU admissions.
- Future research is needed to better understand how reducing unwanted intensive care at the end of life impacts the financial burden placed on family members of dying patients.