Published in final edited form as:

Org Lett. 2014 February 7; 16(3): 724–726. doi:10.1021/ol403402h.

Leveraging the Micellar Effect: Gold-Catalyzed Dehydrative Cyclizations ... *in Water* at Room Temperature

Stefan R. K. Minkler[†], Nicholas A. Isley[‡], Daniel J. Lippincott[‡], Norbert Krause[†], and Bruce H. Lipshutz[‡]

Organic Chemistry, Dortmund University of Technology, Otto-Hahn-Strasse 6, D-44227 Dortmund, Germany, and Department of Chemistry and Biochemistry, University of California, Santa Barbara, California 93106

Norbert Krause: norbert.krause@tu-dortmund.de; Bruce H. Lipshutz: lipshutz@chem.ucsb.edu

Abstract

The first examples of gold-catalyzed cyclizations of diols and triols to the corresponding hetero- or spirocycles in a bulk aqueous medium are presented. These reactions take place within nanomicelles, where the hydrophobic effect is operating, thereby driving the dehydrations, notwithstanding the surrounding water. By the addition of simple salts such as sodium chloride, reaction times and catalyst loadings can be significantly decreased.

Dehydrative reactions that take place in organic media are typically driven by the presence of a dehydrating agent, such as molecular sieves. As an alternative strategy, micellar catalysis offers an opportunity to take advantage of the hydrophobic effect that exists within the lipophilic cores of nanomicelles formed upon dissolution of a surfactant in water.¹ Since the surfactant is providing a very limited amount of an organic medium in which these reactions can take place, reactions occur under typically high internal concentrations and under mild, usually room temperature conditions. Water is expelled for entropic reasons; it cannot solvate the hydrocarbon present on the inside of these nanoparticles. Nonetheless, solvent effects can be large, as with any organic reaction run in an organic medium, and hence, the nature of the amphiphile can play a significant role in the outcome of reactions run under such aqueous conditions. This has been extensively demonstrated in transition metal-catalyzed reactions, including several name reactions that are Nobel-prize winning,

Correspondence to: Norbert Krause, norbert.krause@tu-dortmund.de; Bruce H. Lipshutz, lipshutz@chem.ucsb.edu.

[†]Dortmund University of Technology.

[‡]University of California, Santa Barbara.

Supporting Information Available: Experimental procedures and data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

such as olefin metathesis,² Heck reactions,³ and Suzuki-Miyaura couplings.⁴ The preclusion of water from the inside of such nanoreactors has also been used, contrary to textbook thinking, to effect both net Negishi-type couplings of highly moisture sensitive organozinc reagents,⁵ as well as conjugate additions via water-intolerant organocopper complexes, in the absence of pre-formed organometallics.⁶ These latter two cases provide strong evidence that the surrounding water does not easily penetrate the hydrophobic micellar pockets. On the basis of these observations, it was anticipated that the hydrophobic effect would act to drive gold-catalyzed dehydrative reactions to cyclic products⁷ in bulk water.

Our investigations began with acetylenic diol **1** which undergoes dehydrative cyclization to furan **2**. Previously, cationic or neutral gold(I) catalysts in THF or toluene was used for this purpose, sometimes in the presence of activated molecular sieves for scavenging water.⁸ Treatment of **1** with 5 mol % AuBr₃ in 2 weight % aqueous PTS (poly(oxyethyl)- α -tocopheryl sebacate; global concentration 0.5 M) for 2 h afforded furan **2** in excellent yield (90%; Table 1, entry 1). In accordance with previous work by Aponick et al.^{8a}, a control experiment done “on water” (i.e., without amphiphile) led to no product formation under identical conditions, documenting that these are reactions taking place under micellar catalysis conditions (entry 2). With second generation surfactant TPGS-750-M the yield was slightly higher (entry 3). Other surfactants such as TPGS-1000, Brij 30, and Nok⁹ gave inferior levels of conversion (entries 4–6). It was possible to decrease the gold catalyst loading to 2 mol % but with somewhat elongated reaction times (entries 7 and 8). Addition of silver salts or use of cationic gold catalyst **A** (entry 13) did not lead to higher yields or shorter reaction times (entries 9–13). All subsequent experiments were performed in aqueous TPGS-750-M.

After this first example (**1** to **2**) we explored other diols as well as amino alcohols bearing secondary amino groups. Primary amines have shown poor reactivity in gold catalyzed cyclizations due to deactivation of the gold catalyst by the Lewis basic amine.¹⁰ Under these micellar conditions, however, pyrroles **4a** and **4b** were formed smoothly and quickly; after 1 h, with high yields of 91% and 93%, respectively (Table 2, entries 1, 2). Diols **3c–3e** were also cyclized readily with different gold(I) and gold(III) catalysts (entries 3–14); the latter seem to be best suited for these substrates with reaction times of 5–20 min and result in high yields (83–92%; entries 3, 7, 11). This may reflect a limited stability of gold(I) salts in aqueous media. Diols **3f–3i** with different substitution patterns were subjected to 2 mol % AuBr₃/AgOTf to form furans with various substituents at the 2-, 3-, and 5-positions. With this lower catalyst loading the products could be obtained in 2 h with consistently good yields. It is worth noting that with increasing steric hindrance at the 2- and/or 3-positions, the yield was increased (compare entries 16 and 17). Moreover, these reactions tolerate the presence of a nitrile (entries 11–14) and an olefin (entry 15).

Further optimization focused on decreasing the time associated with these dehydrative cyclization reactions. From previous work it had been shown that dissolution of salts, such as NaCl, in the aqueous reaction medium can lead to beneficial increases in micellar diameter, thereby extending the time for substrate/catalyst exposure to the inner lipophilic core.^{1,11} The outgrowth of this simple modification, in addition to a further reduction in

catalyst loading to 1 mol %, is an enhanced reaction rate leading to shorter reaction times (Table 3, entries 4, 6, 8–10).

Previous experiments run in D₂O have shown that the proton that replaces the gold on the ring derives from the surrounding water, rather than the starting alcohol. If iodine (one equivalent) is present in the reaction of diol **3g**, iodinated furan **5** is formed in 93% isolated yield (Scheme 1). With the iodinated furan in hand, it is possible to use it in palladium-catalyzed cross-coupling reactions.¹²

Further application of this technology beyond diols was made to triols, thereby forming various spirocycles (Table 4). Previously, cationic gold(I) catalysts in THF were used for this transformation, in the presence of activated molecular sieves for scavenging water.¹³ Triol **6a** was smoothly cyclized with Ph₃PAuCl/AgOTf in water containing 2 wt % TPGS-750-M at room temperature over 5 h to the corresponding spirocycle **7a** (89% yield; entry 1). Under the same conditions, increased steric hindrance at the 1- and 3-positions dramatically slowed the conversion (triol **6b**; entry 2). An increased rate of protodeauration after the initial cyclization led to formation of a complex mixture containing unsaturated monocyclization products^{13a} together with the desired spirocycle **7b**. With a cationic gold(I) catalyst formed from **B** and AgOTf the yield of **7b** could be doubled (entry 3), but it was only after the use of precatalyst **C**/AgOTf that a moderate yield of product **7b** could be isolated (46%; entry 4). The related triol **6c** bearing the same steric hindrance at the 1- and 3-positions, but with a longer carbon chain (albeit without phenyl substituents) could be cyclized in 64% yield to the corresponding spirocycle **7c** upon heating to 40 °C for 12 h.

In summary, acetylenic vicinal diols and amino alcohols can be cyclodehydrated to the corresponding furans and pyrroles, respectively, using nanoparticle technology in water. Nanomicelles composed of the designer surfactant TPGS-750-M, which is an item of commerce, enable these gold-catalyzed reactions to take place at ambient temperatures and in high isolated yields. Appropriately substituted alkynes containing three hydroxyl residues can be converted to spirocyclic products in the presence of a Au(I) catalyst. Further work on developing an enantioselective version of this route to spirocycles is currently under study.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial support for our programs in green chemistry provided by the NIH (GM 86485) to BHL is warmly acknowledged with thanks.

References

1. a) Lipshutz BH, Ghorai S. *Aldrichimica Acta*. 2008; 41:59–72. b) Lipshutz BH, Ghorai S. *Aldrichimica Acta*. 2012; 45:3–16. [PubMed: 23807816] c) Lipshutz BH, Isley NA, Fennewald JC, Slack ED. *Angew Chem Int Ed*. 2013; 52:10952–10958.
2. a) Lipshutz BH, Aguinaldo GT, Ghorai S, Voigtritter K. *Org Lett*. 2008; 10:1325–1328. [PubMed: 18335947] b) Lipshutz BH, Ghorai S, Aguinaldo GT. *Adv Synth Catal*. 2008; 350:953–956. c) Lipshutz BH, Ghorai S. *Tetrahedron*. 2010; 66:1057–1063.

3. Lipshutz BH, Taft BR. *Org Lett.* 2008; 10:1329–1332. [PubMed: 18335945]
4. a) Lipshutz BH, Petersen TB, Abela AR. *Org Lett.* 2008; 10:1333–1336. [PubMed: 18335944] b) Lipshutz BH, Abela AR. *Org Lett.* 2008; 10:5329–5332. [PubMed: 18998705] c) Nishikata T, Lipshutz BH. *J Am Chem Soc.* 2009; 131:12103–12105. [PubMed: 19663513]
5. a) Krasovskiy A, Duplais C, Lipshutz BH. *J Am Chem Soc.* 2009; 131:15592–15593. [PubMed: 19827762] b) Krasovskiy A, Duplais C, Lipshutz BH. *Org Lett.* 2010; 12:4742–4744. [PubMed: 20882969]
6. Lipshutz BH, Huang S, Leong WWY, Zhong G, Isley NA. *J Am Chem Soc.* 2012; 134:19985–19988. [PubMed: 23190029]
7. Biannic B, Aponick A. *Eur J Org Chem.* 2011:6605–6617. See also: Manabe K, Iimura S, Sun X-M, Kobayashi S. *J Am Chem Soc.* 2002; 124:11971–11978. [PubMed: 12358542]
8. a) Aponick A, Li CY, Malinge J, Marques EF. *Org Lett.* 2009; 11:4624–4627. [PubMed: 19772313] b) Egi M, Azechi K, Akai S. *Org Lett.* 2009; 11:5002–5005. [PubMed: 19780532] c) Egi M, Azechi K, Akai S. *Adv Synth Catal.* 2011; 353:287–290.
9. Klumphu P, Lipshutz BH. *J Org Chem.* in press.
10. a) Morita N, Krause N. *Org Lett.* 2004; 6:4121–4123. [PubMed: 15496114] b) Morita N, Krause N. *Eur J Org Chem.* 2006:4634–4641.
11. Minkler SRK, Lipshutz BH, Krause N. *Angew Chem Int Ed.* 2011; 50:7820–7823.
12. a) Gockel B, Krause N. *Eur J Org Chem.* 2010:311–316. b) Bew SP, El-Taieb GMM, Jones S, Knight DW, Tan WF. *Eur J Org Chem.* 2007:5759–5770.
13. Aponick A, Li CY, Palmes JA. *Org Lett.* 2009; 11:121–124. [PubMed: 19049406] See also: Fang C, Pang Y, Forsyth CJ. *Org Lett.* 2010; 12:4528–4531. [PubMed: 20849144]

Scheme 1.

Gold-catalyzed dehydrative cyclization of diol **3g** in the presence of iodine.

Table 1

Gold-catalyzed Dehydrative Cyclization of Diol 1.

entry	[Au]	surfactant	time (h)	conv. ^b (%)	yield ^c (%)
1	AuBr ₃	PTS	2	94	90
2	AuBr ₃	none	7	0	nd
3	AuBr ₃	TPGS-750-M	2	87	91 (2.5 h)
4	AuBr ₃	TPGS-1000	2	79	nd
5	AuBr ₃	Brij 30	2	50	nd
6	AuBr ₃	NOK	2	63	nd
7	AuBr ₃ ^c	PTS	2	35	88 (4 h)
8	AuBr ₃ ^d	TPGS-750-M	2	16	86 (6 h)
9	AuCl ₃ /AgOTf	TPGS-750-M	6	nd	70
10	AuCl ₃ /AgSbF ₆	TPGS-750-M	6	nd	68
11	AuBr ₃ /AgOTf	TPGS-750-M	2	nd	92
12	AuBr ₃ /AgSbF ₆	TPGS-750-M	2	nd	90
13	A ^e	TPGS-750-M	3	nd	84

^aPTS = poly(oxyethyl)- α -tocopheryl sebacate; TPGS-750-M = D- α -tocopherol-polyethyleneglycol-750-succinate monomethyl ether; TPGS-1000 = D- α -tocopherol-polyethyleneglycol-1000-succinate; Brij 30 = tetraethyleneglycol-monoiododecyl ether; NOK = β -sitosterol-polyethyleneglycol-600-succinate monomethyl ether.

^bConversion determined by GC-MS.

^cIsolated yield.

^d2 mol %.

^e**A** = [*t*-Bu₂(α -biphenyl)]PAu(MeCN)SbF₆.

Table 2

Gold-catalyzed Dehydrative Cyclization of Diols and Amino Alcohols **3**.

entry	3	X	R ¹	R ²	R ³	[Au]	time	4/yield ^a (%)
1	3a	NBu	Ph	H	H	AuBr ₃	1 h	4a (91)
2	3b	NBn	Ph	H	H	AuBr ₃	1 h	4b (93)
3	3c	O	Ph	H	H	AuBr ₃	20 min	4c (92)
4	3c	O	Ph	H	H	AuCl	2 h	4c (79)
5	3c	O	Ph	H	H	Ph ₃ PAuCl/AgOTf	40 min	4c (82)
6	3c	O	Ph	H	H	Ph ₃ PAuCl/AgBF ₄	4 h	4c (84)
7	3d	O	<i>n</i> -C ₈ H ₁₇	H	H	AuBr ₃	5 min	4d (90)
8	3d	O	<i>n</i> -C ₈ H ₁₇	H	H	AuCl	3 h	4d (76)
9	3d	O	<i>n</i> -C ₈ H ₁₇	H	H	Ph ₃ PAuCl/AgOTf	1 h	4d (85)
10	3d	O	<i>n</i> -C ₈ H ₁₇	H	H	Ph ₃ PAuCl/AgBF ₄	6 h	4d (86)
11	3e	O	(CH ₂) ₃ CN	H	H	AuBr ₃	10 min	4e (83)
12	3e	O	(CH ₂) ₃ CN	H	H	AuCl	2 h	4e (80)
13	3e	O	(CH ₂) ₃ CN	H	H	Ph ₃ PAuCl/AgOTf	1 h	4e (82)
14	3e	O	(CH ₂) ₃ CN	H	H	Ph ₃ PAuCl/AgBF ₄	4 h	4e (80)
15	3f	O	cyclohexen-1-yl	H	H	AuBr ₃ /AgOTf ^b	2 h	4f (83)
16	3g	O	Ph	Ph	Ph	AuBr ₃ /AgOTf ^b	2 h	4g (95)
17	3h	O	Ph	Me	Me	AuBr ₃ /AgOTf ^b	2 h	4h (85)
18	3i	O	<i>n</i> -Hex	Me	Me	AuBr ₃ /AgOTf ^b	2 h	4i (85)

^a Isolated yield.^b 2 mol %.

Table 3

Gold-catalyzed Dehydrative Cyclization of Diols **3** in the Presence of NaCl.

entry	3	R	AuBr₃ (mol %)	additive	time (min)	4/yeild^a (%)
1	3c	Ph	5	-	20	4c (92)
2	3c	Ph	2	-	50	4c (91)
3	3c	Ph	1	-	120	4c (88)
4	3c	Ph	1	NaCl ^b	30	4c (90)
5	3d	<i>n</i> -C ₈ H ₁₇	5	-	5	4d (90)
6	3d	<i>n</i> -C ₈ H ₁₇	1	NaCl ^b	20	4d (87)
7	3e	(CH ₂) ₃ CN	5	-	10	4e (83)
8	3e	(CH ₂) ₃ CN	1	NaCl ^b	40	4e (91)
9	3j	<i>n</i> -Hex	2	NaCl ^b	30	4j (89)
10	3k	(CH ₂) ₃ Cl	2	NaCl ^b	30	4k (86)

^a Isolated yield.

^b 2 M.

Table 4

Gold-catalyzed Dehydrative Spirocyclization of Triols **6**.

entry	6	R¹	R²	n	[Au]	time (h)	7/yeild ^a (%)
1	6a	Ph	H	0	Ph ₃ PAuCl	5	7a (89)
2	6b	Ph	Me	0	Ph ₃ PAuCl	4	7b (11)
3	6b	Ph	Me	0	B^b	4	7b (22) ^c
4	6b	Ph	Me	0	C^d	4	7b (46)
5	6c	H	Me	1	Ph ₃ PAuCl	12	7c (64) ^e

^aIsolated yield.

^b**B** = [*c*-Hex2(*o*-biphenyl)]PAuCl.

^c40 °C.

^d**C** = [*t*-Bu2(*o*-biphenyl)]PAuCl.

^e19% reisolated starting material.