Published in final edited form as:

Curr Heart Fail Rep. 2014 December ; 11(4): 485–499. doi:10.1007/s11897-014-0226-4.

Novel Renal Biomarkers to Assess Cardiorenal Syndrome

Meredith A. Brisco, MD MSCE [Assistant Professor of Medicine] and

Heart Failure and Cardiac Transplantation Medical University of South Carolina 25 Courtenay Drive, ART 7061 MSC 592 Charleston, SC 29425-5920 Phone: (843) 876-4807 Fax: (843) 876-4809 brisco@musc.edu

Jeffrey M. Testani, MD MTR [Assistant Professor of Medicine]

Department of Internal Medicine Program of Applied Translational Research Yale University 60 Temple Street Suite 6C New Haven, CT 06510

Abstract

Renal dysfunction (RD) in heart failure portends adverse outcomes and often limits aggressive medical and decongestive therapies. Despite the high prevalence in this population, not all forms of RD are prognostically or mechanistically equivalent: RD can result from irreversible nephron loss secondary to diabetic or hypertensive kidney disease or it can develop secondary to the HF itself, i.e. the cardiorenal syndrome. Furthermore, filtration is only one aspect of renal performance such that significant renal impairment secondary to cardiorenal syndrome can exist despite a normal glomerular filtration rate. Renal biomarkers have the potential to inform some of the intricacies involved in accurately assessing cardiorenal interactions. This article discusses novel biomarkers for cardiorenal syndrome and their utility in prognosis, diagnosis, and targeted treatment of heart failure-induced RD.

Keywords

cardiorenal syndrome; heart failure; worsening renal function; improvement in renal function; acute kidney injury; kidney injury biomarkers; cystatin C; albuminuria; Neutrophil Gelatinase-Associated Lipocalin; N-Acetyl- β -D-Glucosaminidase; Kidney Injury Molecule-1; Interleukin-18; Blood Urea Nitrogen to Creatinine Ratio; Brain Natriuretic Peptide; C-Type Natriuretic Peptide; diuretic resistance; hepatic congestion; galectin-3

Introduction

The complex interplay between the heart and the kidney in the setting of concomitant impairment of each organ has been referred to as the cardiorenal syndrome (CRS) [1]. The relevance of the kidney in heart failure (HF) is obvious as it serves as the regulator of fluid

(Corresponding Author) Phone: (203) 737-6227 Fax: (203) 746-8373 jeffrey.testani@yale.edu.

Compliance with Ethics Guidelines

Conflict of Interest

Meredith A. Brisco and Jeffrey M. Testani declare that they have no conflict of interest.

Human and Animal Rights and Informed Consent

This article does not contain any studies with human or animal subjects performed by any of the authors.

and sodium balance in the body. Thus perturbations in renal physiology are central to the development of volume retention, which is a hallmark of the HF syndrome [2]. Additionally, a requisite to the use of many lifesaving therapies, such as angiotensin converting enzyme inhibitors, is adequate renal function. As a result of the above, it is not surprising that it has repeatedly been demonstrated that RD is common in HF and is one of the strongest predictors of morbidity and mortality in these patients [3-6].

Despite widespread recognition of the importance of CRS, progress toward an understanding of this “syndrome” has remained a challenge [7]. The true difficulty lies in the fact that not all elevated creatinine levels in HF occur via the same mechanism [8-10]. Although some patients have RD caused by the HF itself, many patients have comorbid intrinsic kidney disease from shared risk factors such as diabetes or hypertension [11]. Unfortunately, assuming that all RD in HF should be treated similarly is akin to treating all anemia with vitamin B-12 supplementation and expecting improvement in hemoglobin levels in all patients. It is therefore not surprising in the face of these challenges that the majority of our “cardio-renal” trials have been disappointingly negative [12-16].

However, novel biomarkers provide many opportunities in overcoming these challenges. There are three important purposes for cardiorenal biomarkers: 1) Prognostication 2) Discrimination between RD phenotypes and 3) Patient identification for targeted therapeutic intervention. Currently, most cardiorenal biomarkers have aided in prognosis, but the ultimate goal is for them to facilitate selection of patients most likely to benefit from specific CRS or HF therapies. The purpose of this review is to provide an overview of biomarkers for CRS highlighting that while in some cases novel means “new”, in other cases novel refers to the repurposing of traditional laboratory metrics to inform on CRS. Each biomarker will be discussed in the context of the potential pathophysiology of CRS they may query (Table 1).

Glomerular Filtration and Integrity

Initial recognition of cardiorenal dysfunction focused on the presence of a reduced glomerular filtration rate (GFR). Serum creatinine-based measures of renal function are significantly influenced by a number of factors unrelated to the GFR [17]. Therefore determining alternative less-biased biomarkers for GFR estimation in addition to parameters that describe glomerular integrity has been the focus of significant research.

Cystatin C (CysC)

CysC is a 13 kDa cysteine protease, ubiquitous in all nucleated cells that is produced at a constant rate, freely filtered, and not secreted in renal tubules. Despite initial enthusiasm that this marker would perform dramatically better than serum creatinine since it is not primarily determined by muscle mass, frequent discrepancies from measured GFR are now well-recognized [18, 19]. Although the data has not necessarily born out that CysC is vastly superior to creatinine with respect to measured GFR, the sources of bias are different and thus significant and context-specific incremental information appears to be present with this marker [20-23].

Prognosis in CRS—The prognostic capability of CysC was first demonstrated in over 4,000 older patients in the Cardiovascular Health Study where CysC was independently and linearly associated with increased cardiovascular (CV) events and mortality, outperforming creatinine [24]. In a subset of patients with chronic HF, the highest quartile of CysC (>1.55 mg/L) was associated with two times the risk of CV mortality adjusted for baseline characteristics [25]. Subsequent larger studies in stable chronic HF confirmed the associated risk of death with increasing CysC even when added to models including eGFR using either the Cockcroft-Gault or Modified Diet in Renal Disease (MDRD) equations [21, 26-28]. Additionally, elevated CysC identified patients at high risk for death without RD as determined by the MDRD equation [28]. In patients presenting with ADHF, CysC performs similarly as a mortality indicator and is associated with repeat HF hospitalization and adjusted risk of death at 30 days and 1 year, again outperforming eGFR, regardless of the estimating equation used [29-32]. In those patients with $\text{eGFR} > 90 \text{ ml/min/1.73m}^2$, CysC remained a robust predictor of survival [29]. CysC also aids in risk stratification alongside other powerful HF biomarkers. Lassus et al. demonstrated in 620 HF patients admitted with volume overload that patients with the highest tertile of N-terminal prohormone brain natriuretic peptide (NT-proBNP) and CysC suffered a 48.7% 1-year mortality compared to 5.2% in those with the lowest tertile of both markers, while Manzano et al. showed that patients with elevations in cardiac troponin T ($>0.011 \text{ ng/mL}$), NT-proBNP ($>3.345 \text{ pg/mL}$) and CysC ($>1.21 \text{ mg/L}$) had a 66.7% risk of mortality [29, 32]. Similar additive prognostication of CysC when added to BNP was illustrated in stable HF outpatients [28].

Diagnosis of CRS—The enhanced value of CysC as a prognostic indicator in HF patients with normal renal function ($\text{eGFR} > 90 \text{ ml/min/1.73m}^2$) begs the question as to whether CysC is merely more accurately reflecting GFR than creatinine-based measures, particularly in populations with significant muscle wasting. Four out of ten patients without RD presenting with HF have elevated CysC levels indicating that traditional estimates of GFR may underdiagnose RD in HF [26]. To that end, Shilpak and colleagues recently proved the superiority of CysC eGFR across 11 large population studies in 90,750 participants; when CysC eGFR was used resulting in reclassification of patients into different CKD stages, the relationship between CKD and death was stronger and more linear across all eGFRs [20]. Furthermore, the increased risk of CV mortality appeared reliably at a CysC-based eGFR $< 85 \text{ ml/min/1.73m}^2$. In HF patients, CysC-based eGFR outperformed all other estimating equations with the lowest bias, great precision and excellent accuracy compared to directly measured GFR using iothalamate clearance [33]. However, despite potential improved accuracy and risk stratification with CysC, it offers no ability to differentiate different mechanisms for CRS.

Albuminuria

The size- and charge-selective glomerular barrier prevents filtration of the majority of large proteins like albumin and this is in part the reason that albumin is not normally found in the urine [34]. In the setting of glomerular capillary damage, glomerular integrity is disrupted and leakage results in increased amounts of albumin entering the renal tubules producing albuminuria. Additionally, damage to the proximal tubule may inhibit albumin reabsorption mechanisms leading to its appearance in the urine at abnormal levels [35]. Perhaps most

commonly associated with diabetes, HTN and both AKI and CKD, albuminuria is a puissant risk factor for both CV death and all-cause mortality, independent of other CV risk factors, and in patients with normal renal function [36, 37]. In chronic HF, microalbuminuria (30-300 mg/g creatinine (Cr)) is present in a third of patients and macroalbuminuria (>300 mg/gCr) occurs in 5-11% of patients [38-40]. Although albuminuria is inversely correlated with eGFR and occurs more commonly in HF patients with diabetes, it remains highly prevalent in nondiabetics and those with normal renal function [39]. The mechanism of albuminuria in HF compared to other disease states has not been elucidated. Inflammation and endothelial dysfunction have been proposed as possible etiologies, yet there was no association between markers like C-reactive protein and von Willebrand Factor and albuminuria in HF patients [41]. However, correlations between albuminuria and reduced renal blood flow, elevated NT-proBNP levels and physical exam findings of volume overload have been demonstrated, evoking potential involvement of “arterial underfilling” and venous congestion [41, 42]. In a study of 115 ADHF patients, in whom spot urine albumin to creatinine ratios (UACR) were measured on days 1 and 7 of hospital admission, the prevalence of microalbuminuria was 42% and a surprising 31% of patients had macroalbuminuria [43]. Interestingly, the mean UACR decreased significantly over 7 days, regardless of changes in renal function, and these improvements in UACR were correlated with similar improvements in NT-proBNP and serum bilirubin [43]. The improvement in albuminuria alongside the clinical improvement of congestive symptoms of HF exacerbation endorses a role for hypervolemia in the pathophysiology of albuminuria in HF.

Prognosis in CRS—Three substudies of major HF trials have revealed the prognostic power of albuminuria. In 2,310 patients enrolled in CHARM, albuminuria was associated with a 40-80% increase in the adjusted risk for all-cause mortality, CV mortality, and admission for HF [44]. In the GISSI-HF study, patients with microalbuminuria (HR=1.42, 95% CI 1.11-1.81, p=0.005) and macroalbuminuria (HR=1.70, 95% CI 1.16-2.50, p=0.006) suffered marked increases in mortality, independent of eGFR, diabetes and HTN [39]. Importantly in both studies, the magnitude of increased risk associated with albuminuria remained in those patients whose urinary albumin excretion was in the normal range. Baseline proteinuria as assessed by urinary dipstick in over 5,000 patients in Val-HeFT was also independently associated with increased mortality as well as morbid events including sudden death, HF hospitalization or administration of inotropic agents [42]. This observed risk did not differ between those patients with (58%) and without baseline RD implying that proteinuria is either an early precursor of RD, or is a result of HF-induced kidney damage via a different mechanism than what leads to reductions in GFR. Still, it is important to remember that patients with significantly decreased eGFR (Cr>2.5 mg/dL) were excluded from this study, so whether existence of more severe RD modifies the survival disadvantage associated with albuminuria is unknown.

Diagnosis of CRS—The prevalence of albuminuria in HF patients without concomitant RD (27% in CHARM) or diabetes and HTN (36% in GISSI-HF) suggests it may be valuable in identifying patients with CRS who would be overlooked using eGFR alone [39, 44]. Examining changes in albuminuria with decompensation may also further distinguish patients with CRS [43]. However, although utilizing albuminuria may enhance the

sensitivity of CRS discrimination, it also impairs specificity. Until there is a means of determining the etiology of albuminuria in those patients with coexisting diabetes or HTN and HF, albuminuria alone in these patients is unlikely to further phenotype CRS.

Targeted treatment in CRS—Albuminuria is one of the few cardiorenal biomarkers that has some data related to targeted treatment, albeit not particularly promising. In CHARM, candesartan had no effect on UACR during treatment, nor did aliskerin in the ALOFT trial but neither trial examined whether patients with albuminuria garnered improved outcomes with additional renin-angiotensin-aldosterone system inhibition despite lack of changes in UACR [38, 44]. Still, secondary analyses of clinical trials, while informative, should be interpreted with caution and currently there have been no trials of specific CRS therapies directed toward patients with albuminuria in HF.

Tubular Kidney Injury

In the case of acute kidney injury (AKI), creatinine often lags far behind the incidence of the actual kidney damage. Furthermore, significant kidney damage may exist without meaningful decrements in GFR secondary to renal reserve. Consequently, timely diagnosis of AKI is impeded and alterations in treatment, if appropriate, are delayed. Still, even with earlier recognition of AKI, all worsening in renal function (WRF) in HF is not created equally, a fact supported by the fact that WRF caused by different provocations (i.e., aggressive diuresis or a drop in blood pressure) have dramatically different prognostic implications[45-48]. Newly developed biomarkers of tubular kidney injury, unlike creatinine, have shown promise in earlier AKI identification and offer some discrimination to the site and degree of insult [49]. As a result, the following biomarkers indicative of tubular kidney injury may be of particular import and utility in CRS.

Neutrophil Gelatinase-Associated Lipocalin (NGAL)

NGAL is a small 25 kDa protein originally found in granules of neutrophils. In steady-state it's concentration is less than 20 ng/mL in both the serum and the urine but it can be elevated in chronic inflammatory conditions. Due to its involvement in iron transport, plasma levels are inversely correlated with indices of anemia [50]. In response to kidney injury, NGAL messenger RNA is transcribed in the kidney and both serum and urine levels are precipitously increased peaking within 24-48 hours following injury [51, 52]. NGAL is freely filtered in the glomerulus but it is nearly completely resorbed in the proximal convoluted tubule unless tubular damage exists. Schmidt-Ott outlined a model of NGAL trafficking to highlight the differences in what serum and urine measurements represent: urine NGAL reflects primarily intrarenal production from the thick ascending loop of Henle and collecting ducts, whereas systemic NGAL reflects extrarenal synthesis and potentially some renal-derived NGAL [53].

Urinary NGAL is increased in chronic HF outpatients compared to controls with only minimal correlation with eGFR and no correlation with NT-proBNP [54]. Serum NGAL is also elevated in chronic HF patients, including those with normal renal function [55]. Serum NGAL is not strongly correlated with EF or NYHA functional class or with other biomarkers like urinary KIM-1 and NAG [56, 57]. In patients presenting with ADHF, serum

NGAL is significantly correlated with serum creatinine ($r=0.68$, $p<0.0001$) and eGFR ($r=-0.69$, $p<0.0001$) whereas urinary NGAL has little relationship with renal function indices [58]. As a result, some have described serum NGAL as “expensive creatinine.”

Prognosis in CRS—In chronic HF, elevated serum and urine NGAL levels were unrelated to all-cause mortality in three smaller single-center studies [56, 57, 59]. However in over 2,000 patients enrolled in the GISSI-HF study, urinary NGAL was weakly associated with increased mortality (adjusted HR=1.23 per SD, 95% CI 1.07-1.41, $p=.003$) but failed to demonstrate a significant independent relationship with HF hospitalization [60]. Unlike other injury biomarkers, NGAL has been examined in multiple studies of ADHF and is nearly uniformly associated with increased mortality [55, 61-63]. Aghel et al. showed that an admission serum NGAL ≥ 215 ng/mL was associated with a 3-fold higher hazard for death and when admission systemic levels of NGAL exceed 100 ng/mL in patients with a BNP ≥ 330 pg/mL, the HR for mortality or HF re-hospitalization was 16.85, $p<0.006$ [61]. In a secondary analysis of the COACH study, systemic NGAL above the median was associated with increased mortality in patients with RD at discharge (HR=1.97, $p<0.001$) and in those with normal renal function (HR=2.01, $p=0.003$) [55]. NGAL outperformed other renal indices (eGFR and cystatin C) in predicting subsequent mortality. Interestingly, elevated urinary NGAL during admission for ADHF fails to confer an increased risk of death [64].

Diagnosis of CRS—Some of NGAL's promise as a CRS biomarker lies in its strong associations with development of WRF in the setting of ADHF regardless of baseline renal function. In one study of ADHF, patients with serum NGAL >140 ng/mL at hospital admission were 7.4 times more likely to develop WRF ($p<0.001$) [61]. Avelos et al. reported a similar relationship using a cut-off for serum NGAL >170 ng/mL resulting in an AUC for WRF during admission of 0.93, $p<0.001$ [65]. Serum NGAL remained associated with incident WRF, albeit less powerfully, in a significantly older population (median age=80) admitted with ADHF [66]. Serial measurements of serum NGAL in ADHF also appear to strengthen its ability to predict WRF where the degree of change in NGAL from baseline to peak produced an AUC for WRF of 0.91 compared to admission NGAL alone with an AUC of 0.69 [67]. However, not all study populations have exhibited positive associations between NGAL and WRF; Verbrugge et al. found no relationship between urinary NGAL and WRF or development of persistent renal impairment after discharge [64]. Although elevated serum NGAL conferred increased risk for WRF in the GISSI-HF study, it lost statistical significance in multivariable analysis [68].

The differing pathophysiology of renal damage indicated by urinary vs. serum levels of NGAL is unique among tubular injury biomarkers and may ultimately prove vital for characterization of the cardiorenal phenotype [53]. Shrestha et al. evaluated the relationship between both urine and serum NGAL and eGFR, effective diuresis and natriuresis in 93 patients admitted with ADHF [58]. Serum NGAL was strongly associated with eGFR and predicted development of WRF yet showed no correlation with metrics of diuresis including weight loss, fluid loss and natriuresis. Although serum and urine NGAL levels correlated moderately with each other, urinary NGAL demonstrated only weak correlations with

eGFR, was unrelated to WRF, and yet elevated levels were associated with decreased diuresis, weight loss and natriuresis with ADHF treatment. The authors proposed that urinary NGAL may identify HF patients with diuretic resistance, a hallmark of CRS. In a subsequent study from the same group in a less advanced and older ADHF population, urinary levels of NGAL only rose modestly in patients with an increase in serum creatinine, and once urinary NGAL levels were indexed to urine creatinine, volume of diuresis was identical between groups [69]. Further research is necessary to understand these findings and to determine whether applying differing treatment strategies based on serum/urinary NGAL profiles translates into clinical benefit [70].

N-Acetyl- β -D-Glucosaminidase (NAG)

NAG is a large lysosomal enzyme originating in proximal tubule cells and when detected in the urine, is indicative of proximal tubular injury with disruption of lysosomal integrity [71]. NAG was initially developed and validated as an early accurate marker of subsequent AKI, and when elevated in patients with existing AKI (normal urinary levels are <3 U/gCr), is associated with dialysis requirement and mortality [72, 73]. In patients with chronic HF, NAG levels are significantly higher compared to healthy controls, a finding that persists after adjustment for baseline eGFR [56]. Furthermore, patients with more significant LV dysfunction ($EF < 40\%$) have greater NAG levels compared to mild dysfunction [57]. Although elevations in NAG are more pronounced in those with HF and concomitant RD, defined as an $eGFR < 60$ ml/min/1.73m², this marker is also elevated in HF patients with preserved renal function [60].

Prognosis in CRS—As a CRS biomarker, NAG provides prognostic value; in multiple HF studies, NAG is associated with increased all-cause mortality and HF hospitalization independent of eGFR [56, 57, 60]. Damman et al. demonstrated in over 2,000 patients participating in the GISSI-HF trial that NAG conferred a significant and independent increased risk for HF hospitalization (adjusted HR=1.17 per SD, $p=0.025$) and mortality (adjusted HR= 1.30 per SD, $p=0.001$) in multivariable models. Patients with both RD and high NAG were two times more likely to die compared to those with low NAG and normal renal function [60].

Diagnosis of CRS—Although baseline NAG predicted subsequent WRF in the GISSI-HF trial in a univariate analysis, this relationship dissipated with covariate adjustment [68]. Interestingly, elevated NAG correlates with congestion as measured by NT-proBNP, brain natriuretic peptide (BNP) and atrial natriuretic peptide (ANP) and decreases similarly in response to diuretic-induced decongestion [74]. As venous congestion plays an important role in CRS, NAG has potential for phenotyping existing RD in HF. Unfortunately, although elevated urinary NAG is sensitive for tubular injury, it is also increased in diabetes and hypertension as well, limiting its specificity for its ability to differentiate CRS from tubular injury secondary to other etiologies [60].

Kidney Injury Molecule-1 (KIM-1)

KIM-1 is a type I cell membrane glycoprotein expressed in regenerating proximal tubular cells and facilitates phagocytosis of neighboring apoptotic tubular epithelial cells; it is not

expressed in the normal kidney [75]. Within 24 hours of tubular injury, KIM-1 increases dramatically (normal <200 ng/gCr) and sheds its ectodomain which is detectable in the urine [72]. In animal models, elevations correlated with severity of AKI by histopathology, outperforming creatinine and NAG [76]. In HF patients, KIM-1 is only modestly correlated with other biomarkers of tubular injury, eGFR, and urinary albumin excretion [56]. Although levels of KIM-1 are increased in those patients with RD, chronic HF patients exhibit higher levels even in the presence of normal kidney function [60]. Similar to urinary NAG, KIM-1 levels increase as EF decreases and as the severity of HF symptoms increase as exemplified by worse NYHA class [57]. However, in ADHF, Verbrugge et al. noted KIM-1 levels were only slightly elevated at time of hospital admission [64]. Interestingly, KIM-1 is the only injury biomarker that has also been associated with development of incident HF in the Framingham Heart Study and incident HF hospitalization in the Uppsala Longitudinal Study of Adult Men [77, 78].

Prognosis in CRS—In chronic HF, KIM-1 is associated with increased risk for HF hospitalization and all-cause mortality [56, 57]. In the GISSI-HF study, KIM-1 was also associated with the combined endpoint of HF hospitalization and death (adjusted HR=1.13, $p=0.018$), but the adjusted relationship lost significance when these endpoints were examined individually [60]. Alternatively, in ADHF there was no relationship between levels of KIM-1 on admission and all-cause mortality (HR=1.29, $p=0.072$) [64]. Further research is necessary to better understand the prognostic ability of KIM-1 and how/why that differs in chronic versus ADHF.

Diagnosis in CRS—Similar to NAG, KIM-1 is strongly correlated with measures of congestion including NT-proBNP, and elevated in patients requiring higher doses of diuretics [57]. The concentration of KIM-1 increases with diuretic withdrawal and subsequently returns to normal with reinstitution of diuretic therapy [74]. Although this pattern was also described for NAG, the rapidity and degree of fluctuations in KIM-1 were significantly more marked than NAG, particularly in those patients with baseline RD. It is important to note that while levels of KIM-1 were fluctuating, serum creatinine remained relatively unchanged. Increased levels of KIM-1 in chronic HF outpatients is significantly associated with development of WRF over 1 year later when adjusted extensively for baseline characteristics (HR 1.23, 95% CI 1.09-1.39 per log increase, $p<0.001$) [68]. Despite its robust association with WRF in chronic HF, levels of KIM-1 at admission for ADHF did not predict subsequent AKI with treatment nor was it associated with persistent renal impairment following discharge [64]. The sensitivity of KIM-1 to changes in fluid status and diuretic use coupled with its strong association with WRF in chronic HF suggests it may be valuable in phenotyping CRS, although the lack of a relationship between KIM-1 and WRF in decompensated HF may in turn limit its utility and therefore deserves further exploration. It is important to note that whether the changes described in tubular function captured with KIM-1 are actually due to HF has yet to be determined.

Interleukin-18 (IL-18)

IL-18 is cytokine that is produced in mononuclear cells, macrophages and non-immune cells and in its active form, mediates inflammation and ischemic injury in multiple organs

including the heart, lungs and colon [49]. In fact, plasma concentrations of IL-18 are elevated in patients with acute coronary syndrome [79]. In the kidney, IL-18 is produced and released from the proximal convoluted tubule within hours of kidney injury, peaking at 12 hours and remaining in the system for 48 hours [80]. Urinary IL-18 elevations are more specific for ATN than other renal diseases including chronic kidney disease and urinary tract infections [81]. In children undergoing cardiac bypass, increases in IL-18 appeared 6 hours prior to the rise in creatinine and improved the AUC for AKI detection from 0.72 to 0.84 [82]. In adults, patients with the highest quartile of urinary IL-18 prior to cardiac bypass experienced significantly increased mortality in those with and without AKI [83]. Interestingly, circulating levels of IL-18 have been reported in patients with HF and in one small study correlated with disease severity [64, 84, 85].

Prognosis in CRS—In HF, there have been few studies examining IL-18, but Verbrugge et al. examined IL-18 levels in 83 patients admitted with ADHF along with other kidney injury biomarkers. Only IL-18 was significantly associated with all-cause mortality in univariate analysis of this population (HR=1.48, 95% CI 1.16-1.87, $p=0.001$) [64]. Given the strong mortality associations already reported in AKI patients and in patients with coronary artery disease, further investigation into the prognostic potential of this biomarker in CRS is warranted.

Diagnosis of CRS—In ADHF, IL-18 is only modestly correlated with other kidney injury biomarkers, and when corrected for urine creatinine production, demonstrates no correlation with eGFR or serum cystatin C [64]. Although IL-18 did not predict WRF during hospitalization, it was the only biomarker significantly associated with persistent renal impairment following discharge; baseline levels >7 pg/gCr were 68% sensitive and 60% specific.

Neurohormonal Activation and Sodium Avidity

HF arises when progressive pump dysfunction can no longer be buffered by endogenous compensatory mechanisms. One of the best studied of these compensatory mechanisms is upregulation of neurohormonal systems such as renin, angiotensin II, aldosterone, vasopressin, and the sympathetic nervous system [2]. In the kidney, the direct effects of neurohormonal activation are analogous to the homeostatic response to acute blood loss, causing a substantial reduction in renal blood flow, GFR, and increased sodium/fluid avidity. In fact, neurohormonal levels have been shown to correlate better with GFR than with left ventricular ejection fraction in HF [4].

Blood Urea Nitrogen to Creatinine Ratio (BUN/Cr)

Although the BUN/Cr has been widely used clinically to distinguish intrinsic kidney disease from pre-renal RD, it has only recently been employed in HF patients with RD, hence its inclusion here as a novel biomarker [86]. Certainly the individual components of this ratio are well known for their powerful prognostic abilities in HF, however the BUN/Cr provides additional diagnostic and prognostic value particularly in CRS given what it represents [87]. Renal clearance of urea is determined by the amount filtered and the degree of tubular reabsorption. While decreases in eGFR from any etiology can lead to decreased urea

filtration, the tubular reabsorption of urea is largely influenced by neurohormonal activation with increases in angiotensin II and vasopressin leading to higher urea concentration in the proximal tubule and increased urea transporters in the collecting duct respectively, thereby enhancing absorption [88, 89]. As such, in diseases of increased neurohormonal activation, like HF, the BUN is elevated out of proportion to serum creatinine as opposed to intrinsic kidney disease where filtration may be reduced, but tubular reabsorption of urea is preserved yielding a lower BUN/Cr [90, 91].

Prognosis in CRS—Increasing levels of BUN/Cr are associated with increased mortality in chronic and ADHF despite adjustment for eGFR, but the enhanced value of BUN/Cr lies in its differentiation of high-risk forms of RD in HF [92, 93]. In four separate populations, ranging from stable HF outpatients to severely decompensated inpatients, we have demonstrated that in patients with an elevated BUN/Cr the adjusted risk of death attributed to RD ranges from 2.2-4.6 compared to those with normal renal function, but in patients with a low BUN/Cr, the presence of RD had no impact on mortality (p-interaction<0.03 for all; Figure 1) [5, 94]. Patients with elevated BUN/Cr also displayed multiple signs of increased neurohormonal activation. These results suggest that the mechanism of eGFR reduction in HF patients (i.e. CRS versus intrinsic kidney disease) is more important than the RD itself.

Diagnosis in CRS—Unlike the irreversible nephron loss associated with intrinsic kidney disease, RD secondary to CRS is often reversible as evidenced by marked improvements in renal function (IRF) in many patients following treatment for ADHF [95, 96]. Our group previously showed in a cohort of 896 patients with ADHF that an admission BUN/Cr ≥ 20 was independently associated with increased incidence of IRF during the hospitalization despite extensive adjustment for baseline characteristics and eGFR (OR=1.66, 95% CI 1.15-2.41, p=0.007) [94]. These results suggest that an elevated BUN/Cr identifies a potentially reversible form of CRS. Further research is required to delineate whether targeted treatment of these high-risk patients yields subsequent benefit.

Diuretic Resistance / Diuretic Efficiency

The failure to realize a significant diuresis or natriuresis following dosing of a loop diuretic signifies a sodium avid state which by in large is driven by neurohormonal activation [97-100]. Importantly, this renal sodium avidity queries a different dimension of renal function than filtration or structural integrity of components of the renal parenchyma. Recently, our group and others have focused on not just the diuretic dose but the *response* to that dose using metrics which attempt to capture the renal response to the diuretic stimulus: 1. Diuretic efficiency defined as net fluid output in mL/40 mg furosemide equivalents 2. Weight change in kg/40 mg furosemide equivalents and 3. Natriuretic response to continuous IV furosemide defined as urine sodium to urine furosemide ratio ($U_{Na}:U_{Furosemide}$) [101-103].

Prognosis in CRS—In two separate cohorts, we demonstrated that while diuretic efficiency was only modestly correlated with diuretic dose, net urine output, and estimated GFR, it was independently associated with significantly increased mortality; patients with

diuretic efficiency below the median in the ESCAPE trial experienced nearly three times the risk of death compared to those patients with diuretic efficiency above the median despite extensive adjustment for baseline and in-hospital characteristics including net fluid output and diuretic dose (HR=2.86, 95% CI 1.53-5.36, $p=0.001$) [101]. Furthermore, the relationship between low diuretic efficiency and increased mortality was present in both patients on low and high dose diuretics reinforcing that diuretic efficiency captures more than just the diuretic dose. Valente et al. found similar results where patients with less weight loss per 40 mg furosemide equivalents experienced an increased hazard for 60-day HF rehospitalization and 180-day mortality [102]. A final metric of diuretic responsiveness, the $U_{Na}:U_{Furosemide}$ was examined by Singh et al who measured spot urine sodium, creatinine and furosemide in 52 ADHF patients hospitalized on continuous IV furosemide infusion [103]. $U_{Na}:U_{Furosemide}$ was not significantly associated with eGFR or diuretic dose but higher $U_{Na}:U_{Furosemide}$ was associated with 3.6 times the likelihood of increased urinary output and 2.7 times the likelihood of increased weight loss in 24 hours. Patients with $U_{Na}:U_{Furosemide} < 2$ mmol/mg (indicative of low diuretic efficiency) experienced less weight loss and fluid removal in the first 24 hours and were at significantly increased risk for death, HF rehospitalization, and cardiac transplantation in an analysis adjusted for age and eGFR (HR=2.2, 95% CI 1.08-4.49, $p=.032$). The results of the aforementioned studies suggest that metrics of diuretic responsiveness are superior to diuretic dose in identifying HF patients with diuretic resistance who in turn are at increased risk for poor outcomes.

Diagnosis in CRS—Diuretic resistance or low DE in HF is believed to result from a number of different mechanisms. First a significant reduction in GFR can decrease drug delivery to the tubules, but as most loop diuretics are bound to albumin and secreted by the proximal tubular cells, adequate drug delivery is far more dependent on renal blood flow than filtration [97]. Diuretics themselves can lead to increases in activation of RAAS subsequently decreasing renal blood flow and tubular secretion [104]. Other mechanisms of resistance that have been proposed are pharmacodynamic like diuretic “braking” during which acute tolerance develops as means to preserve intravascular volume and hypertrophy of distal tubular cells in response to increased sodium delivery downstream induced by diuretics leading to increased sodium reabsorption and decreased responsiveness even in the setting of normal drug delivery [98]. Due to the fact that most of the proposed mechanisms for diuretic resistance are independent of GFR yet indicative of HF-induced RD, accurate assessment of diuretic resistance may identify patients with significant cardiorenal dysfunction. As an example, we found that the presence of a low eGFR did not exclude the possibility of excellent DE and vice versa [101]. Furthermore, patients with $U_{Na}:U_{Furosemide} < 2$ mmol/mg in the study by Singh et al. were also more likely to experience WRF with treatment, a hallmark of CRS [103]. Further research is necessary to explore these metrics of decreased diuretic responsiveness and whether targeted treatment strategies in these patients are beneficial.

Venous Congestion

One of the chief mechanisms believed to be involved in cardiorenal dysfunction is systemic venous congestion. Data from animal models illustrates how increases in renal vein pressures yield decreased renal blood flow, GFR and sodium excretion, abnormalities which

improve with relief of congestion [105]. Furthermore, positive associations between volume overload and RD in HF patients have only strengthened the potential role of venous congestion in CRS [106, 107]. The inherently complicated interplay between venous congestion and CRS is not the focus of this review, however, those biomarkers which represent or result from congestion may prove extremely useful in deciphering the pathophysiology of CRS while simultaneously aiding in its recognition [108].

Brain Natriuretic Peptide (BNP)

BNP is a hormone produced by the myocardium in response to increased wall stretch and elevated filling pressures. Both BNP and its biologically inactive form, NT-proBNP are elevated in HF and are well-established as diagnostic and prognostic indicators [109, 110]. Increases in NT-proBNP during hospitalization for ADHF are associated with a 3-fold increase in mortality [111]. Similarly, increases in BNP in stable HF patients over four months also predicts poor prognosis [112].

Prognosis in CRS—Possibly due to the common misconception that elevations of BNP and NT-proBNP in patients with RD merely reflect a reduction in eGFR, their use in patients with HF and RD was not extensively explored prior to a study by van Kimmenade and colleagues. In 720 patients presenting with ADHF, both RD (eGFR < 60 ml/min/1.73m²) and NT-proBNP above the median (>4,647 pg/mL) were independently associated with mortality, however; those patients with both RD and elevated NT-proBNP suffered the worst mortality (OR=3.46, 95% CI 2.3-5.6, p<0.001). In fact, patients with RD but lower NT-proBNP levels had a similar prognosis to patients with normal renal function suggesting that NT-proBNP may characterize RD in HF instead of simply reflecting it (Figure 1) [113].

Diagnosis of CRS—In the aforementioned study, van Kimmenade et al. also examined the relationship between WRF (rise in creatinine >0.3 mg/dL) and NT-proBNP during treatment of ADHF [113]. Notably, WRF was in fact associated with decreased 60-day survival, but this survival disadvantage was restricted to those patients with an elevated NT-proBNP. Patients with WRF in the setting of lower NT-proBNP had similar survival to those patients with stable renal function. As WRF in HF is heterogeneous, utilization of NT-proBNP in addition to WRF may further distinguish CRS from other forms of RD in HF.

Biochemical Evidence of Hepatic Dysfunction

Physicians have long been aware of the negative impact of HF on the liver: reduced hepatic perfusion can lead to hepatocellular injury with increases in aspartate aminotransferase (AST) and alanine aminotransferase (ALT) while elevated venous pressure and congestion ultimately result in centrilobular necrosis with increases in bilirubin (BIL) and INR [114-116]. Notably, similar pathophysiologic mechanisms like venous congestion and reduced perfusion are thought to contribute to CRS. As both the kidneys and the liver occupy the same circulatory and congestive environment, congestion and abnormal cardiac output secondary to HF likely affects both organs similarly and simultaneously. Therefore laboratory evidence of hepatic dysfunction (i.e. congestive hepatopathy) may identify patients with concomitant CRS.

Prognosis in CRS—Multiple studies have demonstrated that elevated liver function tests in patients with acute and chronic HF, specifically AST, ALT and bilirubin, are associated with increased mortality and HF rehospitalization, but only bilirubin has retained prognostic ability in adjusted analyses [116-119]. Still, whether markers of liver dysfunction are of heightened prognostic value in patients with RD has yet to be explored.

Diagnosis in CRS—Similar to CRS, the centrilobular necrosis signified by elevated liver function tests and characteristic of HF-induced liver dysfunction is reversible with decongestion [114]. Therefore patients presenting with ADHF and markers of hepatic congestion may also be more likely to experience IRF with return to compensation. We previously demonstrated in a cohort of 823 patients admitted with ADHF that those who presented with elevated AST, ALT, BIL or INR (in those not on warfarin) were significantly more likely to experience IRF (defined as a ≥20% improvement in eGFR) during the hospitalization suggesting the presence of CRS at the time of presentation [120]. The strength of the associations between markers of liver dysfunction and subsequent IRF were much stronger in patients who presented with marked RD (eGFR <45 ml/min/1.73m²). Patients with RD and an elevated BIL at admission were 5 times more likely to experience IRF (OR=5.1, p<0.001) while in patients without RD, an elevated admission BIL was not significantly related to IRF (OR=1.6, p=0.06, p-interaction=0.01). Similar results were observed for an elevated admission INR and ALT. In the absence of clear-cut diagnostic criteria for CRS, using evidence of HF-induced dysfunction of another organ to inform on the likely presence of concomitant CRS deserves further investigation.

Tissue Fibrosis

Renal tissue fibrosis can develop secondary to various modes of kidney injury both acute and chronic, while myocardial fibrosis is a sine qua non of the HF syndrome and cardiac remodeling. As tissue fibrosis is involved in the final common pathway for damage to both organs, biomarkers along these pathways may improve our understanding about the cross-talk between the heart and kidneys in CRS.

Galectin-3

Galectin-3 is a β -galactoside-binding lectin, expressed both intracellularly and extracellularly, where it is involved in cell proliferation, apoptosis, inflammation and cell-growth. In the kidney the role of galectin-3 appears to be protective in AKI, attenuating fibrosis, yet also activates kidney fibrosis in the setting of persistent renal injury [121, 122]. In the myocardium, galectin-3 has binding sites on cardiac fibroblasts and induces their proliferation and ultimately collagen deposition leading to ventricular dysfunction and myocardial fibrosis [123]. Galectin-3 is measured in the blood with levels >17.8 ng/mL considered elevated according to the package insert.

Patients with both chronic and decompensated HF demonstrate similarly increased galectin-3 levels, as do patients with both reduced and preserved EF; there is no correlation between galectin-3 and metrics of disease severity including NYHA functional class [124-127]. Although galectin-3 was helpful in diagnosing suspected acute HF, its performance was inferior in comparison to NT-proBNP as a diagnostic biomarker [125].

Prognosis in CRS—There is a strong relationship between galectin-3 and increased mortality in both stable chronic and decompensated HF patients. In the PRIDE substudy, ADHF patients with elevated admission galectin-3 had 10 times the odds of death in 60 days after adjustment for eGFR and baseline characteristics and Lok et al. reported an adjusted HR of 1.24 for galectin-3 in an outpatient HF cohort of 240 patients [125, 126]. When measured at discharge following a HF exacerbation, galectin-3 has also recently been shown to predict 30, 60, 90 and 120-day risk for HF rehospitalization, significantly improving patient reclassification in readmission models [128]. Fluctuations in galectin-3 are also significantly associated with survival [129]. In a secondary analysis of the Controlled Rosuvastatin Multinational Trial in Heart Failure (CORONA, HF outpatients) and COACH (ADHF patients) trials, patients whose galectin-3 levels increased by >15% over 3-6 months had a significantly increased adjusted risk for all cause-mortality and HF hospitalization (CORONA HR=1.54, $p<0.001$; COACH HR=2.04, $p=0.026$) [130]. Although all the aforementioned studies adjusted their comparisons for renal function, whether RD modifies the relationship between galectin-3 and mortality in HF has yet to be examined. However, Tang et al. reported in a single-center study that the magnitude of the survival disadvantage attributed to increased galectin-3 was more marked in HF patients with concomitant elevations in serum CysC compared to those with normal CysC ($p<0.001$) [127]. Further research is required to delineate prognostic value of galectin-3 in CRS.

Diagnosis of CRS—Given the powerful mortality associations reported for galectin-3 in HF, one implication is that it may serve as a surrogate for kidney function. However, as galectin-3 is predominantly hepatically cleared and is only modestly correlated with eGFR, this is less likely. Furthermore, in over 2,000 subjects in the Framingham Offspring Study, patients with normal renal function (mean eGFR~90 ml/min/1.73m²) and baseline galectin-3 in the highest quartile were at significantly increased odds of developing incident RD defined as an eGFR<60 ml/min/1.73m² (adjusted OR=1.47, 95% CI 1.27-1.71; $p<0.0001$) [131]. Therefore, galectin-3 elevations in this cohort were not merely reflecting abnormal filtration. Despite the strong correlations between galectin-3 with incident RD, to date, there are no studies investigating an association between galectin-3 and WRF in HF, leaving its potential to distinguish HF-induced RD unexplored.

Undefined Mechanism

C-Type Natriuretic Peptide (CNP)

CNP is member of the natriuretic peptide family which, similar to BNP, results from cleavage of a precursor and subsequent post processing that yields the mature, active form, CNP22. Although CNP has been detected in cardiac tissue, it is predominantly a vascular endothelial product with vasodilatory properties that indirectly lead to decreased cardiac filling pressures [132]. Unlike BNP and ANP, CNP has much less of a diuretic effect, and a very brief circulatory half-life, potentially limiting its use as a biomarker in the serum. However, CNP is also produced and processed in the kidney, and is detectable in the urine, offering a different method in which measuring CNP may prove beneficial [133]. Given that the stimulation for CNP release in the kidney in heart failure is not completely understood,

CNP's potential role as a CRS biomarker is described here without a specific CRS mechanistic context.

In symptomatic HF patients, plasma concentrations of both CNP and NT-proCNP are significantly elevated compared to controls and are associated with HF disease severity expressed by NYHA functional class and EF [134]. Interestingly, CNP levels are also increased in patients with asymptomatic LV dysfunction [135]. Although elevations of NT-proCNP increased the accuracy diagnosing HF above BNP alone, it was not superior as a diagnostic tool [136].

Prognosis in CRS—Given that CNP is also produced by the renal tubules and correlated with serum creatinine, elevated urinary levels of CNP and NT-proCNP may indicate both renal tubular damage and congestion thereby providing unique prognostic information in CRS over natriuretic peptides or other tubular injury markers alone. In a study of 58 ADHF patients, Zakeri et al. demonstrated that urinary NT-proCNP was independently associated with all-cause rehospitalization and death and outperformed both KIM-1 and NGAL as mortality predictors [137]. Serum CNP was also elevated in patients with ADHF, but was not correlated with urinary forms of CNP and showed no association with mortality or rehospitalization.

Diagnosis of CRS—Although all isoforms of CNP are elevated in the serum and urine of ADHF patients, it has not been studied specifically in HF patients with RD. Urinary CNP may in fact serve as a partial marker of tubular injury, but until it is further explored in normal patients and patients with other forms of RD, its capacity to distinguish CRS is unknown.

Conclusions

As described in this review, there is a growing literature of promising and novel biomarkers for cardio-renal dysfunction. Although the focus of this research thus far has centered on prognostication, the currently available and emerging biomarkers in this area query a diverse spectrum of mechanistic pathways and biology. As a result, it is likely that through continued research in this area, pathways to differentiate specific mechanism for renal dysfunction ultimately allowing patient specific therapeutic approaches will be possible.

REFERENCES

Papers of particular interest, published recently, have been highlighted as:

- Of importance
 - Of major importance
1. Cardio-renal connections in heart failure and cardiovascular disease, national heart-lung and blood institute working group report. 2004
 2. Braunwald, E.; Bonow, RO. Braunwald's heart disease : a textbook of cardiovascular medicine. 9th ed.. Saunders; Philadelphia: 2012.
 3. Bock JS, Gottlieb SS. Cardiorenal syndrome: new perspectives. *Circulation*. 2010; 121:2592–600. [PubMed: 20547939]

4. Hillege HL, Girbes ARJ, de Kam PJ, Boomsma F, de Zeeuw D, Charlesworth A, et al. Renal Function, Neurohormonal Activation, and Survival in Patients With Chronic Heart Failure. *Circulation*. 2000; 102:203–10. [PubMed: 10889132]
5. Testani JM, Coca SG, Shannon RP, Kimmel SE, Cappola TP. Influence of renal dysfunction phenotype on mortality in the setting of cardiac dysfunction: analysis of three randomized controlled trials. *European Journal of Heart Failure*. 2011; 13:1224–30. [PubMed: 21926073]
6. Heywood JT, Fonarow GC, Costanzo MR, Mathur VS, Wigneswaran JR, Wynne J, et al. High prevalence of renal dysfunction and its impact on outcome in 118,465 patients hospitalized with acute decompensated heart failure: a report from the ADHERE database. *J Card Fail*. 2007; 13:422–30. [PubMed: 17675055]
7. Ronco C, Ciccoira M, McCullough PA. Cardiorenal syndrome type 1: pathophysiological crosstalk leading to combined heart and kidney dysfunction in the setting of acutely decompensated heart failure. *Journal of the American College of Cardiology*. 2012; 60:1031–42. [PubMed: 22840531]
8. Testani JM, Kimmel SE, Dries DL, Coca SG. Prognostic importance of early worsening renal function after initiation of angiotensin-converting enzyme inhibitor therapy in patients with cardiac dysfunction. *Circulation Heart failure*. 2011; 4:685–91. [PubMed: 21903907]
9. Testani JM, Chen J, McCauley BD, Kimmel SE, Shannon RP. Potential effects of aggressive decongestion during the treatment of decompensated heart failure on renal function and survival. *Circulation*. 2010; 122:265–72. [PubMed: 20606118]
10. Testani JM, Brisco MA, Han G, Laur O, Kula AJ, Cheng SJ, et al. Influence of age-related versus nonage-related renal dysfunction on survival in patients with left ventricular dysfunction. *Am J Cardiol*. 2014; 113:127–31. [PubMed: 24216124]
11. Fonarow GC, Heywood JT. The confounding issue of comorbid renal insufficiency. *Am J Med*. 2006; 119:S17–25. [PubMed: 17113396]
12. Chen HH, Anstrom KJ, Givertz MM, Stevenson LW, Semigran MJ, Goldsmith SR, et al. Low-dose dopamine or low-dose nesiritide in acute heart failure with renal dysfunction: the ROSE acute heart failure randomized trial. *JAMA: The Journal of the American Medical Association*. 2013; 310:2533–43.
13. Massie BM, O'Connor CM, Metra M, Ponikowski P, Teerlink JR, Cotter G, et al. Rolofylline, an adenosine A1-receptor antagonist, in acute heart failure. *N Engl J Med*. 2010; 363:1419–28. [PubMed: 20925544]
14. O'Connor CM, Starling RC, Hernandez AF, Armstrong PW, Dickstein K, Hasselblad V, et al. Effect of nesiritide in patients with acute decompensated heart failure. *N Engl J Med*. 2011; 365:32–43. [PubMed: 21732835]
15. Gheorghade M, Konstam MA, Burnett JC Jr, Grinfeld L, Maggioni AP, Swedberg K, et al. Short-term clinical effects of tolvaptan, an oral vasopressin antagonist, in patients hospitalized for heart failure: the EVEREST Clinical Status Trials. *JAMA : the journal of the American Medical Association*. 2007; 297:1332–43.
16. Bart BA, Goldsmith SR, Lee KL, Givertz MM, O'Connor CM, Bull DA, et al. Ultrafiltration in decompensated heart failure with cardiorenal syndrome. *N Engl J Med*. 2012; 367:2296–304. [PubMed: 23131078]
17. Smilde TD, van Veldhuisen DJ, Navis G, Voors AA, Hillege HL. Drawbacks and prognostic value of formulas estimating renal function in patients with chronic heart failure and systolic dysfunction. *Circulation*. 2006; 114:1572–80. [PubMed: 17015793]
18. Knight EL, Verhave JC, Spiegelman D, Hillege HL, de Zeeuw D, Curhan GC, et al. Factors influencing serum cystatin C levels other than renal function and the impact on renal function measurement. *Kidney Int*. 2004; 65:1416–21. [PubMed: 15086483]
19. Dupont M, Shrestha K, Singh D, Finucan M, Tang WHW. Lack of concordance in defining worsening renal function by rise in creatinine vs rise in cystatin C. *Congestive heart failure (Greenwich, Conn)*. 2013; 19:E17–21.
- 20•. Shlipak MG, Matsushita K, Ärnlöv J, Inker LA, Katz R, Polkinghorne KR, et al. Cystatin C versus creatinine in determining risk based on kidney function. *The New England journal of medicine*. 2013; 369:932–43. [PubMed: 24004120] [This is a meta-analysis of over 90,000

patients illustrating that using cystatin C to estimate renal function strengthens the association between renal dysfunction and mortality.]

21. Damman K, van der Harst P, Smilde TDJ, Voors AA, Navis G, van Veldhuisen DJ, et al. Use of cystatin C levels in estimating renal function and prognosis in patients with chronic systolic heart failure. *Heart (British Cardiac Society)*. 2012; 98:319–24. [PubMed: 22038544]
22. Zamora E, Lupón J, Vila J, Urrutia A, de Antonio M, Sanz H, et al. Estimated glomerular filtration rate and prognosis in heart failure: value of the Modification of Diet in Renal Disease Study-4, chronic kidney disease epidemiology collaboration, and cockcroft-gault formulas. *Journal of the American College of Cardiology*. 2012; 59:1709–15. [PubMed: 22554602]
23. Laterza OF, Price CP, Scott MG. Cystatin C: an improved estimator of glomerular filtration rate? *Clinical chemistry*. 2002; 48:699–707. [PubMed: 11978596]
24. Shlipak MG, Sarnak MJ, Katz R, Fried LF, Seliger SL, Newman AB, et al. Cystatin C and the risk of death and cardiovascular events among elderly persons. *The New England journal of medicine*. 2005; 352:2049–60. [PubMed: 15901858]
25. Shlipak MG, Katz R, Fried LF, Jenny NS, Stehman-Breen CO, Newman AB, et al. Cystatin-C and mortality in elderly persons with heart failure. *Journal of the American College of Cardiology*. 2005; 45:268–71. [PubMed: 15653026]
26. Tang WHW, Van Lente F, Shrestha K, Troughton RW, Francis GS, Tong W, et al. Impact of myocardial function on cystatin C measurements in chronic systolic heart failure. *Journal of Cardiac Failure*. 2008; 14:394–9. [PubMed: 18514931]
27. Zamora E, Lupón J, de Antonio M, Vila J, Galán A, Gastelurrutia P, et al. Limited value of cystatin-C over estimated glomerular filtration rate for heart failure risk stratification. *PLoS ONE*. 2012; 7:e51234. [PubMed: 23240006]
28. Dupont M, Wu Y, Hazen SL, Tang WHW. Cystatin C identifies patients with stable chronic heart failure at increased risk for adverse cardiovascular events. *Circulation: Heart Failure*. 2012; 5:602–9. [PubMed: 22899766] [This is a study in chronic heart failure patients showing cystatin C improves estimation of risk, particularly in those with normal renal function.]
29. Lassus J, Harjola V-P, Sund R, Siirilä-Waris K, Melin J, Peuhkurinen K, et al. Prognostic value of cystatin C in acute heart failure in relation to other markers of renal function and NT-proBNP. *European Heart Journal*. 2007; 28:1841–7. [PubMed: 17289743]
30. Arimoto T, Takeishi Y, Niizeki T, Takabatake N, Okuyama H, Fukui A, et al. Cystatin C, a novel measure of renal function, is an independent predictor of cardiac events in patients with heart failure. *Journal of Cardiac Failure*. 2005; 11:595–601. [PubMed: 16230262]
31. Naruse H, Ishii J, Kawai T, Hattori K, Ishikawa M, Okumura M, et al. Cystatin C in acute heart failure without advanced renal impairment. *Am J Med*. 2009; 122:566–73. [PubMed: 19393984]
32. Manzano-Fernández S, Boronat-Garcia M, Albaladejo-Otón MD, Pastor P, Garrido IP, Pastor-Pérez FJ, et al. Complementary prognostic value of cystatin C, N-terminal pro-B-type natriuretic Peptide and cardiac troponin T in patients with acute heart failure. *The American Journal of Cardiology*. 2009; 103:1753–9. [PubMed: 19539088]
33. Valente MAE, Hillege HL, Navis G, Voors AA, Dunselman PHJM, van Veldhuisen DJ, et al. The Chronic Kidney Disease Epidemiology Collaboration equation outperforms the Modification of Diet in Renal Disease equation for estimating glomerular filtration rate in chronic systolic heart failure. *European Journal of Heart Failure*. 2014; 16:86–94. [PubMed: 23901055] [This study compared the accuracy and prognostic value of estimating equations of glomerular filtration rate in heart failure patients to iothalamate clearance.]
34. Taal MW.; Brenner, BM.; Rector, FC. Brenner & Rector's the kidney. 9th ed.. Elsevier/Saunders; Philadelphia, PA: 2012.
35. Dickson LE, Wagner MC, Sandoval RM, Molitoris BA. The proximal tubule and albuminuria: really! *J Am Soc Nephrol*. 2014; 25:443–53. [PubMed: 24408874]
36. Agewall S, Wikstrand J, Ljungman S, Fagerberg B. Usefulness of microalbuminuria in predicting cardiovascular mortality in treated hypertensive men with and without diabetes mellitus. Risk Factor Intervention Study Group. *Am J Cardiol*. 1997; 80:164–9. [PubMed: 9230153]

37. Hillege HL, Fidler V, Diercks GF, van Gilst WH, de Zeeuw D, van Veldhuisen DJ, et al. Urinary albumin excretion predicts cardiovascular and noncardiovascular mortality in general population. *Circulation*. 2002; 106:1777–82. [PubMed: 12356629]
38. Jackson CE, MacDonald MR, Petrie MC, Solomon SD, Pitt B, Latini R, et al. Associations of albuminuria in patients with chronic heart failure: findings in the ALiskiren Observation of heart Failure Treatment study. *European Journal of Heart Failure*. 2014; 13:746–54. [PubMed: 21459891] [This secondary analysis of the ALOFT trial illustrated that patients with albuminuria without concomitant diabetes had indices consistent with increased venous congestion.]
39. Masson S, Latini R, Milani V, Moretti L, Rossi MG, Carbonieri E, et al. Prevalence and prognostic value of elevated urinary albumin excretion in patients with chronic heart failure: data from the GISSI-Heart Failure trial. *Circulation Heart failure*. 2010; 3:65–72. [PubMed: 19850697]
40. van de Wal RMA, Asselbergs FW, Plokker HWT, Smilde TDJ, Lok D, van Veldhuisen DJ, et al. High prevalence of microalbuminuria in chronic heart failure patients. *Journal of Cardiac Failure*. 2005; 11:602–6. [PubMed: 16230263]
41. Smilde TDJ, Damman K, Harst P, Navis G, Daan Westenbrink B, Voors AA, et al. Differential associations between renal function and “modifiable” risk factors in patients with chronic heart failure. *Clinical Research in Cardiology*. 2008; 98:121–9. [PubMed: 18979056]
42. Anand IS, Bishu K, Rector TS, Ishani A, Kuskowski MA, Cohn JN. Proteinuria, chronic kidney disease, and the effect of an angiotensin receptor blocker in addition to an angiotensin-converting enzyme inhibitor in patients with moderate to severe heart failure. *Circulation*. 2009; 120:1577–84. [PubMed: 19805651]
43. Koyama S, Sato Y, Tanada Y, Fujiwara H, Takatsu Y. Early evolution and correlates of urine albumin excretion in patients presenting with acutely decompensated heart failure. *Circulation: Heart Failure*. 2013; 6:227–32. [PubMed: 23395932] [This single-center study established a form of albuminuria secondary to heart failure that improves with decongestion, suggesting its high utility as a marker of cardiorenal dysfunction.]
44. Jackson CE, Solomon SD, Gerstein HC, Zetterstrand S, Olofsson B, Michelson EL, et al. Albuminuria in chronic heart failure: prevalence and prognostic importance. *Lancet*. 2009; 374:543–50. [PubMed: 19683640]
45. Testani JM, McCauley BD, Kimmel SE, Shannon RP. Characteristics of patients with improvement or worsening in renal function during treatment of acute decompensated heart failure. *Am J Cardiol*. 2010; 106:1763–9. [PubMed: 21055713]
46. Testani JM, Brisco MA, Chen J, McCauley BD, Parikh CR, Tang WH. Timing of hemoconcentration during treatment of acute decompensated heart failure and subsequent survival: importance of sustained decongestion. *J Am Coll Cardiol*. 2013; 62:516–24. [PubMed: 23747773]
47. Testani JM, Cappola TP, McCauley BD, Chen J, Shen J, Shannon RP, et al. Impact of worsening renal function during the treatment of decompensated heart failure on changes in renal function during subsequent hospitalization. *Am Heart J*. 2011; 161:944–9. [PubMed: 21570527]
48. Testani JM, Coca SG, McCauley BD, Shannon RP, Kimmel SE. Impact of changes in blood pressure during the treatment of acute decompensated heart failure on renal and clinical outcomes. *Eur J Heart Fail*. 2011; 13:877–84. [PubMed: 21693504]
49. Charlton JR, Portilla D, Okusa MD. A basic science view of acute kidney injury biomarkers. *Nephrology, Dialysis, Transplantation*. 2014; 29:1301–11.
50. Shrestha K, Borowski AG, Troughton RW, Klein AL, Tang WHW. Association between systemic neutrophil gelatinase-associated lipocalin and anemia, relative hypochromia, and inflammation in chronic systolic heart failure. *Congestive heart failure (Greenwich, Conn)*. 2012; 18:239–44.
51. Mishra J, Dent C, Tarabishi R, Mitsnefes MM, Ma Q, Kelly C, et al. Neutrophil gelatinase-associated lipocalin (NGAL) as a biomarker for acute renal injury after cardiac surgery. *Lancet*. 2005; 365:1231–8. [PubMed: 15811456]
52. Mishra J, Ma Q, Prada A, Mitsnefes M, Zahedi K, Yang J, et al. Identification of neutrophil gelatinase-associated lipocalin as a novel early urinary biomarker for ischemic renal injury. *J Am Soc Nephrol*. 2003; 14:2534–43. [PubMed: 14514731]
53. Schmidt-Ott KM. Neutrophil gelatinase-associated lipocalin as a biomarker of acute kidney injury--where do we stand today? *Nephrology, Dialysis, Transplantation*. 2011; 26:762–4.

54. Damman K, van Veldhuisen DJ, Navis G, Voors AA, Hillege HL. Urinary neutrophil gelatinase associated lipocalin (NGAL), a marker of tubular damage, is increased in patients with chronic heart failure. *European journal of heart failure : journal of the Working Group on Heart Failure of the European Society of Cardiology*. 2008; 10:997–1000.
55. van Deursen VM, Damman K, Voors AA, van der Wal MH, Jaarsma T, van Veldhuisen DJ, et al. Prognostic Value of Plasma Neutrophil Gelatinase–Associated Lipocalin for Mortality in Patients With Heart Failure. *Circulation: Heart Failure*. 2014; 7:35–42. [PubMed: 24347663]
56. Damman K, van Veldhuisen DJ, Navis G, Vaidya VS, Smilde TDJ, Westenbrink BD, et al. Tubular damage in chronic systolic heart failure is associated with reduced survival independent of glomerular filtration rate. *Heart (British Cardiac Society)*. 2010; 96:1297–302. [PubMed: 20659949]
57. Jungbauer CG, Birner C, Jung B, Buchner S, Lubnow M, von Bary C, et al. Kidney injury molecule-1 and N-acetyl- β -D-glucosaminidase in chronic heart failure: possible biomarkers of cardiorenal syndrome. *European Journal of Heart Failure*. 2014; 13:1104–10. [PubMed: 21846754]
58. Shrestha K, Shao Z, Singh D, Dupont M, Tang WHW. Relation of systemic and urinary neutrophil gelatinase-associated lipocalin levels to different aspects of impaired renal function in patients with acute decompensated heart failure. *The American Journal of Cardiology*. 2012; 110:1329–35. [PubMed: 22835414]
59. Shrestha K, Borowski AG, Troughton RW, Thomas JD, Klein AL, Tang WH. Renal dysfunction is a stronger determinant of systemic neutrophil gelatinase-associated lipocalin levels than myocardial dysfunction in systolic heart failure. *J Card Fail*. 2011; 17:472–8. [PubMed: 21624735]
60. Damman K, Masson S, Hillege HL, Maggioni AP, Voors AA, Opasich C, et al. Clinical outcome of renal tubular damage in chronic heart failure. *European Heart Journal*. 2011; 32:2705–12. [PubMed: 21666249]
61. Aghel A, Shrestha K, Mullens W, Borowski A, Tang WH. Serum neutrophil gelatinase-associated lipocalin (NGAL) in predicting worsening renal function in acute decompensated heart failure. *J Card Fail*. 2010; 16:49–54. [PubMed: 20123318]
62. Maisel AS, Mueller C, Fitzgerald R, Brikhan R, Hiestand BC, Iqbal N, et al. Prognostic utility of plasma neutrophil gelatinase-associated lipocalin in patients with acute heart failure: the NGAL Evaluation Along with B-type NaTriuretic Peptide in acutely decompensated heart failure (GALLANT) trial. *European Journal of Heart Failure*. 2011; 13:846–51. [PubMed: 21791540]
63. Palazzuoli A, Ruocco G, Beltrami M, Franci B, Pellegrini M, Lucani B, et al. Admission plasma neutrophil gelatinase associated lipocalin (NGAL) predicts worsening renal function during hospitalization and post discharge outcome in patients with acute heart failure. *Acute Card Care*. 2014
64. Verbrugge FH, Dupont M, Shao Z, Shrestha K, Singh D, Finucan M, et al. Novel urinary biomarkers in detecting acute kidney injury, persistent renal impairment, and all-cause mortality following decongestive therapy in acute decompensated heart failure. *Journal of Cardiac Failure*. 2013; 19:621–8. [PubMed: 24054338] [This illustrates the predictive potential of urinary biomarkers for kidney injury in decompensated heart failure.]
65. Alvelos M, Pimentel R, Pinho E, Gomes A, Lourenço P, Teles MJ, et al. Neutrophil gelatinase-associated lipocalin in the diagnosis of type 1 cardio-renal syndrome in the general ward. *Clinical Journal of the American Society of Nephrology*. 2011; 6:476–81. [PubMed: 21115620]
66. Macdonald S, Arendts G, Nagree Y, Xu X-F. Neutrophil Gelatinase-Associated Lipocalin (NGAL) predicts renal injury in acute decompensated cardiac failure: a prospective observational study. *BMC cardiovascular disorders*. 2012; 12:8. [PubMed: 22340728]
67. Mortara A, Bonadies M, Mazzetti S, Fracchioni I, Delfino P, Chioffi M, et al. Neutrophil gelatinase-associated lipocalin predicts worsening of renal function in acute heart failure: methodological and clinical issues. *Journal of cardiovascular medicine (Hagerstown, Md)*. 2013; 14:629–34.
68. Damman K, Masson S, Hillege HL, Voors AA, van Veldhuisen DJ, Rossignol P, et al. Tubular damage and worsening renal function in chronic heart failure. *JACC: Heart Failure*. 2013; 1:417–

24. [PubMed: 24621973] [This secondary analysis of the GISSI-HF study suggests renal biomarkers may help predict worsening renal function in heart failure.]
69. Dupont M, Shrestha K, Singh D, Awad A, Kovach C, Scarpino M, et al. Lack of significant renal tubular injury despite acute kidney injury in acute decompensated heart failure. *European Journal of Heart Failure*. 2012; 14:597–604. [PubMed: 22505396] [Significant worsening in renal function with diuresis was associated with increases in urinary NGAL but not the magnitude of other forms of kidney injury suggesting different pathophysiology of renal dysfunction in heart failure.]
70. Testani JM, Tang WH. Biomarkers of acute kidney injury in chronic heart failure: what do the signals mean? *JACC Heart failure*. 2013; 1:425–6. [PubMed: 24621974]
71. Waring WS, Moonie A. Earlier recognition of nephrotoxicity using novel biomarkers of acute kidney injury. *Clinical toxicology*. 2011; 49:720–8. [PubMed: 21970770]
72. Han WK, Wagener G, Zhu Y, Wang S, Lee HT. Urinary biomarkers in the early detection of acute kidney injury after cardiac surgery. *Clinical Journal of the American Society of Nephrology*. 2009; 4:873–82. [PubMed: 19406962]
73. Liangos O, Perianayagam MC, Vaidya VS, Han WK, Wald R, Tighiouart H, et al. Urinary N-acetyl-beta-(D)-glucosaminidase activity and kidney injury molecule-1 level are associated with adverse outcomes in acute renal failure. *J Am Soc Nephrol*. 2007; 18:904–12. [PubMed: 17267747]
74. Damman K, Ng Kam Chuen MJ, MacFadyen RJ, Lip GYH, Gaze D, Collinson PO, et al. Volume status and diuretic therapy in systolic heart failure and the detection of early abnormalities in renal and tubular function. *Journal of the American College of Cardiology*. 2011; 57:2233–41. [PubMed: 21616283]
75. Bonventre JV, Yang L. Kidney injury molecule-1. *Current opinion in critical care*. 2010; 16:556–61. [PubMed: 20930626]
76. Vaidya VS, Ozer JS, Dieterle F, Collings FB, Ramirez V, Troth S, et al. Kidney injury molecule-1 outperforms traditional biomarkers of kidney injury in preclinical biomarker qualification studies. *Nature biotechnology*. 2010; 28:478–85.
77. O'Seaghda CM, Hwang S-J, Larson MG, Meigs JB, Vasan RS, Fox CS. Analysis of a urinary biomarker panel for incident kidney disease and clinical outcomes. *J Am Soc Nephrol*. 2013; 24:1880–8. [PubMed: 23990678]
78. Carlsson AC, Larsson A, Helmersson-Karlqvist J, Lind L, Ingelsson E, Larsson TE, et al. Urinary kidney injury molecule 1 and incidence of heart failure in elderly men. *European Journal of Heart Failure*. 2013; 15:441–6. [PubMed: 23220287]
79. Mallat Z, Henry P, Fressonnet R, Alouani S, Scoazec A, Beaufils P, et al. Increased plasma concentrations of interleukin-18 in acute coronary syndromes. *Heart (British Cardiac Society)*. 2002; 88:467–9. [PubMed: 12381634]
80. Parikh CR, Mishra J, Thiessen-Philbrook H, Dursun B, Ma Q, Kelly C, et al. Urinary IL-18 is an early predictive biomarker of acute kidney injury after cardiac surgery. *Kidney Int*. 2006; 70:199–203. [PubMed: 16710348]
81. Parikh CR, Jani A, Melnikov VY, Faubel S, Edelstein CL. Urinary interleukin-18 is a marker of human acute tubular necrosis. *American Journal of Kidney Diseases*. 2004; 43:405–14. [PubMed: 14981598]
82. Krawczeski CD, Goldstein SL, Woo JG, Wang Y, Piyaphanee N, Ma Q, et al. Temporal relationship and predictive value of urinary acute kidney injury biomarkers after pediatric cardiopulmonary bypass. *Journal of the American College of Cardiology*. 2011; 58:2301–9. [PubMed: 22093507]
83. Coca SG, Garg AX, Thiessen-Philbrook H, Koyner JL, Patel UD, Krumholz HM, et al. Urinary Biomarkers of AKI and Mortality 3 Years after Cardiac Surgery. *J Am Soc Nephrol*. 2014; 25:1063–71. [PubMed: 24357673]
84. Naito Y, Tsujino T, Fujioka Y, Ohyanagi M, Okamura H, Iwasaki T. Increased circulating interleukin-18 in patients with congestive heart failure. *Heart (British Cardiac Society)*. 2002; 88:296–7. [PubMed: 12181230]

85. Driver TH, Katz R, Ix JH, Magnani JW, Peralta CA, Parikh CR, et al. Urinary Kidney Injury Molecule 1 (KIM-1) and Interleukin 18 (IL-18) as Risk Markers for Heart Failure in Older Adults: The Health, Aging, and Body Composition (Health ABC) Study. *American Journal of Kidney Diseases : The Official Journal of the National Kidney Foundation*. 2014; 64:49–56. [PubMed: 24656453]
86. Cecil, RL.; Goldman, L.; Schafer, AI. *Goldman's Cecil medicine*. 24th ed.. Elsevier/Saunders/; Philadelphia: 2012.
87. Fonarow GC, Adams KF, Abraham WT, Yancy CW, Boscardin WJ, ADHERE Scientific Advisory Committee SG, and Investigators. Risk stratification for in-hospital mortality in acutely decompensated heart failure: classification and regression tree analysis. *JAMA: The Journal of the American Medical Association*. 2005; 293:572–80.
88. Fenton RA. Essential role of vasopressin-regulated urea transport processes in the mammalian kidney. *Pflugers Archiv : European journal of physiology*. 2009; 458:169–77. [PubMed: 19011892]
89. Kazory A. Emergence of blood urea nitrogen as a biomarker of neurohormonal activation in heart failure. *Am J Cardiol*. 2010; 106:694–700. [PubMed: 20723648]
90. Schrier RW. Blood Urea Nitrogen and Serum Creatinine: Not Married in Heart Failure. *Circulation: Heart Failure*. 2008; 1:2–5. [PubMed: 19808263]
91. Lindenfeld J, Schrier RW. Blood urea nitrogen a marker for adverse effects of loop diuretics? *Journal of the American College of Cardiology*. 2011; 58:383–5. [PubMed: 21757115]
92. Lin H-J, Chao C-L, Chien K-L, Ho Y-L, Lee C-M, Lin Y-H, et al. Elevated blood urea nitrogen-to-creatinine ratio increased the risk of hospitalization and all-cause death in patients with chronic heart failure. *Clinical Research in Cardiology*. 2009; 98:487–92. [PubMed: 19468780]
93. Aronson D, Mittleman MA, Burger AJ. Elevated blood urea nitrogen level as a predictor of mortality in patients admitted for decompensated heart failure. *The American Journal of Medicine*. 2004; 116:466–73. [PubMed: 15047036]
- 94•. Brisco MA, Coca SG, Chen J, Owens AT, McCauley BD, Kimmel SE, et al. Blood urea nitrogen/creatinine ratio identifies a high-risk but potentially reversible form of renal dysfunction in patients with decompensated heart failure. *Circulation Heart failure*. 2013; 6:233–9. [PubMed: 23325460] [Suggests that BUN/Cr may be useful in differentiating forms of renal dysfunction in heart failure.]
95. Testani JM, Khera AV, St John Sutton MG, Keane MG, Wieggers SE, Shannon RP, et al. Effect of Right Ventricular Function and Venous Congestion on Cardio-Renal Interactions during the Treatment of Decompensated Heart Failure. *The American Journal of Cardiology*. 2010; 105:511–6. [PubMed: 20152246]
96. Testani JM, McCauley BD, Chen J, Coca SG, Cappola TP, Kimmel SE. Clinical characteristics and outcomes of patients with improvement in renal function during the treatment of decompensated heart failure. *J Card Fail*. 2011; 17:993–1000. [PubMed: 22123361]
97. Brater DC, Seiwel R, Anderson S, Burdette A, Dehmer GJ, Chennavasin P. Absorption and disposition of furosemide in congestive heart failure. *Kidney Int*. 1982; 22:171–6. [PubMed: 7132061]
98. Brater DC. Update in diuretic therapy: clinical pharmacology. *Seminars in Nephrology*. 2011; 31:483–94. [PubMed: 22099505]
99. Felker GM, Mentz RJ. Diuretics and ultrafiltration in acute decompensated heart failure. *Journal of the American College of Cardiology*. 2012; 59:2145–53. [PubMed: 22676934]
100. Braunwald E. Responsiveness to loop diuretics in heart failure. *European Heart Journal*. 2014; 35:1235–7. [PubMed: 24685713]
- 101•. Testani JM, Brisco MA, Turner JM, Spatz ES, Bellumkonda L, Parikh CR, et al. Loop diuretic efficiency: a metric of diuretic responsiveness with prognostic importance in acute decompensated heart failure. *Circulation: Heart Failure*. 2014; 7:261–70. [PubMed: 24379278] [This is an analysis of two decompensated heart failure cohorts suggesting diuretic efficiency may serve as a metric of diuretic resistance with distinct prognostic information, further identifying cardiorenal dysfunction.]

102. Valente MAE, Voors AA, Damman K, van Veldhuisen DJ, Massie BM, O'Connor CM, et al. Diuretic response in acute heart failure: clinical characteristics and prognostic significance. *European Heart Journal*. 2014; 35:1284–93. [PubMed: 24585267]
103. Singh D, Shrestha K, Testani JM, Verbrugge FH, Dupont M, Mullens W, et al. Insufficient natriuretic response to continuous intravenous furosemide is associated with poor long-term outcomes in acute decompensated heart failure. *Journal of Cardiac Failure*. 2014; 20:392–9. [PubMed: 24704538] [This is a single-center study describing the prognostic and potentially therapeutic utility of urine sodium measurements during diuretic treatment.]
104. Mulder H, Schopman W Jr. van der Lely AJ, Schopman W Sr. Acute changes in plasma renin activity, plasma aldosterone concentration and plasma electrolyte concentrations following furosemide administration in patients with congestive heart failure--interrelationships and diuretic response. *Hormone and metabolic research*. 1987; 19:80–3. [PubMed: 3549504]
105. Winton FR. The influence of venous pressure on the isolated mammalian kidney. *The Journal of physiology*. 1931; 72:49–61. [PubMed: 16994199]
106. Testani JM, Khera AV, St John Sutton MG, Keane MG, Wiegers SE, Shannon RP, et al. Effect of right ventricular function and venous congestion on cardiorenal interactions during the treatment of decompensated heart failure. *Am J Cardiol*. 2010; 105:511–6. [PubMed: 20152246]
107. Damman K, Navis G, Smilde TDJ, Voors AA, van der Bij W, van Veldhuisen DJ, et al. Decreased cardiac output, venous congestion and the association with renal impairment in patients with cardiac dysfunction. *European journal of heart failure : journal of the Working Group on Heart Failure of the European Society of Cardiology*. 2007; 9:872–8.
108. Testani JM, Damman K. Venous congestion and renal function in heart failure ... it's complicated. *Eur J Heart Fail*. 2013; 15:599–601. [PubMed: 23603089]
109. Rothenburger M, Wichter T, Schmid C, Stypmann J, Tjan TD, Berendes E, et al. Aminoterminal pro type B natriuretic peptide as a predictive and prognostic marker in patients with chronic heart failure. *The Journal of heart and lung transplantation : the official publication of the International Society for Heart Transplantation*. 2004; 23:1189–97.
110. Hartmann F, Packer M, Coats AJS, Fowler MB, Krum H, Mohacsi P, et al. Prognostic impact of plasma N-terminal pro-brain natriuretic peptide in severe chronic congestive heart failure: a substudy of the Carvedilol Prospective Randomized Cumulative Survival (COPERNICUS) trial. *Circulation*. 2004; 110:1780–6. [PubMed: 15381643]
111. Bettencourt P, Azevedo A, Pimenta J, Friões F, Ferreira S, Ferreira A. N-terminal-pro-brain natriuretic peptide predicts outcome after hospital discharge in heart failure patients. *Circulation*. 2004; 110:2168–74. [PubMed: 15451800]
112. Latini R, Masson S, Wong M, Barlera S, Carretta E, Staszewsky L, et al. Incremental prognostic value of changes in B-type natriuretic peptide in heart failure. *Am J Med*. 2006; 119:70, e23–30. [PubMed: 16431191]
113. van Kimmenade RR, Januzzi J, James L, Ellinor PT, Sharma UC, Bakker JA, Low AF, et al. Utility of Amino-Terminal Pro-Brain Natriuretic Peptide, Galectin-3, and Apelin for the Evaluation of Patients With Acute Heart Failure. *Journal of the American College of Cardiology*. 2006; 48:1217–24. [PubMed: 16979009]
114. Sherlock S. The liver in heart failure; relation of anatomical, functional, and circulatory changes. *British Heart Journal*. 1951; 13:273–93. [PubMed: 14848381]
115. Giallourakis CC, Rosenberg PM, Friedman LS. The liver in heart failure. *Clinics in liver disease*. 2002; 6:947–67, viii-ix. [PubMed: 12516201]
116. van Deursen VM, K Damman MDP, H L Hillege MDP, A P Van Beek MDP, D J Van Veldhuisen MDP, A A Voors MDP. Abnormal Liver Function in Relation to Hemodynamic Profile in Heart Failure Patients. *Journal of Cardiac Failure*. 2010; 16:84–90. [PubMed: 20123323]
117. Kim MS, Kato TS, Farr M, Wu C, Givens RC, Collado E, et al. Hepatic dysfunction in ambulatory patients with heart failure: application of the MELD scoring system for outcome prediction. *Journal of the American College of Cardiology*. 2013; 61:2253–61. [PubMed: 23563127]

- 118•. van Deursen VM, Edwards C, Cotter G, Davison BA, Damman K, Teerlink JR, et al. Liver function, in-hospital, and post-discharge clinical outcome in patients with acute heart failure—results from the relaxin for the treatment of patients with acute heart failure study. *Journal of Cardiac Failure*. 2014; 20:407–13. [PubMed: 24642379] [This study examines the association between liver function tests and mortality, rehospitalization and worsening heart failure.]
119. Allen LA, Felker GM, Pocock S, McMurray JJV, Pfeffer MA, Swedberg K, et al. Liver function abnormalities and outcome in patients with chronic heart failure: data from the Candesartan in Heart Failure: Assessment of Reduction in Mortality and Morbidity (CHARM) program. *European journal of heart failure : journal of the Working Group on Heart Failure of the European Society of Cardiology*. 2009; 11:170–7.
- 120•. Brisco MA, McCauley BD, Chen J, Parikh CR, Testani JM. Biochemical evidence of mild hepatic dysfunction identifies decompensated heart failure patients with reversible renal dysfunction. *J Card Fail*. 2013; 19:739–45. [PubMed: 24263117] [This study suggests that markers of hepatic congestion in patients with renal dysfunction may identify patients with heart failure induced renal dysfunction at presentation.]
121. Henderson NC, Mackinnon AC, Farnworth SL, Kipari T, Haslett C, Iredale JP, et al. Galectin-3 expression and secretion links macrophages to the promotion of renal fibrosis. *The American journal of pathology*. 2008; 172:288–98. [PubMed: 18202187]
122. Okamura DM, Pasichnyk K, Lopez-Guisa JM, Collins S, Hsu DK, Liu FT, et al. Galectin-3 preserves renal tubules and modulates extracellular matrix remodeling in progressive fibrosis. *Am J Physiol Renal Physiol*. 2011; 300:F245–53. [PubMed: 20962111]
123. Sharma UC, Pokharel S, van Brakel TJ, van Berlo JH, Cleutjens JPM, Schroen B, et al. Galectin-3 marks activated macrophages in failure-prone hypertrophied hearts and contributes to cardiac dysfunction. *Circulation*. 2004; 110:3121–8. [PubMed: 15520318]
124. Gopal DM, Kommineni M, Ayalon N, Koelbl C, Ayalon R, Biolo A, et al. Relationship of plasma galectin-3 to renal function in patients with heart failure: effects of clinical status, pathophysiology of heart failure, and presence or absence of heart failure. *Journal of the American Heart Association*. 2012; 1:e000760. [PubMed: 23316284]
125. van Kimmenade RR, Januzzi J, James L, Ellinor PT, Sharma UC, Bakker JA, Low AF, et al. Utility of Amino-Terminal Pro-Brain Natriuretic Peptide, Galectin-3, and Apelin for the Evaluation of Patients With Acute Heart Failure. *Journal of the American College of Cardiology*. 2006; 48:1217–24. [PubMed: 16979009]
126. Lok DJA, Meer P, Porte PWB-A, Lipsic E, Wijngaarden J, Hillege HL, et al. Prognostic value of galectin-3, a novel marker of fibrosis, in patients with chronic heart failure: data from the DEAL-HF study. *Clinical Research in Cardiology*. 2010; 99:323–8. [PubMed: 20130888]
127. Tang WHW, Shrestha K, Shao Z, Borowski AG, Troughton RW, Thomas JD, et al. Usefulness of plasma galectin-3 levels in systolic heart failure to predict renal insufficiency and survival. *The American Journal of Cardiology*. 2011; 108:385–90. [PubMed: 21600537]
- 128•. Meijers WC, Januzzi JL, deFilippi C, Adourian AS, Shah SJ, van Veldhuisen DJ, et al. Elevated plasma galectin-3 is associated with near-term rehospitalization in heart failure: a pooled analysis of 3 clinical trials. *American Heart Journal*. 2014; 167:853–60.e4. [PubMed: 24890535] [This study suggested galectin-3 may aid in predicting heart failure rehospitalization.]
129. Shah RV, Chen-Tournoux AA, Picard MH, van Kimmenade RRJ, Januzzi JL. Galectin-3, cardiac structure and function, and long-term mortality in patients with acutely decompensated heart failure. *European Journal of Heart Failure*. 2010; 12:826–32. [PubMed: 20525986]
- 130•. van der Velde AR, Gullestad L, Ueland T, Aukrust P, Guo Y, Adourian A, et al. Prognostic value of changes in galectin-3 levels over time in patients with heart failure: data from CORONA and COACH. *Circulation: Heart Failure*. 2013; 6:219–26. [PubMed: 23395934] [This analysis of two cohorts revealed that galectin-3 may identify both acute and chronic heart failure patients at risk for mortality.]
131. O'Seaghdha CM, Hwang S-J, Ho JE, Vasan RS, Levy D, Fox CS. Elevated galectin-3 precedes the development of CKD. *J Am Soc Nephrol*. 2013; 24:1470–7. [PubMed: 23766533]
132. Davidson NC, Barr CS, Struthers AD. C-type natriuretic peptide. An endogenous inhibitor of vascular angiotensin-converting enzyme activity. *Circulation*. 1996; 93:1155–9. [PubMed: 8653836]

133. Mattingly MT, Brandt RR, Heublein DM, Wei CM, Nir A, Burnett JC. Presence of C-type natriuretic peptide in human kidney and urine. *Kidney Int.* 1994; 46:744–7. [PubMed: 7996796]
134. Del Ry S, Passino C, Maltinti M, Emdin M, Giannessi D. C-type natriuretic peptide plasma levels increase in patients with chronic heart failure as a function of clinical severity. *European Journal of Heart Failure.* 2005; 7:1145–8. [PubMed: 15922659]
135. Del Ry S, Giannessi D, Maltinti M, Prontera C, Iervasi A, Colotti C, et al. Increased levels of C-type natriuretic peptide in patients with idiopathic left ventricular dysfunction. *Peptides.* 2007; 28:1068–73. [PubMed: 17428580]
136. Wright SP, Prickett TCR, Doughty RN, Frampton C, Gamble GD, Yandle TG, et al. Amino-terminal pro-C-type natriuretic peptide in heart failure. *Hypertension.* 2004; 43:94–100. [PubMed: 14656955]
137. Zakeri R, Sangaralingham SJ, Sandberg SM, Heublein DM, Scott CG, Burnett JC. Urinary C-type natriuretic peptide: a new heart failure biomarker. *JACC: Heart Failure.* 2013; 1:170–7. [PubMed: 24244916] [This study revealed the potential for urinary CNP as a prognostic biomarker in decompensated heart failure.]

Figure 1. Survival of heart failure patients as a function of GFR and BUN/Cr (left panel) and GFR and NT-proBNP (right panel) in two clinical trial populations of decompensated heart failure

Patients with renal dysfunction (estimated glomerular filtration rate less than 60 ml/min/1.73m²) and elevated BUN/Cr or NT-proBNP are at the highest risk for mortality. Evaluation of markers of congestion (NT-proBNP) or neurohormonal activation (BUN/Cr) in heart failure patients with renal dysfunction may aid in distinguishing CRS from other forms of renal impairment.

Left panel reproduced from Testani JM, Coca SG, Shannon RP, Kimmel SE, Cappola TP. Influence of renal dysfunction phenotype on mortality in the setting of cardiac dysfunction: analysis of three randomized controlled trials. *European Journal of Heart Failure*. 2011;13:1224-30, with permission from John Wiley and Sons.

Right panel reproduced from van Kimmenade RRJ, Januzzi J, James L, Baggish AL, Lainchbury JG, Bayes-Genis A, Richards AM, et al. Amino-terminal pro-brain natriuretic Peptide, renal function, and outcomes in acute heart failure: redefining the cardiorenal interaction? *Journal of the American College of Cardiology*. 2006;48:1621-7, with permission from Elsevier.

Table 1

Novel renal biomarkers and their utility in prognosis, diagnosis, and targeted treatment of cardiorenal syndrome

Biomarker	Prognosis in CRS	Diagnosis of CRS	Targeted Treatment
Glomerular Filtration and Integrity			
Cystatin C	***	*	None
Albuminuria	***	*	*
Tubular Kidney Injury			
NGAL	***	*	None
NAG	***	*	None
KIM-1	***	*	None
IL-18	*	*	None
Neurohormonal Activation and Sodium Avidity			
BUN/Cr	***	**	None
Diuretic Efficiency	***	**	None
Venous Congestion			
BNP	***	**	None
Biochemical Evidence of	*	*	None
Hepatic Dysfunction			
Tissue Fibrosis			
Galectin-3	*	None	None
Mechanism Undefined			
CNP	**	None	None

*** Utility strongly demonstrated by evidence

** Utility moderately demonstrated by evidence

* Utility only suggested by evidence; "None" No evidence; NGAL: Neutrophil Gelatinase-Associated Lipocalin, NAG:N-Acetyl- β -D-Glucosaminidase, KIM-1: Kidney Injury Molecule-1, IL-18:Interleukin-18, BUN/Cr: Blood Urea Nitrogen/Creatinine Ratio, BNP:Brain Natriuretic Peptide, CNP:C-Type Natriuretic Peptide