Published in final edited form as:

Angew Chem Int Ed Engl. 2014 October 6; 53(41): 11023–11026. doi:10.1002/anie.201405696.

A Multitasking Vanadium-Dependent Chloroperoxidase as an Inspiration for the Chemical Synthesis of the Merochlorins**

Dr. Stefan Diethelm,

Center for Marine Biotechnology and Biomedicine, Scripps Institution of Oceanography,
University of California San Diego, La Jolla, CA 92093

Dr. Robin Teufel,

Center for Marine Biotechnology and Biomedicine, Scripps Institution of Oceanography,
University of California San Diego, La Jolla, CA 92093

Dr. Leonard Kaysser, and

Center for Marine Biotechnology and Biomedicine, Scripps Institution of Oceanography,
University of California San Diego, La Jolla, CA 92093

Prof. Dr. Bradley S. Moore

Center for Marine Biotechnology and Biomedicine, Scripps Institution of Oceanography,
University of California San Diego, La Jolla, CA 92093, Homepage: <http://scrippsscholars.ucsd.edu/bsmoore>. Skaggs School of Pharmacy and Pharmaceutical Sciences,
University of California San Diego, La Jolla, CA 92093

Bradley S. Moore: bsmoore@ucsd.edu

Abstract

The vanadium-dependent chloroperoxidase Mcl24 was discovered to mediate a complex series of unprecedented transformations in the biosynthesis of the merochlorin meroterpenoid antibiotics. In particular, a site-selective naphthol chlorination is followed by a sequence of oxidative dearomatization/terpene cyclization reactions to build up the stereochemically complex carbon framework of the merochlorins in one step. Inspired by the enzyme reactivity, we developed a chemical chlorination protocol paralleling the biocatalytic process. These chemical studies led to the identification of previously overlooked merochlorin natural products.

Keywords

biosynthesis; vanadium-dependent haloperoxidase; oxidative dearomatization; biomimetic synthesis; chlorination

**We are grateful to Dr. Brendan Duggan for assistance with NMR measurements. This research was supported by the US National Institute of Health (NIH, Grant No. R01-AI047818) and by postdoctoral fellowships to R.T. from the Deutsche Forschungsgemeinschaft (TE 931/1-1), to L.K. from the Alexander von Humboldt Foundation, and to S.D. from the Swiss National Science Foundation (SNF).

© 2013 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

Correspondence to: Bradley S. Moore, bsmoore@ucsd.edu.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201405696>

In the past decades, marine microorganisms have emerged as valuable sources for structurally diverse secondary metabolites.^[1] In this context, the isolation and characterization of natural products from marine bacteria incorporating unusual carbon skeletons has been of particular interest. Merochlorin A (**1**) and B (**2**) were recently isolated from marine *Streptomyces* sp. CNH-189, constituting a new class of structurally unprecedented meroterpenoid secondary metabolites.^[2,3] The merochlorins are composed of a tetrahydroxynaphthalene fragment decorated with a rare terpenoid appendage. Furthermore, chlorination at C(11)/C(12) adds to the structural complexity of these compounds. These unusual structural features along with the antimicrobial activity of the merochlorins attracted the interest of the synthetic community, culminating in elegant total syntheses of both merochlorin A and B.^[4] In order to shed light on the biosynthetic origin of these natural products, our group recently identified the gene cluster associated with the production of the merochlorins.^[3] The genomic data suggested a biosynthetic pathway starting with the prenylation of 1,3,6,8-tetrahydroxynaphthalene (THN, **3**) to produce triene **4** (pre-merochlorin) as a key intermediate (Scheme 1). In the preceding publication, we reported on the investigation of these early steps in the merochlorin biosynthesis, leading to the identification of a highly unusual pathway for the synthesis and transfer of an isosqualavandulyl terpene unit **5** onto THN (**3**).^[5]

We herein report the biochemical characterization of Mcl24, a multitasking vanadium-dependent chloroperoxidase involved in the late stage of merochlorin biosynthesis (Scheme 1). The enzyme was found to mediate a complex series of transformations, including a site-selective naphthol chlorination, followed by a sequence of oxidative dearomatization/terpene cyclization. Furthermore, the enzyme's reactivity served as a blueprint for the development of a parallel chemical chlorination protocol, enabling the oxidative dearomatization/cyclization of the enzyme substrate and leading to the production of various merochlorin analogues. Besides the identification of previously overlooked natural merochlorin isomers, the chemical transformation suggests a mechanistic rationale for the enzymatic reaction cascade. Most importantly, the applied concept showcases the yet largely unexplored potential of unusual biosynthesis pathways to serve as an inspiration for the development of novel synthetic reagent systems based on biosynthetic logic.^[6]

The identification of prenylated naphthalene derivative **4** as a key intermediate towards merochlorin A and B^[5] suggested an oxidative cyclization event to account for the formation of the polycyclic core structure of both natural products. Surprisingly, although numerous secondary metabolites have been suggested to arise from oxidative cyclization involving an aromatic core, no enzymatic system capable of mediating such a transformation has been characterized to date.^[7] We identified a number of oxidative enzymes in the merochlorin gene cluster that were possible candidates to promote such a reaction, including an iron-sulfur cluster containing protein.^[3] Interestingly though, deletion of the gene *mcl30*, annotated to code for this protein, did not affect merochlorin production in the mutant strain. Moreover, the gene cluster contained the vanadium-dependent haloperoxidase (VHPO) Mcl24, which we suspected was likely responsible for introduction of the chlorine substituent. This enzyme particularly attracted our interest, as VHPO's represent a class of biocatalysts still poorly characterized in the context of complex molecule biosynthesis.^[8,9]

At the outset of our studies directed towards investigation of the final steps in the merochlorin biosynthesis, the nature of the oxidative cyclization event and the exact role of the haloperoxidase were largely unclear.

We commenced our efforts with the heterologous expression of octahistidyl-tagged Mcl24 in *Escherichia coli* in order to address the proposed C(2)-chlorination of intermediate **4**.^[5] Accordingly, purified Mcl24 was subjected to a standard VHPO enzyme assay (pH 6.0 MES buffer, 50 mM KCl, 1 mM H₂O₂, 100 μ M Na₃VO₄) using chemically synthesized pre-merochlorin **4**^[10] as the putative enzyme substrate. To our surprise, analysis of the Mcl24 reaction suggested the direct formation of the natural products merochlorin A (**1**) and B (**2**) by the chloroperoxidase enzyme, as indicated by reversed phase HPLC retention time as well as UV and MS spectral data of the product peaks (Figure 1A). Isolation of the major reaction components and ¹H-NMR analysis confirmed the identity of the products formed by comparison with spectral data of the isolated natural products.^[3] The unexpected finding that Mcl24 is able to catalyze multiple transformations including a site-selective chlorination at C(2), along with a highly unusual oxidative cyclization event, raised questions about the exact functional role of the enzyme in this reaction cascade. To gain further insight, we next tested various substrate analogues closely resembling the physiological substrate **4**, as well as the simple terpene alcohols geraniol and nerolidol. Surprisingly, none of these alternative substrates were accepted by the enzyme (see SI for details). We further subjected Mcl24 to a monochlorodimedone (MCD) assay, a standard protocol to assess the activity of halonium-generating halogenating enzymes.^[11] As indicated in Figure 1B, we found that Mcl24 is not able to oxidize MCD under standard assay conditions in the presence of chloride only.

However, when KBr was added to the reaction mixture, rapid disappearance of the MCD signal was observed. We recently reported a similar observation with the vanadium-dependent chloroperoxidase NaphH1, involved in a highly selective chloroether formation in the biosynthesis of the napyradiomycins.^[9c] As such, these results raise questions about the feasibility of the MCD assay to properly reflect the natural role of halogenating enzymes.

On the basis of these initial observations, we hypothesized that an active chloronium species could trigger an oxidative dearomatization of the highly electron rich naphthol moiety in **4**. The thus generated reactive intermediate would then initiate a terpene cyclization to arrive at the polycyclic frameworks of the merochlorins. To gain further insight into the details of such an enzyme catalyzed oxidative naphthol dearomatization/cyclization cascade, we opted for the development of a parallel chemical protocol inducing oxidative cyclization of **4** through a chlorination event. Although oxidative dearomatization/cyclization cascades initiated by a halogenating species are, to our knowledge, unprecedented in the chemical literature,^[12] we were curious, if chemical chlorination conditions could be identified that would enable generating cyclized products starting with pre-merochlorin **4**. Accordingly, various chlorination conditions were evaluated towards this end as outlined in Table 1 (see SI for further details). In particular, we sought protocols mediating selective *ortho*-chlorination of phenol and naphthol derivatives, thus paralleling the chlorination reactivity of Mcl24.^[13] 2,3,4,5,6,6-Hexachloro-2,4-cyclohexadien-1-one has been reported to effect such transformations in the presence of DMF as a co-solvent. Subjecting naphthol **4** to these conditions led to complete decomposition of the starting material (entry 1).^[14] Interestingly

though, addition of 5 equiv. of triethylamine to this reaction effected formation of trace quantities of a set of products with very similar retention times on reversed phase HPLC as well as identical UV and MS spectra to the enzymatic products **1** and **2** (entry 2).

Following this initial result, we set out to further study the oxidative cyclization of **4**. A number of standard chlorinating reagents failed to give any of the previously observed products (entries 3–6). Interestingly, the use of hypochlorous acid, the generally postulated reactive species of chloroperoxidases, only led to decomposition of triene **4** (entry 7).^[15] At this point we began to suspect that the amine additive in the previously successful conditions might play a key role in the reaction, possibly leading to the formation of a reactive chloramine intermediate.^[16] Accordingly, substrate **4** was subjected to *N*-chlorosuccinimide in the presence of 2 equiv. of *i*-Pr₂NH.^[17] To our delight, formation of the previously observed product mixture ensued in a total yield of 30% (entry 8). Moreover, the use of independently prepared *N,N*-dichloro-*tert*-butylamine produced the cyclized products, albeit in lower yield (entry 9).^[18] Finally, the use of stronger bases was tested. Addition of sodium hydride equally led to the formation of the characteristic product pattern in 22% yield when conducting the reaction at –78 °C (entry 10).

With an efficient protocol for chemical oxidative cyclization of **4** in hand, we turned to a closer investigation of the individual components observed in the product mixture. As shown in Figure 2, HR-LCMS comparison of the enzymatic Mcl24 reaction with the chemical oxidation of **4** using NCS/*i*-Pr₂NH, indicated a markedly different product distribution pattern. Upon preparative HPLC purification of both, the enzymatic and the chemical reaction mixtures, the individual products were characterized by detailed 2D NMR and MS analysis. As depicted in Figure 2, a full spectrum of differently chlorinated merochlorin A and B analogues were identified. The NCS/*i*-Pr₂NH reaction thereby predominantly produced deschloro-merochlorin B (**6**), isochloro-merochlorin B (**7**), deschloro-merochlorin A (**8**) and minor amounts of dichloromerochlorin B (**9**) (coelutes with **8**). On the other hand, the enzymatic reaction preferentially provides the previously described merochlorin A (**1**) and B (**2**), along with only minor amounts of isomers **6**, **7** and **8**. Reinvestigation of the natural merochlorin producer strain *Streptomyces* sp. CNH-189 revealed that all of the observed merochlorin isomers were in fact produced by this organism, albeit in only minor amounts (see SI for details).

The inverted selectivity profiles of the chemical and the enzymatic oxidative cyclization of **4** suggested implications on the reaction path Mcl24 follows. Remarkably, the enzyme seems to control both, the site-selectivity of halogenation as well as the timing of the chlorination reaction in respect to the oxidative cyclization step. In contrast to the NCS/*i*-Pr₂NH reaction, Mcl24 preferentially chlorinates at C(2) in **4** prior to cyclization. On the other hand, treatment of **4** with NCS/*i*-Pr₂NH leads to an initial oxidative cyclization followed by optional chlorination at C(7). In fact, conducting the chemical chlorination at low temperature (–78 °C) prevents this chlorination step, leading to the exclusive production of isomers **6** and **8** (Table 1, entry 10 and SI). The origin of the distinct reactivity profile of the enzymatic reaction remains unclear at this point.

The studies undertaken towards chemical oxidative cyclization of **4** led us to propose a mechanistic rationale for merochlorin production. As mentioned before, we surmise that an *in situ* generated chloramine intermediate is the active oxidant for the chemical transformation. Interestingly, a ϵ -lysine chloramine intermediate is postulated for the rebeccamyin flavin-dependent chlorinase RebH.^[19] The chlorinated amino acid side chain was proposed to transfer its halogen atom in a highly selective manner to the enzyme substrate. It seems attractive to invoke a similar intermediate with a chlorinated amino acid side chain to account for the unusual selectivity pattern observed for Mcl24. We further hypothesize that the active chloronium species initiates the enzymatic reaction by selective C(2) chlorination of substrate **4** (Scheme 2).^[20] A second chlorination step would then likely produce an aromatic hypochlorite species **10**, and upon loss of chloride would give benzylic carbocation **11**.^[21] The proposed formation of intermediate **10** is also consistent with the observation that a NaH/NCS reagent combination can promote cyclization of **4** (Table 1, entry 10). Finally, a cation-induced terpene cyclization would explain formation of the merochlorins *via* intermediate **12**.

In conclusion, the characterization of the vanadium-dependent chloroperoxidase Mcl24 led to the discovery of an unprecedented enzymatic reaction cascade. Mcl24 was found to catalyze a site-selective naphthol chlorination prior to the initiation of an oxidative dearomatization event that further triggers a terpene cyclization. We were able to shed light on the characteristics of the enzymatic reaction by the development of a synthetic chlorination protocol mimicking Mcl24 reactivity. In addition, this novel reagent system enabled the total synthesis of previously unnoticed merochlorin natural products in only five steps. The applied concept stresses the potential of unusual biosynthetic transformations to serve as an inspiration for the development of synthetic reagent systems.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

1. Moore BS. Nat Prod Rep. 2005; 22:580–593. [PubMed: 16193157]
2. Sakoulas G, Nam SJ, Loesgen S, Fenical W, Jensen PR, Nizet V, Hensler M. PLoS ONE. 2012; 7:e29439. [PubMed: 22279537]
3. Kaysser L, Bernhardt P, Nam SJ, Loesgen S, Ruby JG, Skewes-Cox P, Jensen PR, Fenical W, Moore BS. J Am Chem Soc. 2012; 134:11988–11991. [PubMed: 22784372]
4. a) Pepper HP, George JH. Angew Chem Int Ed. 2013; 52:12170–12173. b) Meier R, Strych S, Trauner D. Org Lett. 2014; 16:2634–2637. [PubMed: 24804897]
5. Teufel R, Kaysser L, Villaume MT, Diethelm S, Carbullido MK, Baran PS, Moore BS. preceding publication.
6. For a couple of recent examples of biosynthetically inspired synthetic reagents, see: Gui J, Zhou Q, Pan CM, Yabe Y, Burns AC, Collins MR, Ornelas MA, Ishihara Y, Baran PS. J Am Chem Soc. 2014; 136:4853–4856. [PubMed: 24611732] Chen MS, White MC. Science. 2007; 318:783–787. [PubMed: 17975062]
7. a) Singh IP, Bharate SB. Nat Prod Rep. 2006; 23:558–591. [PubMed: 16874390] b) Roche SP, Proco JA Jr. Angew Chem Int Ed. 2011; 50:4068–4093.
8. a) Littlechild J. Curr Opin Chem Biol. 1999; 3:28–34. [PubMed: 10021409] b) Butler A, Carter-Franklin JN. Nat Prod Rep. 2004; 21:180–188. [PubMed: 15039842] c) Vaillancourt FH, Yeh E,

- Vosburg DA, Garneau-Tsodikova S, Walsh CT. Chem Rev. 2006; 106:3364–3378. [PubMed: 16895332] d) Winter JM, Moore BS. J Biol Chem. 2009; 284:18577–18581. [PubMed: 19363038]
9. a) Carter-Franklin JN, Parrish JD, Tschirret-Guth RA, Little RD, Butler A. J Am Chem Soc. 2003; 125:3688–3689. [PubMed: 12656585] b) Carter-Franklin JN, Butler A. J Am Chem Soc. 2004; 126:15060–15066. [PubMed: 15548002] c) Bernhardt P, Okino T, Winter JM, Miyanaga A, Moore BS. J Am Chem Soc. 2011; 133:4268–4270. [PubMed: 21384874]
10. The synthesis of **4** is reported in the preceding publication [5].
11. Morris DR, Hager LP. J Biol Chem. 1966; 241:1763–1768. [PubMed: 5949836]
12. For a comprehensive review on oxidative dearomatization in chemistry, see: Quideau S, Pouységu L, Deffieux D. Synlett. 2008:467–495. See also ref. [7b].
13. For a review on selective phenol chlorination, see: Pearson DE, Buehler CA. Synthesis. 1971:455–477.
14. a) Guy A, Lemaire M, Guetté JP. Tetrahedron. 1982; 38:2339–2346. b) Guy A, Lemaire M, Guetté JP. Tetrahedron. 1982; 38:2347–2354.
15. Slow addition over 1 hour (syringe pump) of only 1 equiv. of HOCl to a solution of **4** in acetone equally led to complete decomposition of the starting material.
16. Literature precedence suggested that chloramines can mediate ortho-selective phenol chlorinations: Lindsay Smith JR, McKeer LC, Taylor JM. J Chem Soc, Perkin Trans. 1988; 2:385–391. Smith K, Butters M. Tetrahedron Lett. 1988; 29:1319–1322. Gnaïm JM, Sheldon RA. Tetrahedron Lett. 1995; 36:3893–3896. Saper NI, Snider BB. J Org Chem. 2014; 79:809–813. [PubMed: 24328120]
17. The use of primary amines, such as *t*-BuNH₂, proved equally effective (see supporting information for further detail).
18. Ogata Y, Takagi K, Kondo Y. J Chinese Chem Soc. 1983; 30:261–266. For the preparation of *t*-BuNCl₂, see: De Luca L, Giacomelli G. Synlett. 2004:2180–2184.
19. Yeh E, Blasiak LC, Koglin A, Drennan CL, Walsh CT. Biochemistry. 2007; 46:1284–1292. [PubMed: 17260957]
20. C(2)-chlorinated pre-merochlorin could not be detected by HPLC and HR-LCMS analysis in either the Mcl24 or the NCS/*i*-Pr₂NH reaction. This can likely be attributed to the high oxygen sensitivity of pre-merochlorin compounds and resulting rapid oxidative decomposition of such intermediates.
21. For a report on the generation of an aromatic hypochlorite by the action of an N-Cl species, see: Dietze PE, Underwood GR. J Org Chem. 1984; 49:2491–2494.

Figure 1.

Reactivity of Mcl24. (A) Reversed phase HPLC chromatogram (UV at 254 nm) of the reaction of Mcl24 with **4**: a) Mcl24 (50 $\mu\text{g/mL}$), KCl (50 mM), Na_3VO_4 (100 μM), H_2O_2 (1 mM), **4** (100 μmol), pH 6.0 MES buffer (50 mM). **1** = merochlorin A; **2** = merochlorin B; * = non-enzymatic oxidized degradation product. b) Negative control: no substrate **4**. c) Negative control: no enzyme. (B) Spectrophotometric MCD assay for Mcl24. The reaction was initiated by addition of H_2O_2 . As no decrease in absorbance was observed in the presence of chloride, KBr was added to the mixture after 7 min.

Figure 2.

HR-LCMS comparison (UV at 254 nm shown) of the Mcl24 reaction (top) with the NCS/*i*-Pr₂NH reaction (bottom) of **4**.

Scheme 1.

Biosynthetic pathway towards merochlorin A (1) and B (2). DMAPP = dimethylallyl diphosphate, GPP = geranyl diphosphate.

Scheme 2.

Proposed mechanism for the oxidative dearomatization/cyclization towards merochlorin A (path *a*) and B (path *b*).

Table 1

Chemical chlorination of pre-merochlorin **4**.

Entry ^[a]	Chlorinating Agent	Additive	Solvent	Combined Yield ^[b]
1	HCDO	DMF (100 equiv.)	CCl ₄	-
2	HCDO	NEt ₃ (5 equiv.)	CCl ₄	< 5%
3	NCS	-	CH ₂ Cl ₂	-
4	TCCA	-	CH ₂ Cl ₂	-
5	<i>t</i> -BuOCl	-	CH ₂ Cl ₂	-
6	SOCl ₂	<i>i</i> -Pr ₂ NH (2 equiv.)	toluene	-
7	HOCl	-	acetone	-
8 ^[c]	NCS	<i>i</i> -Pr ₂ NH (2 equiv.)	CH ₂ Cl ₂	30%
9	<i>t</i> -BuNCl ₂	-	CCl ₄	~5%
10 ^{[d],[e]}	NCS	NaH (2 equiv.)	CH ₂ Cl ₂	22%

^[a] Reaction conditions: substrate (5 mg, 12.6 μmol), chlorinating agent (2.0 equiv.), additive, solvent (1 mL), 25 °C, 2 h.

^[b] combined yield of product mixture after silica gel purification.

^[c] at 0 °C.

^[d] at -78 °C.

^[e] only non-chlorinated isomers **6** and **8** produced. HCDO = 2,3,4,5,6,6-hexachloro-2,4-cyclohexadiene-1-one; TCCA = trichloroisocyanuric acid; NCS = *N*-chlorosuccinimide; DMF = *N,N*-dimethylformamide.