Published in final edited form as:

New Horiz Transl Med. 2014 September 1; 2(1): 16–19. doi:10.1016/j.nhtm.2014.08.003.

Small Molecule PAI-1 Functional Inhibitor Attenuates Vascular Smooth Muscle Cell Migration and Survival: Implications for the Therapy of Vascular Disease

Tessa M. Simone and Paul J. Higgins

Center for Cell Biology & Cancer Research, Albany Medical College, Albany, New York 12208

Keywords

Vascular disease; PAI-1; SERPIN; pericellular proteolysis; tiplaxtinin; vascular smooth muscle cells; migration; apoptosis

Introduction: PAI-1 in Vascular Pathology

Vascular restenosis, the pathologic re-narrowing of a blood vessel after percutaneous coronary intervention, involves increased vascular smooth muscle cell (VSMC) migration, elevated proliferation and decreased VSMC apoptosis (1, 2). Few treatment options for vascular restenosis exist aside from re-catheterization. Among several factors implicated in the pathophysiologic vascular response to injury (balloon angioplasty is one type of trauma), studies in animal models and the available clinical evidence suggest that plasminogen activator inhibitor-1 (PAI-1), a member of the serine protease inhibitor (SERPIN) superfamily and the major physiologic regulator of the plasmin-based pericellular proteolytic cascade, is perhaps the most prominent. PAI-1 attenuates fibrinolysis and promotes tissue fibrosis by inhibition of the plasminogen→plasmin-generating system (Figure 1).

Elevated PAI-1 expression is a significant causative factor in vascular disease and a major contributor to the pathophysiology of a number of significant human disorders including diabetes, pulmonary/renal fibrosis, metabolic syndrome, intravascular septic coagulopathy, atherosclerosis and restenosis, particularly in the setting of increased tissue TGF- β 1 levels. PAI-1 exerts spatial and temporal control over the integrated processes of pericellular proteolysis and extracellular matrix (ECM) deposition/turnover that impact stromal remodeling, inflammation, cell migration, proliferation and apoptosis, each of which are critical determinants in tissue fibrosis and vascular disease (Figure 2).

© 2014 European Society for Translational Medicine. Published by Elsevier Ltd. All rights reserved.

Correspondence: Dr. Paul J. Higgins, Center for Cell Biology & Cancer Research, Albany Medical College, 47 New Scotland Avenue, MC-165, Albany, New York 12208, Tel: 518-262-5168, FAX: 518-262-5669, higginp@mail.amc.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

PAI-1 Structure/Function

During the interaction of PAI-1 with its target proteases, the isopeptide bond in the reactive center loop (RCL) is cleaved by the target protease to form a covalent ester bond between a serine hydroxyl group of the enzyme and a PAI-1 carboxyl group. Upon PAI-1 cleavage, the N-terminus of the RCL inserts into β -sheet A, while the RCLC-terminus forms strands 1C in β -sheet C producing a 70Å separation of the P1 and P1' residues, thereby deforming the protease and rendering it inactive. This cleavage also renders PAI-1 anti-proteolytically inactive, giving rise to its designation as a “suicide inhibitor”. A substrate form of PAI-1 exists as well in which PAI-1 is cleaved by its target proteases without covalent formation of a PAI-1:protease complex (3,4–6).

Due to the complexity of PAI-1 structure and function, several low-molecular weight antagonists of PAI-1 have been developed to evaluate specific contributions of this SERPIN to disease pathologies (7). Tiplaxtinin (PAI-039), one of the most well-studied small-molecule inhibitors, attenuates asthmatic episodes, hyperlipidemia, hyperglycemia and angiogenesis (7–14). The specific mechanism by which tiplaxtinin antagonizes the anti-fibrinolytic activity of PAI-1 involves promotion of a substrate-like conformation resulting in PAI-1 cleavage and impaired uPA and tPA inhibition (15,16). The relative abundance of the different conformational states of PAI-1 within the immediate pericellular microenvironment likely dictates, in fact, whether VSMC migrate, proliferate or undergo apoptosis in response to injury. Indeed, in the context of a spectrum of cardiovascular diseases, it appears that PAI-1 is both pro- and anti-restenotic (17) depending on the nature of the wound model, level of both active or available PAI-1 and vessel TGF- β 1 expression. Global PAI-1 deletion in knock-out mice, over-reliance on application of only the full-length, active form of PAI-1 to assess vascular remodeling, and uncertainties as to the role of PAI-1 conformation-dependent processes, however, are major confounders.

PAI-1: Function Beyond Protease inhibition

It is increasingly evident that aside from its anti-proteolytic role, PAI-1 also functions as a multifunctional signaling “ligand” where it impacts cellular responses at the site of injury. All three forms of PAI-1 (full-length, latent and cleaved) interact with the low-density lipoprotein receptor-related protein 1 (LRP-1) and stimulate JAK/STAT1-mediated VSMC migration (18). Outcomes, however, are clearly concentration-dependent. Low dose (2 nM), acute exposure (3 h) to cleaved PAI-1 stimulates VSMC migration (18); chronic exposure (24 h) to high dose (40 nM) cleaved PAI-1 (via application of tiplaxtinin), in contrast, attenuated motility. Since both tiplaxtinin and cleaved PAI-1 stimulate apoptosis after 24 h, chronic exposure to cleaved PAI-1 appears to switch VSMC from the pro-migratory to a pro-apoptotic phenotype.

While full-length, active PAI-1 reduces both spontaneous and stimulated prostate cancer cell apoptosis, latent PAI-1 was unable to rescue either response. The ability of active PAI-1 to inhibit apoptosis, furthermore, is not due to its urokinase PA (uPA) binding or uPA receptor (uPAR) signaling roles (19) suggesting that LRP-1 may be the more relevant “survival” receptor. Truncated PAI-1 (PAI-1₂₃), a mutant with deletions in much of the heparin-

binding domain and the RCL, also stimulated endothelial cell apoptosis (20,21). It appears that an intact RCL is required for the pro-survival function of PAI-1 and disruption of this structure promotes an apoptotic phenotype. This is consistent with findings that cleaved PAI-1 stimulates VSMC apoptosis (Simone et al., submitted). Indeed, unpublished recent data from this laboratory clearly indicate that application of an elastase-cleaved PAI-1 promotes VSMC apoptosis, both with and without additional death stimuli (e.g., chemotherapeutic drugs), while tiplaxtinin (10 μ M), in the presence of recombinant PAI-1 (40 nM), promotes apoptosis to a greater extent than tiplaxtinin alone. Since 20 μ M tiplaxtinin completely cleaves 400 nM PAI-1 in 30 min, it is likely under the latter conditions (24 h) tiplaxtinin effectively cleaves both exogenous recombinant and endogenous PAI-1 pools.

Importantly, elastase levels increase immediately after balloon angioplasty, peak at one week and then decline (22) suggesting that lowered elastase activity levels, might contribute to a decrease in cleaved PAI-1 and, therefore, reduced apoptosis and increased VSMC persistence. As elastase cleaves PAI-1 at the peptide bond between Val³⁴³-Ser³⁴⁴ (P4-P3) and, whereas, tiplaxtinin promotes substrate behavior of PAI-1, allowing for uPA/tPA protease attack at Arg³⁴⁶-Met³⁴⁷ (P1-P1') (23), cleavage at either of these sites may be sufficient for promotion of apoptosis. Tiplaxtinin could be a useful therapeutic option in the context of elevated vessel uPA and PAI-1, as it promotes a substrate-like form of PAI-1, a uPA target (24–30).

Conclusions

In the absence of PAI-1, the increase inactive plasmin releases a pro-apoptotic Fas ligand that stimulates endothelial cell death (31). Unlike endothelial cells, however, VSMC are relatively resistant to Fas ligand-stimulated apoptosis (32) suggesting that, when cleaved, PAI-1-stimulated VSMC apoptosis, is likely to be independent of Fas. Interestingly, PAI-1 expression is stimulated by both TNF- α (33,34) and tumor necrosis factor-like weak inducer of apoptosis (TWEAK)(35). Furthermore, full-length PAI-1 binds to, and inhibits, caspase-3 in TNF- α -treated cells (36), implicating PAI-1 as a VSMC pro-survival factor in response to and dependent on members of the TNF- α -superfamily. Importantly, both tiplaxtinin and elastase-cleaved PAI-1 down-regulate TWEAK, NF- κ B, and TNFSR1b, (a TNF- α -binding receptor) transcripts in VSMC. Cleaved PAI-1 might potentiate VSMC apoptosis by deregulating these pro-survival signaling pathways (37–40) inducing death receptor-initiated apoptosis and the executioner caspase cascade. Collectively, these data suggest that PAI-1 overexpression is both a causative factor in vascular and non-vascular fibrosis as well as a biomarker of cardiovascular disease mortality. The adaptation of PAI-1 function-disrupting strategies may have eventual translational utility for the treatment of the pathophysiology of vascular disorders, thrombosis and tissue fibrosis. Targeted manipulation of the underlying PAI-1-dependent pro-survival pathways, with function-blocking PAI-1 mutants or small molecule pharmacologic inhibitors, is likely to have significant clinical implications in the context of PAI-1-dependent vascular pathologies.

Acknowledgments

This work was supported by NIH Grant GM507242 to PJ Higgins and AHA Pre-Doctoral Fellowship 14PRE18170012 to TM Simone.

References

1. Bochaton-Piallat ML, Gabbiani F, Redard M, Desmouliere A, Gabbiani G. Apoptosis participates in cellularity regulation during rat aortic intimal thickening. *Am J Pathol.* 1995; 146:1059–1064. [PubMed: 7747800]
2. Schwartz SM. Perspectives series: cell adhesion in vascular biology. Smooth muscle migration in atherosclerosis and restenosis. *J Clin Invest.* 1997; 99:2814–2816. [PubMed: 9185501]
3. Declerck PJ, De Mol M, Vaughan DE, Collen D. Identification of a conformationally distinct form of plasminogen activator inhibitor-1, acting as a noninhibitory substrate for tissue-type plasminogen activator. *J Biol Chem.* 1992; 267:11693–11696. [PubMed: 1601844]
4. Patston PA, Gettins P, Beechem J, Schapira M. Mechanism of serpin action: evidence that C1 inhibitor functions as a suicide substrate. *Biochemistry.* 1991; 30:8876–8882. [PubMed: 1888745]
5. Gils A, Declerck PJ. Proteinase specificity and functional diversity in point mutants of plasminogen activator inhibitor 1. *J Biol Chem.* 1997; 272:12662–12666. [PubMed: 9139722]
6. Aertgeerts K, De Bondt HL, De Ranter CJ, Declerck PJ. Mechanisms contributing to the conformational and functional flexibility of plasminogen activator inhibitor-1. *Nat Struct Biol.* 1995; 2:891–897. [PubMed: 7552714]
7. Lee SH, Eren M, Vaughan DE, Schleimer RP, Cho SH. A plasminogen activator inhibitor-1 inhibitor reduces airway remodeling in a murine model of chronic asthma. *Am J Respir Cell Mol Biol.* 2012; 46:842–846. [PubMed: 22323366]
8. Krause MP, Moradi J, Nissar AA, Riddell MC, Hawke TJ. Inhibition of plasminogen activator inhibitor-1 restores skeletal muscle regeneration in untreated type 1 diabetic mice. *Diabetes.* 2011; 60:1964–1972. [PubMed: 21593201]
9. Lijnen HR, Alessi MC, Frederix L, Collen D, Juhan-Vague I. Tiplaxtinin impairs nutritionally induced obesity in mice. *Thromb Haemost.* 2006; 96:731–737. [PubMed: 17139361]
10. Schalkwijk CG, Stehouwer CD. PAI-1 inhibition in obesity and the metabolic syndrome: a promising therapeutic strategy. *Thromb Haemost.* 2006; 96:698–699. [PubMed: 17139361]
11. Leik CE, Su EJ, Nambi P, Crandall DL, Lawrence DA. Effect of pharmacologic plasminogen activator inhibitor-1 inhibition on cell motility and tumor angiogenesis. *J Thromb Haemost.* 2006; 4:2710–2715. [PubMed: 17010152]
12. Crandall DL, Quinet EM, El Ayachi S, Hreha AL, Leik CE, Savio DA, Juhan-Vague I, Alessi MC. Modulation of adipose tissue development by pharmacological inhibition of PAI-1. *Arterioscler Thromb Vasc Biol.* 2006; 26:2209–2215. [PubMed: 16825598]
13. Lijnen HR, Alessi MC, Van Hoef B, Collen D, Juhan-Vague I. On the role of plasminogen activator inhibitor-1 in adipose tissue development and insulin resistance in mice. *J Thromb Haemost.* 2005; 3:1174–1179. [PubMed: 15946208]
14. Weisberg AD, Albornoz F, Griffin JP, Crandall DL, Elokda H, Fogo AB, Vaughan DE, Brown NJ. Pharmacological inhibition and genetic deficiency of plasminogen activator inhibitor-1 attenuates angiotensin II/salt-induced aortic remodeling. *Arterioscler Thromb Vasc Biol.* 2005; 25:365–371. [PubMed: 15576638]
15. Gorlatova NV, Cale JM, Elokda H, Li D, Fan K, Warnock M, Crandall DL, Lawrence DA. Mechanism of inactivation of plasminogen activator inhibitor-1 by a small molecule inhibitor. *J Biol Chem.* 2007; 282:9288–9296. [PubMed: 17276980]
16. Elokda H, Abou-Gharbia M, Hennen JK, McFarlane G, Mugford CP, Krishnamurthy G, Crandall DL. Tiplaxtinin, a novel, orally efficacious inhibitor of plasminogen activator inhibitor-1: design, synthesis, and preclinical characterization. *J Med Chem.* 2004; 47:3491–3494. [PubMed: 15214776]
17. Diebold I, Kraicun D, Bonello S, Gorlach A. The 'PAI-1 paradox' in vascular remodeling. *Thromb Haemost.* 2008; 100:984–991. [PubMed: 19132221]

18. Degryse B, Neels JG, Czekay RP, Aertgeerts K, Kamikubo Y, Loskutoff DJ. The low density lipoprotein receptor-related protein is a motogenic receptor for plasminogen activator inhibitor-1. *J Biol Chem.* 2004; 279:22595–22604. [PubMed: 15001579]
19. Kwaan HC, Wang J, Svoboda K, Declerck PJ. Plasminogen activator inhibitor 1 may promote tumour growth through inhibition of apoptosis. *Br J Cancer.* 2000; 82:1702–1708. [PubMed: 10817507]
20. Drinane M, Walsh J, Mollmark J, Simons M, Mulligan-Kehoe MJ. The anti-angiogenic activity of rPAI-1(23) inhibits fibroblast growth factor-2 functions. *J Biol Chem.* 2006; 281:33336–33344. [PubMed: 16950776]
21. Mulligan-Kehoe MJ, Wagner R, Wieland C, Powell R. A truncated plasminogen activator inhibitor-1 protein induces and inhibits angiostatin (kringles 1–3), a plasminogen cleavage product. *J Biol Chem.* 2001; 276:8588–8596. [PubMed: 11113116]
22. Barolet AW, Nili N, Cheema A, Robinson R, Natarajan MK, O'Blenes S, Li J, Eskandarian MR, Sparkes J, Rabinovitch M, Strauss BH. Arterial elastase activity after balloon angioplasty and effects of elafin, an elastase inhibitor. *Arterioscler Thromb Vasc Biol.* 2001; 21:1269–1274. [PubMed: 11498452]
23. Lindahl TL, Ohlsson PI, Wiman B. The mechanism of the reaction between human plasminogen-activator inhibitor 1 and tissue plasminogen activator. *Biochem J.* 1990; 265:109–113. [PubMed: 2105714]
24. Lennartz MR, Aggarwal A, Michaud TM, Feustel PJ, Jones DM, Brosnan MJ, Keller RS, Loegering DJ, Kreienberg PB. Ligation of macrophage Fcγ receptors recapitulates the gene expression pattern of vulnerable human carotid plaques. *PLoS One.* 2011; 6:e21803. [PubMed: 21814555]
25. Falkenberg M, Tom C, DeYoung MB, Wen S, Linnemann R, Dichek DA. Increased expression of urokinase during atherosclerotic lesion development causes arterial constriction and lumen loss, and accelerates lesion growth. *Proc Natl Acad Sci U S A.* 2002; 99:10665–10670. [PubMed: 12149463]
26. Peng L, Bhatia N, Parker AC, Zhu Y, Fay WP. Endogenous vitronectin and plasminogen activator inhibitor-1 promote neointima formation in murine carotid arteries. *Arterioscler Thromb Vasc Biol.* 2002; 22:934–939. [PubMed: 12067901]
27. DeYoung MB, Tom C, Dichek DA. Plasminogen activator inhibitor type 1 increases neointima formation in balloon-injured rat carotid arteries. *Circulation.* 2001; 104:1972–1971. [PubMed: 11602503]
28. Chomiki N, Henry M, Alessi MC, Anfosso F, Juhan-Vague I. Plasminogen activator inhibitor-1 expression in human liver and healthy or atherosclerotic vessel walls. *Thromb Haemost.* 1994; 72:44–53. [PubMed: 7974374]
29. Schneiderman J, Sawdey MS, Keeton MR, Bordin GM, Bernstein EF, Dilley RB, Loskutoff DJ. Increased type 1 plasminogen activator inhibitor gene expression in atherosclerotic human arteries. *Proc Natl Acad Sci U S A.* 1992; 89:6998–7002. [PubMed: 1495992]
30. Olofsson BO, Dahlen G, Nilsson TK. Evidence for increased levels of plasminogen activator inhibitor and tissue plasminogen activator in plasma of patients with angiographically verified coronary artery disease. *Eur Heart J.* 1989; 10:77–82. [PubMed: 2495219]
31. Bajou K, Peng H, Laug WE, Maillard C, Noel A, Foidart JM, Martial JA, DeClerck YA. Plasminogen activator inhibitor-1 protects endothelial cells from FasL-mediated apoptosis. *Cancer Cell.* 2008; 14:324–334. [PubMed: 18835034]
32. Sata M, Suhara T, Walsh K. Vascular endothelial cells and smooth muscle cells differ in expression of Fas and Fas ligand and in sensitivity to Fas ligand-induced cell death: implications for vascular disease and therapy. *Arterioscler Thromb Vasc Biol.* 2000; 20:309–316. [PubMed: 10669625]
33. Sawdey MS, Loskutoff DJ. Regulation of murine type 1 plasminogen activator inhibitor gene expression in vivo. Tissue specificity and induction by lipopolysaccharide, tumor necrosis factor-α, and transforming growth factor-β. *J Clin Invest.* 1991; 88:1346–1353. [PubMed: 1918385]

34. Schleef RR, Bevilacqua MP, Sawdey M, Gimbrone MA Jr, Loskutoff DJ. Cytokine activation of vascular endothelium. Effects on tissue-type plasminogen activator and type 1 plasminogen activator inhibitor. *J Biol Chem.* 1988; 263:5797–5803. [PubMed: 3128548]
35. Munoz-Garcia B, Madrigal-Matute J, Moreno JA, Martin-Ventura JL, Lopez-Franco O, Sastre C, Ortega L, Burkly LC, Egido J, Blanco-Colio LM. TWEAK-Fn14 interaction enhances plasminogen activator inhibitor 1 and tissue factor expression in atherosclerotic plaques and in cultured vascular smooth muscle cells. *Cardiovasc Res.* 2011; 89:225–233. [PubMed: 20810696]
36. Chen Y, Kelm RJ Jr, Budd RC, Sobel BE, Schneider DJ. Inhibition of apoptosis and caspase-3 in vascular smooth muscle cells by plasminogen activator inhibitor type-1. *J Cell Biochem.* 2004; 92:178–188. [PubMed: 15095413]
37. Sobel BE, Woodcock-Mitchell J, Schneider DJ, Holt RE, Marutsuka K, Gold H. Increased plasminogen activator inhibitor type 1 in coronary artery atherectomy specimens from type 2 diabetic compared with nondiabetic patients: a potential factor predisposing to thrombosis and its persistence. *Circulation.* 1998; 97:2213–2221. [PubMed: 9631870]
38. Badrichani AZ, Stroka DM, Bilbao G, Curiel DT, Bach FH, Ferran C. Bcl-2 and Bcl-XL serve an anti-inflammatory function in endothelial cells through inhibition of NF-kappaB. *J Clin Invest.* 1999; 103:543–553. [PubMed: 10021463]
39. Tran NL, McDonough WS, Savitch BA, Sawyer TF, Winkles JA, Berens ME. The tumor necrosis factor-like weak inducer of apoptosis (TWEAK)-fibroblast growth factor-inducible 14 (Fn14) signaling system regulates glioma cell survival via NFkappaB pathway activation and BCL-XL/BCL-W expression. *J Biol Chem.* 2005; 280:3483–3492. [PubMed: 15611130]
40. Vince JE, Pantaki D, Feltham R, Mace PD, Cordier SM, Schmukle AC, Davidson AJ, Callus BA, Wong WW, Gentle IE, Carter H, Lee EF, Walczak H, Day CL, Vaux DL, Silke J. TRAF2 must bind to cellular inhibitors of apoptosis for tumor necrosis factor (tnf) to efficiently activate nf- κ b and to prevent tnf-induced apoptosis. *J Biol Chem.* 2009; 284:35906–35915. [PubMed: 19815541]

Focal Points

Bedside

Targeted pharmacologic disruption of PAI-1 function with small molecule inhibitors may have general applicability for the treatment of fibroproliferative disorders, in general, and vascular disease, in particular. The development orally compatible drugs would likely simplify delivery and patient compliance.

Benchside

It is apparent that PAI-1 is a multi-functional SERPIN, affecting such diverse physiological and pathophysiological processes as smooth muscle cell growth, migration, survival and pericellular proteolysis activity. Clearly, PAI-1 functional blockade has widespread clinical implications that are not restricted to one aspect of the tissue response to injury.

Industry

The original small molecule panel of PAI-1 inhibitors derived from a collaboration between basic scientists and, largely, cardiovascular disease-oriented medical chemists as well as pharmaceutical investigators. This collegial relationship is expected to expand as it is evident that PAI-1 inhibition has generally utility in the treatment of cardiovascular diseases, tissue fibrosis (in particular, the renal and pulmonary systems) and cancer.

Executive Summary

Among factors implicated in the tissue response to injury, plasminogen activator inhibitor-1 (PAI-1, the major negative regulator of the plasminogen→plasmin-generating cascade, is perhaps the most pathophysiologically-relevant.

PAI-1 exerts spatial and temporal control over the integrated processes of pericellular proteolysis and matrix turnover that, collectively, impact stromal remodeling, the inflammatory response, cellular migration, proliferation and apoptosis; each of which are critical determinants in tissue fibrosis and vascular disease.

Targeted blockade of PAI-1 function with small molecule inhibitors effectively inhibits the initiation and progression of pulmonary, renal, vascular and cutaneous fibrotic disease.

Tiplaxtinin is one of the most well-studied small-molecule PAI-1 inhibitors. Tiplaxtinin attenuates asthmatic episodes, hyperlipidemia, hyperglycemia and angiogenesis.

The specific mechanism by which tiplaxtinin antagonizes the anti-fibrinolytic activity of PAI-1 involves promotion of a substrate-like conformation resulting in PAI-1 cleavage and impaired uPA and tPA inhibition, and increased vascular smooth muscle cell apoptosis.

Development of PAI-1 function-disrupting strategies may have clinical utility for the treatment of the pathophysiology of vascular disorders, thrombosis and tissue fibrosis.

Figure 1. The plasmin/matrix metalloproteinase (MMP) axis in pericellular proteolytic control
 Urokinase plasminogen activator (uPA), tethered to its cell surface receptor (uPAR), converts plasminogen receptor (PlgR)-bound plasminogen to the broad-spectrum protease plasmin that, in turn, activates several MMP family members. Collectively, plasmin and MMPs regulate the extent, duration and locale of stromal remodeling.

Figure 2. The repertoire of PAI-1 functions in the context of normal and pathologic tissue injury repair

The findings in animal model systems as well as the available clinical evidence implicate PAI-1 as a critical element in normal and dysfunctional injury repair. Use of small molecule PAI-1 inhibitors has led to the realization that PAI-1 targeting may have therapeutic implications for patients with vascular disease and various fibrotic disorders.