Published in final edited form as:

J Psychopathol Behav Assess. 2014 June ; 36(2): 201–210. doi:10.1007/s10862-013-9388-8.

Higher Levels of Psychopathy Predict Poorer Motor Control: Implications for Understanding the Psychopathy Construct

Michael D. Robinson and Konrad Bresin

Department of Psychology, North Dakota State University, Fargo, ND 58108

Abstract

A review of the literature suggests that higher levels of psychopathy may be linked to less effective behavioral control. However, several commentators have urged caution in making statements of this type in the absence of direct evidence. In two studies (total $N = 142$), moment-to-moment accuracy in a motor control task was examined as a function of dimensional variations in psychopathy in an undergraduate population. As hypothesized, motor control was distinctively worse at higher levels of psychopathy relative to lower levels, both as a function of primary and secondary psychopathy and particularly their shared variance. These novel findings provide support for the idea that motor control systematically varies by psychopathy, in a basic manner, consistent with views of psychopathy emphasizing lesser control.

Keywords

Psychopathy; Behavior; Control; Self-Control; Motor; Executive

Delete

On the basis of his clinical observations, Cleckley (1941) characterized a group of people as intelligent and non-neurotic, but behaviorally somewhat aimless. He further suggested that such psychopathic individuals could be successful, but seemed less capable of modifying their behavior on the basis of feedback or punishment. Also on the basis of clinical observations, Karpman (1941) suggested that there are several features of psychopathy, some of which are likely genetic in origin (primary psychopathy) and some of which appear to be more due to socialization factors (secondary psychopathy). Hare (1991) created a structured interview for the assessment of psychopathy among prison inmates, one that emphasized overlap among primary and secondary factors, but also linked the former to interpersonal and affective features (e.g., narcissism, callousness) and the latter to an antisocial lifestyle (e.g., earlier criminal offenses, antisocial behaviors). Subsequently, a number of additional assessments of psychopathy have been developed (e.g., Levenson, Kiehl, & Fitzpatrick, 1995; Lilienfeld & Andrews, 1996) and there is now general agreement that psychopathy encompasses several related traits and tendencies such as narcissism, impulsivity, and superficial affect.

The present studies pursue the idea that higher levels of psychopathy may be associated with lesser behavioral control. Several theories of psychopathy emphasize related attributes. For example, Cleckley (1964) characterized the erratic nature of the behavior of psychopaths, Hart and Dempster (1997) suggested that poor behavioral control is a cardinal feature of psychopathy, and Hare (1993) made a similar suggestion. In fact, poor behavioral control is an item on the Psychopathy Checklist-Revised (PCL-R: Hare, 2003), a common psychopathy assessment tool. The correlates of psychopathy, moreover, make a case for the idea that higher levels of psychopathy seem to be linked to lesser behavioral control. Individual differences in psychopathy are predictive of criminality (Lee & Salekin, 2010), classic behaviors thought to be uncontrolled (Baumeister, Heatherton, & Tice, 1994) such as substance abuse and gambling (Miller & Lynam, 2003), personality disorders of an “erratic” type (Miller, Gaughan, & Pryor, 2008), violent behaviors (Porter & Woodworth, 2006), impulsive or short-term forms of sexuality (Jonason, Li, Webster, & Schmitt, 2009), and susceptibility to temptations such as academic cheating (Nathanson, Paulhus, & Williams, 2006). Moreover, higher levels of psychopathy are marked by lower levels of agreeableness and conscientiousness (Miller, Jones, & Lynam, 2011), traits theoretically and empirically linked to better “effortful control” in developmental and personality literatures (Rothbart, Ellis, & Posner, 2011).

Although it may be intuitive to suggest that higher levels of psychopathy are inversely predictive of behavioral control, several commentators have cautioned against such assertions. For example, Poythress and Hall (2011) examined relations between psychopathy and self-reported impulsivity in a meta-analysis. Such relations were inconsistent and heterogeneous, partly due to the multi-faceted nature of self-report scales of impulsivity (Whiteside & Lynam, 2001). Reidy, Shelley-Tremblay, and Lilienfeld (2011) raised similar cautions concerning aggression. Although psychopathy predicts aggression (Jones & Paulhus, 2010), and although aggression is often thought of in terms of uncontrolled behavior (Berkowitz, 1993), there are actually two types of aggression – reactive and instrumental – only the former of which can be considered uncontrolled (Wilkowski & Robinson, 2010). Data suggest that psychopathy predicts intended or instrumental aggression to a much greater extent than reactive aggression and therefore data are not especially strong in supporting the point that psychopaths engage in uncontrolled aggression to a greater extent than non-psychopaths (Reidy et al., 2011).

There is a more basic issue here, however. Baumeister et al. (1994) considered the possibility that several of the correlates of psychopathy – such as aggression, gambling, and substance abuse – reflect lesser control over one’s behaviors. Contrary to this idea, they concluded that such behaviors *could* be controlled if the individual was motivated enough to do so. For example, substance abuse often appears partially or fully willed by the individual (West, 2006). What is needed to circumvent such ambiguities is a method of assessing control in a more basic and unambiguous manner. Fortunately, such paradigms exist in the kinesiology literature. When assessed in terms of moment-to-moment performance, it is clear that motor control is imperfect at best (Slifkin & Newell, 1998). Furthermore, moment-to-moment assessments of motor control are sensitive to factors implicated in self-control. For example, people exhibit better motor control to the extent that they receive more

frequent visual feedback (Ranganathan & Newell, 2009), consistent with an executive attention perspective of how self-control should operate (Shallice, 1988). Additionally, it has been shown that groups of people who are thought to have difficulties controlling their behavior perform more poorly in motor control tasks, including young children (Getchell, 2006), old-aged adults (Kovacs, 2005), and brain-damaged individuals (Winstein, Merians, & Sullivan, 1999). Such basic tasks can additionally be used to examine dimensional, rather than group-based, variations in controlled performance (Bresin, Fetterman, & Robinson, 2012), the focus of the present studies.

Motor control tasks may assess the extent to which one can control one's behaviors in a manner that is ambiguous when people self-report on outcomes such as aggression or substance abuse (Baumeister, Vohs, & Funder, 2007). Nonetheless, we have and will use the phrase "behavioral control" to characterize the outcomes typically examined in the psychopathy literature and the phrase "motor control" to characterize what we assess. It is our hypothesis that these two different levels of analysis can be linked – i.e., that people self-reporting higher levels of psychopathy should exhibit lesser motor control in a task designed to assess it.

Overview of Investigation

Based on several theoretical perspectives of psychopathy (Cleckley, 1964; Hare, 1993; Hart & Demster, 1997) and its correlates (Lee & Salekin, 2010; Lynam et al., 2011; Miller & Lynam, 2003), a basic deficit in motor control was hypothesized to characterize higher levels of psychopathy relative to lower levels of it. A basis for this idea is that motor control difficulties may contribute to, or at least serve as a marker of (Shallice, 1988), less effective behavioral control in more general terms (Getchell, 2006; Winstein et al., 1999). To ensure replicability, two studies were conducted. They focus on dimensional perspectives of psychopathy rather than diagnostic groups. Such studies have been recommended in that psychopathy is dimensional (Walters, Brinkley, Magaletta, & Diamond, 2008) and a full range of psychopathy scores can result in a better understanding of how this trait functions relative to a restricted range (Sadeh & Verona, 2008). Non-forensic samples are also useful in guarding against factors such as extensive drug use or a history of incarceration (Lilienfeld, 1994) that, we suggest, could quite likely affect motor control. Given the design, further work would be necessary to extend the present work to diagnostic levels of psychopathy or forensic populations.

There are two agreed-upon types of psychopathy and we sought to assess both. Primary psychopathy (sometimes termed Factor 1) taps emotional and interpersonal features of psychopathy such as narcissism, callousness, and deficient empathy. Secondary psychopathy (sometimes termed Factor 2) focuses more on behavioral aspects of psychopathy such as impulsivity, irresponsibility, and conduct problems. Primary and secondary psychopathy are moderately correlated (Hare, 1991), both predict antisocial behavior (Lee & Salekin, 2010), and both are equally marked by low levels of agreeableness (Lynam & Derefinko, 2006). In addition, Lynam, Whiteside, and Jones (1999) found both forms of psychopathy to predict commission errors in a go/no go task, results at least suggestive of the idea that motor control may be poorer for both forms of psychopathy (also see Zeier, Baskin-Sommers,

Hiatt Racer, & Newman, 2012). Secondary psychopathy, however, is a better predictor of seemingly uncontrolled behaviors such as substance abuse, aggression, and criminality (Benning, Patrick, Hicks, Blonigen, & Krueger, 2003; Falkenbach, Poythress, Falki, & Manchak, 2007), results at least suggestive of the idea that secondary psychopathy may be the better predictor of motor control difficulties. These alternative perspectives will be tested using basic probes of motor control.

We also deemed it potentially informative to manipulate aversive experiences in examining the psychopathy/motor control relationship. Possible interactions involving this manipulated factor were uncertain, however. On the one hand, there are data to suggest that psychopathic individuals are less reactive to aversive experiences in physiological terms (Patrick, 1994). On the other hand, some have suggested that links between psychopathy and controlled performance might be more evident under high stress conditions (e.g., Zeier et al., 2012). Given such discrepant ideas and findings, we could not make a directional prediction. Nonetheless, potential moderating effects seemed useful to consider, at the very least in establishing a (motor control) phenomenon that is apparent in both aversive and non-aversive contexts.

Study 1

Study 1 assessed primary and secondary psychopathy using the well-validated Levenson Self-Report Psychopathy Scale (Levenson et al., 1995). Aversive experiences were manipulated in a trial-specific manner in terms of anticipated noise blasts. Motor control was assessed using the joystick task developed by Bresin et al. (2012), as described below.

Method

Participants and Procedures—Sixty-four (37 female; 95% Caucasian) undergraduates from an upper Midwest University received psychology course credit upon completion of the study. They reported to the laboratory in groups of 6 or less and were generally told that they would complete several tasks on computer. Individuals were then seated at personal computers with dividers. The motor control task was programmed with E-Prime software and the psychopathy questionnaire was programmed with Medialab software. The motor control task occurred first to preclude possible influences of thinking about one's personality tendencies on implicit performance (Robinson & Neighbors, 2006). For conceptual reasons, though, the psychopathy assessment is described first.

Psychopathy Assessment—To ensure a degree of comparability across forensic and non-forensic samples, it may be useful to administer a psychopathy test modeled after the PCL-R (Hare, 2003), often considered the gold standard of psychopathy assessment (Lee & Salekin, 2010). Levenson et al. (1995) created such an instrument, one that is now termed the Levenson Self-Report Psychopathy (LSRP) scale. Items were modeled after the PCL-R, evidence was found for correlated yet distinct primary and secondary factors, and convergent and predictive validity for the scales was further established (Levenson et al., 1995). Subsequent investigations have provided support for the LSRP in relation to model-based cluster analyses (Falkenbach, Poythress, & Creevy, 2008), convergence with other self-report scales of psychopathy (Lynam et al., 2011), the ability to predict personality

disorder symptoms and traits (Miller et al., 2008), and sensitivity to dimensional rather than taxonomic features of psychopathy (Walters et al., 2008).

In specific terms, participants indicated the extent to which (1 = disagree strongly; 4 = agree strongly) 26 statements encompassing psychopathic attitudes and behaviors were true of the self (Levenson et al., 1995). Sixteen of these statements were designed to assess primary psychopathy (e.g., “People that are stupid enough to get ripped off usually deserve it”) and 10 were designed to assess secondary psychopathy (e.g., “I find myself in the same kinds of trouble time after time”). Means were computed by averaging across the relevant items and descriptive statistics for the scales were practically identical to those reported by Levenson et al. (1995) for both primary ($M = 1.99$; $SD = 0.44$; $\alpha = .86$) and secondary ($M = 2.06$; $SD = 0.40$; $\alpha = .63$) psychopathy. We further note that there was a good range of scores for psychopathy in both its primary (1.00–3.12) and secondary (1.40–2.90) forms and that the scales correlated with each other to a moderate degree, $r = .41$, $p < .01$.

Motor Control Task

Equipment: The E-prime program used a 32-bit distribution of Windows XP and monitors had a screen height of 13.64 inches. The screen resolution was set to 1280×1024 pixels. During the task, participants wore headphones that allowed us to present sounds of equal strength to both headphone speakers. Computers were equipped with Saitek Aviator-01 Dual Throttle joysticks connected to a USB port.

Task: To examine potential relations between psychopathy and motor control, we programmed the following task (also see Bresin et al., 2012). On each of 60 trials, a spatial target (a white + sign) was presented on the computer screen. Its position was randomly varied across trials, but excluded centered presentations along either the x- or y-axis, which would have rendered a trial easier than desired. The joystick cursor was initially presented at center screen and it was also a white + sign. Participants were told to move the joystick cursor to the presented target and hold it steadily on the stationary target thereafter. When the joystick cursor was within 5 pixels of the target, both horizontally and vertically, the target turned yellow, indicating that the motor control portion of the trial had begun. For each trial, we then sampled joystick position every 50 ms for 2000 ms (i.e., there were 40 samples collected per trial). After this 2000 ms interval, participants were instructed to return to center screen. After they had done so, and after the manipulation described next, there was a 250 ms delay until the next trial began.

A Trial-Specific Manipulation of Aversive Experience: At the beginning of each trial, participants were informed that they would (“you will receive noise”; 80 dB of white noise for 4500 ms) or would not (“you will receive silence”; 0 dB of white noise for 4500 ms) encounter a noise blast over headphones at the end of the trial, a message presented for 2000 ms to ensure its registration. Anticipating an aversive experience is very stressful according to the results of Monat, Averill, and Lazarus (1972). In addition, a pilot study established that people find white noise blasts to be quite a bit more (1 = not at all; 5 = extremely) “irritating” ($M = 3.13$) than silence ($M = 1.32$), $F(1, 84) = 192.73$, $p < .01$. An equal number of noise and silence trials occurred, randomized by trial number.

Quantification of Motor Control Difficulties: The task required participants to keep their joystick cursor as close to the target position as possible. We quantified success in maintaining this proximity in Euclidean pixel-based terms. For each sample of each trial, the Pythagorean Theorem ($c^2 = a^2 + b^2$, with a & b being position discrepancies along the x - and y -axes of the computer screen) was used to calculate the shortest distance from the sampled joystick position to the stationary target. These squared distance estimates were then averaged across samples for a given trial. Subsequently, trial scores were averaged by participant and condition such that each participant had one mean distance score for the aversive condition and one for the silence condition. As is true of many performance-based measures (Sanders, 1998), these target distance means were positively skewed. We therefore log-transformed them to reduce skew (Robinson, 2007), though results are reported in terms of non-transformed means for ease of interpretation.

Results

Overview—To simultaneously examine possible effects involving condition and psychopathy, General Linear Model (GLM) procedures were used. The GLM is a parent to both analysis of variance and multiple regression procedures, but is more flexible in handling mixed designs of the present type (Robinson, 2007). In the present case, condition was a discrete, within-subject predictor of motor control and psychopathy was a continuous, z -scored (Robinson, 2007) between-subjects predictor. Initial GLMs focus on each form of psychopathy separately, omitting sex for the sake of parsimony. Subsequently, potential interactions by sex will be examined and regressions will be performed to refine our understanding of the findings.^{1,2}

It should be emphasized that variations in primary and secondary psychopathy were treated as continuous predictors in all analyses. No analysis, that is, dichotomized people into low and high psychopathy groups. When we refer to people low and high in psychopathy, we do so on the basis of estimated (rather than actual) means (± 1 SD) along the relevant psychopathy continuum, following the procedures advocated by Aiken and West (1991).

Primary Psychopathy Findings—Motor control performance was first examined in a GLM crossing variations in primary psychopathy with the condition manipulation. As hypothesized, there was a main effect for Primary Psychopathy, $F(1, 62) = 5.55, p < .05$. To understand this effect, estimated means were calculated for prototypically low (-1 SD) versus high ($+1$ SD) levels of primary psychopathy and these estimated means are reported in the top panel of Figure 1. As shown there, motor control was 183% worse at the high relative to low level of primary psychopathy. We had thought it possible that motor control would be worse when anticipating noise, but there was no main effect for Condition, $F < 1$.

¹Aiken and West (1991) advocate standardizing continuous predictors in multiple regression analyses and Robinson (2007) extends such considerations to the GLM. Using nonstandardized personality scores can result in erroneous findings involving the repeated-measures variable (here, anticipated noise condition), though will not alter the main effects for psychopathy that are reported (Maxwell & Delaney, 2004). This is because z -scoring does not alter the shape of the distribution for a continuous predictor (Maxwell & Delaney, 2004). Thus, for example, participants' untransformed primary psychopathy scores correlated at $r = 1$ with their z -scored primary psychopathy scores.

²Separate GLMs for primary and secondary psychopathy were performed to examine whether each form of psychopathy, as an independent entity, predicted motor control. Entering both forms of psychopathy simultaneously would eliminate their common variance and the consequences of doing so are saved for a subsequently reported multiple regression.

Of potential interest, there was a Primary Psychopathy by Condition interaction, $F(1, 62) = 4.93, p < .05$. The interaction was subtle, however, in that estimated means as well as simple regressions indicated a marked effect for primary psychopathy in both the control (estimated $M_s = 62.18$ versus 190.57), $t = 2.51, p < .05$, and anticipated noise (estimated $M_s = 66.69$ versus 177.69), $t = 2.20, p < .05$, conditions, with perhaps a slightly larger effect in the control condition. Given the subtle nature of the interaction, we emphasize the main effect for primary psychopathy that occurred, though Study 2 will allow us to reexamine the possibility of an interaction with another manipulation of aversive experience.

Secondary Psychopathy Findings—To determine whether motor control varies by secondary psychopathy, a parallel GLM analysis replaced primary psychopathy scores with secondary psychopathy scores. In this analysis, there was a main effect for Secondary Psychopathy, $F(1, 62) = 4.67, p < .05$, no main effect for Condition, $F < 1$, and no Secondary Psychopathy by Condition interaction, $F < 1$. In other words, a basic, condition-invariant, relationship between secondary psychopathy and motor control was evident. Estimated means for the main effect are reported in the bottom panel of Figure 1. Motor control performance was 166% worse at the prototypically high (+1 SD) relative to low (−1 SD) level of the secondary psychopathy continuum.

Potential Interactions by Participant Sex—Men often score moderately higher than women on dimensional assessments of psychopathy (Paulhus & Williams, 2002). Nonetheless, the correlates of both primary and secondary psychopathy – in terms of personality facets and personality disorder symptoms – are highly similar for men and women (Miller et al., 2008). Accordingly, there were reasons for thinking that relations between psychopathy and motor control might be invariant across participant sex. To formally examine this possibility, we added participant sex as a factor to the GLM analyses reported above. More specifically, one GLM included the factors of primary psychopathy, sex, condition, and all two-way and three-way interactions involving these predictors; the other replaced primary psychopathy with secondary psychopathy. No effect involving sex was significant, all $p_s > .25$. Thus, higher levels of psychopathy predicted poorer motor control equally so for males and females.

Poorer Motor Control as a General Feature of Psychopathy—Irrespective of the instrument used, primary and secondary psychopathy assessments are highly correlated (Hare, 2003; Miller et al., 2008). Additionally, given that both primary and secondary psychopathy predicted motor control, it seemed likely that poorer motor control characterizes psychopathy in general terms. This point was substantiated in a multiple regression. When removing overlapping variance, neither Primary Psychopathy, $t = 1.61, p > .10$, nor Secondary Psychopathy, $t = 1.33, p > .15$, predicted motor control performance. By contrast, when averaging across primary and secondary psychopathy scores, overall levels of psychopathy *did* predict motor control difficulties in a simple regression, $t = 2.73, p < .01$, Beta = .33. At least with respect to the data collected in Study 1, then, poorer motor control appears to reflect what is common to primary and secondary psychopathy.

Study 2

Given the novel nature of the findings, a more or less direct replication effort seemed desirable. In addition, though, Study 2 sought to instantiate a stronger manipulation of aversive experience in that aversive or neutral sound clips were played over headphones *while* participants attempted motor control. Such procedures will allow us to more directly determine the relevance of emotional factors to relations between psychopathy and motor control.

Method

Participants, Procedures, and Psychopathy Assessment—A different sample of 78 (40 female; 75% Caucasian) undergraduates from an upper Midwest University received course credit. They completed the study in groups of 6 or less on personal computers, with an order of assessment parallel to Study 1. Individual differences in psychopathy were again assessed by the LSRP (Levenson et al., 1995). Descriptive statistics for primary ($M = 2.05$; $SD = 0.47$; $\alpha = .86$) and secondary ($M = 2.15$; $SD = 0.41$; $\alpha = .63$) psychopathy were very similar to Study 1. In addition, the range of scores was substantial for both primary (1.06–3.18) and secondary (1.20–3.40) psychopathy. The correlation between these two forms of psychopathy was $r = .29$, $p < .01$.

Motor Control Task—Equipment and programming for the motor control task were identical to Study 1, as was the number of trials (60). There were two key differences, however. Motor control had been assessed for 2000 ms in Study 1. To obtain more extensive data, each trial of Study 2 consisted of 6000 ms of motor control efforts. The sampling rate was 50 ms, resulting in 120 samples per trial. The other major change was the emotion-related manipulation. In Study 2, it was decided to manipulate aversive experiences during motor control efforts. Instructions stated that we were interested in abilities to perform two tasks simultaneously, one involving listening to sounds for a later memory test (which was not administered) and the other involving motor control. These instructions provided a rationale for the manipulation that occurred.³

Sound clips (6000 ms each) from the well-validated database of Bradley and Lang (1999) were used. Fifteen (e.g., screaming, vomiting) were aversive (IAD-2 #s 115, 242, 255, 260, 261, 276, 277, 279, 282, 284, 292, 296, 422, 712, & 719) and 15 (e.g., cattle sounds, yawning) were neutral (IAD-2 #s 114, 130, 246, 262, 358, 364, 368, 373, 382, 425, 627, 701, 708, 722, & 723). Along the 1–9 rating scales of Bradley and Lang (1999), the aversive sounds were more unpleasant ($M = 2.24$) than the neutral sounds ($M = 4.84$), $F(1, 28) = 370.14$, $p < .01$, and also more arousing ($M = 7.06$) than the neutral sounds ($M = 4.68$), $F(1, 28) = 75.66$, $p < .01$. Sound clips of this type have been effectively used in other psychopathy studies (e.g., Verona, Patrick, Curtin, Bradley, & Lang, 2004), though in relation to different dependent measures than ours.

³Participants were not probed for awareness of the possible influence of the manipulation on motor control. When we have probed for awareness using similar “dual task” instructions in the past, people have not spontaneously mentioned that one task involved some sort of condition difference (i.e., a manipulation) that might affect what they were subsequently asked to do.

Trial procedures consisted of the following. A white + sign appeared in a random location on the computer screen, with the constraint that it could not be centered. A white + sign, representing the joystick cursor, appeared at center screen. Participants were to move the cursor to the target and hold it as steadily as possible on the target for 6000 ms. When the cursor was within 5 pixels of the target, both horizontally and vertically, two events occurred. First, the cursor turned yellow, indicating that the motor control portion of the trial had begun. Second, a sound clip was selected and played over headphones at a loud, but reasonable volume (~ 80 dB). Sounds were chosen randomly without replacement, such that each sound was played twice and there were 30 trials of each type – neutral and aversive. At the end of 6000 ms, the sound stopped and participants were instructed to return the cursor to center screen. After they had done so, there was a 250 ms blank delay until the next trial started.

Motor control performance was quantified as in Study 1, but in the context of 120 samples per trial. After scoring each trial, means were then calculated across the 30 trials of each condition, separately so by participant. The latter means were log-transformed for analysis purposes because this transformation normalizes the skew typical of cognitive measures (Robinson, 2007), but results are reported in terms of original (non-transformed) units because such units are more intuitive in appreciating a pattern of findings (Robinson, 2007).

Results

Motor control performance was examined in two GLMs, with condition (unpleasant versus neutral) as a within-subject predictor and z-scored variations in a type of psychopathy as a continuous, between-subjects predictor. In the GLM involving primary psychopathy, there was a main effect for Primary Psychopathy, $F(1, 76) = 4.28, p < .05$, no effect for Condition, $F(1, 76) = 1.38, p > .20$, and no Primary Psychopathy by Condition interaction, $F < 1$. Estimated means for the primary psychopathy main effect are reported in the top panel of Figure 2. Motor control was 200% worse at the high relative to low level of primary psychopathy. Note that the subtle interaction of Study 1 was not replicated and is therefore not robust.

Results were similar for secondary psychopathy. In this analysis, there was a main effect for Secondary Psychopathy, $F(1, 76) = 5.02, p < .05$, no main effect for Condition, $F(1, 76) = 1.38, p > .20$, and no Secondary Psychopathy by Condition interaction, $F < 1$. Estimated means for the secondary psychopathy main effect are reported in the bottom panel of Figure 2. Motor control was 254% worse at the high relative to low level of secondary psychopathy.

As in Study 1, follow-up GLMs were performed in which sex was added to the more streamlined GLMs reported above. One, for example, examined the effects of primary psychopathy, sex, condition, and all two-way and three-way interactions among these predictors. No effect involving sex – whether main effect or interaction – was significant, all $ps > .25$.

Study 1 established that motor control varies by what is common to primary and secondary psychopathy rather than what is unique to one type of psychopathy relative to the other. We

performed two regression-based analyses to determine whether the same was true in Study 2. We first conducted a multiple regression in which the common variance between primary and secondary psychopathy was controlled for. When removing this common variance, neither Primary Psychopathy, $t = 1.40, p > .15$, nor Secondary Psychopathy, $t = 1.82, p > .05$, predicted motor control deficits. When averaging primary and secondary psychopathy, the overall psychopathy score was a robust predictor of motor control, $t = 2.72, p < .01$, Beta = .30. The Study 2 results therefore closely replicate those of Study 1 in suggesting that people higher in psychopathy, regardless of its form, have poorer motor control.

General Discussion

Two studies pursued the idea that higher levels of psychopathy can be linked to less effective motor control. To circumvent concerns about self-reports (Baumeister et al., 2007), an objective performance task was used. The task was one in which control cannot be perfect, rendering it sensitive to individual differences in the executive control of the motoric system (Newell, Mayer-Kress, & Liu, 2009). Both studies found inverse relations between psychopathy and motor control. Estimated means differed greatly by psychopathy and motor control deficits were common to its primary and secondary forms. The relationships were general in nature in that they were not exacerbated by manipulations of aversive experience. The findings are discussed in terms of their implications for understanding individual differences in psychopathy and future research directions are also presented.

Executive Functions and Motor Control in Psychopathy

Individual differences in “executive functions” – i.e., those mediated by the frontal cortex (Goldberg, 2009) – have primarily been investigated in tasks that might be viewed as inhibition-related in nature. For example, the classic Stroop task (MacLeod, 1991) requires people to inhibit distracting sources of information present on some trials relative to other trials. The motor control tasks administered in the present studies were quite different. First, motor output rather than cognition was to be controlled. Second, there were no congruent or incongruent conditions. Rather, every trial assessed the extent to which control was effectively administered. Third, we could quantify moment-to-moment variations in control on individual trials in a manner consistent with the kinesiology literature (Slifkin & Newell, 1998) relative to quantifications in which there is only one data-point per trial (Sanders, 1998).

Such considerations aside, it is useful to discuss the present findings in relation to work on psychopathy and executive processing deficits owing to this work’s similar focus on deficits in control. Two studies (Blair et al., 2006; Lapierre, Braun, & Hodgins, 1995) have linked psychopathy to poorer performance in tasks thought to be mediated by the orbitofrontal cortex (OFC), but not to poorer performance on other tasks thought to be mediated by the dorsolateral prefrontal cortex (dlPFC). The OFC plays a role in reversing previously reinforced associations (Rolls, 2005), but it also plays a role in mitigating impulsive responding (Lapierre et al., 1995), including in tasks such as the go/no go task (Roussy & Toupin, 2000). Along the latter lines, Lynam et al. (1999) found that both primary and secondary psychopathy were predictive of commission errors in go/no go tasks. Our

findings, we think, should be viewed as compatible with the Lynam et al. (1999) findings and with this OFC perspective of psychopathy.

By contrast, predictions for more traditional interference-related dlPFC tasks such as the flanker effect (Kerns et al., 2004) are more complicated (Zeier et al., 2012). There are sources of data to suggest that psychopaths may “gate out” irrelevant perceptual inputs at an early level of processing (e.g., Hiatt, Schmitt, & Newman, 2004). Accordingly, it may not be surprising that links of psychopathy to such traditional executive processing tasks are not particularly consistent across studies (Ogilvie, Stewart, Chan, & Shum, 2011). Our task was not of this dlPFC type in that cognitive sources of interference were not manipulated. It is likely in part for this reason that robust links of psychopathy to motor control deficits were found.

Secondary psychopathy, relative to primary psychopathy, is often the better predictor of seemingly dysregulated outcomes such as substance abuse, aggression, and criminality (Benning et al., 2003; Falkenbach et al., 2007). Yet, both forms of psychopathy have been linked to traits thought to reflect lower effortful control (Miller et al., 2011), personality disorder symptoms of an “erratic” type (Miller et al., 2008), and antisocial behaviors and aggression (Sadeh & Verona, 2008). In addition, there are theories (e.g., Moffitt, 1993) and sources of data (e.g., Morgan & Lilienfeld, 2000) indicating that antisocial behavior, which is common to primary and secondary psychopathy (Sadeh & Verona, 2008), is predictive of deficits in executive functioning. On the basis of our findings, something like poorer effortful control, which is common to both primary and secondary psychopathy (Miller et al., 2008), appears to underlie problems with motor control. This would explain why controlling for overlapping variance among primary and secondary psychopathy resulted in a non-significant prediction of motor control deficits.

Even so, the psychopathy inventory administered may have a limitation in that the primary psychopathy scale of the LSRP (Levenson et al., 1995) appears more closely aligned with its secondary psychopathy scale than is the case when using the revised Psychopathic Personality Inventory, in which the primary factor has been termed fearless-dominance (PPI-R: Poythress et al., 2010). It is therefore possible that dissociations involving primary and secondary psychopathy could be found with the PPI-R scales, though it should also be mentioned that there are concerns related to the operationalization of primary psychopathy in the PPI-R (Miller & Lynam, 2012) and the LSRP scales continue to have broad support (Falkenbach et al., 2007; Miller et al., 2008). Nonetheless, we suggest that motor control could be revisited in the context of alternative assessments of the various factors of psychopathy.

Aversive experiences did not interact with psychopathy to predict motor control. The results are therefore consistent with the idea that the psychopathy/control interface can be a basic one (e.g., Morgan & Lilienfeld, 2000) that is independent of the emotional processing features that have sometimes been emphasized in this literature (e.g., Lykken, 1995). It should be recognized, though, that the manipulations of aversive experience did not affect motor control in main effect terms, despite the within-subject power and strength (e.g., concurrent audio clips: Study 2) of this feature of the design. It is possible that affective

manipulations that are more self-relevant (e.g., error feedback: Moeller & Robinson, 2010) might have a larger influence on motor control. Alternatively, motor control may primarily vary by factors that are chronic and trait-related rather than temporary and state-related. Future research using other manipulations will be necessary to decide among these two possibilities.

Potential Clinical Significance and Future Research Directions

Our focus was on basic processes, but the findings possess potential clinical significance. Meyer and Kurtz (2006) have highlighted the limitations of self-reports in the clinical realm. Implicit assessments such as ours can bypass some such limitations. For example, it is doubtful that one could “fake” higher levels of motor control when one lacks it. With appropriate norms, motor control performance may aid case conceptualization much as intelligence tests currently do (Kazdin, 2008). Further, difficulties in motor control may alert the client and the assessor or therapist to targets of intervention (Shedler & Westen, 2007). In turn, changes in motor control might inform treatment progress, in that successful treatment may be expected to improve motor control (Wiers & Stacy, 2006). Finally, it is conceivable that training motor control might have some therapeutic benefit. This suggestion is plausible given current theories of self-regulation (Carver & Scheier, 1998; Muraven & Baumeister, 2000) and it is also consistent with cognitive retraining results in the mood disorder literature (Yiend, 2004). At the very least, our findings highlight a robust correlate of psychopathy that deserves applied research attention.

There are further limitations to the present results and therefore several additional directions of research that can be advocated. It would be useful to conceptually replicate the present results using other tasks in which motor control can be assessed such as pinch grip force tasks or circle tracing tasks (see Slifkin & Newell, 1998, for an overview of some of these tasks). Especially high levels of psychopathy, or at least its criminal manifestations, might not be especially prevalent among college undergraduates and replication with other – e.g., forensic (Hare, 1993) – samples would be desirable. We focused on the dimension of psychopathy rather than its correlates and it would therefore be useful to investigate whether motor control difficulties can predict psychopathy-linked outcomes such as aggression (Reidy et al., 2011) and/or substance use (Miller & Lynam, 2003). In this connection, we suspect that poorer motor control may be a robust predictor of externalizing behaviors of multiple types (e.g., those reviewed by Krueger et al., 2009) though establishing this point requires additional research. The predictive value of motor control might be further established by using longitudinal designs as the prediction of future externalizing problems may be of more clinical value than the prediction of problematic behaviors that have already occurred. Altogether, research of the type outlined would be useful in further validating the idea that people who display poorer motor control in basic tasks are likely to display self-control problems more generally considered.

References

- Aiken, L.S.; West, S.G. Multiple regression: Testing and interpreting interactions. Thousand Oaks, CA: Sage Publications; 1991.

- Baumeister, RF.; Heatherton, TF.; Tice, DM. *Losing control: How and why people fail at self-regulation*. San Diego, CA: Academic Press; 1994.
- Baumeister RF, Vohs KD, Funder DC. Psychology as the science of self-reports and finger movements: What happened to actual behavior? *Perspectives in Psychological Science*. 2007; 2:396–403.
- Benning SD, Patrick CJ, Hicks BM, Blonigen DM, Krueger RF. Factor structure of the Psychopathic Personality Inventory: Validity and implications for clinical assessment. *Psychological Assessment*. 2003; 15:340–350. [PubMed: 14593834]
- Berkowitz, L. *Aggression: Its causes, consequences, and control*. New York: McGraw-Hill Book Company; 1993.
- Blair KS, Newman C, Mitchell DGV, Richell RA, Leonard A, Morton J, et al. Differentiating among prefrontal substrates in psychopathy: Neuropsychological test findings. *Neuropsychology*. 2006; 20:153–165. [PubMed: 16594776]
- Bradley, MM.; Lang, PJ. *International affective digitized sounds (IADS): Stimuli, instruction manual and affective ratings*. Gainesville, FL: The Center for Research in Psychopathology, University of Florida; 1999.
- Bresin K, Fetterman AK, Robinson MD. Motor control accuracy: A consequential probe of individual differences in emotion regulation. *Emotion*. 2012; 12:479–486. [PubMed: 22023366]
- Carver, CS.; Scheier, MF. *On the self-regulation of behavior*. New York: Cambridge University Press; 1998.
- Cleckley, H. *The mask of sanity*. St Louis, MO: Mosby; 1941.
- Cleckley, H. *The mask of sanity*. 4. St. Louis, MO: Mosby; 1964.
- Falkenbach D, Poythress N, Creevy C. The exploration of subclinical psychopathic subtypes and the relationship with types of aggression. *Personality and Individual Differences*. 2008; 44:821–832.
- Falkenbach D, Poythress N, Falki M, Manchak S. Reliability and validity of two self-report measures of psychopathy. *Assessment*. 2007; 14:341–350. [PubMed: 17986652]
- Getchell N. Age and task-related differences in timing stability, consistency, and natural frequency of children's rhythmic, motor coordination. *Developmental Psychobiology*. 2006; 48:675–685. [PubMed: 17111404]
- Goldberg, E. *The new executive brain: Frontal lobes in a complex world*. New York: Oxford University Press; 2009.
- Hare, RD. *The Hare Psychopathy Checklist-Revised*. Toronto, Ontario, Canada: Multi-Health Systems; 1991.
- Hare, RD. *Without conscience: The disturbing world of the psychopaths among us*. New York: Guilford; 1993.
- Hare, RD. *Manual for the Hare Psychopathy Checklist-Revised*. 2. Toronto, Ontario, Canada: Multi-Health Systems; 2003.
- Hart, SD.; Dempster, RJ. Impulsivity and psychopathy. In: Webster, CD.; Jackson, MA., editors. *Impulsivity: Theory, assessment, and treatment*. New York: Guilford; 1997. p. 212–232.
- Hiatt KD, Schmitt WA, Newman JP. Stroop tasks reveal abnormal selective attention among psychopathic offenders. *Neuropsychology*. 2004; 18:50–59. [PubMed: 14744187]
- Jonason PK, Li NP, Webster GD, Schmitt DP. The dark triad: Facilitating a short-term mating strategy in men. *European Journal of Personality*. 2009; 23:5–18.
- Jones DN, Paulhus DL. Different provocations trigger aggression in narcissists and psychopaths. *Social Psychological and Personality Science*. 2010; 1:12–18.
- Karpman B. On the need of separating psychopathy into two distinct clinical types: The symptomatic and the idiopathic. *Journal of Criminal Psychopathology*. 1941; 3:112–137.
- Kazdin AE. Evidence-based treatment and practice: New opportunities to bridge clinical research and practice, enhance the knowledge base, and improve patient care. *American Psychologist*. 2008; 63:146–159. [PubMed: 18377105]
- Kerns JG, Cohen JD, MacDonald AW, Cho RY, Stenger VA, Carter CS. Anterior cingulate conflict monitoring and adjustments in control. *Science*. 2004; 303:1023–1026. [PubMed: 14963333]

- Kovacs CR. Age-related changes in gait and obstacle avoidance capabilities in older adults: A review. *Journal of Applied Gerontology*. 2005; 24:21–34.
- Krueger, RF.; Hicks, BM.; Patrick, CJ.; Carlson, SR.; Iacono, WG.; McGue, M. Etiologic connections among substance dependence, antisocial behavior, and personality: Modeling the externalizing spectrum. In: Marlatt, GA.; Witkiewitz, K., editors. *Addictive behaviors: New readings on etiology, prevention, and treatment*. Washington, DC: American Psychological Association; 2009. p. 59-88.
- Lapierre, Braun; Hodgins, S. Ventral frontal deficits in psychopathy: Neuropsychological test findings. *Neuropsychologia*. 1995; 33:139–151. [PubMed: 7746360]
- Lee Z, Salekin RT. Psychopathy in a noninstitutional sample: Differences in primary and secondary subtypes. *Personality Disorders: Theory, Research, and Treatment*. 2010; 1:153–169.
- Levenson MR, Kiehl KA, Fitzpatrick CM. Assessing psychopathic attributes in a noninstitutionalized population. *Journal of Personality and Social Psychology*. 1995; 68:151–158. [PubMed: 7861311]
- Lilienfeld SO. Conceptual problems in the assessment of psychopathy. *Clinical Psychology Review*. 1994; 14:17–38.
- Lilienfeld SO, Andrews BP. Development and preliminary validation of a self-report measure of psychopathic personality traits in noncriminal populations. *Journal of Personality Assessment*. 1996; 66:488–524. [PubMed: 8667144]
- Lykken, DT. *The antisocial personalities*. Hillsdale, NJ: Lawrence Erlbaum Associates; 1995.
- Lynam, DR.; Derefinko, KJ. Psychopathy and personality. In: Patrick, CJ., editor. *Handbook of psychopathy*. New York: Guilford; 2006. p. 133-155.
- Lynam DR, Gaughan ET, Miller JD, Miller DJ, Mullins-Sweatt S, Widiger TA. Assessing the basic traits associated with psychopathy: Development and validation of the Elementary Psychopathy Assessment. *Psychological Assessment*. 2011; 23:108–124. [PubMed: 21171784]
- Lynam DR, Whiteside S, Jones S. Self-reported psychopathy: A validation study. *Journal of Personality Assessment*. 1999; 73:110–132. [PubMed: 10497804]
- MacLeod CM. Half a century of research on the Stroop effect: An integrative review. *Psychological Bulletin*. 1991; 109:163–203. [PubMed: 2034749]
- Maxwell, SE.; Delaney, HD. *Designing experiments and analyzing data: A model comparison perspective*. 2. Mahwah, NJ: Lawrence Erlbaum Associates; 2004.
- Meyer GJ, Kurtz JE. Advancing personality assessment technology: Time to retire ‘objective’ and ‘projective’ as personality test descriptors. *Journal of Personality Assessment*. 2006; 87:223–225. [PubMed: 17134328]
- Miller JD, Gaughan ET, Pryor LR. The Levenson Self-Report Psychopathy Scale: An examination of the personality traits and disorders associated with the LSRP factors. *Assessment*. 2008; 15:450–464. [PubMed: 18480502]
- Miller JD, Jones SE, Lynam DR. Psychopathic traits from the perspective of self and informant reports: Is there evidence for a lack of insight? *Journal of Abnormal Psychology*. 2011; 120:758–764. [PubMed: 21381803]
- Miller JD, Lynam DR. Psychopathy and the Five-factor model of personality: A replication and extension. *Journal of Personality Assessment*. 2003; 81:168–178. [PubMed: 12946923]
- Miller JD, Lynam DR. An examination of the Psychopathic Personality Inventory’s nomological network: A meta-analytic review. *Personality Disorders: Theory, Research, and Treatment*. 2012; 3:305–326.
- Moeller SK, Robinson MD. Cognitive sources of evidence for neuroticism’s link to punishment-sensitivity processes. *Cognition and Emotion*. 2010; 24:741–759.
- Moffitt TE. The neuropsychology of conduct disorder. *Development and Psychopathology*. 1993; 5:135–152.
- Monat A, Averill JR, Lazarus RS. Anticipatory stress and coping reactions under various conditions of uncertainty. *Journal of Personality and Social Psychology*. 1972; 24:237–253. [PubMed: 5081195]
- Morgan AB, Lilienfeld SO. A meta-analytical review of the relation between antisocial behavior and neuropsychological measures of executive function. *Clinical Psychology Review*. 2000; 20:113–156. [PubMed: 10660831]

- Muraven M, Baumeister RF. Self-regulation and depletion of limited resources: Does self-control resemble a muscle? *Psychological Bulletin*. 2000; 126:247–259. [PubMed: 10748642]
- Nathanson C, Paulhus DL, Williams KM. Predictors of a behavioral measure of scholastic cheating: Personality and competence but not demographics. *Contemporary Educational Psychology*. 2006; 31:97–122.
- Newell KM, Mayer-Kress G, Liu Y. Aging, time scales, and sensorimotor variability. *Psychology and Aging*. 2009; 24:809–818. [PubMed: 20025397]
- Ogilvie JM, Stewart AL, Chan RCK, Shum DHK. Neuropsychological measures of executive function and antisocial behavior: A meta-analysis. *Criminology: An Interdisciplinary Journal*. 2011; 49:1063–1107.
- Patrick CJ. Emotion and psychopathy: Startling new insights. *Psychophysiology*. 1994; 31:319–330. [PubMed: 10690912]
- Paulhus DL, Williams KM. The Dark Triad of personality: Narcissism, Machiavellianism and psychopathy. *Journal of Research in Personality*. 2002; 36:556–563.
- Porter, S.; Woodworth, M. Psychopathy and aggression. In: Patrick, CJ., editor. *Handbook of psychopathy*. New York: Guilford; 2006. p. 481–494.
- Poythress NG, Edens JF, Skeem JL, Lilienfeld SO, Douglas KS, Frick PJ, et al. Identifying subtypes among offenders with antisocial personality disorder: A cluster-analytic study. *Journal of Abnormal Psychology*. 2010; 119:389–400. [PubMed: 20455611]
- Poythress NG, Hall JR. Psychopathy and impulsivity reconsidered. *Aggression and Violent Behavior*. 2011; 16:120–134.
- Ranganathan R, Newell KM. Influence of augmented feedback on coordination strategies. *Journal of Motor Behavior*. 2009; 41:317–330. [PubMed: 19508958]
- Reidy DE, Shelley-Tremblay JF, Lilienfeld SO. Psychopathy, reactive aggression, and precarious proclamations: A review of behavioral, cognitive, and biological research. *Aggression and Violent Behavior*. 2011; 16:512–524.
- Robinson, MD. Lives lived in milliseconds: Using cognitive methods in personality research. In: Robins, RW.; Fraley, RC.; Krueger, R., editors. *Handbook of research methods in personality psychology*. New York: Guilford Press; 2007. p. 345–359.
- Robinson, MD.; Neighbors, C. Catching the mind in action: Implicit methods personality research and assessment. In: Eid, M.; Diener, E., editors. *Handbook of multimethod measurement in psychology*. Washington, DC: American Psychological Association; 2006. p. 115–125.
- Rolls, ET. What are emotions, why do we have emotions, and what is their computational basis the brain?. In: Fellous, J.; Arbib, MA., editors. *Why need emotions? The brain meets the robot*. New York: Oxford University Press; 2005. p. 117–146.
- Rothbart, MK.; Ellis, LK.; Posner, MI. Temperament and self-regulation. In: Vohs, KD.; Baumeister, RF., editors. *Handbook of self-regulation: Research, theory, and applications*. 2. New York: Guilford; 2011. p. 441–460.
- Roussy S, Toupin J. Behavioral inhibition deficits in juvenile psychopaths. *Aggressive Behavior*. 2000; 26:413–424.
- Sadeh N, Verona E. Psychopathic personality traits associated with abnormal selective attention and impaired cognitive control. *Neuropsychology*. 2008; 22:669–680. [PubMed: 18763886]
- Sanders, AF. *Elements of human performance: Reaction processes and attention in human skill*. Mahwah, NJ: Lawrence Erlbaum Associates; 1998.
- Shallice, T. *From neuropsychology to mental structure*. New York: Cambridge University Press; 1988.
- Shedler J, Westen D. The Shedler-Westen Assessment Procedure (SWAP): Making personality diagnosis clinically meaningful. *Journal of Personality Assessment*. 2007; 89:41–55. [PubMed: 17604533]
- Slifkin AB, Newell KM. Is variability in human performance a reflection of system noise? *Current Directions in Psychological Science*. 1998; 7:170–177.
- Verona E, Patrick CJ, Curtin JJ, Bradley MM, Lang PJ. Psychopathy and physiological response to emotionally evocative sounds. *Journal of Abnormal Psychology*. 2004; 113:99–108. [PubMed: 14992662]

- Walters GD, Brinkley CA, Magaletta PR, Diamond PM. Taxometric analysis of the Levenson Self-Report Psychopathy Scale. *Journal of Personality Assessment*. 2008; 90:491–498. [PubMed: 18704808]
- West, R. *Theory of addiction*. Oxford, England: Blackwell Publishing; 2006.
- Whiteside SP, Lynam DR. The Five Factor Model and impulsivity: Using a structural model of personality to understand psychopathy. *Personality and Individual Differences*. 2001; 30:669–689.
- Wiers, RW.; Stacy, AW. *Handbook of implicit cognition and addiction*. Thousand Oaks, CA: Sage Publications; 2006.
- Wilkowski BM, Robinson MD. The anatomy of anger: An integrative cognitive model of trait anger and reactive aggression. *Journal of Personality*. 2010; 78:9–38. [PubMed: 20433611]
- Winstein CJ, Merians AS, Sullivan KJ. Motor learning after unilateral brain damage. *Neuropsychologia*. 1999; 37:975–987. [PubMed: 10426521]
- Yiend, J. *Cognition, emotion and psychopathology: Theoretical, empirical and clinical directions*. New York: Cambridge University Press; 2004.
- Zeier JD, Baskin-Sommers AR, Hiatt Racer KD, Newman JP. Cognitive control deficits associated with antisocial personality disorder and psychopathy. *Personality Disorders: Theory, Research, and Treatment*. 2012; 3:283–293.

Figure 1. Motor Control Difficulty Estimated Means as a Function of Primary Psychopathy (Top Panel) and Secondary Psychopathy (Bottom Panel), Study 1

Figure 2. Motor Control Difficulty Estimated Means as a Function of Primary Psychopathy (Top Panel) and Secondary Psychopathy (Bottom Panel), Study 2