Published in final edited form as:

Am J Clin Dermatol. 2014 December ; 15(6): 537–542. doi:10.1007/s40257-014-0090-8.

Isotretinoin Use and Celiac disease: a Population-Based Cross-Sectional Study

Benjamin Lebwohl, MD, MS^{1,2}, Anders Sundström³, Bana Jabri⁴, Sonia S. Kupfer⁴, Peter HR Green, MD¹, and Jonas F. Ludvigsson, MD, PhD^{2,5,*}

¹Celiac Disease Center, Department of Medicine, Columbia University College of Physicians and Surgeons, New York, New York, USA

²Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden

³Clinical Epidemiology Unit, Karolinska Institutet, Stockholm, Sweden

⁴University of Chicago, Celiac Disease Center, Chicago, IL, USA

⁵Department of Paediatrics, Örebro University Hospital, Sweden

Abstract

Background and aim—Isotretinoin, a vitamin A analogue, can promote a pro-inflammatory milieu in the small intestine in response to dietary antigens. We hypothesized that oral isotretinoin exposure would increase the risk of celiac disease (CD).

Methods—We contacted all 28 pathology departments in Sweden, and through biopsy reports identified 26,739 individuals with CD. We then compared the prevalence of ever using oral isotretinoin to the prevalence in 134,277 matched controls through conditional logistic regression. Data on isotretinoin exposure were obtained from the national Swedish Prescribed Drug Registry.

Correspondence and reprint requests: Jonas F Ludvigsson, Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Sweden, jonasludvigsson@yahoo.com.

Authors' contributions

ICMJE criteria for authorship read and met: BL, AS, BJ, SSK, PG, JFL.

Agree with the manuscript's results and conclusions: BL, AS, BJ, SSK, PG, JFL.

Designed the experiments/the study: BL, JFL.

Collected data: JFL

Analyzed the data: BL

Wrote the first draft of the paper: BL and JFL.

Contributed to study design, interpretation of data and writing: AS, PG.

Interpretation of data; approved the final version of the manuscript: BL, AS, BJ, SSK, PG, JFL

Responsible for data integrity: JFL.

Obtained funding: JFL.

Details of ethics approval: This project (2006/633-31/4) was approved by the Research Ethics Committee of the Karolinska Institute, Sweden on June 14, 2006.

Competing interests

Dr. Lebwohl: The author declares that he has no conflict of interest.

Dr. Sundström: The author declares that he has no conflict of interest.

Dr. Jabri: The author declares that she has no conflict of interest.

Dr. Kupfer: The author declares that she has no conflict of interest.

Dr. Green: The author declares that he has no conflict of interest.

Dr. Ludvigsson: The author declares that he has no conflict of interest.

As the only indication for isotretinoin use in Sweden is acne, we also examined its relationship to CD. Data on acne were obtained from the Swedish Patient Registry.

Results—93 (0.35%) individuals with CD and 378 (0.28%) matched controls had a prescription of isotretinoin. This corresponded to an odds ratio (OR) of 1.22 (95%CI=0.97-1.54). Risk estimates were similar in men and women, and when we restricted our data to individuals diagnosed after the start of the Prescribed Drug Registry. Restricting our analyses to individuals diagnosed aged 12-45 years did not influence the risk estimates (OR=1.38; 0.97-1.97). Meanwhile, having a diagnosis of acne was positively associated with CD (OR 1.34, 95%CI 1.20-1.51).

Conclusions—This study found no association between isotretinoin use and CD, but a small excess risk of CD in patients with a diagnosis of acne.

Keywords

autoimmunity; celiac; inflammation; isotretinoin

INTRODUCTION

Celiac disease (CD) is characterized by small intestinal inflammation and occurs in about 1% of the Western population[1, 2]. Individuals with CD are at increased risk of a number of disorders including lymphoproliferative malignancy[3], neuropathy[4], type 1 diabetes[5] and thyroid disease[5].

Other than gluten exposure and DQ2/DQ8[6], short duration of breastfeeding [7] may be an important risk factor. Breastfeeding is known to influence the microbiota[8] in the child, and patients with CD have an aberrant duodenal microbiota composition[9, 10]. But breastfeeding may also protect against intestinal inflammation[11], and we hypothesized that substances that induce intestinal inflammation may also increase the risk of CD.

Retinoic acid plays an important role for the regulation of the intestinal immune system[12]. Together with IL-15[13], retinoic acid seems to promote inflammatory responses to dietary antigens[14]. Oral isotretinoin, a form of retinoic acid, has been implicated in one study to trigger inflammatory bowel disease,[15] though this association has not been confirmed[16, 17].

In this study we examined for an association between oral isotretinoin (13-cis-retinoic acid) and CD. To do so we matched data from the Swedish Prescribed Drug Registry to histopathology data from 26,739 individuals with CD. A secondary aim was to examine the association between acne and CD.

METHODS

Data Sources

During the years spanning 2006-2008, we obtained computerized biopsy reports from all 28 Swedish pathology departments[19]. These biopsies had been carried out between 1969 and 2008. Searches for biopsy reports were carried out by local information technology personnel who delivered data on personal identity number[20], morphology (for a list of

relevant morphology codes according to the Swedish SnoMed system[21]), topography (duodenum or jejunum), and date of biopsy. Cases were determined based on these biopsies. Controls were provided by the government agency, Statistics Sweden, using the matching parameters described below. After identifying cases and controls, we cross-checked these individuals with the Swedish Prescribed Drug Registry. The Prescribed Drug Registry began on July 1, 2005[18], and accounts for some 84% of the total drug utilization (>99% of all dispensed drugs) in Sweden.

Study Design

We performed a population-based cross-sectional study of patients in Sweden with biopsy-diagnosed CD (i.e. cases) and non-CD controls. We measured for an association between isotretinoin prescription and case status. In this cross-sectional study, we included isotretinoin exposure that took place both before and after CD diagnosis (and the corresponding date of matching in controls). Exposure was measured regardless of temporal relationship with case/control ascertainment due to the fact that the Swedish Prescribed Drug Registry commenced in 2005, while the CD patients and controls were ascertained during the years spanning 1969-2008. A sensitivity analysis that incorporated temporality (see *Statistical Analysis* below) resulted in fewer than 50 prescriptions for isotretinoin in the whole cohort.

Case Status - Celiac disease

We defined CD as having villous atrophy (histopathology stage Marsh 3)[19] according to a biopsy report. As these were the first instances of villous atrophy for individuals, these were regarded as incident cases. In a patient chart validation of 114 randomly selected individuals with villous atrophy, 108 (95%) had CD.[21] The biopsy reports were on average based on three tissue specimens,[22] which should rule in approximately 95% of all CD cases[23]. Some 88% of patients with villous atrophy has positive celiac serology at time of celiac diagnosis (defined as date of first biopsy with villous atrophy).[21]

Controls

The government agency Statistics Sweden matched each CD individual with up to 5 controls, using the following matching criteria, on the date of CD diagnosis of their corresponding case: age, sex, county of residence at time of biopsy, and calendar year. Controls were sampled from the Swedish Total population register with the exception that they should not have an earlier record of small intestinal biopsy prior to the date of CD diagnosis of their corresponding case.

Sample Size

After removal of duplicates and other data irregularities our data file was identical to that of our study on mortality in CD [24]. We then excluded patients who died or emigrated prior to July 1, 2005, since the Prescribed Drug Registry started on this date, and therefore our final dataset consisted of 26,739 individuals with CD and 134,277 matched controls.

Exposure variable

Exposure information was derived from the Swedish Prescribed Drug Registry. Isotretinoin (oral use) was identified through the relevant anatomical therapeutic chemical (ATC) code (D10BA01) in Patients who died or emigrated prior to July 1, 2005 (the inception of the Swedish Prescribed Drug Registry) were excluded from this analysis.

Statistical Analysis

We used conditional logistic regression to calculate odds ratios (ORs) for the association between CD and ever oral isotretinoin use. Each individual with CD was compared to his or her matched controls within the same stratum, with regard to isotretinoin exposure. This eliminated the effect of age, sex, county and calendar year on our risk estimate. We then summarized stratum-specific data and calculated an overall OR for CD.

In *a priori* secondary analyses, we tested for the association between CD and isotretinoin use stratified by sex and restricted to the age group spanning 12-45 years, as this group is most likely to be exposed to isotretinoin[15]. So as to detect a possible dose-response effect, we measured point estimates based on the number of prescriptions of isotretinoin (none, <3 prescriptions, or 3 prescriptions).

In an additional pre-planned analysis, we adjusted for education according to four *a priori*-defined categories.[26] We used data on education as proxy for socioeconomic status (Four groups: 9 years of primary and secondary school, 2 years of high school (usually programs for manual, clerical or assistance work), 3 years of high school (theoretical programs) and college/university studies)[25]. These data were obtained from the Swedish Education Register maintained by Statistics Sweden. Some 4% of individuals had no data on education and were fitted into a separate fifth category in the multivariate analysis.

In a sensitivity analysis, we subsequently restricted our calculations to those patients with CD and corresponding controls who were diagnosed subsequent to July 1, 2005, the date of inception of the Swedish Prescription Drug Register.

A post-hoc power analysis found that we had an 80% (at 0.05 significance level) to detect a relative risk of 1.39 for CD among isotretinoin users.

Although it is not known whether acne is more common in patients with CD, given the possibility that an observed relationship between isotretinoin use and CD could be influenced by acne, in our model we also examined the OR for CD among patients with a diagnosis of acne. Data on acne obtained from the Swedish patient registry and defined according to the following ICD codes: (ICD-7: 714,1; ICD-8: 706; ICD-9: 706; ICD-10: L70). While we did measure the association between an acne diagnosis and CD, we did not adjust our isotretinoin data for acne since acne per se is the only medical indication for isotretinoin in Sweden.

We used SAS version 9.2 (Cary, NC, USA) for all analyses. We report ORs with corresponding 95% confidence intervals. All p values reported are 2-sided.

Ethics

The Ethics Review board of Stockholm, Sweden, approved this study and deemed that no individual informed consent was required since data were strictly register-based. The study and analysis did not deviate from the approved protocol.

RESULTS

Some 63% of the study participants were female (Table 1). The median age at CD diagnosis was 27 years (interquartile range 5-50 years). As reported previously, the median year of diagnosis was 1998 and the vast majority of both cases (CD: 96.7%) and controls (94.3%) were born in the Nordic countries.[24] The median age at first exposure to isotretinoin was 18 years, with the youngest subject being 5 years, and the eldest 69 years.

Some 93 (0.35%) individuals with CD and 378 (0.28%) matched controls had a prescription of oral isotretinoin. Individuals with isotretinoin were at a non-significantly increased risk of CD (Odds ratio (OR)=1.22; 95% CI=0.97-1.54). Additional adjustment for socioeconomic status did not influence our risk estimates (Table 2).

ORs were similar in men and women (Table 2), and when we restricted our data to individuals diagnosed after the start of the Prescribed Drug Registry (July 1, 2005).

In a subanalysis we examined the association between amount of isotretinoin and risk of CD. Few individuals had received 3 prescriptions during this time period (2 individuals with CD and 3 controls). This corresponded to an OR (compared to no exposure) of 3.33 (95% CI=0.56-19.95)(Table 2).

A diagnosis of acne was present in 363 (1.4%) patients with CD and 1,354 (1.0%) controls (OR 1.34, 95% CI 1.20-1.51, $p<0.0001$).

DISCUSSION

In this nationwide case-control study we linked data from biopsy records with CD from all Swedish pathology departments with data from the Swedish Prescribed Drug Registry. We found no association between oral isotretinoin exposure and CD, but a statistically significant positive association with acne, and we cannot rule out that this latter association has influenced the OR for CD among isotretinoin-users in our study.

We are not aware of any studies on isotretinoin and CD but one earlier study has suggested that this substance could trigger ulcerative colitis[15]. However, this positive association with inflammatory bowel disease has been contradicted by one case-control study[16], and by a recent meta-analysis based on 2159 inflammatory bowel disease cases and more than 40,000 controls (Etminan *et al*: relative risk=0.99 with a 95% CI=0.52-1.90)[17].

While the vitamin A metabolite, retinoic acid, can induce intestinal regulatory immune responses in concert with transforming growth factor β (TGF- β)[12, 27], recent work has shown that, under conditions of intestinal stress such as in CD with high levels of IL-15, Retinoic acid prevents rather promotes oral tolerance[14]. Specifically, retinoic acid

activates dendritic cells to induce the release of pro-inflammatory cytokines IL12p70 and IL-23 that lead to decreased regulatory T cell differentiation. Intriguingly, children with vitamin A deficiency show less efficient response to oral vaccine therapy[28, 29] and studies in mice have shown that in the absence of vitamin A mice cannot mount efficient mucosal vaccine responses and protective immunity against pathogens [30]. Altogether, these findings suggest that Retinoic acid may have adjuvant effects and promote inflammation. In line with this hypothesis, a link between isotretinoin and inflammatory bowel disease was proposed [31]. Importantly, it is the context in which retinoic acid operates that determines whether retinoic acid will promote inflammation or tolerance in the gut. Hence, retinoic acid would promote CD only in individuals who already have upregulation of IL-15 in the gut such as those individuals who experience an episode of gastroenteritis (rotavirus or campylobacter) while taking isotretinoin. Whereas in others retinoic acid may promote immune-regulatory effects in the absence of IL-15 upregulation. Unfortunately we lack data on IL-15 in this study.

Albeit not statistically significant, isotretinoin users were at a 22% increased risk of CD. We cannot rule out that the underlying reason for isotretinoin treatment, i.e. the presence of acne, may have contributed to the association with CD especially since the risk for CD among patients with acne was increased 34%. Acne has been linked to increased mortality[32], and severe acne is likely to result in frequent contact with dermatologists. It is possible that, on some occasions, this may either have lead to the discovery of dermatitis herpetiformis[33] or of CD-related symptoms. Both these conditions may have resulted in a small intestinal biopsy showing villous atrophy and inclusion in our cohort. Finally, Bowe et al have suggested that acne can be associated with altered microbiota[34]. It has recently been acknowledged that dysbiosis may play a role in the pathogenesis of CD[9, 10], and this might serve as a link between acne and CD. However, we cannot rule out that the association of CD and acne is due to surveillance bias as patients with acne more likely to attend a doctor more than controls and hence more likely to get a any formal diagnosis.

We used biopsy record data to ascertain CD. This has several advantages. Firstly, it allowed us to identify average patients with CD, as earlier research has shown that Swedish patients with CD identified through inpatient data[35] may have significantly more comorbidity than patients identified through pathology departments[36]. Secondly, villous atrophy has a higher positive predictive value for CD (95%)[21] than having a CD diagnosis in the Swedish Inpatient Registry (86%)[37]. Our earlier patient chart validation has also shown that symptoms in patients with villous atrophy (diarrhea (36%) and anemia (35%)[21]) are similar to those in CD[38, 39]. While we did not use the Inpatient Registry to identify CD, inpatient and outpatient data (combined they are called the “Patient Registry”) were used to identify individuals with acne. We are unaware of any study validating the sensitivity and specificity of acne in the Patient Registry but for most other disorders the positive predictive value lies in the range 85-95%.[40]

In Sweden, villous atrophy has few other causes than CD and when two examiners manually screened more than 1500 biopsy reports with villous atrophy or inflammation, the second most common cause of villous atrophy, IBD, only occurred in 0.3% of the biopsy records with villous atrophy. Finally, our use of biopsy records to detect diagnosed CD should have

a high sensitivity since >96% of all Swedish gastroenterologists and paediatricians require a biopsy before CD diagnosis.[21]

This study has several additional limitations. The Prescribed Drug Registry only began in July 2005, and we therefore chose to examine the association between isotretinoin and CD independent of temporality. For isotretinoin to play a role in the pathogenesis of CD, it would need to be instituted before diagnosis, and as the majority of patients were diagnosed before July 2005, causality cannot be inferred. Despite the large number of individuals with CD, and more than 140,000 controls, this study may have suffered from insufficient power due to the low number of exposures to isotretinoin. This is especially so in our analysis of amount of exposure (OR for CD was 3.33) where only 2 individuals with CD vs. 3 controls had received 3 prescriptions with isotretinoin prior to CD diagnosis.

Finally we did not have data on other small intestinal inflammatory disorders such as autoimmune enteropathy and we cannot rule out that autoimmune enteropathy is linked to isotretinoin.

Although not the primary aim of this paper, we also examined the relationship between acne and CD. We urge caution when interpreting those results. The diagnosis of acne is most probably underestimated in the Patient Registry, as acne diagnosed by a general practitioner will not be registered.

CONCLUSIONS

This study found no association between isotretinoin use and CD, but a small excess risk of CD in patients with a diagnosis of acne.

Acknowledgments

BL was funded by the National Center for Advancing Translational Sciences, National Institutes of Health, through Grant Number UL1 TR000040

JFL was supported by grants from The Swedish Society of Medicine, the Swedish Research Council – Medicine (522-2A09-195), and the Swedish Celiac Society.

Guarantor: JFL had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analyses.

Independence (role of the sponsors): None of the funders had any role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; and preparation, review, or approval of the manuscript.

References

1. Maki M, Mustalahti K, Kokkonen J, Kulmala P, Haapalahti M, Karttunen T, et al. Prevalence of Celiac disease among children in Finland. *N Engl J Med*. 2003 Jun 19; 348(25):2517–24. [PubMed: 12815137]
2. Rubio-Tapia A, Ludvigsson JF, Brantner TL, Murray JA, Everhart JE. The prevalence of celiac disease in the United States. *Am J Gastroenterol*. 2012 Oct; 107(10):1538–44. quiz 7, 45. [PubMed: 22850429]

3. Elfstrom P, Granath F, Ekstrom Smedby K, Montgomery SM, Askling J, Ekbom A, et al. Risk of Lymphoproliferative Malignancy in Relation to Small Intestinal Histopathology Among Patients With Celiac Disease. *J Natl Cancer Inst.* 2011 Mar 2; 103(5):436–44. [PubMed: 21289299]
4. Ludvigsson JF, Olsson T, Ekbom A, Montgomery SM. A population-based study of coeliac disease, neurodegenerative and neuroinflammatory diseases. *Aliment Pharmacol Ther.* 2007 Jun 1; 25(11): 1317–27. [PubMed: 17509100]
5. Collin P, Reunala T, Pukkala E, Laippala P, Keyrilainen O, Pasternack A. Coeliac disease--associated disorders and survival. *Gut.* 1994 Sep; 35(9):1215–8. [PubMed: 7959226]
6. Sollid LM. Molecular basis of celiac disease. *Annu Rev Immunol.* 2000; 18:53–81. [PubMed: 10837052]
7. Akobeng AK, Ramanan AV, Buchan I, Heller RF. Effect of breast feeding on risk of coeliac disease: a systematic review and meta-analysis of observational studies. *Arch Dis Child.* 2006 Jan; 91(1):39–43. [PubMed: 16287899]
8. Penders J, Thijs C, Vink C, Stelma FF, Snijders B, Kummeling I, et al. Factors influencing the composition of the intestinal microbiota in early infancy. *Pediatrics.* 2006 Aug; 118(2):511–21. [PubMed: 16882802]
9. Nadal I, Donat E, Ribes-Koninckx C, Calabuig M, Sanz Y. Imbalance in the composition of the duodenal microbiota of children with coeliac disease. *J Med Microbiol.* 2007 Dec; 56(Pt 12):1669–74. [PubMed: 18033837]
10. Ou G, Hedberg M, Horstedt P, Baranov V, Forsberg G, Drobni M, et al. Proximal small intestinal microbiota and identification of rod-shaped bacteria associated with childhood celiac disease. *Am J Gastroenterol.* 2009 Dec; 104(12):3058–67. [PubMed: 19755974]
11. Hanson LA. Breastfeeding provides passive and likely long-lasting active immunity. *Ann Allergy Asthma Immunol.* 1998; 81(6):523–33. quiz 33–4, 37. [PubMed: 9892025]
12. Mora JR, Iwata M, von Andrian UH. Vitamin effects on the immune system: vitamins A and D take centre stage. *Nat Rev Immunol.* 2008 Sep; 8(9):685–98. [PubMed: 19172691]
13. Mention JJ, Ben Ahmed M, Begue B, Barbe U, Verkarre V, Asnafi V, et al. Interleukin 15: a key to disrupted intraepithelial lymphocyte homeostasis and lymphomagenesis in celiac disease. *Gastroenterology.* 2003 Sep; 125(3):730–45. [PubMed: 12949719]
14. DePaolo RW, Abadie V, Tang F, Fehner-Peach H, Hall JA, Wang W, et al. Co-adjuvant effects of retinoic acid and IL-15 induce inflammatory immunity to dietary antigens. *Nature.* 2011 Mar 10; 471(7337):220–4. [PubMed: 21307853]
15. Crockett SD, Porter CQ, Martin CF, Sandler RS, Kappelman MD. Isotretinoin use and the risk of inflammatory bowel disease: a case-control study. *Am J Gastroenterol.* 2010 Sep; 105(9):1986–93. [PubMed: 20354506]
16. Bernstein CN, Nugent Z, Longobardi T, Blanchard JF. Isotretinoin is not associated with inflammatory bowel disease: a population-based case-control study. *Am J Gastroenterol.* 2009 Nov; 104(11):2774–8. [PubMed: 19623167]
17. Etminan M, Bird ST, Delaney JA, Bressler B, Brophy JM. Isotretinoin and risk for inflammatory bowel disease: a nested case-control study and meta-analysis of published and unpublished data. *JAMA Dermatol.* 2013 Feb; 149(2):216–20. [PubMed: 23426479]
18. Wettermark B, Hammar N, Fored CM, Leimanis A, Otterblad Olausson P, Bergman U, et al. The new Swedish Prescribed Drug Register--opportunities for pharmacoepidemiological research and experience from the first six months. *Pharmacoepidemiol Drug Saf.* 2007 Jul; 16(7):726–35. [PubMed: 16897791]
19. Marsh MN. Gluten, major histocompatibility complex, and the small intestine. A molecular and immunobiologic approach to the spectrum of gluten sensitivity 'celiac sprue'. *Gastroenterology.* 1992; 102(1):330–54. [PubMed: 1727768]
20. Ludvigsson JF, Otterblad-Olausson P, Pettersson BU, Ekbom A. The Swedish personal identity number: possibilities and pitfalls in healthcare and medical research. *Eur J Epidemiol.* 2009; 24(11):659–67. [PubMed: 19504049]
21. Ludvigsson JF, Brandt L, Montgomery SM, Granath F, Ekbom A. Validation study of villous atrophy and small intestinal inflammation in Swedish biopsy registers. *BMC Gastroenterol.* 2009 Mar 11; 9(1):19. [PubMed: 19284576]

22. Ludvigsson JF, Brandt L, Montgomery SM. Symptoms and signs in individuals with serology positive for celiac disease but normal mucosa. *BMC Gastroenterol.* 2009; 9:57. [PubMed: 19624815]
23. Pais WP, Duerksen DR, Pettigrew NM, Bernstein CN. How many duodenal biopsy specimens are required to make a diagnosis of celiac disease? *Gastrointest Endosc.* 2008 Jun; 67(7):1082–7. [PubMed: 18308317]
24. Ludvigsson JF, Montgomery SM, Ekbom A, Brandt L, Granath F. Small-intestinal histopathology and mortality risk in celiac disease. *JAMA.* 2009 Sep 16; 302(11):1171–8. [PubMed: 19755695]
25. Olen O, Bihagen E, Rasmussen F, Ludvigsson JF. Socioeconomic position and education in patients with coeliac disease. *Dig Liver Dis.* 2012 Jun; 44(6):471–6. [PubMed: 22341742]
26. Olen O, Bihagen E, Rasmussen F, Ludvigsson JF. Socioeconomic position and education in patients with coeliac disease. *Dig Liver Dis.* 2012 Feb 14.
27. Mucida D, Park Y, Kim G, Turovskaya O, Scott I, Kronenberg M, et al. Reciprocal TH17 and regulatory T cell differentiation mediated by retinoic acid. *Science.* 2007 Jul 13; 317(5835):256–60. [PubMed: 17569825]
28. Holmgren J, Czerkinsky C. Mucosal immunity and vaccines. *Nat Med.* 2005 Apr; 11(4 Suppl):S45–53. [PubMed: 15812489]
29. Stephensen CB, Livingston KA. Vitamin supplements and vaccines: maximize benefits, evaluate potential risks. *Am J Clin Nutr.* 2009 Sep; 90(3):457–8. [PubMed: 19640957]
30. Hall JA, Cannons JL, Grainger JR, Dos Santos LM, Hand TW, Naik S, et al. Essential role for retinoic acid in the promotion of CD4(+) T cell effector responses via retinoic acid receptor alpha. *Immunity.* 2011 Mar 25; 34(3):435–47. [PubMed: 21419664]
31. Reddy D, Siegel CA, Sands BE, Kane S. Possible association between isotretinoin and inflammatory bowel disease. *Am J Gastroenterol.* 2006 Jul; 101(7):1569–73. [PubMed: 16863562]
32. Sundstrom A, Alfredsson L, Sjolín-Forsberg G, Gerden B, Bergman U, Jokinen J. Association of suicide attempts with acne and treatment with isotretinoin: retrospective Swedish cohort study. *BMJ.* 2010; 341:c5812. [PubMed: 21071484]
33. Ludvigsson JF, Leffler DA, Bai JC, Biagi F, Fasano A, Green PH, et al. The Oslo definitions for coeliac disease and related terms. *Gut.* 2013 Jan; 62(1):43–52. [PubMed: 22345659]
34. Bowe WP, Patel NB, Logan AC. Acne vulgaris, probiotics and the gut-brain-skin axis: from anecdote to translational medicine. *Benef Microbes.* 2013 Jul 25.:1–15. [PubMed: 23434947]
35. Ludvigsson JF, Montgomery SM, Ekbom A. Risk of pancreatitis in 14,000 individuals with celiac disease. *Clin Gastroenterol Hepatol.* 2007 Nov; 5(11):1347–53. [PubMed: 17702659]
36. Sadr-Azodi O, Sanders DS, Murray JA, Ludvigsson JF. Patients with celiac disease have an increased risk for pancreatitis. *Clin Gastroenterol Hepatol.* 2012 Oct; 10(10):1136–42. e3. [PubMed: 22801059]
37. Smedby KE, Akerman M, Hildebrand H, Glimelius B, Ekbom A, Askling J. Malignant lymphomas in coeliac disease: evidence of increased risks for lymphoma types other than enteropathy-type T cell lymphoma. *Gut.* 2005 Jan; 54(1):54–9. [PubMed: 15591504]
38. Ciacci C, Cirillo M, Sollazzo R, Savino G, Sabbatini F, Mazzacca G. Gender and clinical presentation in adult celiac disease. *Scand J Gastroenterol.* 1995 Nov; 30(11):1077–81. [PubMed: 8578167]
39. Fasano A. Clinical presentation of celiac disease in the pediatric population. *Gastroenterology.* 2005 Apr; 128(4 Suppl 1):S68–73. [PubMed: 15825129]
40. Ludvigsson JF, Andersson E, Ekbom A, Feychting M, Kim JL, Reuterwall C, et al. External review and validation of the Swedish national inpatient register. *BMC Public Health.* 2011 Jun 9; 11(1):450. [PubMed: 21658213]

List of abbreviations

CD	Celiac disease
CI	Confidence Interval

HR Hazard Ratio

Key message

- In certain circumstances, retinoic acid seems to promote inflammatory responses to dietary antigens. Retinoic acid has been linked to inflammatory bowel disease
- Oral isotretinoin does not seem to be associated with celiac disease
- However, the main indication for oral isotretinoin, acne, may be more common in patients with celiac disease

Table 1
Characteristics of Celiac Disease Patients and Controls

Characteristics	Cases (n=26,739)	Controls (n=134,277)
Age at diagnosis / study entry (years)		
Mean/median/SD	29.3/27/24.4	29.8/27/24.6
<20	11,712 (44)	57,947 (43)
20-39	5,187 (19)	25,750 (19)
40-59	6,008 (22)	30,501 (23)
60-79	3,517 (13)	18,487(14)
80	315 (1)	1,592 (1)
Gender		
Male	9,863 (37)	49,922 (37)
Female	16,876 (63)	84,355 (63)

Table 2
Isotretinoin Use and Celiac Disease

	Cases	Controls	Odds ratio	p value
Any Isotretinoin (regardless of temporality)	93 (0.35)	378 (0.28)	1.22 (0.97-1.54)	0.08
Restrict to patients diagnosed after July 1, 2005	10 (0.34)	39 (0.27)	1.32 (0.66-2.64)	0.44
Restricted to ages 12-45	39 (0.42)	141 (0.31)	1.38 (0.97-1.97)	0.08
Any Isotretinoin, stratified by gender				
Men	41 (0.42)	162 (0.32)	1.27 (0.90-1.79)	0.18
Women	52 (0.31)	216 (0.26)	1.19 (0.88-1.62)	0.26
Adjusted for SES	--	--	1.22 (0.97-1.54)	0.08
Amount of Exposure				
None	26,646 (99.7)	133,899 (99.7)	1.0	0.38
<3 prescriptions prior to CD diagnosis	91 (0.34)	375 (0.28)	1.21 (0.96-1.52)	0.22
3 prescriptions prior to CD diagnosis	2 (0.01)	3 (0.00002)	3.33 (0.56-19.95)	