Growing Tissues in Real and Simulated Microgravity: New Methods for Tissue Engineering

Daniela Grimm, MD,¹ Markus Wehland, PhD,² Jessica Pietsch, PhD,² Ganna Aleshcheva, MSc,²
Petra Wise, PhD,³ Jack van Loon, PhD,⁴ Claudia Ulbrich, DVM,⁵ Nils E. Magnusson, PhD,⁶
Manfred Infanger, MD,² and Johann Bauer, PhD⁷

Tissue engineering in simulated (s-) and real microgravity (r- μ g) is currently a topic in Space medicine contributing to biomedical sciences and their applications on Earth. The principal aim of this review is to highlight the advances and accomplishments in the field of tissue engineering that could be achieved by culturing cells in Space or by devices created to simulate microgravity on Earth. Understanding the biology of three-dimensional (3D) multicellular structures is very important for a more complete appreciation of *in vivo* tissue function and advancing *in vitro* tissue engineering efforts. Various cells exposed to r- μ g in Space or to s- μ g created by a random positioning machine, a 2D-clinostat, or a rotating wall vessel bioreactor grew in the form of 3D tissues. Hence, these methods represent a new strategy for tissue engineering of a variety of tissues, such as regenerated cartilage, artificial vessel constructs, and other organ tissues as well as multicellular cancer spheroids. These aggregates are used to study molecular mechanisms involved in angiogenesis, cancer development, and biology and for pharmacological testing of, for example, chemotherapeutic drugs or inhibitors of neoangiogenesis. Moreover, they are useful for studying multicellular responses in toxicology and radiation biology, or for performing coculture experiments. The future will show whether these tissue-engineered constructs can be used for medical transplantations. Unveiling the mechanisms of microgravity-dependent molecular and cellular changes is an up-to-date requirement for improving Space medicine and developing new treatment strategies that can be translated to *in vivo* models while reducing the use of laboratory animals.

Introduction

A LONG-TERM SPACEFLIGHT induces a variety of health problems, such as cardiovascular and immune disorders, osteoporosis, muscle atrophy, disruption of the biological clock, and others.^{1–7} For a long time, it has been clear that microgravity induces alterations in human cells.^{8–10} However, the corresponding mechanisms are still unknown.

Spaceflight missions are very rare and costly. Therefore, researchers have developed techniques to simulate microgravity on Earth and to prepare their future Space missions.^{11–14} Several cell culture technologies simulating microgravity can be applied in tissue engineering and serve as ground-based facilities for biomedical research. It has been shown that gravitational unloading induces three-dimensional (3D) growth and assembly of cells into functional tissues.¹⁵

Weightlessness provides a very special environment for the cells because of its lack of sedimentation and convection. Using the fast-rotating 2D clinostat microscope at the German Aerospace Center in Cologne, Germany, we have seen that after 24 h follicular thyroid cancer cells of the FTC-133 cell line start to form 3D aggregates (unpublished data). A similar finding was observed by investigating endothelial cells (ECs) on the random positioning machine (RPM). After 24 h, small, 3D aggregates were detectable.¹⁶ Three-dimensional aggregates of cancer cells have become an important topic in cancer research, because they represent a very simple model of a tumor. Multicellular tumor spheroids (MCTSs) resemble the *in vivo* situation more than monolayer cells, but they are not as complex as natural tumors. They mimic small metastases and areas of solid tumors *in vivo*. Therefore, they

¹Institute of Biomedicine, Pharmacology, Aarhus University, Aarhus, Denmark.

²Clinic for Plastic, Aesthetic and Hand Surgery, Otto-von-Guericke-University Magdeburg, Magdeburg, Germany.

³Department of Hematology/Oncology, Children's Hospital Los Angeles, University of Southern California, Los Angeles, California.

⁴Department of Oral and Maxillofacial Surgery/Oral Pathology, VU University Medical Center Amsterdam & Department of Oral Cell Biology, Academic Centre for Dentistry Amsterdam (ACTA), University of Amsterdam and VU University Amsterdam, Amsterdam, The Netherlands & European Space Agency Technology Center, Gravity Lab (ESA-ESTEC-TEC-MMG), Noordwijk, The Netherlands.

⁵Department of Membrane Physiology, Institute of Physiology, University of Hohenheim, Stuttgart, Germany.

⁶Medical Research Laboratory, Institute of Clinical Medicine, Aarhus University, Aarhus, Denmark.

⁷Max-Planck Institute for Biochemistry, Martinsried, Germany.

FIG. 1. Devices suitable for tissue engineering. **(A)** Two-dimensional clinostat in an incubator constructed by the German Aerospace Center (DLR), Institute of Aerospace Medicine, Biomedical Science Support Center, Gravitational Biology, Cologne, Germany. **(B)** Example of a fast-rotating 2D clinostat manufactured by CCM (Neunen, The Netherlands). The system holds three static and three rotating tubes of about 10 mL volume. Rotation speed can be adjusted between 30 and 150 rpm. (Image: J. van Loon, DESC, Amsterdam, The Netherlands). **(C)** The desktop random-positioning machine (RPM). In this picture the automated fluid managing system COBRA is mounted on the platform together with a standard 12-well tissue culture plate (Image: Dutch Space, Leiden, The Netherlands). **(D)** Rotating wall vessel (RWV) with two 50-mL vessels in an incubator from Synthecon. **(E)** The rotary cell culture system (RCCS) from Synthecon dedicated for stem cell research with different sized culture vessels (Image from company Web site; Synthecon, Houston, TX).

appeared suitable for developing *in vitro* anticancer drug test systems and can be used for sparing of animal experiments.¹⁷

In plastic reconstructive surgery small vessels or pieces of tissue are often needed. Therefore, surgeons are looking for alternative sources of tissue. One source is engineered tissues or vessels. If we can go from single cells to 3D tissues, then we should aim as well as be able to construct a transplantable organ or part of an organ in the future. Tissues or vessels can be engineered under weightlessness at low shear stress.¹⁸ They are formed in the absence of an artificial matrix or

scaffolds.^{10–12} Hence, these types of constructs have the advantage that they are not afflicted with foreign materials.

This review will introduce available devices to simulate microgravity on Earth: the RPM, the rotating wall vessel (RWV), and the clinostat (Fig. 1). We will review the available articles that demonstrate research in the field of tissue engineering under microgravity. Moreover, we will focus on the current knowledge about the involved signaling processes for 3D growth in Space and on Earth and new methods for tissue engineering of intima constructs, cartilage, and thyroid cancer spheroids (Table 1).

TABLE 1. SUMMARY OF ARTICLES ADDRESSING TISSUE ENGINEERING ORDERED BY TISSUE

<i>Tissue of</i>	<i>Experimental goal (Ref.)</i>
Endothelial cells	3D growth, ^{12,53,55} altered gene expression, ^{20,97,105} actin polymerization, ³⁸ increase in apoptosis, ^{53,77} influence of bFGF, ¹⁶ VEGF and bFGF influence, ^{60,104} influence on caveolae and caveolae constituents, ⁷³ angiogenic response ⁷⁷
Cartilage	Characterization of human chondrocytes, ^{54,109,114,117,118,119} changes in morphology, gene expression, and protein content, ^{34,110,111} tissue engineering ^{106–108,112,113,115,116}
Thyroid	Differentiation and increase in apoptosis, ¹⁰ spheroid formation, ^{11,52} neoangiogenesis, ¹⁷ thyroid cell growth, ¹²⁰ interaction of proteins, ¹²⁸ influence of keratinocyte growth factor, ¹²⁹ gene expression profile, and altered cytokine secretion ¹³³
MCSs	Human follicular thyroid carcinoma cells, ^{10,11,17,52,80,129} prostate cell cultures ⁷⁹

MCSs, multicellular spheroids; 3D, three-dimensional; VEGF, vascular endothelial growth factor; bFGF, basic fibroblast growth factor.

Devices to Generate Simulated Microgravity

Exposing cells to real microgravity ($r\text{-}\mu\text{g}$) is either very expensive or provides time periods of microgravity that last only seconds or minutes, which are too short for tissue engineering studies, although such durations can be useful to explore various intra- and inter-cellular processes under altered gravity conditions.^{19–21} Therefore, devices that generate simulated microgravity ($s\text{-}\mu\text{g}$) conditions are most often used for tissue engineering.

Fast-rotating clinostat

Wolfgang Briegleb from the German Aerospace Center (DLR) initiated the construction of fast-rotating clinostat (FRC)^{22,23} typically consisting of a pipette-like tube filled with a cell suspension. The tubes have an internal diameter of some millimeters and rotate at a speed between 60 and 90 rpm (Fig. 1A). It is argued that the coupling between the cells and their respective static surrounding liquid boundary layers is the main reason for this microgravity simulation paradigm.^{24,25} FRCs (Fig. 1B) are widely used but they have their limitations. A balance between fluid density, viscosity, and cell-specific density is necessary. To have a high quality of microgravity simulation and a minimal residual acceleration, a proper FRC functionality with larger, that is, heavier, samples such as tissue constructs should be guaranteed. Also, a clinostat with its tube diameter of several millimeters has a too limited volume to accommodate a wide range of samples. Tubes cannot be made too wide in order to not increase the centrifugal forces driving the cells to the wall while rotating at higher speeds. This limits studies that require larger samples and volumes as needed in tissue engineering.

Random-positioning machine

A 3D clinostat²⁶ consists of two frames, each driven by a dedicated motor. In the initial designs the speed of the motors was such that each frame rotated with a constant speed. Later a similar system was developed where the speeds and the direction of rotation of each frame were different and randomized, the so-called RPM^{26–29} (Fig. 1C).

In an RPM one could principally work with larger liquid volumes. Cells and larger multicellular constructs are not exposed to a constant, solid body fluid rotation but rather mainly freely and randomly floating within the liquid phase. An initial study demonstrated that such a movement does generate small fluid shear forces at the container wall in the order of 6 mPa (0.06 dyne/cm²).³⁰ Cells in an RPM are not only oriented randomly with respect to the gravity vector, but are also more likely to interact with other cells within the experimental volume. Hence, there is an increased opportunity to form multicellular structures. The slight fluid flow around cells and constructs also facilitates an increased provision of nutrients and gasses while waste products are more efficiently removed, compared to static batch cell culture conditions where nutrients are delivered by limited convective flows and the very inefficient process of diffusion.²⁵ The ample nutrient and gas supply to the samples also permits to work with higher cell densities, again increasing the probabilities of cell–cell interactions. Due to the construction of an RPM a broad variety of sample enclosures are

applied in RPM studies. This ranges from regular T25 flasks^{31,32} to multiwell plates,³³ flask on slides,³⁴ or more dedicated devices.³⁵ Performing RPM studies, where the best simulation of microgravity is in the center of rotation of the two axes, has limits for the size of the preferred volume for the samples. Depending on the speed of rotation and the distance from center, an acceptable residual gravity can be obtained in the order of $10^{-4} g$ by a maximum speed of 60°/s at a radial distance of 10 cm.²⁵ This allows for a maximum working volume of more than 4 L. Adding drugs or fixing samples during rotation is cumbersome, if not impossible, in an RPM. However, recently tools have been developed to enable fluid managing while rotating.²⁹ RPMs are commercially available by Mitsubishi Heavy Industries (Kobe, Japan) and Dutch Space (Leiden, The Netherlands), while various academic groups developed similar systems of 3D clinostats and RPMs.^{35–40}

Rotating wall vessel

The RWV (Fig. 1D, E) developed by NASA⁴¹ is now commercially available through Synthecon, Inc. (Houston, TX) and is a slow-rotating, relatively large liquid-filled container. It has some overlaps with a 2D clinostat in such a way that cells within the container are prevented from settling by a constant rotation. The rotation speed should be matched to the terminal velocity of the cells, nodules, or other samples within the fluid. The proper rotation speed is mainly determined by the specific weight of the cells, the fluid density, and viscosity, although tissue surface properties and shape also play a role. The cells and tissues in the RWV are constantly falling within the fluid. The settling velocity and direction combined with the rotation of the fluid generated spiral trajectories within the vessel.⁴² This motion of the samples relative to the fluid generates fluid shear forces on a particle surface ranging from 180–320 mPa (0.18–0.32 dyne/cm²) for 50- μm beads⁴³ and ~ 500 mPa (0.5 dyne/cm²) with 3D aggregates of BHK-21 cells⁴⁴ to 520–780 mPa (5.2–7.8 dynes/cm²) for a 200- or 300- μm spherical object.⁴⁵ Over the years various models based on the initial RWV have been developed, differing in vessel geometry and aspect ratio and gas supply (perfusion), such as the 250-mL, slow-turning lateral vessel (Synthecon)⁴⁶ or the 50-mL high aspect ratio vessel (HARV; Synthecon)⁴⁷ or the rotating-wall perfused vessel (Synthecon)⁴⁸ (Fig. 1D, E).

Known Mechanisms Involved in Spheroid Formation

In 3D aggregation processes, it does not play a role whether gravity forces are absent during a spaceflight or whether the gravity vector is randomized only by the RPM or the FRC.^{2,11,49–51}

Only a part of the cells of a monolayer change their growth behavior, when exposed to microgravity, while the other part continues to grow adherently.^{52–54} However, when exposed to microgravity, both the adherently and the 3D growing cells alter their gene expression patterns.⁵⁵ Very often the decrease or increase of the protein content found in a cell population under microgravity is accompanied by a change in the number of cells bearing the respective antigen.^{10,56} It is interesting to explore whether cellular heterogeneities found in cell lines before exposition to microgravity^{57,58} are further expanded under microgravity. In addition, the signals

delivered by removal or randomization of gravity forces cause the formation of different types of aggregates depending on the kind of cells cultured. While cancer cells mainly assemble to spheres,⁵² for example, ECs form tubes⁸ and chondrocytes form pieces of cartilage,⁵⁴ *in vitro* like they naturally do *in vivo*. Although all these cells sense the lack of gravity within seconds,^{20,59} it takes at least 12 h until 3D spheroids are observed^{52,53} and up to 7 days until constructs like tubes resembling an intima can be recognized floating in a culture flask.^{12,60} These observations are not understood yet by far. Recent research, however, has unveiled a few pieces of the mosaic.

At first, the interaction between extracellular matrix (ECM) and cytoskeletal proteins is affected, when gravity forces are randomized or removed. Most cells studied so far adjust to exposure to microgravity by altering the expression patterns of ECM and cytoskeletal proteins at gene and protein levels.^{60–64} Preliminary indications of specific proteins involved in this process came from a comparison of the spheroid formation of the two thyroid cancer cell lines FTC-133 and CGTH-W1. The study revealed that speed of formation and size of resulting spheroids is related to the expression of certain proteins that bind fibronectin.⁶ In this context, integrin- $\alpha 5$ was repeatedly determined as important for spheroid formation of human thyroid cells as well as of porcine hepatocytes.^{52,65} Based on ECM–cytoskeleton interaction studies, Ingber postulated that gravity sensation may be due to a change of the forces, which the cells need for their interaction with the environment under the different conditions.⁶⁶ This hypothesis is in conformance with recent results that reveal that cells can receive information through integrin-mediated cell–matrix adhesions, which connect the cytoskeleton with the ECM.⁶⁷ In this system vinculin plays a

decisive role. It is activated when a mechanical stimulus is translated into a biochemical cell response.⁶⁸ We found a clear upregulation of vinculin protein in the ML-1 thyroid cancer cells during spheroid formation¹⁰ and Gershovich *et al.* observed an intracellular redistribution of vinculin in mesenchymal stem cells cultured on the RPM.⁶⁹ How signals, raised by cell–ECM interaction, are transduced to the cell interior remains to be determined. They may open Ca^{2+} channels allowing an increase of intracellular Ca^{2+} as described by Luo *et al.* for osteoblasts cultured under s- μg .⁷⁰ They may be directly transduced to distinct growth factors via the change of the cytoskeleton architecture^{71–73} or trigger the phosphorylation of caveolae.⁷³

Each system (RPM and RWV) that generates tissue-like cell assemblies under microgravity conditions, initially induces an enhanced rate of apoptosis together with an increase in proapoptotic factors.^{10,53,60,74–76} In EC, apoptosis seems to impair the angiogenic response to s- μg produced on the RPM.⁷⁷ NF- κB 1 plays a central role in bone reduction under microgravity conditions.⁷⁸ Clejan *et al.* showed that the activation of apoptotic pathways during an early phase of culturing DU-145 human prostate carcinoma cells in a HARV is essential for the subsequent formation of 3D organoids.⁷⁹ Our experiments on the RPM on FTC-133 thyroid cancer cells suggested that initiation of an early phase of apoptosis, followed by an escape from the late phase, could be a step in cells that transit from a 2D to a 3D kind of growth.⁸⁰ In this process the signaling elements IL-6, IL-8, OPN, TLN1, and CTGF are involved with NF- κB p65. IL-6 and IL-8 together with the von Willebrand factor also play a role in tube formation. After ECs have been cultured on the RPM, precursors of the tubes with single-layered walls could be observed (Fig. 2), in which IL-8 gene expression was

FIG. 2. Tissue engineering of intima constructs. (A) Microvascular endothelial cells (MVECs) grown under normal static 1 g conditions. (B) MVECs grown for 24 h on the RPM. (C) MVECs grown for 7 days on the RPM. A tubular construct is visible. (D) Coculture of EA.hy926 cells, vascular smooth muscle cells, and fibroblasts for 21 days on the RPM (phase-contrast microscopy). (E, F) Sirius Red staining of cocultures of EA.hy926 cells, vascular smooth muscle cells, and fibroblasts, 21 days on the RPM.

TABLE 2. PROTEINS OR GENES SHOWING ALTERATIONS DURING MICROGRAVITY-INDUCED TISSUE FORMATION

<i>Cells</i>	<i>Tissue type</i>	<i>Gene/protein</i>	<i>Ref.</i>
ML-1 ^a	Spheroids	FAS, BCL-2, vinculin	10
FTC-133 ^a	Spheroids	Integrin- α 5, myosin-10, filamin B	52
Porcine hepatocytes	Liver morphology	Integrin- α 5	65
FTC-133 ^a	Spheroids	NF- κ B1, interleukin-6	78
FTC-133 ^a	Spheroids	<i>Interleukin-6, interleukin-8, osteopontin</i>	80
FTC-133 ^a	Spheroids	<i>EGF, CTGF</i>	11
EA.hy926 ^b	Vessels	<i>Interleukin-8, von Willebrand factor</i>	55
EA.hy926 ^b	Vessels	von Willebrand factor, bFGF	60
EA.hy926 ^b	Vessels	Matrix metalloproteinase 2	12
HUVEC	—	Caveolin-1	73
Chondrocytes ^c	Cartilage	Collagen type I and X	54

^aHuman thyroid cancer cells.^bHuman endothelial cells.^cHuman chondrocytes.

HUVEC, human umbilical vein endothelial cell.

fivefold upregulated and *VWF* gene expression was sixfold downregulated.⁵⁵ IL-6, IL-8, and NF- κ B p65 are differently regulated by basic fibroblast growth factor depending on whether this growth factor is applied at 1 g or s- μ g.¹⁶ A brief overview of these observations is given in Table 2. In addition to growth factors, it should also be considered that miRNA expression is influenced by s- μ g on the RPM.⁸¹ It is known that several miRNAs have an influence on tube formation *in vitro*.^{82,83}

Endothelial Cells

Blood vessel replacements are needed for bypass or reconstructive surgery. Under standard culture conditions, however, ECs grow in a monolayer (Fig. 2A, B) and do not fully exhibit the characteristics they have when they are part of an actual vessel. To obtain a vessel-like structure, endothelial and other cells need to be seeded on preformed scaffolds consisting of biodegradable polymers. While large-caliber constructs generated with this approach usually work well, diameters below 6 mm remain critical due to their increased thrombogenicity.^{84–86}

Cultivation of ECs under s- μ g might provide a solution for these problems, as ECs are highly sensitive to diverse external stimuli, triggering their activation and modulating their function,^{87–91} and strongly react to changes of the influence of gravity. Exposure to r- μ g during parabolic flights (22 s) induced dramatic changes in the microtubule network of ECs in the form of tubulin downregulation and rearrangement and altered the expression of 3605 genes.^{20,92} Further, gravitational unloading helps the cells to form 3D structures without the need for scaffolds. On the clinostat, the formation of 3D cell aggregates has not been published so far. However, after 24 h of clinorotation, tube formation and migration as well as endothelial nitric oxide synthase (eNOS) activity of human umbilical vein endothelial cells were significantly increased, suggesting an s- μ g-mediated promotion of angiogenesis via the PI3K-Akt-eNOS pathway.⁹³ In addition, Wang *et al.* showed that clinorotation also upregulated inducible NOS (iNOS) in an AP-1-dependent manner and suggested a possible involvement of this pathway in the development of postflight orthostatic intolerance.⁹⁴ In parallel to observations

in r- μ g, the cytoskeleton of ECs grown on the clinostat was remodeled. The expression of various surface adhesion factors was changed, but despite these effects, no critical physiological processes were compromised.⁹⁵ Zhang *et al.* reported earlier that mesenchymal stem cells showed an increased endothelial differentiation potential when cultured on a clinostat for 72 h.⁹⁶

Further, clinorotation also has some detrimental consequences for ECs. Kang *et al.* observed that prolonged growth on a clinostat induced among others apoptosis in microvascular ECs.⁹⁷

Moreover, several studies performed on the RWV reported cytoskeletal remodeling, actin downregulation, increased NO release, altered cytokine excretion, and increased caveolin-1 and -2 expression concordantly.^{38,72,98–102} Interestingly, reports about the detrimental effects of cultivation on the RWV showed that they depend on the origin of the investigated cells. Mariotti and Maier reported that only microvascular cells suffer from inhibited growth and angiogenesis, while macrovascular cells proliferated faster when grown on an RWV.¹⁰³

Exposing the cells to altered gravity conditions on an RPM led an increased apoptosis rate,^{17,53} which was significantly reduced by the addition of vascular endothelial growth factor to the culture medium.¹⁰⁴ Similar effects have also been reported in further studies.¹⁰⁵ More importantly, Ea.hy926 cells readily formed 3D multicellular spheroids similar to those observed on RWVs within the first 24 h of incubation on the RPM.^{53,104} Between the 5th and the 7th day of growth on the RPM, double-row cell assemblies were forming. In the course of the next weeks, these assemblies developed into tube-like structures with walls mostly consisting of single-cell layers. They were found to have a lumen and resembled the intima of blood vessels *in vivo*, reaching a length of up to several centimeters.^{12,60} These structures are unlike any other constructs generated by s- μ g tissue engineering either on clinostats or RWVs and are presently the most promising candidates on the way to the engineering of a complete, functional vessel (Fig. 2C–F). First investigations in both short- and long-term s- and r- μ g have shown that this process involves a complex interplay of a multitude of molecules belonging to the cytoskeleton,

FIG. 3. Cartilage formation under simulated microgravity. (A) Control sample after 14 days of cultivation under normal gravity conditions. (B) Three-dimensional aggregates of human chondrocytes after 14 days of cultivation on the RPM. (C) Three-dimensional aggregate grown for 21 days in an RWV and seeded out on a slide flask. Cells are growing out from the spheroid.

ECM, apoptotic processes, cytokines, angiogenesis, and cell cycle with a total number of over 1600 genes identified as differentially regulated on the RPM.^{20,55,92,105}

Cartilage

Interest in tissue engineering of cartilage is motivated by the inability of adult articular cartilage to repair itself after damage by disease or injury.^{106,107} Chondrocytes are the most obvious cell source for the tissue engineering of cartilage.¹⁰⁸ However, chondrocyte expansion in a monolayer causes cell dedifferentiation, characterized by decreased proteoglycan synthesis and type II collagen expression, and increased type I collagen expression.^{109–111}

Cartilage development is achieved *in vitro* by cultivating living cells on biodegradable polymer scaffolds.¹¹² The scaffold should (1) be biodegradable in a controlled way without cytotoxic, tumorigenic, nephrotoxic, or other undesirable effects; (2) have a porosity that allows diffusion of nutrients and waste products; (3) support cell viability, proliferation, differentiation, and ECM production; (4) be able to fix to and integrate with the tissue at the defect site; and (5) give mechanical support to the engineered tissue.^{104,106} Ideally, as the ECM develops, the scaffold dissolves and disappears.¹⁰⁷

A comparison of chondrocyte cells cultured in rotating bioreactors in Space (Mir Space Station) and on Earth was reported by Freed *et al.*¹¹³ Freed and his team performed experiments with rotating cultures of bovine chondrocytes in polyglycolic acid scaffolds and concluded that after 7 months of cultivating the culture on Earth (static 1 g control) it produced cartilage tissues closer to the natural form than that in Space. Very recently, the scaffold-free chondrogenesis of human cartilage in an RWV bioreactor has been reported.¹¹⁴ Good quality cartilage tissue was formed by a rotating culture from aged human articular cartilage after 90 days of cultivation. The report of Ohyabu *et al.*¹¹⁵ was the first on the production of cartilaginous tissue from bone-marrow-derived cells by a rotating culture. A comparison of porcine chondrocyte cells cultured on the International Space Station (ISS) and on the RPM was firstly done by Stamenković *et al.*¹¹⁶ Neocartilage formed in normal gravity was “continuous in shape and form that corresponded to the shape of the cylindrical bioreactor,”¹¹⁶ compared to ISS and RPM tissues, which consisted of irregular cell aggregates.¹¹⁶

RPM-cultivated chondrocytes created tissue with matrix composition characteristics in between the control 1 g and the ISS-produced neocartilages, but produced neocartilage with more evenly dispersed and further spaced apart cells.⁵⁴ The number of cells that can be obtained from a clinically relevant-sized biopsy is not sufficient for cell-based therapies such as autologous cartilage implantation or to produce a functional neocartilage implant. Thus, the classic expansion phase in monolayer is needed to increase cell numbers, adding to the probability of reduced chondrogenic capacities.¹¹⁶ This point is especially a factor in scaffold-free methods that rely on the volume of cells to act as their own scaffold support and tissue volume.¹¹⁴ The RPM therefore offers both opportunities: a reduction in the amount of cells and the prefabrication of implants without the complications associated with scaffolds (Fig. 3). Thus, the RPM has proved to be suitable for tissue engineering of cartilage.³⁴

Thyroid

The first observations of altered thyroid function, that is, thyroid hormones, in astronauts were already made in the late 1970s¹¹⁷ and could be confirmed in later studies on humans as well as mammals.^{117–119} Since the beginning of the new millennium, three Space missions were carried out that either analyzed thyroid rat cells¹²⁰ or thyroids of mice¹²¹ flown in Space. Based on the ratio of sphingomyelinase and sphingomyelin synthase, two enzymes responsible for the sphingomyelin metabolism, Albi *et al.* were able to determine the state of rat thyroid cells (FRTL-5) returning from r-μg as proapoptotic.¹²⁰ Similar changes of the sphingomyelin metabolism could be detected in a single mouse returning after 91 days to Earth. Further, it could be shown that beside the activity of the proteins their location in the thyroid cells was altered.¹²¹ Overall, obvious differences in the thyroid tissue structure could be detected.^{122,123}

Investigating the effects generated by an RPM, changes in the cytoskeleton, the ECM, and in apoptosis of human thyroid cells were observed.^{10,124,125} By proteome analyses, normal and cancerous thyroid-cell-dependent changes in the content of regulatory, metabolic, and structural proteins could be detected.^{52,126,127} The proteome analyses were carried out with samples of human thyroid cells cultured on an RPM for several days, during which they formed small spherical aggregates (spheroids, Fig. 4). Attempts to identify

FIG. 4. Formation of multicellular tumor spheroids. (A) Static control at start of the experiment with the poorly differentiated follicular thyroid carcinoma cell line CGTH-W1. (B) Static control at day 4 at the end of the experiment, CGTH-W1 cell line. (C) RPM experiment, 4 days, adherent cells were visible as well as multicellular tumor spheroids (CGTH-W1 cell line) floating in the supernatant.

proteins that may be involved in spheroid formation hinted to various isoforms of the adhesion molecule integrin.^{52,128} A change of the molecule, however, could not be detected on the RNA level in human thyroid cells cultivated for 14 days in an RWV and thus was deemed not responsible for spheroid formation.¹²⁹ It may be that the integrin protein is essential in the early stages of spheroid formation but later on other proteins are responsible for sustaining the spheroids. Parabolic flights revealed that the cytoskeleton of thyroid carcinoma cells begins to change already after 22 s of exposure to $r\text{-}\mu\text{g}$ for 22 s.⁵⁹

Summary

The preferred environment for tissue engineering studies would be a long-term flight in the microgravity environment, but devices, such as clinostats, the RPM, or RVW, are very useful tools in this area.

Recent research revealed that the formation of 3D aggregates under $s\text{-}\mu\text{g}$ can be divided into three phases of regulation that cells pass through when they transit from a 2D to a 3D kind of growth under gravitational unloading: (1) cells sense a change of the forces required for interaction with the ECM, (2) they face an initiation of apoptosis, and (3) they transduce the signals generated during phases 1 and 2 to the nucleus via kinase-dependent pathways.

Growing cells under $r\text{-}$ or $s\text{-}\mu\text{g}$, particularly on the RPM, became a very promising approach to generate tissue constructs in a scaffold-free manner. Using human ECs to form intima constructs might be a first step to the generation of fully functional vessels usable in both *in vitro* research and (reconstructive) surgery.

In addition, thyroid cells, which can also be stimulated by $s\text{-}\mu\text{g}$ to grow in form of 3D aggregates, can in future be grown up to a state to resemble thyroid tissue. Currently, studies that analyze the molecular mechanisms behind spheroid formation have increased the knowledge of the complex regulation of 3D growth in microgravity.^{52,55,80,130–132}

Further, cartilage may be formed on the RPM and RVW. The advantage of this procedure is that cell aggregation and cell proliferation are simultaneously going on so that the number of cells required at the beginning can be kept low even if an enlarged piece of cartilage is desired. However, the biomechanical stability of 3D chondrocyte aggregates produced under conditions of $s\text{-}\mu\text{g}$ has still to be tested.

Taken together, the use of microgravity to study human cells is very important to understand the exact cellular changes occurring under these special conditions. Hence,

Space medicine has an important contribution to the progress in basic research. The tissues, engineered from single cell layers cultured in $s\text{-}\mu\text{g}$, might be used in the future for medical transplantations. Tissue-engineered cartilage may be important for regenerative medicine. It is also important that with the help of these 3D structures, which are similar to the *in vivo* situation, studies on the molecular mechanisms of processes such as angiogenesis or tumor cell apoptosis are possible without animal testing.

Experiments conducted in Space or in $s\text{-}\mu\text{g}$ environment are not typically the studies a molecular or cellular biologist is able to perform. We have learnt from Space research that the compensation of gravity impacts on the cells enormously and dramatically changes the genome, proteome, and secretome of these cells. However, a spaceflight and a stay on the ISS provides physical conditions that are not achievable on Earth, but can be exploited to study mechanisms and pathways that control cell growth and function. Over the last 40 years, many experiments have shown that exposure to microgravity alters biological processes that may be relevant to the growth behavior and function of human cells. Several spaceflights have demonstrated 3D growth in Space and similar results were detected with the help of devices simulating microgravity in our laboratories.^{11,113,116,133} Therefore, this promising research area can help to develop new strategies for tissue engineering.

Acknowledgments

This article was supported by the German Aerospace Center DLR (D.G.) (BMW grants 50WB0824 and 50WB1124), the European Space Agency (ESA; CORA-GBF-2013-001; CORA-GBF-2013-004), and Aarhus University, Denmark (D.G.). The authors would like to thank Mr. Peter Lindborg, Los Angeles, CA, for reviewing the article for language and grammar errors.

Disclosure Statement

No competing financial interests exist.

References

1. White, R.J., and Averner, M. Humans in space. *Nature* **409**, 1115, 2001.
2. Pietsch, J., Bauer, J., Egli, M., Infanger, M., Wise, P., Ulbrich, C., and Grimm, D. The effects of weightlessness on the human organism and mammalian cells. *Curr Mol Med* **11**, 350, 2011.

3. Grimm, D., Wise, P., Lebert, M., Richter, P., and Baatout, S. How and why does the proteome respond to microgravity? *Expert Rev Proteomics* **8**, 13, 2011.
4. Fritsch-Yelle, J.M., Leuenberger, U.A., D'Aunno, D.S., Rossum, A.C., Brown, T.E., Wood, M.L., Josephson, M.E., and Goldberger, A.L. An episode of ventricular tachycardia during long-duration spaceflight. *Am J Cardiol* **81**, 1391, 1998.
5. Perhonen, M.A., Franco, F., Lane, L.D., Buckey, J.C., Blomqvist, C.G., Zerwekh, J.E., Peshock, R.M., Weatherall, P.T., and Levine, B.D. Cardiac atrophy after bed rest and spaceflight. *J Appl Physiol* **91**, 645, 2001.
6. Pavy-Le-Traon, A., Vasseur, P., Maillet, A., Güell, A., Bes, A., and Gharib, C. Orthostatic hypotension after space flight. *Presse Med* **23**, 1031, 1994.
7. Meck, J.V., Reyes, C.J., Perez, S.A., Goldberger, A.L., and Ziegler, M.G. Marked exacerbation of orthostatic intolerance after long- vs. short-duration spaceflight in veteran astronauts. *Psychosom Med* **63**, 865, 2001.
8. Lewis, M.L., Reynolds, J.L., Cubano, L.A., Hatton, J.P., Lawless, B.D., and Piepmeier, E.H. Spaceflight alters microtubules and increases apoptosis in human lymphocytes (Jurkat). *FASEB J* **12**, 1007, 1998.
9. Cogoli, A., Tschopp, A., and Fuchs-Bislin, P. Cell sensitivity to gravity. *Science* **225**, 228, 1984.
10. Grimm, D., Bauer, J., Kossmehl, P., Shakibaei, M., Schonberger, J., Pickenhahn, H., Schulze-Tansil, G., Vetter, R., Eilles, C., Paul, M., and Cogoli, A. Simulated microgravity alters differentiation and increases apoptosis in human follicular thyroid carcinoma cells. *FASEB J* **16**, 604, 2002.
11. Pietsch, J., Ma, X., Wehland, M., Aleshcheva, G., Schwarzwälder, A., Segerer, J., Birlen, M., Horn, A., Bauer, J., Infanger, M., and Grimm, D. Spheroid formation of human thyroid cancer cells in an automated culturing system during the Shenzhou-8 Space mission. *Biomaterials* **34**, 7694, 2013.
12. Grimm, D., Infanger, M., Westphal, K., Ulbrich, C., Pietsch, J., Kossmehl, P., Vadrucchi, S., Baatout, S., Flick, B., Paul, M., and Bauer, J. A delayed type of three-dimensional growth of human endothelial cells under simulated weightlessness. *Tissue Eng Part A* **15**, 2267, 2009.
13. Conza, N., Mainil-Varlet, P., Rieser, F., Kraemer, J., Bittmann, P., Huijser, R., van den Bergh, L., and Cogoli, A. Tissue engineering in space. *J Gravit Physiol* **8**, 17, 2001.
14. Herranz, R., Anken, R., Boonstra, J., Braun, M., Christianen, P., de Geest, M., Hauslage, J., Hilbig, R., Hill, R., Lebert, M., Medina, F., Vagt, N., Ullrich, O., van Loon, J., and Hemmersbach, R. Ground-based facilities for simulation of microgravity: organism-specific recommendations for their use, and recommended terminology. *Astrobiology* **13**, 1, 2013.
15. Unsworth, B.R., and Lelkes, P.I. Growing tissues in microgravity. *Nat Med* **4**, 901, 1998.
16. Ulbrich, C., Westphal, K., Baatout, S., Wehland, M., Bauer, J., Flick, B., Infanger, M., Kreutz, R., Vadrucchi, S., Egli, M., Cogoli, A., Derradji, H., Pietsch, J., Paul, M., and Grimm, D. Effects of basic fibroblast growth factor on endothelial cells under conditions of simulated microgravity. *J Cell Biochem* **104**, 1324, 2008.
17. Grimm, D., Bauer, J., and Schoenberger, J. Blockade of neoangiogenesis, a new and promising technique to control the growth of malignant tumors and their metastases. *Curr Vasc Pharmacol* **7**, 347, 2009.
18. Nickerson, C.A., Ott, C.M., Wilson, J.W., Ramamurthy, R., and Pierson, D.L. Microbial responses to microgravity and other low-shear environments. *Microbiol Mol Biol Rev* **68**, 345, 2004.
19. Albi, E., Ambesi-Impimbato, F.S., Peverini, M., Damaskopoulou, E., Fontanini, E., Lazzarini, R., Curcio, F., and Perrella, G. Thyrotropin receptor and membrane interactions in FRTL-5 thyroid cell strain in microgravity. *Astrobiology* **11**, 57, 2011.
20. Grosse, J., Wehland, M., Pietsch, J., Ma, X., Ulbrich, C., Schulz, H., Saar, K., Hübner, N., Hauslage, J., Hemmersbach, R., Braun, M., van Loon, J., Vagt, N., Infanger, M., Eilles, C., Egli, M., Richter, P., Baltz, T., Einspanier, R., Sharbati, S., and Grimm, D. Short-term weightlessness produced by parabolic flight maneuvers altered gene expression patterns in human endothelial cells. *FASEB J* **26**, 639, 2012.
21. Hemmersbach, R., Krause, M., Bräucker, R., and Ivanova, K. Gravitopercption in ciliates: steps in the transduction chain. *Adv Space Res* **35**, 296, 2005.
22. Briegleb, W. Some qualitative and quantitative aspects of the fast-rotating clinostats as a research tool. *ASGSB Bull* **5**, 23, 1992.
23. Briegleb, W. Ein modell fuer schwerlosigkeits-simulation an mikroorganismen. *Naturwissenschaften* **54**, 67, 1967.
24. Klaus, D. Clinostats and bioreactors. *ASGSB Bull* **14**, 55, 2001.
25. van Loon, J.J.W.A. Biology in space and life on earth. In: Brinckmann, E., ed. *Effects of Spaceflight on Biological Systems*. Weinheim: Wiley-VCH, 2007, p. 17.
26. Murakami, S., and Yamada, M. Architecture of statocytes and chloroplasts under the microgravity environment. *Jpn Soc Biol Sci Space* **2**, 301, 1988.
27. Hoson, T., Kamisaka, S., Masuda, Y., and Yamashita, M. Changes in plant growth process under microgravity conditions simulated by a three-dimensional clinostat. *Bot Mag* **105**, 53, 1992.
28. Hoson, T., Kamisaka, S., Masuda, Y., Yamashita, M., and Buchen, B. Evaluation of the three dimensional clinostat as a simulator of weightlessness. *Planta* **203**, 197, 1997.
29. Borst, A.G., and van Loon, J.J.W.A. Technology and developments for the random positioning machine, RPM. *Microgravity Sci Technol* **21**, 287, 2009.
30. Leguy, C.A.D., Delfos, R., Pourquie, M.J.B.M., Poelma, C., Krooneman, J., Westerweel, J., and van Loon, J.J.W.A. Fluid motion for microgravity simulations in a random positioning machine. *Gravit Space Biol* **25**, 36, 2011.
31. Wakayama, S., Kawahara, Y., Li, C., Yamagata, K., Yuge, L., and Wakayama, T. Detrimental effects of microgravity on mouse pre-implantation development *in vitro*. *PLoS One* **4**, e6753, 2009.
32. Luo, M.Z., Meng, R., Li, S.S., Di, S.M., Han, J., Zhang, W., and Shang, P. Weightlessness simulated with random positioning machine influences the cytoskeleton and migration of MC3T3-E1 cells. *J Jpn Soc Microgravity Appl* **28**, S41, 2011.
33. Loesberg, W.A., Walboomers, X.F., Bronkhorst, E.M., van Loon, J.J., and Jansen, J.A. The effect of combined simulated microgravity and microgrooved surface topography on fibroblasts. *Cell Motil Cytoskeleton* **64**, 174, 2007.
34. Aleshcheva, G., Sahana, J., Ma, X., Hauslage, J., Hemmersbach, R., Egli, M., Infanger, M., Bauer, J., and Grimm, D. Changes in morphology, gene expression and

- protein content in chondrocytes cultured on a random positioning machine. *PLoS One* **8**, e79057, 2013.
35. Prodanov, L., van Loon, J.J.W.A., te Riet, J., Jansen, J.A., and Walboomers, X.F. Nanostructured substrate conformation can decrease osteoblast-like cell dysfunction in simulated microgravity conditions. *J Tissue Eng Regen Med* 2012.
 36. Ichigi, J., and Asashima, M. Dome formation and tubule morphogenesis by *Xenopus* kidney A6 cell cultures exposed to microgravity simulated with a 3D-clinostat and to hypergravity. *In Vitro Cell Dev Biol Anim* **37**, 31, 2001.
 37. Jiang, Y.D., Li, W.N., Wang, L.F., Zhang, Z.Y., Zhang, B.M., and Wu, H.J. Several new types of clinostats. *Space Med Med Eng* **21**, 368, 2008.
 38. Siamwala, J.H., Reddy, S.H., Majumder, S., Kolluru, G.K., Muley, A., Sinha, S., and Chatterjee, S. Simulated microgravity perturbs actin polymerization to promote nitric oxide-associated migration in human immortalized Eahy926 cells. *Protoplasma* **242**, 3, 2010.
 39. Xiang, L., Qim, F., Daim, D., Li, C., and Jiang, Y. Simulated microgravity affects growth of *Escherichia coli* and recombinant beta-D-glucuronidase production. *Appl Biochem Biotech* **162**, 654, 2010.
 40. Hemmersbach, R., Strauch, S.M., Seibt, D., and Schuber, M. Comparative studies on gravisensitive protists on ground (2D and 3D clinostats) and in microgravity. *Microgravity Sci Technol* **18**, 257, 2006.
 41. Goodwin, T.J., Jessup, J.M., and Wolf, D.A. Morphological differentiation of colon carcinoma cell lines HT-29 and HT-29KM in rotating-wall vessels. *In Vitro Cell Dev Biol* **28A**, 47, 1992.
 42. Hammond, T.G., and Hammond, J.M. Optimized suspension culture: the rotating-wall vessel. *Am J Physiol Renal Physiol* **281**, F12, 2001.
 43. Liu, T., Li, X., Sun, X., Ma, X., and Cui, Z. Analysis on forces and movement of cultivated particles in a rotating wall vessel bioreactor. *Biochem Eng J* **18**, 97, 2004.
 44. Goodwin, T.J., Prewett, T.L., Wolf, D.A., and Spaulding, G.F. Reduced shear stress: a major component in the ability of mammalian tissues to form three-dimensional assemblies in simulated microgravity. *J Cell Biochem* **51**, 301, 1993.
 45. Nauman, E.A., Ott, C.M., Sander, E., Tucker, D.L., Pierson, D., Wilson, J.W., and Nickerson, C.A. Novel quantitative biosystem for modeling physiological fluid shear stress on cells. *Appl Environ Microbiol* **73**, 699, 2007.
 46. Guo, X.M., Zhao, Y.S., Chang, H.X., Wang, C.Y., Zhang, X.A., Duan, C.M., Dong, L.Z., Jiang, H., Li, J., Song, Y., and Yang, X.J. Creation of engineered cardiac tissue *in vitro* from mouse embryonic stem cells. *Circulation* **113**, 2229, 2006.
 47. Tucker, D.L., Ott, C.M., Huff, S., Fofanov, Y., Pierson, D.L., Willson, R.C., and Fox, G.E. Characterization of *Escherichia coli* MG1655 grown in a low-shear modeled microgravity environment. *BMC Microbiol* **7**, 15, 2007.
 48. Cinbiz, M.N., Tigli, R.S., Beşkardeş, I.G., Gümüşderelioğlu, M., and Colak, U. Computational fluid dynamics modeling of momentum transport in rotating wall perfused bioreactor for cartilage tissue engineering. *J Biotech* **150**, 389, 2010.
 49. Jin, F., Zhang, Y.J., Xuan, K., He, D.N., Deng, T.Z., Tang, L., Lu, W., and Duan, Y.Z. Establishment of three-dimensional tissue-engineered bone constructs under microgravity-simulated conditions. *Artif Organs* **34**, 118, 2010.
 50. Hou, Y., Song, C., Xie, W.J., Wei, Z., Huang, R.P., Liu, W., Zhang, Z.L., and Shi, Y.B. Excellent effect of three-dimensional culture condition on pancreatic islets. *Diabetes Res Clin Pract* **86**, 11, 2009.
 51. Schrader, S., Kremling, C., Klinger, M., Laqua, H., and Geerling, G. Cultivation of lacrimal gland acinar cells in a microgravity environment. *Br J Ophthalmol* **93**, 1121, 2009.
 52. Pietsch, J., Sickmann, A., Weber, G., Bauer, J., Egli, M., Wildgruber, R., Infanger, M., and Grimm, D. A proteomic approach to analysing spheroid formation of two human thyroid cell lines cultured on a random positioning machine. *Proteomics* **11**, 2095, 2011.
 53. Infanger, M., Kossmehl, P., Shakibaei, M., Baatout, S., Witzing, A., Grosse, J., Bauer, J., Cogoli, A., Faramarzi, S., Derradji, H., Neefs, M., Paul, M., and Grimm, D. Induction of three-dimensional assembly and increase in apoptosis of human endothelial cells by simulated microgravity: impact of vascular endothelial growth factor. *Apoptosis* **11**, 749, 2006.
 54. Ulbrich, C., Westphal, K., Pietsch, J., Winkler, H.D.F., Leder, A., Bauer, J., Kossmehl, P., Grosse, J., Schoenberger, J., Infanger, M., Egli, M., and Grimm, D. Characterization of human chondrocytes exposed to simulated microgravity. *Cell Physiol Biochem* **25**, 551, 2010.
 55. Ma, X., Wehland, M., Schulz, H., Saar, K., Hübner, N., Infanger, M., Bauer, J., and Grimm, D. Genomic approach to identify factors that drive the formation of three-dimensional structures by EA.hy926 endothelial cells. *PLoS One* **8**, e64402, 2013.
 56. Konstantinova, N.A., Buravkova, L.B., Manuilova, E.S., and Grivennikov, I.A. The effects of gravity vector randomization on mouse embryonic stem cells *in vitro*. *J Gravit Physiol* **13**, 149, 2006.
 57. Schönberger, J., Bauer, J., Spruß, T., Weber, G., Chahoud, I., Eilles, C., and Grimm, D. Establishment and characterization of the follicular thyroid carcinoma cell line ML-1. *J Mol Med* **78**, 102, 2000.
 58. Weber, G., Grimm, D., and Bauer, J. Application of binary buffer systems to free flow cell electrophoresis. *Electrophoresis* **21**, 325, 2000.
 59. Ulbrich, C., Pietsch, J., Grosse, J., Wehland, M., Schulz, H., Saar, K., Hübner, N., Hauslage, J., Hemmersbach, R., Braun, M., Van Loon, J., Vagt, N., Egli, M., Richter, P., Einspanier, R., Sharbaty, S., Baltz, T., Infanger, M., Ma, X., and Grimm, D. Differential gene regulation under altered gravity conditions in follicular thyroid cancer cells: relationship between the extracellular matrix and the cytoskeleton. *Cell Physiol Biochem* **28**, 185, 2011.
 60. Grimm, D., Bauer, J., Ulbrich, C., Westphal, K., Wehland, M., Infanger, M., Aleshcheva, G., Pietsch, J., Ghardi, M., Beck, M., El-Saghire, H., De Saint-Georges, L., and Baatout, S. Different responsiveness of endothelial cells to vascular endothelial growth factor and basic fibroblast growth factor added to culture media under gravity and simulated microgravity. *Tissue Eng Part A* **16**, 1559, 2010.
 61. Qian, A.R., Zhang, W., Xie, L., Weng, Y., Yang, P., Wang, Z., Hu, L., Xu, H., Tian, Z., and Shang, P. Simulated weightlessness alters biological characteristics of human breast cancer cell line MCF-7. *Acta Astronaut* **63**, 947, 2008.

62. Uva, B.M., Masini, M.A., Sturla, M., Prato, P., Passalacqua, M., Giuliani, M., Tagliaferro, G., and Strollo, F. Clinorotation-induced weightlessness influences the cytoskeleton of glial cells in culture. *Brain Res* **934**, 132, 2002.
63. Rösner, H., Wassermann, T., Möller, W., and Hanke, W. Effects of altered gravity on the actin and microtubule cytoskeleton of human SH-SY5Y neuroblastoma cells. *Protoplasma* **229**, 225, 2006.
64. Hughes-Fulford, M. Function of the cytoskeleton in gravisensing during spaceflight. *Adv Space Res* **32**, 1585, 2003.
65. Nelson, L.J., Walker, S.W., Hayes, P.C., and Plevris, J.N. Low-shear modelled microgravity environment maintains morphology and differentiated functionality of primary porcine hepatocyte cultures. *Cells Tissues Organs* **192**, 125, 2010.
66. Ingber, D. How cells (might) sense microgravity, *FASEB J* **13**, S3, 1999.
67. Schiller, H.B., and Fässler, R. Mechanosensitivity and compositional dynamics of cell-matrix adhesions. *EMBO Rep* **14**, 509, 2013.
68. del Rio, A., Perez-Jimenez, R., Liu, R.C., Roca-Cusachs, P., Fernandez, J.M., and Sheetz, M.P. Stretching single talin rod molecules activates vinculin binding. *Science* **323**, 638, 2009.
69. Gershovich, P.M., Gershovich, J.G., and Buravkova, L.B. Simulated microgravity alters actin cytoskeleton and integrin-mediated focal adhesions of cultured human mesenchymal stromal cells. European Space Agency (Special Publication) ESA SP 663 SP. Noordwijk: European Space Agency, 2008.
70. Luo, M., Yang, Z., Li, J., Xu, H., Li, S., Zhang, W., Qian, A., and Shang, P. Calcium influx through stretch-activated channels mediates microfilament reorganization in osteoblasts under simulated weightlessness. *Adv Space Res* **51**, 2058, 2013.
71. Dai, Z.Q., Wu, F., Chen, J., Xu, H.J., Wang, H.H., Guo, F.M., Tan, Y.J., Ding, B., Wang, J.F., Wan, Y.M., and Li, Y.H. Actin microfilament mediates osteoblast Cbfa1 responsiveness to BMP2 under simulated microgravity. *PLoS One* **8**, e63661, 2013.
72. Chang, D., Guo, Y., and Liu, C. Effects of microgravity environment on intracellular signal transduction pathways. *Med J Chin Peoples Liberation Army* **37**, 749, 2012.
73. Spisni, E., Toni, M., Strillacci, A., Galleri, G., Santi, S., Griffoni, C., and Tomasi, V. Caveolae and caveolae constituents in mechanosensing: effect of modeled microgravity on cultured human endothelial cells. *Cell Biochem Biophys* **46**, 155, 2006.
74. Meng, R., Xu, H.Y., Di, S.M., Shi, D.Y., Qian, A.R., Wang, J.F., and Shang, P. Human mesenchymal stem cells are sensitive to abnormal gravity and exhibit classic apoptotic features. *Acta Biochim Biophys Sin (Shanghai)* **43**, 133, 2011.
75. Takahashi, A., Suzuki, H., Omori, K., Seki, M., Hashizume, T., Shimazu, T., Ishioka, N., and Ohnishi, T. The expression of p53-regulated genes in human cultured lymphoblastoid TSCE5 and WTK1 cell lines during spaceflight. *Int J Radiation Biol* **86**, 669, 2010.
76. Uva, B.M., Masini, M.A., Sturla, M., Bruzzone, F., Giuliani, M., Tagliaferro, G., and Strollo, F. Microgravity-induced apoptosis in cultured glial cells. *Eur J Histochem* **46**, 209, 2002.
77. Morbidelli, L., Monici, M., Marziliano, N., Cogoli, A., Fusi, F., Waltenberger, J. and Ziche, M. Simulated hypogravity impairs the angiogenic response of endothelium by up-regulating apoptotic signals. *Biochem Biophys Res Commun* **334**, 491, 2005.
78. Nakamura, H., Aoki, K., Masuda, W., Alles, N., Nagano, K., Fukushima, H., Osawa, K., Yasuda, H., Nakamura, I., Mikuni-Takagaki, Y., Ohya, K., Maki, K., and Jimi, E. Disruption of NF- κ B1 prevents bone loss caused by mechanical unloading. *J Bone Miner Res* **28**, 1457, 2013.
79. Clejan, S., O'Connor, K., and Rosensweig, N. Tri-dimensional prostate cell cultures in simulated microgravity and induced changes in lipid second messengers and signal transduction. *J Cell Mol Med* **5**, 60, 2001.
80. Grosse, J., Wehland, M., Pietsch, J., Schulz, H., Saar, K., Hubner, N., Eilles, C., Bauer, J., Abou-El-Ardat, K., Baatout, S., Ma, X., Infanger, M., Hemmersbach, R., and Grimm, D. Gravity-sensitive signaling drives 3-dimensional formation of multicellular thyroid cancer spheroids. *FASEB J* **26**, 5124, 2012.
81. Xu, D., Guo, S., and Liu, M. Identification of miRNAs involved in long-term simulated microgravity response in *Solanum lycopersicum*. *Plant Physiol Biochem* **66**, 10, 2013.
82. Fang, L., Deng, Z., Shatseva, T., Yang, J., Peng, C., Du, W.W., Yee, A.J., Ang, L.C., He, C., Shan, S.W., and Yang, B.B. MicroRNA miR-93 promotes tumor growth and angiogenesis by targeting integrin-beta 8. *Oncogene* **30**, 806, 2011.
83. Shilo, S., Roy, S., Khanna, S., and Sen, C.K. Evidence for the involvement of miRNA in redox regulated angiogenic response of human microvascular endothelial cells. *Arterioscler Thromb Vasc Biol* **28**, 471, 2008.
84. Pearson, J.D. Normal endothelial cell function. *Lupus* **9**, 183, 2000.
85. Alliegro, M.C. Pigpen and endothelial cell differentiation. *Cell Biol Int* **220**, 2077, 2001.
86. Dvorak, H.F. Vascular permeability factor/vascular endothelial growth factor: a critical cytokine in tumor angiogenesis and a potential target for diagnosis and therapy. *J Clin Oncol* **20**, 4368, 2002.
87. Schiffrin, E.L. The endothelium and control of blood vessel function in health and disease. *Clin Invest Med* **17**, 602, 1994.
88. Mantovani, A., Bussolino, F., and Introna, M. Cytokine regulation of endothelial cell function: from molecular level to the bedside. *Immunol Today* **18**, 231, 1997.
89. Wehland, M., Ma, X., Braun, M., Hauslage, J., Hemmersbach, R., Bauer, J., Grosse, J., Infanger M., and Grimm, D. The impact of altered gravity and vibration on endothelial cells during a parabolic flight. *Cell Physiol Biochem* **31**, 432, 2013.
90. Shinoka, T., Shum-Tim, D., Ma, P.X., Tanel, R.E., Isogai, N., Langer, R., Vacanti, J.P., and Mayer, J.E. Creation of viable pulmonary artery autografts through tissue engineering. *J Thorac Cardiovasc Surg* **115**, 536, 1998.
91. Chlupáč, J., Filová, E., and Bacáková, L. Blood vessel replacement: 50 years of development and tissue engineering paradigms in vascular surgery. *Physiol Res* **58(Suppl 2)**, S119, 2009.
92. Naito, Y., Shinoka, T., Duncan, D., Hibino, N., Solomon, D., Cleary, M., Rathore, A., Fein, C., Church, S., and Breuer, C. Vascular tissue engineering: towards the next generation vascular grafts. *Adv Drug Deliv Rev* **63**, 312, 2011.

93. Shi, F., Wang, Y.C., Zhao, T.Z., Zhang, S., Du, T.Y., Yang, C.B., Li, Y.H., and Sun, X.Q. Effects of simulated microgravity on human umbilical vein endothelial cell angiogenesis and role of the PI3K-Akt-eNOS signal pathway. *PLoS One* **7**, e40365, 2012.
94. Wang, Y.C., Zhang, S., Du, T.Y., Wang, B., and Sun, X.Q. Clinorotation upregulates inducible nitric oxide synthase by inhibiting AP-1 activation in human umbilical vein endothelial cells. *J Cell Biochem* **107**, 357, 2009.
95. Buravkova, L., Romanov, Y., Rykova, M., Grigorieva, O., and Merzlikina, N. Cell-to-cell interactions in changed gravity: ground-based and flight experiments. *Acta Astronaut* **57**, 67, 2005.
96. Zhang, X., Nan, Y., Wang, H., Chen, J., Wang, N., Xie, J., Ma, J., and Wang, Z. Model microgravity enhances endothelium differentiation of mesenchymal stem cells. *Naturwissenschaften* **100**, 125, 2013.
97. Kang, C.Y., Zou, L., Yuan, M., Wang, Y., Li, T.Z., Zhang, Y., Wang, J.F., Li, Y., Deng, X.W., and Liu, C.T. Impact of simulated microgravity on microvascular endothelial cell apoptosis. *Eur J Appl Physiol* **111**, 2131, 2011.
98. Versari, S., Villa, A., Bradamante, S., and Maier, J.A. Alterations of the actin cytoskeleton and increased nitric oxide synthesis are common features in human primary endothelial cell response to changes in gravity. *Biochim Biophys Acta* **1773**, 1645, 2007.
99. Cotrupi, S., Ranzani, D., and Maier, J.A. Impact of modeled microgravity on microvascular endothelial cells. *Biochim Biophys Acta* **1746**, 163, 2005.
100. Carlsson, S.I., Bertilaccio, M.T., Ballabio, E., and Maier, J.A. Endothelial stress by gravitational unloading: effects on cell growth and cytoskeletal organization. *Biochim Biophys Acta* **1642**, 173, 2003.
101. Grenon, S.M., Jeanne, M., Aguado-Zuniga, J., Conte, M.S., and Hughes-Fulford, M. Effects of gravitational mechanical unloading in endothelial cells: association between caveolins, inflammation and adhesion molecules. *Sci Rep* **3**, 1494, 2013.
102. Carlsson, S.I., Bertilaccio, M.T., Ascari, I., Bradamante, S., and Maier, J.A. Modulation of human endothelial cell behaviour in simulated microgravity. *J Gravit Physiol* **9**, P273, 2002.
103. Mariotti, M., and Maier, J.A. Gravitational unloading induces an anti-angiogenic phenotype in human microvascular endothelial cells. *J Cell Biochem* **104**, 129, 2008.
104. Infanger, M., Kossmehl, P., Shakibaei, M., Cogoli, A., Witzing, A., Faramarzi, S., Wittfoht, W., Paul, M., and Grimm, D. Vascular endothelial growth factor inhibits programmed cell death of endothelial cells induced by clinorotation. *J Gravit Physiol* **11**, P199, 2004.
105. Infanger, M., Ulbrich, C., Baatout, S., Wehland, M., Kreutz, R., Bauer, J., Grosse, J., Vadrucci, S., Cogoli, A., Derradji, H., Neefs, M., Küsters, S., Spain, M., Paul, M., and Grimm, D. Modeled gravitational unloading induced downregulation of endothelin-1 in human endothelial cells. *J Cell Biochem* **101**, 1439, 2007.
106. Chang, C.H., Lin, F.H., Kuo, T.F., and Liu, H.C. Cartilage tissue engineering. *Biomed Eng Appl Basis Commun* **17**, 61, 2005.
107. Mahmoudifar, N., and Doran, P.M. Tissue engineering of human cartilage in bioreactors using single and composite cell-seeded scaffolds. *Biotech Bioeng* **91**, 338, 2005.
108. Kock, L., van Donkelaar, C.C., and Ito, K. Tissue engineering of functional articular cartilage: the current status. *Cell Tissue Res* **347**, 613, 2012.
109. Darling, E.M., and Athanasiou, K.A. Rapid phenotypic changes in passaged articular chondrocyte subpopulations. *J Orthop Res* **23**, 425, 2005.
110. Goessler, U.R., Bugert, P., Bieback, K., Baisch, A., Sadick, H., Verse, T., Kluter, H., Hormann, K., and Riedel, F. Expression of collagen and fiber-associated proteins in human septal cartilage during *in vitro* dedifferentiation. *Int J Mol Med* **14**, 1015, 2004.
111. Goessler, U.R., Bieback, K., Bugert, P., Naim, R., Schaffer, C., Sadick, H., Hormann, K., and Riedel, F. Human chondrocytes differentially express matrix modulators during *in vitro* expansion for tissue engineering. *Int J Mol Med* **16**, 509, 2005.
112. Saltzman, W.M. Weaving cartilage at zero g: the reality of tissue engineering in space. *Proc Natl Acad Sci U S A* **94**, 13380, 1997.
113. Freed, L.E., Langer, R., Martin, I., Pellis, N.R., and Vunjak-Novakovic, G. Tissue engineering of cartilage in space. *Proc Natl Acad Sci U S A* **94**, 13885, 1997.
114. Marlovits, S., Tichy, B., Truppe, M., Gruber, D., and Vécsei, V. Chondrogenesis of aged human articular cartilage in a scaffold-free bioreactor. *Tissue Eng Part A* **9**, 1215, 2003.
115. Ohyabu, Y., Kida, N., Kojima, H., Taguchi, T., Tanaka, J., and Uemura, T. Cartilaginous tissue formation from bone marrow cells using rotating wall vessel (RWV) bioreactor. *Biotechnol Bioeng* **95**, 1003, 2006.
116. Stamenković, V., Keller, G., Nesic, D., Cogoli, A., and Grogan, S.P. Neocartilage formation in 1 g, simulated, and microgravity environments: implications for tissue engineering. *Tissue Eng Part A* **16**, 1729, 2010.
117. National Research Council. Recapturing a Future for Space Exploration: Life and Physical Sciences Research for a New Era. Washington, DC: The National Academies Press, 2011.
118. Strollo, F. Hormonal changes in humans during spaceflight. *Adv Space Biol Med* **7**, 99, 1999.
119. Grindeland, R.E., Popova, I.A., Vasques, M., and Arnaud, S.B. Cosmos 1887 mission overview: effects of microgravity on rat body and adrenal weights and plasma constituents. *FASEB J* **41**, 105, 1990.
120. Albi, E., Ambesi-Impiombato, F.S., Villani, M., De Pol, I., Spelat, R., Lazzarini, R., and Perrella, G. Thyroid cell growth: sphingomyelin metabolism as non-invasive marker for cell damage acquired during spaceflight. *Astrobiology* **10**, 811, 2010.
121. Albi, E., Curcio, F., Spelat, R., Lazzarini, A., Lazzarini, R., Loreti, E., Ferri, I., and Ambesi-Impiombato, F.S. Observing the mouse thyroid sphingomyelin under space conditions: a case study from the MDS mission in comparison with hypergravity conditions. *Astrobiology* **12**, 1035, 2012.
122. Masini, M.A., Albi, E., Barmo, C., Bonfiglio, T., Bruni, L., Cenesi, L., Cataldi, S., Curcio, F., D'Amora, M., Ferri, I., Goto, K., Kawano, F., Lazzarini, R., Loreti, E., Nakai, N., Ohira, T., Ohira, Y., Palmero, S., Prato, P., Ricci, F., Scrabelli, L., Shibaguchi, T., Spelat, R., Strollo, F., and Ambesi-Impiombato, F.S. The impact of long-term exposure to space environment on adult mammalian organisms: a study on mouse thyroid and testis. *PLoS One* **7**, e35418, 2012.

123. Albi, E., Curcio, F., Spelat, R., Lazzarini, A., Lazzarini, R., Loreti, E., Ferri, I., and Ambesi-Impimbato, F.S. Loss of parafollicular cells during gravitational changes (microgravity, hypergravity) and the secret effect of pleiotrophin. *PLoS One* **7**, e48518, 2012.
124. Infanger, M., Kossmehl, P., Shakibaei, M., Bauer, J., Kossmehl-Zorn, S., Cogoli, A., Curcio, F., Oksche, A., Wehland, M., Kreutz, R., Paul, M., and Grimm, D. Simulated weightlessness changes the cytoskeleton and extracellular matrix proteins in papillary thyroid carcinoma cells. *Cell Tissue Res* **324**, 267, 2006.
125. Kossmehl, P., Shakibaei, M., Cogoli, A., Infanger, M., Curcio, F., Schönberger, J., Eilles, C., Bauer, J., Pickenhahn, H., Schulze-Tanzil, G., Paul, M., and Grimm, D. Weightlessness induced apoptosis in normal thyroid cells and papillary thyroid carcinoma cells via extrinsic and intrinsic pathways. *Endocrinology* **144**, 4172, 2003.
126. Pietsch, J., Kussian, R., Sickmann, A., Bauer, J., Weber, G., Nissum, M., Westphal, K., Egli, M., Grosse, J., Schönberger, J., Wildgruber, R., Infanger, M., and Grimm, D. Application of free-flow IEF to identify protein candidates changing under microgravity conditions. *Proteomics* **10**, 904, 2010.
127. Pietsch, J., Sickmann, A., Weber, G., Bauer, J., Egli, M., Wildgruber, R., Infanger, M., and Grimm, D. Metabolic enzyme diversity in different human thyroid cell lines and their sensitivity to gravitational forces. *Proteomics* **12**, 2539, 2012.
128. Pietsch, J., Riwaltdt, S., Bauer, J., Sickmann, A., Weber, G., Grosse, J., Infanger, M., Eilles, C., and Grimm, D. Interaction of proteins identified in human thyroid cells. *Int J Mol Sci* **14**, 1164, 2013.
129. Martin, A., Zhou, A., Gordon, R.E., Henderson, S.C., Schwartz, A.E., Friedmann, E.W., and Davies, T.F. Thyroid organoid formation in simulated microgravity: influence of keratinocyte growth factor. *Thyroid* **10**, 481, 2000.
130. Konstantinov, S.M., Mindova, M.M., Gospodinov, P.T., and Genova, P.I. Three-dimensional bioreactor cultures: a useful dynamic model for the study of cellular interactions. *Ann N Y Acad Sci* **1030**, 103, 2004.
131. Chang, T.T., and Hughes-Fulford, M. Monolayer and spheroid culture of human liver hepatocellular carcinoma cell line cells demonstrate distinct global gene expression patterns and functional phenotypes. *Tissue Eng Part A* **15**, 559, 2009.
132. Weaver, V.M., Petersen, O.W., Wang, F., Larabell, C.A., Briand, P., Damsky, C., and Bissell, M.J. Reversion of the malignant phenotype of human breast cells in three-dimensional culture and *in vivo* by integrin blocking antibodies. *J Cell Biol* **137**, 231, 1997.
133. Ma, X., Pietsch, J., Wehland, M., Schulz, H., Saar, K., Hübner, N., Bauer, J., Braun, M., Schwarzwälder, A., Segerer, J., Birlem, M., Horn, A., Hemmersbach, R., Waßer, K., Grosse, J., Infanger, M., and Grimm, D. Differential gene expression profile and altered cytokine secretion of thyroid cancer cells in space. *FASEB J* **28**, 813, 2014.

Address correspondence to:

Daniela Grimm, MD
 Institute of Biomedicine, Pharmacology
 Aarhus University
 Wilhelm Meyers Allé 4
 Aarhus C DK-8000
 Denmark

E-mail: daniela.grimm@farm.au.dk

Received: November 20, 2013

Accepted: February 19, 2014

Online Publication Date: April 3, 2014