Published in final edited form as:

Eur J Neurosci. 2013 March ; 37(6): 850–859. doi:10.1111/ejn.12098.

GABA_A, NMDA, AND mGlu2 RECEPTORS TONICALLY REGULATE INHIBITION AND EXCITATION IN THE THALAMIC RETICULAR NUCLEUS

John W. Crabtree¹, David Lodge¹, Zafar I. Bashir¹, and John T.R. Isaac²

¹Medical Research Council Centre for Synaptic Plasticity, School of Physiology and Pharmacology, Medical Sciences Building, University of Bristol, Bristol BS8 1TD, UK

²Eli Lilly and Company, Erl Wood Manor, Windlesham, Surrey GU20 6PH, UK

Abstract

Traditionally, neurotransmitters are associated with a fast, or phasic, type of action on neurons in the CNS. However, accumulating evidence indicates that GABA and glutamate can also have a continual, or tonic, influence on these cells. Here, in voltage- and current-clamp recordings in rat brain slices, we identify three types of tonically active receptors in a single CNS structure, the thalamic reticular nucleus (TRN). Thus, TRN contains constitutively active GABA_A receptors (GABA_ARs), which are located on TRN neurons and generate a persistent outward Cl[−] current. When TRN neurons are depolarized, blockade of this current increases their action potential output in response to current injection. Furthermore, TRN contains tonically active GluN2B-containing N-methyl-D-aspartate receptors (NMDARs). These are located on reticuloreticular GABAergic terminals in TRN and generate a persistent facilitation of vesicular GABA release from these terminals. In addition, TRN contains tonically active metabotropic glutamate type 2 receptors (mGlu2Rs). These are located on glutamatergic cortical terminals in TRN and generate a persistent reduction of vesicular glutamate release from these terminals. Although tonically active GABA_ARs, NMDARs, and mGlu2Rs operate through different mechanisms, we propose that the continual and combined activity of these three receptor types ultimately serves to hyperpolarize TRN neurons, which will differentially affect the output of these cells depending upon the current state of their membrane potential. Thus, when TRN cells are relatively depolarized, their firing in single-spike tonic mode will be reduced, whereas, when these cells are relatively hyperpolarized, their ability to fire in multispikes burst mode will be facilitated.

Keywords

constitutively active receptors; miniature inhibitory postsynaptic currents; miniature excitatory postsynaptic currents; thalamic reticular neurons; Wistar rat

Corresponding author: John W. Crabtree, MRC Centre for Synaptic Plasticity, School of Physiology and Pharmacology, Medical Sciences Building, University Walk, University of Bristol, Bristol BS8 1TD, United Kingdom, Fax: +44 (0)117 3312288, j.w.crabtree@bristol.ac.uk.

The other authors declare no conflicts of interest, financial or otherwise.

Introduction

Neurotransmitters are usually associated with a fast (phasic) type of action on neurons in the CNS, but mounting evidence indicates that GABA and glutamate can also have a persistent (tonic) influence on these cells (Semyanov *et al.*, 2004; Cavelier *et al.*, 2005; Farrant & Nusser, 2005; Glykys and Mody, 2007; Featherstone & Shippy, 2008; Brickley & Mody, 2012). Unlike receptors located at the synapse that respond phasically to relatively high concentrations of synaptically released neurotransmitter, tonically active receptors are located at perisynaptic or extrasynaptic sites and continually respond to relatively low concentrations of ambient GABA and glutamate in the extracellular space. Postsynaptically, tonically active receptors will affect neuronal excitability by continually generating a membrane conductance. Presynaptically, tonically active receptors will affect this excitability by continually modulating the inhibitory or excitatory ‘tone’ that results from vesicular release of GABA or glutamate, respectively. In the thalamus, control of neuronal excitability is thought to depend on the interactions among intrinsic membrane conductances and relatively long-lasting changes in membrane potentials due to activation of postsynaptic receptors (Sherman & Guillery, 2001). However, this view should be broadened to take into account the effects of tonically active receptors in the thalamus (Belelli *et al.*, 2005; Cope *et al.*, 2005; Jia *et al.*, 2005; Bright *et al.*, 2007; Bright & Brickley, 2008).

The thalamic reticular nucleus (TRN) is a sheet of GABAergic neurons (Houser *et al.*, 1980) that surrounds much of the lateral and rostral margins of the main body of thalamic relay nuclei. TRN neurons receive glutamatergic inputs from collaterals of axons arising from thalamocortical (TC) neurons (Jones, 1975) and layer VI corticothalamic (CT) neurons (Bourassa *et al.*, 1995). The main targets of TRN GABAergic outputs are TC neurons (Jones, 1975). Thus, TRN neurons provide a strong inhibitory innervation of TC neurons (Cox *et al.*, 1997; Kim & McCormick, 1998), which will strongly influence their transfer of sensory information to cortex (Salt, 1989; Lee *et al.*, 1994a b; Warren & Jones, 1994; Hartings & Simons, 2000; Cotillon-Williams *et al.*, 2008). Therefore, mechanisms regulating the excitability of TRN neurons will make important contributions to the control of information flow through the thalamus.

Several mechanisms intrinsic to TRN have been identified that regulate the excitability of its neurons, including chemical (Sanchez-Vives *et al.*, 1997; Deleuze & Huguenard, 2006; Lam *et al.*, 2006; Mistry *et al.*, 2008) and electrical (Landisman *et al.*, 2002; Long *et al.*, 2004) synapses, depolarization-induced suppression of inhibition (Sun *et al.*, 2011), and short-term depression at glutamatergic synapses (Mistry *et al.*, 2008). However, demonstration of additional regulatory mechanisms in TRN that are mediated by tonically active receptors has been inconclusive (Belelli *et al.*, 2005; Cope *et al.*, 2005; Jia *et al.*, 2005). Here we show that TRN contains three types of tonically active receptors, postsynaptic GABA_A receptors (GABA_ARs), presynaptic N-methyl-D-aspartate receptors (NMDARs), and presynaptic metabotropic glutamate type 2 receptors (mGlu2Rs). Although operating through different mechanisms, the combined and continual activity of these three types of receptors will be to hyperpolarize TRN neurons.

Materials and Methods

Thalamic slice preparation

All experimental procedures were performed in accordance with the UK Animals (Scientific Procedures) Act 1986 and ethically approved by the University of Bristol. Young (14- to 21-d-old) Wistar rats of either sex were killed by cervical dislocation. The brain was rapidly removed and placed in ice-cold oxygenated (95% O₂, 5% CO₂) artificial cerebrospinal fluid (ACSF). Horizontal slices (500 µm thick) were cut through the thalamus on a vibratome and were allowed to recover in oxygenated ACSF for at least 1 h at room temperature (19–22°C). Slices were then transferred to a recording chamber and submerged beneath continuously perfusing oxygenated ACSF. The ACSF contained (in mM): 119 NaCl, 2.5 KCl, 1.0 NaH₂PO₄, 26.2 NaHCO₃, 11.0 glucose, 2.5 CaCl₂, and 1.3 MgSO₄.

Electrophysiology

Using 3–5 MΩ glass electrodes, whole-cell voltage- or current-clamp recordings were made from neurons in a centrovventral sector of TRN (Crabtree & Isaac, 2002). Transilluminating a slice created contrast interference patterns that allowed visualization of TRN and adjacent thalamic nuclei (Crabtree & Isaac, 2002; Mistry *et al.*, 2008). During experiments a recording electrode was advanced ‘blind’ through a slice and only one neuron per slice was recorded. For voltage-clamp recordings, the whole-cell solution contained (in mM): 135 Cs methane sulphonate, 10 HEPES, 0.5 EGTA, 3.0 NaCl, 5.0 QX-314, 4.0 Mg-ATP, and 0.3 Na-GTP, pH 7.2 using CsOH (275–280 mOsm). For current-clamp recordings, the whole-cell solution contained (in mM): 130 K methane sulphonate, 5.0 HEPES, 0.2 EGTA, 8.5 NaCl, 4.0 Mg-ATP, and 0.5 Na-GTP, pH 7.2 using KOH (275–280 mOsm). During voltage-clamp recordings, different combinations of 6-imino-3(4-methoxyphenyl)-1-(6H)-pyridazinebutanoic acid hydrobromide (gabazine; 50 µM or 100 µM) or picrotoxin (50 µM), 2,3-Dioxo-6-nitro-,1,2,3,4-tetrahydrobenzo[f]quinoxaline-7-sulfonamide (NBQX; 2 µM), D-(–)-2-Amino-5-phosphonopentanoic acid (D-AP5; 50 µM), and tetrodotoxin (TTX; 1 µM) were bath applied in the ACSF to block synaptic currents mediated by GABA_ARs, α-amino-3-hydroxy-5-methylisoxazole-4-propionic acid receptors (AMPA_Rs), NMDARs, and action potentials presynaptic to a recorded cell, respectively, as specified for the various experimental procedures. During voltage-clamp recordings, cesium ions and QX-314 in the whole-cell solution blocked postsynaptic currents mediated by GABA_B receptors (GABA_BRs). In some of these recordings, (5R,10S)-(–)-5-methyl-10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5,10-imine maleate (MK-801; 3 mM) was included in the whole-cell solution (iMK-801) to block postsynaptic NMDAR-mediated currents. To ensure blockade of these currents, neurons were depolarized to –10 mV for 10 s once every min for 10 min after whole-cell access (Woodhall *et al.*, 2001). In other voltage-clamp recordings, guanosine 5′-[β-thio]diphosphate trilithium salt (GDPβS; 1 mM) was included in the whole-cell solution to block postsynaptic mGluR-mediated currents (Lee & McCormick, 1997; Cox & Sherman, 1999; Alexander & Godwin, 2006) and data acquisition began 20 min after whole-cell access. All drugs were obtained from Ascent Scientific except for RO 25-6981, which was obtained from Tocris Bioscience, and picrotoxin and GDPβS, which were obtained from Sigma-Aldrich. During voltage-clamp recordings, cells were held at 0 mV or –60 mV as specified for the various experimental procedures. During current-clamp

recordings, the membrane potential was close to -55 mV. Using an in-line heater (Scientifica, Uckfield, UK), all recordings were made at close to physiological temperature (32 – 35°C).

Data acquisition and analysis

Recordings were made using an Axopatch 200B amplifier (Axon Instruments, Foster City, CA, USA). Data were filtered at 5 kHz, digitized at 10 kHz, and stored on computer. Holding current (DC), series resistance (R_s), input resistance (R_m), inhibitory postsynaptic currents (IPSCs), miniature inhibitory (or excitatory) postsynaptic currents (mI(E)PSCs), and action potentials were displayed on-line using WinLTP (Anderson & Collingridge, 2007). Data were acceptable from neurons with a stable whole-cell access (on average about 25 M Ω) or were discarded if the R_s changed by $>20\%$. DC was measured at the beginning of 2 -s periods sampled every 4 s. DC measurements were discarded if they fell on IPSCs. R_m was measured at the beginning of 5 -s periods sampled every 20 s or at the beginning of each of 6 5 -s periods sampled every 2 min. R_m measurements were discarded if they fell on mIPSCs or mEPSCs. During a 5 -min baseline period, 100 – 120 IPSCs were sampled and averaged for each neuron using Clampfit (Axon Instruments). The total IPSC current, or IPSC charge transfer (integral of the average IPSC) multiplied by IPSC frequency, was then determined for each cell. mIPSCs were recorded over 5 -s periods sampled every 20 s and 18 – 24 trials were given before and after a 10 min wash-in of an agonist or antagonist. mEPSCs were recorded over 5 -s periods for 6 trials sampled every 2 min. Amplitudes and frequencies of mI(E)PSCs were analyzed using Mini-Analysis (Synaptosoft, Decatur, GA, USA). Detection threshold for mI(E)PSCs was 10 pA. Action potentials were recorded over 200 ms current-pulse periods sampled every 10 s and 72 trials were given before and after a 10 min wash-in of picrotoxin (50 μM). Pooled data are expressed as cumulative probabilities or as means $\pm$ SEM. Statistical significance was assessed using the Wilcoxon matched-pair test or Kolmogorov-Smirnov test. Differences were considered significant when $p < 0.05$.

Results

TRN contains tonically active GABA_ARs

Blockade of intra-TRN inhibition increased the tonic spike output of TRN neurons to stimulation of their glutamatergic inputs (Mistry *et al.*, 2008). During such blockade, we observed that the amplitudes of the first evoked excitatory EPSPs in response trains increased compared to baseline amplitudes. This suggested that, in addition to intra-TRN inhibition, another source of inhibition was present in TRN because, under control conditions, the first evoked EPSP in a response train will occur much earlier than any influence intra-TRN inhibition will have on neuronal responses. Tonically active GABA_ARs were a likely candidate to account for this other source of inhibition (Semyanov *et al.*, 2004; Cavalier *et al.*, 2005; Farrant & Nusser, 2005; Glykys & Mody, 2007; Belelli *et al.*, 2009; Brickley & Mody, 2012). Therefore, we tested the hypothesis that tonically active GABA_ARs are present on TRN neurons.

TRN neurons were recorded in voltage-clamp mode and held at 0 mV in the presence of NBQX (2 μM) and D-AP5 (50 μM). Bath-applied gabazine (50 μM ; $n = 11$, or 100 μM ; $n =$

3) abolished phasic IPSCs and had no effect on the holding current (Fig. 1A). However, bath-applied picrotoxin (50 μ M) not only abolished phasic IPSCs but also caused a reduction in an outward holding current (Fig. 1B); in 15 of 18 TRN neurons (83.3%), there was a significant shift in holding current averaged over the last 5 min compared to the holding current averaged over the first 5 min of each experiment (Fig. 1C). The amplitude of this current shift was 21.1 ± 8.9 pA. These findings are consistent with the blockade of constitutively active postsynaptic GABA_ARs that mediate a tonic Cl⁻ current. Because of this persistent current, constitutively active postsynaptic GABA_ARs would provide a mechanism for reducing the excitability of TRN neurons.

The proportions of the total intra-TRN GABA_AR inhibition mediated by phasic and tonic currents were estimated. For the 15 TRN neurons in which a tonic current was observed, IPSCs contributed $5.1 \pm 4.7\%$ (1.0 ± 0.5 of 22.1 ± 9.4 pA) of the total GABA_AR-mediated current, whereas the tonic component contributed $94.9 \pm 4.7\%$ (21.1 ± 8.9 of 22.1 ± 9.4 pA) of this total current. Thus, most (>90%) of the intra-TRN GABA_AR-mediated inhibition is carried by tonically active GABA_ARs.

In the cerebellum (Brickley *et al.*, 1996; Hamann *et al.*, 2002), hippocampus (Semyanov *et al.*, 2003), and thalamic dorsal lateral geniculate nucleus (Cope *et al.*, 2005), tonically active GABA_ARs reduce action potential output of neurons. Therefore, we tested the effect of tonic postsynaptic GABA_ARs on the excitability of TRN neurons. TRN neurons (n=10) were recorded in current-clamp mode and held close to -55 mV before and during the presence of bath-applied picrotoxin (50 μ M). Under control conditions, these cells fired action potentials (spikes) in response to current injections (200 ms pulses) and the spike frequency increased with increasing levels of current injection (Fig. 1D). Compared to control, the spike frequency significantly increased at all levels of current injection used, except the lowest level, in the presence of picrotoxin (Fig. 1D). Thus, removal of the tonic GABA_AR-mediated current by picrotoxin increases the excitability of TRN neurons.

TRN contains tonically active NMDARs

Axon terminals in neocortex (Berretta & Jones, 1996; Corlew *et al.*, 2007; Brasier & Feldman, 2008), hippocampus (Breukel *et al.*, 1998), and cerebellum (Glitsch & Marty, 1999) exhibit tonically active NMDARs. These receptors continually facilitate the vesicular release of GABA or glutamate onto a neuron depending on whether the recipient neuron is GABAergic or glutamatergic, respectively. Thus, an additional mechanism possibly involved in affecting the excitability of TRN neurons is the tonic activation of NMDARs on GABAergic terminals. Therefore, we next investigated whether such receptors are also present on GABAergic terminals in TRN.

TRN neurons were recorded in voltage-clamp mode and held at 0 mV. By electrically stimulating TC and CT axons (see Mistry *et al.*, 2008), we initially recorded EPSCs evoked in TRN neurons (n=3) in the presence of bath-applied NBQX (2 μ M) and confirmed that intracellularly applied MK-801 (iMK-801; 3 mM) was effective in blocking postsynaptic NMDARs (Fig. 2A insert). In control conditions, outward mIPSCs (Fig. 2A-C) were recorded in the presence of TTX (1 μ M), NBQX (2 μ M), and iMK-801 (3 mM). The presence of tonically active presynaptic NMDARs was then tested by recording mIPSCs

under two conditions: during bath application of an NMDAR agonist N-methyl-D-aspartate (NMDA) or an NMDAR antagonist D-AP5. In 6 of 6 TRN neurons, there was no change in the amplitudes of mIPSCs (Figs. 2A left and 2D left) but the frequency of mIPSCs significantly increased (Figs. 2A right and 2D right) following wash-in of NMDA (40 μ M). This increase in mIPSC frequency was $22.0 \pm 3.6\%$. In 8 of 8 TRN neurons, there was again no change in the amplitudes of mIPSCs (Figs. 2B left and 2D left) but the frequency of mIPSCs now significantly decreased (Figs. 2B right and 2D right) following wash-in of D-AP5 (50 μ M). This decrease in mIPSC frequency was $39.5 \pm 1.4\%$. Additional wash-in of picrotoxin (50 μ M) abolished the mIPSCs indicating that they were mediated by GABA_ARs (Fig. 2B insert). That there were changes only in the frequencies of mIPSCs in the presence of NMDA or D-AP5 is indicative of the presence of presynaptic NMDARs. That the frequency of mIPSCs decreased below the baseline level in the presence of D-AP5 indicates that these presynaptic NMDARs are tonically active and facilitate the release of GABA from terminals in TRN. Tonically active NMDARs are reported to contain a GluN2B subunit (Woodhall *et al.*, 2001; Brasier & Feldman, 2008; Larsen *et al.*, 2011). We then tested whether the tonic NMDARs in TRN contained this subunit by recording mIPSCs during bath application of a GluN2B antagonist (α R, β S)- α -(4-hydroxyphenyl)- β -methyl-4-(phenylmethyl)-1-piperidinepropanol maleate (Ro 25-6981). In 6 of 6 TRN neurons, there was no change in the amplitudes of mIPSCs (Figs. 2C left and 2D left) but the frequency of mIPSCs significantly decreased (Figs. 2C right and 2D right) following wash-in of Ro 25-6981 (3 μ M). This decrease in mIPSC frequency was only $28.1 \pm 2.2\%$, suggesting that tonically active NMDARs on GABAergic terminals in TRN may contain a subunit other than GluN2B. For the neurons in the above experiments, there was no change in Rm following wash-in of NMDA (40 μ M) (baseline, 101.6 ± 9.7 M Ω ; NMDA, 98.4 ± 8.4 M Ω ; $p > 0.05$; Wilcoxon matched-pair test), D-AP5 (50 μ M) (baseline, 125.1 ± 7.0 M Ω ; D-AP5, 128.7 ± 6.9 M Ω ; $p > 0.05$; Wilcoxon matched-pair test), or Ro 25-6981 (3 μ M) (baseline, 123.7 ± 8.0 M Ω ; Ro 25-6981, 126.4 ± 8.3 M Ω ; $p > 0.05$; Wilcoxon matched-pair test).

For the baseline periods, the cumulative probability plots for inter-mIPSC intervals for the NMDA (Fig. 2A right), D-AP5 (Fig. 2B right), and Ro 25-6981 (Fig. 2C right) conditions indicate substantial between-cell variability in mIPSC frequency. Although this variability was evident when the inter-mIPSC interval data were plotted for each neuron in the NMDA (Fig. 3A), D-AP5 (Fig. 3B), and Ro 25-6981 (Fig. 3C) conditions, all the cells in each condition showed the same downward or upward shift in mean inter-mIPSC interval following agonist or antagonist application, respectively. Furthermore, there was no indication that these applications had stronger or weaker effects on cells with substantially different basal mIPSC frequencies.

TRN contains tonically active mGlu2Rs

Axon terminals in hippocampus (Losonczy *et al.*, 2003; Lauri *et al.*, 2006) and hypothalamus (Boudaba *et al.*, 2003; Acuna-Goycolea *et al.*, 2004) exhibit tonically active mGluRs. These receptors continually reduce the vesicular release of glutamate onto a neuron. Thus, another possible mechanism involved in affecting the excitability of TRN neurons is the tonic activation of mGluRs on glutamatergic terminals. Although presynaptic Group II mGluRs (mGlu2R and mGlu3R) do not appear to be present at thalamoreticular

synapses (Ohishi *et al.*, 1993ab), such presynaptic mGluRs are present at corticoreticular synapses and are presumably mGlu2Rs (Alexander & Godwin, 2006). Therefore, we next investigated whether these presumptive mGlu2Rs are tonically active.

TRN neurons were recorded in voltage-clamp mode and held at -60 mV. In control conditions, inward mEPSCs (Fig. 4A,C) were recorded in the presence of TTX (1 μ M), D-AP5 (50 μ M), and picrotoxin (50 μ M). In addition, GDP β S (1 mM) was included in the intracellular recording solution to block any postsynaptic effects of mGluRs (Lee & McCormick, 1997; Cox & Sherman, 1999; Alexander & Godwin, 2006). The presence of tonically active presynaptic mGlu2Rs was then tested by recording mEPSCs under two conditions: during bath application of a selective mGlu2R agonist (1SR,2SR,4RS,5RS,6SR)-2-amino-4-methylbicyclo[3.1.0]-hexane-2,6-dicarboxylic acid (LY395756) (Dominguez *et al.*, 2005; Ceolin *et al.*, 2011) or a selective Group II mGluR antagonist (2S,1'S,2'S)-2-(9-xanthylmethyl)-2-(2'-carboxycyclopropylglycine (LY341495) (Kingston *et al.*, 1998). Bath application of LY395756 (3 μ M) caused a reversible decrease in the frequency of mEPSCs (Fig. 4A). In 3 of 3 TRN neurons, there was a significant decrease in mEPSC frequency averaged over the points at 18–24 min compared to the mEPSC frequency averaged over the points at 2–10 min (Fig. 4B). This decrease was $36.3 \pm 3.4\%$. In comparing these two time periods, there was no change in the amplitudes of mEPSCs (baseline, 16.4 ± 0.6 pA; LY395756, 16.1 ± 1.1 pA; $p > 0.05$; Wilcoxon matched-pair test). In contrast, bath application of LY341495 (300 nM) caused a reversible increase in the frequency of mEPSCs (Fig. 4C). In 8 of 8 TRN neurons, there was a significant increase in mEPSC frequency averaged over the points at 18–24 min compared to the mEPSC frequency averaged over the points at 2–10 min (Fig. 4D). This increase was $52.9 \pm 4.9\%$. In comparing these two time periods, there was no change in the amplitudes of mEPSCs (Fig. 4E left) but the frequency of mEPSCs significantly increased (Fig. 4E right). Additional wash-in of NBQX (2 μ M) abolished the mEPSCs indicating that they were mediated by AMPARs (Fig. 4E insert). That there were changes only in the frequency of mEPSCs in the presence of LY341495 is indicative of the presence of presynaptic mGlu2Rs. That the frequency of mEPSCs increased above the baseline level in the presence of LY341495 indicates that these presynaptic mGlu2Rs are tonically active and reduce the release of glutamate from terminals in TRN. During bath application of LY395756 (3 μ M) or LY341495 (300 nM) there was no change in R_m averaged over the points at 18–24 min compared to the baseline R_m averaged over the points at 2–10 min (baseline, 162.8 ± 16.6 M Ω ; LY395756, 163.8 ± 16.1 M Ω ; $p > 0.05$; Wilcoxon matched-pair test; baseline, 184.2 ± 10.1 M Ω ; LY341495, 180.1 ± 10.6 M Ω ; $p > 0.05$; Wilcoxon matched-pair test).

Discussion

Here we show the presence of three types of tonically active receptors in a single CNS structure, the thalamic reticular nucleus (TRN). Figure 5 summarizes our findings: each of the three receptors that we have studied is indicated by an asterisk. Thus, TRN contains constitutively active postsynaptic GABA_ARs, which generate in TRN neurons a persistent outward Cl[−] current approximately 21 pA in amplitude. Furthermore, TRN exhibits tonically active GluN2B-containing NMDARs on reticuloreticular GABAergic terminals, which generate a continual facilitation of GABA release as indicated by an approximately 40% or

28% decrease in the frequency of mIPSCs when these NMDARs are blocked by D-AP5 or R0 25-6981, respectively. Moreover, TRN contains tonically active mGlu2Rs on corticoreticular glutamatergic terminals, which generate a continual reduction of glutamate release as indicated by an approximately 53% increase in the frequency of mEPSCs when these mGlu2Rs are blocked by LY341495. Therefore, each of these receptor types serves to hyperpolarize TRN neurons. It remains to be determined whether individual TRN neurons are influenced by all three of these tonically active receptor types.

The presence of tonically active GABA_ARs appears ubiquitous in the CNS. Neurons containing these receptors are found in the neocortex (Salin & Prince, 1996; Krook-Magnuson *et al.*, 2008), dentate gyrus (Nusser & Mody, 2002; Stell & Mody, 2002), hippocampus (Bai *et al.*, 2001; Semyanov *et al.*, 2003), dorsal striatum (Ade *et al.*, 2008), amygdala (Marowsky *et al.*, 2012), thalamic relay nuclei (Porcello *et al.*, 2003; Belelli *et al.*, 2005; Cope *et al.*, 2005; Jia *et al.*, 2005), cerebellum (Kaneda *et al.*, 1995; Brickley *et al.*, 1996; Wall & Usowicz, 1997), and spinal cord (Wang *et al.*, 2008). However, in contrast to previous studies in rodents (Belelli *et al.*, 2005; Cope *et al.*, 2005; Jia *et al.*, 2005), we now find tonically active GABA_ARs in TRN. The discrepancy between our findings and these previous studies is most likely due to the types of GABA_AR antagonists used. Whereas the earlier studies used only competitive GABA_AR antagonists (bicuculline or gabazine), we used both a competitive (gabazine) and a noncompetitive (picrotoxin) antagonist. But it is only when picrotoxin is used that a reduction in holding current in TRN neurons occurs. This different result when using a noncompetitive antagonist is entirely consistent with the presence of constitutively active GABA_ARs in TRN. Such receptors do not require the endogenous ligand (GABA) to open the channel and therefore are not sensitive to orthosteric antagonists but are sensitive to the open channel blocker picrotoxin (Sigel *et al.*, 1989; Krishek *et al.*, 1996; Bouairi *et al.*, 2006; McCartney *et al.*, 2007).

In rodents, functional GABA_ARs are pentameric structures that are usually composed of combinations of at least 1 of 6 α subunits, 1 of 3 β subunits, and 1 of 3 γ subunits, although a δ subunit can replace a γ subunit (Pirker *et al.*, 2000). Different combinations of subunits will impart distinct pharmacological and electrophysiological properties onto GABA_ARs (Sieghart *et al.*, 1999; Mody & Pearce, 2004; Semyanov *et al.*, 2004; Farrant & Nusser, 2005; Glykys & Mody, 2007; Belelli *et al.*, 2009; Eyre *et al.*, 2012). GABA_ARs in TRN express the $\alpha 3$, $\beta 3$, and $\gamma 2$ subunits but not the $\alpha 1$, $\alpha 4$, $\beta 2$, and δ subunits (Pirker *et al.*, 2000; Eyre *et al.*, 2012). The combinations of subunits that make up phasic (synaptic) and tonic (constitutively active) GABA_ARs in TRN are currently unknown.

GABA_ARs containing $\beta 1$ subunits are also expressed in the rodent TRN (Pirker *et al.*, 2000; Huntsman & Huguenard, 2006). These subunits impart a much greater heterogeneity in the electrophysiological properties of GABA_ARs (Huntsman & Huguenard, 2006). Homomeric $\beta 1$ subunit-containing GABA_ARs form functional Cl⁻ selective channels (Sigel *et al.*, 1989; Krishek *et al.*, 1996). These receptors exhibit channel opening in the absence of GABA, are insensitive to GABA and orthosteric antagonists, and are blocked by picrotoxin. Therefore, homomeric $\beta 1$ GABA_ARs would be ideally suited to fulfil the role of constitutively active receptors that generate a persistent outward Cl⁻ current in TRN neurons. If these receptors

are present in TRN, they could be regulated intracellularly by postsynaptic activation of G-protein-coupled receptors.

Picrotoxin (50 μ M) can act as a partial antagonist at homomeric α subunit-containing glycine receptors (GlyRs) (Chattipakorn & McMahon, 2002). Therefore, the presence of these receptors in TRN could account for the persistent outward Cl^- current that we attribute to tonically active GABA_ARs. GlyR $\alpha 2$ and β subunits are expressed early in development in the rat TRN (Malosio *et al.*, 1991). Although nonligand-binding β subunits are still expressed here in the adult, there is no expression of ligand-binding $\alpha 2$ subunits in TRN on postnatal day 15 or thereafter. Because our data investigating tonic Cl^- currents are from 2- to 3-week-old rats, it is unlikely that blockade of GlyRs explains the picrotoxin-induced reduction in holding current of TRN neurons.

Our data showing changes in postsynaptic miniature event frequency indicate that TRN contains tonically active presynaptic NMDARs that continually facilitate GABA release from reticuloreticular terminals. Such presynaptic facilitation of transmitter release agrees with previous findings in the neocortex (Berretta and Jones, 1996; Corlew *et al.*, 2007; Brasier & Feldman, 2008), hippocampus (Breukel *et al.*, 1998), and cerebellum (Glitsch & Marty, 1999). Possible presynaptic NMDARs in the somatodendritic compartment appear to play only a small role in changing the frequency of postsynaptic miniature events (Glitsch & Marty, 1999). Thus, the presynaptic NMDARs responsible for these changes would be mostly located in the axonal domain. These receptors contain a GluN2B subunit that is selectively antagonized by RO 25-6981. However, this compound is only a partial antagonist of such subunits (Bartlett *et al.*, 2007; Paoletti & Neyton, 2007) so that, in our experiments, the reduction in frequency of mIPSCs by RO 25-6981 is only about 70% of that produced by D-AP5 . Therefore, the possibility remains that not all tonically active NMDARs in TRN necessarily contain GluN2B subunits.

Our data showing changes in postsynaptic miniature event frequency further indicate that TRN contains tonically active presynaptic mGlu2Rs that continually reduce glutamate release from corticoreticular terminals. Such presynaptic mGluR-mediated reduction of transmitter release agrees with previous findings in the hippocampus (Losonczy *et al.*, 2003; Lauri *et al.*, 2006) and hypothalamus (Boudaba *et al.*, 2003; Acuna-Goycolea *et al.*, 2004). However, 3 μ M LY395756 not only acts as an agonist at mGlu2Rs but can also act as an antagonist at mGlu3Rs (Dominguez *et al.*, 2005; Ceolin *et al.*, 2011). Thus, in TRN, the mGlu2R-mediated decrease in mEPSC frequency of approximately 36% may be an underestimation due to an opposing effect mediated by mGlu3Rs. Furthermore, 300nM LY341495 is not only a Group II mGluR antagonist but can also act as an antagonist at mGlu8Rs (Kingston *et al.*, 1998). Because TRN does not appear to express mGlu8Rs (Saugstad *et al.*, 1997), the increase in mEPSC frequency of approximately 53% that we see during LY341495 application may accurately reflect the antagonist effect of this compound on mGlu2/3Rs regulating glutamate release from corticoreticular terminals.

Presynaptic Group II mGluRs do not appear to be present at reticuloreticular synapses (Alexander and Godwin, 2006). However, the tonically active presynaptic NMDARs that we have identified in TRN may be expressed at other locations in this structure or TRN may

express other types of tonically active presynaptic receptors. Thus, adenosine A1 receptors are present on GABAergic and glutamatergic terminals (Ulrich and Huguenard, 1995), GABA_BRs are present on GABAergic (Ulrich and Huguenard, 1996) and thalamoreticular (Sanchez-Vives et al., 1997) terminals, the Group III mGlu4Rs are present on thalamoreticular terminals (Snead et al., 2000), and kainate receptors are present on corticoreticular terminals (Miyata and Imoto, 2009). All of these receptors modulate the release of neurotransmitter in TRN. Although not yet demonstrated, all of the above-mentioned receptors could be tonically active. Whether TRN contains tonically active NMDARs on glutamatergic axons/terminals was not tested in the present study and remains to be determined.

Like the types of tonically active receptors that are present throughout the CNS (Mody & Pearce, 2004; Semyanov *et al.*, 2004; Cavelier *et al.*, 2005; Farrant & Nusser, 2005; Glykys & Mody, 2007; Corlew *et al.*, 2008; Featherstone & Shippy, 2008; Belelli *et al.*, 2009; Brickley & Mody, 2012), those in TRN are presumably located at perisynaptic or extrasynaptic sites. Furthermore, tonically active NMDARs and mGlu2Rs on terminals in TRN are presumably activated by relatively low concentrations of ambient glutamate in the extracellular space. Possible sources of this endogenous agonist, with estimated basal ambient concentrations in the low μM range (Cavelier *et al.*, 2005; Featherstone & Shippy, 2008), include activity-dependent homo- and heterosynaptic release and ‘spillover’ of neurotransmitter, activity-independent synaptic release of neurotransmitter, observed as mEPSCs, exocytosis of neurotransmitter from axons, dendrites, and glial cells, and the cystine-glutamate transmembrane exchange system. Whatever the sources, the basal ambient concentration of glutamate, and hence the strength of tonically active NMDARs and mGlu2Rs, is most likely under the control of glutamate transport systems (Danbolt, 2001; Cavelier *et al.*, 2005; Featherstone & Shippy, 2008). High frequency synaptic activity could temporarily increase the strength of these tonically active receptors by momentarily increasing the ambient concentration of glutamate.

Functional implications of tonically active receptors in TRN

Our findings suggest that the three types of tonically active receptors we have identified in TRN will act together harmoniously to hyperpolarize the membrane potential of TRN neurons. Thus, the Cl^- current continually generated by constitutively active postsynaptic GABA_ARs in TRN neurons will directly hyperpolarize the membrane potential of these cells. Furthermore, the persistent increase in frequency of mIPSCs generated by tonically active presynaptic NMDARs will increase the inhibitory tone impinging on TRN neurons by increasing the amount of GABA that is available to bind at postsynaptic GABA_ARs and GABA_BRs. In addition, the persistent decrease in frequency of mEPSCs generated by tonically active presynaptic mGlu2Rs will decrease the excitatory tone impinging on TRN neurons by decreasing the amount of glutamate that is available to bind at postsynaptic AMPARs and NMDARs. However, because TRN neurons exhibit tonic and burst firing modes (Steriade *et al.*, 1986; Avanzini *et al.*, 1989; Hartings *et al.*, 2003), which occur at relatively depolarized or hyperpolarized membrane potentials, respectively, the hyperpolarizing effects of the tonically active receptors will differentially affect the action potential output of these cells depending on the moment-to-moment state of their membrane

potential. Thus, when TRN neurons are depolarized, the tonically active receptors will act to reduce firing in single-spike tonic mode. Indeed, we show that blockade of the persistent Cl^- current, which accounts for more than 90% of the intra-TRN inhibition, increases the action potential output of TRN neurons (cf. Brickley *et al.*, 1996; Hamann *et al.*, 2002; Semyanov *et al.*, 2003; Cope *et al.*, 2005). Furthermore, when TRN neurons are hyperpolarized, the tonically active receptors will act to increase the tendency to fire in multispikes burst mode through depolarizing intra-TRN GABAergic inputs and activation of T-type Ca^{2+} channels (Sun *et al.*, 2012). Therefore, the presence of tonically active GABA_ARs, NMDARs, and mGlu2Rs in TRN will affect its constituent neurons by decreasing their inhibitory output when they are relatively depolarized or increasing their inhibitory output when they are relatively hyperpolarized.

Acknowledgments

We thank Bill Anderson for supplying the data-acquisition software and Eli Lilly, Co. for generously supplying the LY341495 and LY395756. The support of The Wellcome Trust (J.W.C, Z.I.B, and J.T.R.I.) and NINDS (J.T.R.I.) is gratefully acknowledged. John Isaac is a full-time employee of Eli Lilly and Company.

Abbreviations

AMPA	α -amino-3-hydroxy-5-methylisoxazole-4-propionic acid receptor
CT	corticothalamic
EPSP	excitatory postsynaptic potential
GABA_AR	GABA _A receptor
GlyR	glycine receptor
IPSC	inhibitory postsynaptic current
NMDAR	N-methyl-D-aspartate receptor
mEPSC	miniature excitatory postsynaptic current
mGlu2R	metabotropic glutamate type 2 receptor
mIPSC	miniature inhibitory postsynaptic current
TC	thalamocortical
TRN	thalamic reticular nucleus

References

- Acuna-Goycolea C, Li Y, van den Pol AN. Group III metabotropic glutamate receptors maintain tonic inhibition of excitatory synaptic input to hypocretin/orexin neurons. *J. Neurosci.* 2004; 24:3013–3022. [PubMed: 15044540]
- Ade KK, Janssen MJ, Ortinski PI, Vicini S. Differential tonic GABA conductances in striatal medium spiny neurons. *J. Neurosci.* 2008; 28:1185–1197. [PubMed: 18234896]
- Alexander GM, Godwin DW. unique presynaptic and postsynaptic roles of group II metabotropic glutamate receptors in the modulation of thalamic network activity. *Neuroscience.* 2006; 141:501–513. [PubMed: 16690217]

- Anderson WW, Collingridge GL. Capabilities of the WinLTP data acquisition program extending beyond basic LTP experimental functions. *J. Neurosci. Methods.* 2007; 162:346–356. [PubMed: 17306885]
- Avanzini G, de Curtis M, Panzica F, Spreafico R. Intrinsic properties of nucleus reticularis thalami neurones of the rat studied *in vitro*. *J. Physiol.* 1989; 416:111–122. [PubMed: 2558172]
- Bai D, Zhu G, Pennefather P, Jackson MF, MacDonald JF, Orser BA. Distinct functional and pharmacological properties of tonic and quantal inhibitory postsynaptic currents mediated by γ -aminobutyric acid_A receptors in hippocampal neurons. *Mol. Pharmacol.* 2001; 59:814–824. [PubMed: 11259626]
- Bartlett TE, Bannister NJ, Collett VJ, Dargan SL, Massey PV, Bortolotto ZA, Fitzjohn SM, Bashir ZI, Collingridge GL, Lodge D. Differential roles of NR2A and NR2B-containing NMDA receptors in LTP and LTD in the CA1 region of two-week old rat hippocampus. *Neuropharmacology.* 2007; 52:60–70. [PubMed: 16904707]
- Belelli D, Harrison NL, Maguire J, Macdonald RL, Walker MC, Cope DW. Extrasynaptic GABA_A receptors: Form, pharmacology, and function. *J. Neurosci.* 2009; 29:12757–12763. [PubMed: 19828786]
- Belelli D, Peden DR, Rosahl TW, Wafford KA, Lambert JJ. Extrasynaptic GABA_A receptors of thalamocortical neurons: A molecular target for hypnotics. *J. Neurosci.* 2005; 25:11513–11520. [PubMed: 16354909]
- Berretta N, Jones RSG. Tonic facilitation of glutamate release by presynaptic N-methyl-D-aspartate autoreceptors in the entorhinal cortex. *Neuroscience.* 1996; 75:339–344. [PubMed: 8931000]
- Bouairi E, Kamendi H, Wang X, Gorini C, Mendelowitz D. Multiple types of GABA_A receptors mediate inhibition in brain stem parasympathetic cardiac neurons in the nucleus ambiguus. *J. Neurophysiol.* 2006; 96:3266–3272. [PubMed: 16914614]
- Boudaba C, Linn DM, Halmos K.Cs, Tasker JG. Increased tonic activation of presynaptic metabotropic glutamate receptors in the rat supraoptic nucleus following chronic dehydration. *J. Physiol.* 2003; 551:815–823. [PubMed: 12824445]
- Bourassa J, Pinault D, Deschênes M. Corticothalamic projections from the cortical barrel field to the somatosensory thalamus in rats: a single-fibre study using biocytin as an anterograde tracer. *Eur. J. Neurosci.* 1995; 7:19–30. [PubMed: 7711933]
- Brasier DJ, Feldman DE. Synapse-specific expression of functional presynaptic NMDA receptors in rat somatosensory cortex. *J. Neurosci.* 2008; 28:2199–2211. [PubMed: 18305253]
- Breukel AIM, Besselsen E, Lopes da Silva FH, Ghijsen WEJM. A presynaptic N-methyl-D-aspartate autoreceptor in rat hippocampus modulating amino acid release from a cytoplasmic pool. *Eur. J. Neurosci.* 1998; 10:106–114. [PubMed: 9753118]
- Brickley SG, Cull-Candy SG, Farrant M. Development of a tonic form of synaptic inhibition in rat cerebellar granule cells resulting from persistent activation of GABA_A receptors. *J. Physiol.* 1996; 497:753–759. [PubMed: 9003560]
- Brickley SG, Mody I. Extrasynaptic GABA_A receptors: Their function in the CNS and implications for disease. *Neuron.* 2012; 73:23–34. [PubMed: 22243744]
- Bright DP, Aller MI, Brickley SG. Synaptic release generates a tonic GABA_A receptor-mediated conductance that modulates burst precision in thalamic relay neurons. *J. Neurosci.* 2007; 27:2560–2569. [PubMed: 17344393]
- Bright DP, Brickley SG. Acting locally but sensing globally: impact of GABAergic synaptic plasticity on phasic and tonic inhibition in the thalamus. *J. Physiol.* 2008; 586:5091–5099. [PubMed: 18772202]
- Cavelier P, Hamann M, Rossi D, Mobbs P, Attwell D. Tonic excitation and inhibition of neurons: ambient transmitter sources and computational consequences. *Prog. Biophys. Mol. Biol.* 2005; 87:3–16. [PubMed: 15471587]
- Ceolín L, Kantamneni S, Barker GRI, Hanna L, Murray L, Warburton EC, Robinson ESJ, Monn JA, Fitzjohn SM, Collingridge GL, Bortolotto ZA, Lodge D. Study of novel selective mGlu2 agonist in the temporo-ammonic input to CA1 neurons reveals reduced mGlu2 receptor expression in a Wistar substrain with an anxiety-like phenotype. *J. Neurosci.* 2011; 31:6721–6731. [PubMed: 21543601]

- Chattipakorn SC, McMahon LL. Pharmacological characterization of glycine-gated chloride currents recorded in rat hippocampal slices. *J. Neurophysiol.* 2002; 87:1515–1525. [PubMed: 11877523]
- Cope DW, Hughes SW, Crunelli V. GABA_A receptor-mediated tonic inhibition in thalamic neurons. *J. Neurosci.* 2005; 25:11553–11563. [PubMed: 16354913]
- Corlew R, Brasier DJ, Feldman DE, Philpot BD. Presynaptic NMDA receptors: Newly appreciated roles in cortical synaptic function and plasticity. *Neuroscientist.* 2008; 14:609–625. [PubMed: 19029059]
- Corlew R, Wang Y, Ghermazien H, Erisir A, Philpot BD. Developmental switch in the contribution of presynaptic and postsynaptic NMDA receptors to long-term depression. *J. Neurosci.* 2007; 27:9835–9845. [PubMed: 17855598]
- Cotillon-Williams N, Huetz C, Hennevin E, Edeline J-M. Tonotopic control of auditory thalamus frequency tuning by reticular thalamic neurons. *J. Neurophysiol.* 2008; 99:1137–1151. [PubMed: 18160422]
- Cox CL, Huguenard JR, Prince DA. Nucleus reticularis neurons mediate diverse inhibitory effects in thalamus. *Proc. Natl. Acad. Sci. U. S. A.* 1997; 94:8854–8859. [PubMed: 9238067]
- Cox CL, Sherman SM. Glutamate inhibits thalamic reticular neurons. *J. Neurosci.* 1999; 19:6694–6699. [PubMed: 10414998]
- Crabtree JW, Isaac JTR. New intrathalamic pathways allowing modality-related and cross-modality switching in the dorsal thalamus. *J. Neurosci.* 2002; 22:8754–8761. [PubMed: 12351751]
- Danbolt NC. Glutamate uptake. *Prog. Neurobiol.* 2001; 65:1–105. [PubMed: 11369436]
- Deleuze C, Huguenard JR. Distinct electrical and chemical connectivity maps in the thalamic reticular nucleus: Potential roles in synchronization and sensation. *J. Neurosci.* 2006; 26:8633–8645. [PubMed: 16914689]
- Dominguez C, Prieto L, Valli MJ, Massey SM, Bures M, Wright RA, Johnson BG, Andis SL, Kingston A, Schoepp DD, Monn JA. Methyl substitution of 2-aminobicyclo[3.1.0]hexane 2,6-dicarboxylate (LY354740) determines functional activity at metabotropic glutamate receptors: identification of a subtype selective mGlu2 receptor agonist. *J. Med. Chem.* 2005; 48:3605–3612. [PubMed: 15887967]
- Eyre MD, Renzi M, Farrant M, Nusser Z. Setting the time course of inhibitory synaptic currents by mixing multiple GABA_A receptor α subunit isoforms. *J. Neurosci.* 2012; 32:5853–5867. [PubMed: 22539847]
- Farrant M, Nusser Z. Variations on an inhibitory theme: Phasic and tonic activation of GABA_A receptors. *Nat. Rev. Neurosci.* 2005; 6:215–229. [PubMed: 15738957]
- Featherstone DE, Shippy SA. Regulation of synaptic transmission by ambient extracellular glutamate. *Neuroscientist.* 2008; 14:171–181. [PubMed: 17947494]
- Glitsch M, Marty A. Presynaptic effects of NMDA in cerebellar Purkinje cells and interneurons. *J. Neurosci.* 1999; 19:511–519. [PubMed: 9880571]
- Glykys J, Mody I. Activation of GABA_A receptors: Views from outside the synaptic cleft. *Neuron.* 2007; 56:763–770. [PubMed: 18054854]
- Hamann M, Rossi DJ, Attwell D. Tonic and spillover inhibition of granule cells control information flow through cerebellar cortex. *Neuron.* 2002; 33:625–633. [PubMed: 11856535]
- Hartings JA, Simons DJ. Inhibition suppresses transmission of tonic vibrissae-evoked activity in the rat ventrobasal thalamus. *J. Neurosci.* 2000; 20 RC100:1–5. [PubMed: 10627575]
- Hartings JA, Temereanca A, Simons DJ. State-dependent processing of sensory stimuli by thalamic reticular neurons. *J. Neurosci.* 2003; 23:5264–5271. [PubMed: 12832551]
- Huntsman MM, Huguenard JR. Fast IPSCs in rat thalamic reticular nucleus require the GABA_A receptor β 1 subunit. *J. Physiol.* 2006; 572:459–475. [PubMed: 16469775]
- Jia F, Pignataro L, Schofield CM, Yue M, Harrison NL, Goldstein PA. An extrasynaptic GABA_A receptor mediates tonic inhibition in thalamic VB neurons. *J. Neurophysiol.* 2005; 94:4491–4501. [PubMed: 16162835]
- Jones EG. Some aspects of the organization of the thalamic reticular complex. *J. Comp. Neurol.* 1975; 162:285–308. [PubMed: 1150923]

- Kaneda M, Farrant M, Cull-Candy SG. Whole-cell and single-channel currents activated by GABA and glycine in granule cells of the rat cerebellum. *J. Physiol.* 1995; 485:419–435. [PubMed: 7545231]
- Kim U, McCormick DA. The functional influence of burst and tonic firing mode on synaptic interactions in the thalamus. *J. Neurosci.* 1998; 18:9500–9516. [PubMed: 9801387]
- Kingston AE, Ornstein PL, Wright RA, Johnson BG, Mayne NG, Burnett JP, Belagaje R, Wu S, Schoepp DD. LY341495 is a nanomolar potent and selective antagonist of group II metabotropic glutamate receptors. *Neuropharmacology.* 1998; 37:1–12. [PubMed: 9680254]
- Krishek BJ, Moss SJ, Smart TG. Homomeric $\beta 1$ γ -aminobutyric acid_A receptor-ion channels: evaluation of pharmacological and physiological properties. *Mol. Pharmacol.* 1996; 49:494–504. [PubMed: 8643089]
- Krook-Magnuson EI, Li P, Paluszkiwicz SM, Huntsman MM. Tonically active inhibition selectively controls feedforward circuits in mouse barrel cortex. *J. Neurophysiol.* 2008; 100:932–944. [PubMed: 18509076]
- Lam Y-W, Nelson CS, Sherman SM. Mapping of the functional interconnections between thalamic reticular neurons using photostimulation. *J. Neurophysiol.* 2006; 96:2593–2600. [PubMed: 16855107]
- Landisman CE, Long MA, Beierlein M, Deans MR, Paul DL, Connors BW. Electrical synapses in the thalamic reticular nucleus. *J. Neurosci.* 2002; 22:1002–1009. [PubMed: 11826128]
- Larsen RS, Corlew RJ, Henson MA, Roberts AC, Mishina M, Watanabe M, Lipton SA, Nakanishi N, Pérez-Otaño I, Weinberg RJ, Philpot BD. NR3A-containing NMDARs promote neurotransmitter release and spike timing-dependent plasticity. *Nat. Neurosci.* 2011; 14:338–344. [PubMed: 21297630]
- Lauri SE, Vesikansa A, Segerstråle M, Collingridge GL, Isaac JTR, Taira T. Functional maturation of CA1 synapses involves activity-dependent loss of tonic kainate receptor-mediated inhibition of glutamate release. *Neuron.* 2006; 50:415–429. [PubMed: 16675396]
- Lee KH, McCormick DA. Modulation of spindle oscillations by acetylcholine, cholecystokinin and 1S, 3R-ACPD in the ferret lateral geniculate and perigeniculate nuclei in vitro. *Neuroscience.* 1997; 77:335–350. [PubMed: 9472394]
- Lee SM, Friedberg MH, Ebner FF. The role of GABA-mediated inhibition in the rat ventral posterior medial thalamus. I. Assessment of receptive field changes following thalamic reticular nucleus lesions. *J. Neurophysiol.* 1994a; 71:1702–1715. [PubMed: 8064343]
- Lee SM, Friedberg MH, Ebner FF. The role of GABA-mediated inhibition in the rat ventral posterior medial thalamus. II. Differential effects of GABA_A and GABA_B receptor antagonists on responses of VPM neurons. *J. Neurophysiol.* 1994b; 71:1716–1726. [PubMed: 8064344]
- Long MA, Landisman CE, Connors BW. Small clusters of electrically coupled neurons generate synchronous rhythms in the thalamic reticular nucleus. *J. Neurosci.* 2004; 24:341–349. [PubMed: 14724232]
- Losonczy A, Somogyi P, Nusser Z. Reduction of excitatory postsynaptic responses by persistently active metabotropic glutamate receptors in the hippocampus. *J. Neurophysiol.* 2003; 89:1910–1919. [PubMed: 12686572]
- Malosio M-L, Marquère-Pouey B, Kuhse J, Betz H. Widespread expression of glycine receptor subunit mRNAs in the adult and developing rat brain. *EMBO J.* 1991; 10:2401–2409. [PubMed: 1651228]
- Marowsky A, Rudolph U, Fritschy J-M, Arand M. Tonic inhibition in principal cells of the amygdala: A central role for $\alpha 3$ subunit-containing GABA_A receptors. *J. Neurosci.* 2012; 32:8611–8619. [PubMed: 22723702]
- McCartney MR, Deeb TZ, Henderson TN, Hales TG. Tonically active GABA_A receptors in hippocampal pyramidal neurons exhibit constitutive GABA-independent gating. *Mol. Pharmacol.* 2007; 71:539–548. [PubMed: 17090706]
- Mistry RB, Isaac JTR, Crabtree JW. Two differential frequency-dependent mechanisms regulating tonic firing of thalamic reticular neurons. *Eur. J. Neurosci.* 2008; 27:2643–2656. [PubMed: 18547248]

- Miyata M, Imoto K. Contrary roles of kainate receptors in transmitter release at corticothalamic synapses onto thalamic relay and reticular neurons. *J. Physiol.* 2009; 587:999–1012. [PubMed: 19124541]
- Mody I, Pearce RA. Diversity of inhibitory neurotransmission through GABA_A receptors. *TINS.* 2004; 27:569–575. [PubMed: 15331240]
- Nusser Z, Mody I. Selective modulation of tonic and phasic inhibition in dentate gyrus granule cells. *J. Neurophysiol.* 2002; 87:2624–2628. [PubMed: 11976398]
- Ohishi H, Shigemoto R, Nakanishi S, Mizuno N. Distribution of the messenger RNA for a metabotropic glutamate receptor, mGluR2, in the central nervous system of the rat. *Neuroscience.* 1993a; 53:1009–1018. [PubMed: 8389425]
- Ohishi H, Shigemoto R, Nakanishi S, Mizuno N. Distribution of the mRNA for a metabotropic glutamate receptor (mGluR3) in the rat brain: An in situ hybridization study. *J. Comp. Neurol.* 1993b; 335:252–266. [PubMed: 8227517]
- Paoletti P, Neyton J. NMDA receptor subunits: function and pharmacology. *Curr. Opin. Pharmacol.* 2007; 7:39–47. [PubMed: 17088105]
- Pirker S, Schwarzer C, Wieselthaler A, Sieghart W, Sperk G. GABA_A receptors: Immunocytochemical distribution of 13 subunits in the adult rat brain. *Neuroscience.* 2000; 101:815–850. [PubMed: 11113332]
- Porcello DM, Huntsman MM, Mihalek RM, Homanics GE, Huguenard JR. Intact synaptic GABAergic inhibition and altered neurosteroid modulation of thalamic relay neurons in mice lacking δ subunit. *J. Neurophysiol.* 2003; 89:1378–1386. [PubMed: 12626617]
- Salin PA, Prince DA. Spontaneous GABA_A receptor-mediated inhibitory currents in adult rat somatosensory cortex. *J. Neurophysiol.* 1996; 75:1573–1588. [PubMed: 8727397]
- Salt TE. Gamma-aminobutyric acid and afferent inhibition in the cat and rat ventrobasal thalamus. *Neuroscience.* 1989; 28:17–26. [PubMed: 2668781]
- Sanchez-Vives MV, Bal T, McCormick DA. Inhibitory interactions between perigeniculate GABAergic neurons. *J. Neurosci.* 1997; 17:8894–8908. [PubMed: 9348356]
- Saugstad JA, Kinzie JM, Shinohara MM, Segerson TP, Westbrook GL. Cloning and expression of rat metabotropic glutamate receptor 8 reveals a distinct pharmacological profile. *Mol. Pharmacol.* 1997; 51:119–125. [PubMed: 9016353]
- Semyanov A, Walker MC, Kullmann DM. GABA uptake regulates cortical excitability via cell type-specific tonic inhibition. *Nat. Neurosci.* 2003; 6:484–490. [PubMed: 12679782]
- Semyanov A, Walker MC, Kullmann DM, Silver RA. Tonically active GABA_A receptors: modulating gain and maintaining the tone. *TINS.* 2004; 27:262–269. [PubMed: 15111008]
- Sherman, SM.; Guillery, RW. Exploring the thalamus. Academic; San Diego, CA: 2001.
- Sieghart W, Fuchs K, Tretter V, Ebert V, Jechlinger M, Höger H, Adamiker D. Structure and subunit composition of GABA_A receptors. *Neurochem. Int.* 1999; 34:379–385. [PubMed: 10397365]
- Sigel E, Baur R, Malherbe P, Möhler H. The rat β 1-subunit of the GABA_A receptor forms a picrotoxin-sensitive anion channel open in the absence of GABA. *FEBS Lett.* 1989; 257:377–379. [PubMed: 2479580]
- Snead OC, Banerjee PK, Burnham M, Hampson D. Modulation of absence seizures by the GABA_A receptor: A critical role for metabotropic glutamate receptor 4 (mGluR4). *J. Neurosci.* 2000; 20:6218–6224. [PubMed: 10934271]
- Stell BM, Mody I. Receptors with different affinities mediate phasic and tonic GABA_A conductances in hippocampal neurons. *J. Neurosci.* 2002; 22 RC223:1–5. [PubMed: 11756482]
- Steriade M, Domich L, Oakson G. Reticularis thalami neurons revisited: activity changes during shifts in states of vigilance. *J. Neurosci.* 1986; 6:68–81. [PubMed: 3944624]
- Sun Y-G, Wu C-S, Lu H-C, Beierlein M. Target-dependent control of synaptic inhibition by endocannabinoids in the thalamus. *J. Neurosci.* 2011; 31:9222–9230. [PubMed: 21697372]
- Sun Y-G, Wu C-S, Renger JJ, Uebele VN, Lu H-C, Beierlein M. GABAergic synaptic transmission triggers action potentials in thalamic reticular nucleus neurons. *J. Neurosci.* 2012; 32:7782–7790. [PubMed: 22674255]

- Ulrich D, Huguenard JR. Purinergic inhibition of GABA and glutamate release in the thalamus: Implications for thalamic network activity. *Neuron*. 1995; 15:909–918. [PubMed: 7576639]
- Ulrich D, Huguenard JR. GABA_B receptor-mediated responses in GABAergic projection neurones of rat nucleus reticularis thalami *in vitro*. *J. Physiol.* 1996; 493:845–856. [PubMed: 8799904]
- Wall MJ, Usowicz MM. Development of action potential-dependent and independent spontaneous GABA_A receptor-mediated currents in granule cells of postnatal rat cerebellum. *Eur. J. Neurosci.* 1997; 9:533–548. [PubMed: 9104595]
- Wang L, Spary E, Deuchars J, Deuchars SA. Tonic GABAergic inhibition of sympathetic preganglionic neurons: A novel substrate for sympathetic control. *J. Neurosci.* 2008; 28:12445–12452. [PubMed: 19020037]
- Warren RA, Jones EG. Glutamate activation of cat thalamic reticular nucleus: effects on response properties of ventroposterior neurons. *Exp. Brain Res.* 1994; 100:215–226. [PubMed: 7813660]
- Woodhall G, Evans DI, Cunningham MO, Jones RSG. NR2B-containing NMDA autoreceptors at synapses on entorhinal cortical neurons. *J. Neurophysiol.* 2001; 86:1644–1651. [PubMed: 11600627]

Fig. 1.

A tonic GABA_AR-mediated conductance in TRN neurons. (A) Holding current (DC) versus time recorded from a representative TRN neuron at a holding potential of 0 mV. Bath applied gabazine (50 μ M; black bar) abolished outward spontaneous IPSCs but did not affect the holding current. The individual traces at the top were taken at the times indicated (1, 2). (B) Holding current (DC) versus time recorded from a representative TRN neuron at a holding potential of 0 mV. Bath applied picrotoxin (50 μ M; black bar) abolished outward spontaneous IPSCs and reduced an outward holding current indicating that a tonic

GABA_AR-mediated conductance was present in this cell. The individual traces at the top were taken at the times indicated (1, 2). The scale bars for these traces also apply to those in A. (C) Summary graph of the effect of picrotoxin (50 μ M; black bar) on DC over time in 15 TRN neurons. Data are expressed as changes from the average baseline response before picrotoxin application. Compared to this baseline, the shift in DC averaged over the last five minutes was significant ($p < 0.05$; Wilcoxon matched-pair test). The dashed line indicates the baseline response level. (D) In an example TRN neuron, depolarizing current pulses (200-ms steps) produced firing of action potentials (spikes) under control conditions (upper traces) and in the presence of picrotoxin (50 μ M; lower traces). The membrane potential of this cell is shown to the left of each row of traces. The scale bar for the trace at the upper left applies to all the traces. For the example TRN neuron, the number of spikes versus injected current (lower left plot). Each point represents the number of spikes over a 200-ms period averaged over 12 trials. Pooled data from 10 TRN neurons for the number of spikes versus injected current (lower right plot). Compared to control, the number of spikes significantly increased at all levels of injected current, except the lowest level, after wash-in of picrotoxin ($*p < 0.05$; Wilcoxon matched-pair tests).

Fig. 2.

A tonic presynaptic NMDAR facilitates release of GABA from terminals in TRN. (A) Pooled data from 6 TRN neurons before (solid lines) and after (dashed red lines) wash-in of NMDA (40 μ M). Data were obtained by collecting 200 mIPSCs under each condition from each of the cells. Comparing before and after NMDA wash-in, the mIPSC amplitudes did not significantly differ ($p > 0.05$; Kolmogorov-Smirnov test) but the inter-mIPSC interval was significantly shifted to the left ($p < 0.05$; Kolmogorov-Smirnov test). The individual traces at the top were taken under the conditions indicated. The insert shows over 200-ms

periods an evoked NMDAR-mediated EPSC recorded from a TRN neuron (top trace) and the abolition of this EPSC (bottom trace) after the cell was dialyzed with MK-801. (B) Pooled data from 8 TRN neurons before (solid lines) and after (dashed red lines) wash-in of D-AP5 (50 μ M). Data were obtained by collecting 200 mIPSCs under each condition from each of the cells. Comparing before and after D-AP5 wash-in, the mIPSC amplitudes did not significantly differ ($p > 0.05$; Kolmogorov-Smirnov test) but the inter-mIPSC interval was significantly shifted to the right ($p < 0.05$; Kolmogorov-Smirnov test). The individual traces at the top were taken under the conditions indicated. The scale bars for the trace on the right applies to all the traces in A-C. The insert shows over 5-s periods mIPSCs recorded from a TRN neuron (top trace) and the abolition of mIPSCs (bottom trace) after bath application of picrotoxin (50 μ M). (C) Pooled data from 6 TRN neurons before (solid lines) and after (dashed red lines) wash-in of Ro 25-6981 (3 μ M). Data were obtained by collecting 200 mIPSCs under each condition from each of the cells. Comparing before and after RO 25-6981 wash-in, the mIPSC amplitudes did not significantly differ ($p > 0.05$; Kolmogorov-Smirnov test) but the inter-mIPSC interval was significantly shifted to the right ($p < 0.05$; Kolmogorov-Smirnov test). The individual traces at the top were taken under the conditions indicated. (D) Pooled data from the 6 neurons shown in A, the 8 neurons shown in B, and the 6 neurons shown in C before (Baseline) and after wash-in of NMDA (40 μ M), D-AP5 (50 μ M), or Ro 25-6981 (3 μ M). The bars represent the amplitude or frequency of mIPSCs over a 5-s period averaged over 18 trials for each cell under the various conditions and are expressed as percentages of the average baseline response before NMDA, D-AP5, or Ro 25-6981 application. Compared to baseline, the mIPSC amplitudes did not significantly differ after the various agonist or antagonist wash-ins ($p > 0.05$; Wilcoxon matched-pair tests) but the mIPSC frequency significantly increased after NMDA wash-in ($*p < 0.05$; Wilcoxon matched-pair test) or decreased after D-AP5 or Ro 25-6981 wash-in ($*p < 0.05$; Wilcoxon matched-pair tests).

Fig. 3.

Change in mean inter-mIPSC interval for each neuron. Bars represent the population mean. (A) Data from 6 TRN neurons before (Baseline) and after wash-in of NMDA (40 μ M). (B) Data from 8 TRN neurons before (Baseline) and after wash-in of D-AP5 (50 μ M). (C) Data from 6 TRN neurons before (Baseline) and after wash-in of Ro 25-6981 (3 μ M).

Fig. 4.

A tonic presynaptic mGlu2R reduces release of glutamate from terminals in TRN. (A) mEPSC frequency versus time recorded from a representative TRN neuron at a holding potential of -60 mV. Each point represents the number of mEPSCs over a 5-s period averaged over 6 trials. Bath applied LY395756 (3 μ M; black bar) reversibly reduced mEPSC frequency. The individual traces at the top were taken at the times indicated (1, 2, 3). (B) Summary graph of the effect of LY395756 (3 μ M; black bar) on mEPSC frequency over time in 3 TRN neurons. Data are expressed as percentages of the average baseline response

over the points at 2-10 min before LY395756 application. Compared to this baseline, the decrease in mEPSC frequency averaged over the points at 18-24 min was significant ($p < 0.05$; Wilcoxon matched-pair test). The dashed line indicates the baseline response level (100%). (C) mEPSC frequency versus time recorded from a representative TRN neuron at a holding potential of -60 mV. Each point represents the number of mEPSCs over a 5-s period averaged over 6 trials. Bath applied LY341495 (300 nM; black bar) reversibly increased mEPSC frequency. The individual traces at the top were taken at the times indicated (1, 2, 3). (D) Summary graph of the effect of LY341495 (300 nM; black bar) on mEPSC frequency over time in 8 TRN neurons. Data are expressed as percentages of the average baseline response over the points at 2-10 min before LY341495 application. Compared to this baseline, the increase in mEPSC frequency averaged over the points at 18-24 min was significant ($p < 0.05$; Wilcoxon matched-pair test). The dashed line indicates the baseline response level (100%). (E) Pooled data from the 8 TRN neurons shown in D before (solid lines) and after (dashed red lines) wash-in of LY341495 (300 nM). Data were obtained by collecting 200 mEPSCs each at the 2-10 min points and the 18-24 min points from each of the cells. Comparing before and after LY341495 wash-in, the mIPSC amplitudes did not significantly differ ($p > 0.05$; Kolmogorov-Smirnov test) but the inter-mIPSC interval was significantly shifted to the left ($p < 0.05$; Kolmogorov-Smirnov test). The insert shows over 5-s periods mEPSCs in a TRN neuron (top trace) and the abolition of mEPSCs (bottom trace) after bath application of NBQX ($2 \mu\text{M}$).

Fig.5.

Summary of postsynaptic and presynaptic receptors in TRN. The schematic shows a GABAergic neuron in TRN (R). For clarity, only a single unbranched dendrite is shown. The receptors are those that we took into account in this study. Receptors indicated by an asterisk, either on the dendrite, the GABAergic (GABA) axon terminal, or the glutamatergic (GLUT) axon terminal, summarize our findings regarding the presence of 3 types of tonically active receptors in TRN. Receptor and terminal locations are only approximate.