Published in final edited form as:

Cloning Transgenes. 2013 ; 2: . doi:10.4172/2168-9849.1000116.

Cryopreservation of transgenic Huntington's disease rhesus macaque sperm-A Case Report

Kittiphong Putkhao, Ph.D^{1,4}, Anthony W.S. Chan, DVM Ph.D², Yuksel Agca, DVM Ph.D⁴, and Rangsun Parnpai, Ph.D^{1,3}

¹Yerkes National Primate Research Center Emory University School of Medicine 954 Gatewood Rd., N.E. Atlanta, Georgia 30329, USA. ²Department of Veterinary Pathobiology, College of Veterinary Medicine, University of Missouri 1600 East Rollins Rd., Columbia, MO 65211 agca@missouri.edu Phone: 573-884-0311 Fax: 573-884-7521 ³Yerkes National Primate Research Center Emory University School of Medicine 954 Gatewood Rd., N.E. Atlanta, Georgia 30329, USA. ⁴Embryo Technology and Stem Cell Research Center, School of Biotechnology, Suranaree University of Technology, Nakhon Ratchasima 30000, Thailand

Abstract

The cryoprotective effects of glycerol in three different semen freezing extenders, Tris-citrate (TRIS), TEST, and Tes-Tris-Egg yolk (TTE), on wild-type (WT) rhesus monkey (*Macaca mulatta*) sperm cryopreservation have been tested. Sperm motility and viability were examined to evaluate the integrity of frozen-thawed sperm, and the best extender was selected to cryopreserve sperm from transgenic Huntington's disease (HD) rhesus monkey. The results showed no post-thaw motility difference among the freezing extender tested ($P>0.05$). However, sperm membrane integrity in TEST and TTE were significantly better than in TRIS extender ($P<0.05$). TEST was chosen for HD rhesus monkey sperm cryopreservation. The results showed that post-thawed HD sperm motility and viability was not different compared with WT control group ($P>0.05$). The present study demonstrates that TEST and TTE were excellent extenders and suitable for rhesus monkey sperm cryopreservation and no detectable differences of post-thaw sperm motility and viability between HD and WT rhesus monkeys resulted from TEST extender.

Keywords

Transgenic monkey model; Huntington's disease; Sperm cryopreservation; Freezing extender

Introduction

Sperm cryopreservation technology is one of the most important assisted reproductive techniques (ART) that is used for genome banking of biomedically important laboratory animals, farm animals and human infertility programs.¹ Since, the discovery of glycerol as a

Correspondence should be addressed to: Anthony W.S. Chan DVM Ph.D awchan@emory.edu Phone: 404-712-8347 Fax: 404-727-5289.
putkhao_kp@hotmail.com Phone: 66-44-224-244 Fax: 66-44-224-150 rangsun@g.sut.ac.th Phone: 66-44-224-244 Fax: 66-44-224-150

cryoprotectant agent (CPA) for sperm cryopreservation,³⁹ glycerol have been the major CPA for sperm cryopreservation in many species including; nonhuman primate (NHP) such as lemurs,⁵ marmosets,^{18,33,34} squirrel monkeys,⁹ African greens,⁴⁴ vervet monkeys,^{6,48} cynomolgus monkeys,^{3, 13, 28, 29, 30, 31, 35, 47, 54} Japanese monkeys,⁴⁶ stump tail monkeys,⁴⁴ baboons,²² chimpanzees,^{15, 24, 44, 57} and gorillas.^{25, 26, 40} Among NHP, rhesus monkeys are the commonly used research model in the development of sperm cryopreservation due to their high physiologic and genetic similarity with humans.^{7, 14, 17, 19, 23, 36, 37, 41, 52, 53, 55} There have been extensive studies on rhesus monkey sperm cryopreservation^{10, 11, 27, 44, 45, 49, 51}. However, sperm cryopreservation efficacy varies tremendously among laboratories, because they often use different freezing methods, extenders types, and species. Thus, it is difficult to compare results among institutions, and there is still the need for an optimal protocol for sperm cryopreservation.

Huntington's disease (HD) is an inherited neurodegenerative disorder resulting from an expanded polyglutamine (CAG) tract in the Exon 1 of the Huntington gene IT15 (*HTT*)¹⁶ that leads to progressive impairment on motor, cognitive and psychiatric functions as an individual ages.^{8, 18, 38, 42, 43} The size of the polyglutamine (CAG) expansion is significantly influenced by gender, as paternal instability has shown to be greater than maternal inheritance in mouse HD model and human patients.^{4, 21} While the mechanism of paternal instability in HD remains largely unknown, HD monkey sperm provide a unique resource for investigating CAG stability and paternal inheritance. Besides aiding the understanding of HD pathogenesis and its correlation with CAG instability, cryopreservation of HD monkey sperm preserves the unique genetics necessary for future propagation of next generation HD monkeys since fertility in HD monkey declines as disease progresses. Thus, the goal of the present study was to develop and optimize a cryopreservation protocol of HD monkey sperm which will be used for future banking of HD monkey sperm.

Materials and Methods

A total of five ejaculates were collected from a single male of wild-type (WT) rhesus monkey and used to study the freezing efficiency of three freezing extenders (TRIS, TEST, and TTE). A total of four ejaculates were collected from a single male HD monkey and compared with a WT control. The extender that offered satisfactory results would be selected for HD monkey sperm cryopreservation.

Animal model

The rhesus monkeys (WT and HD) were housed and maintained at Yerkes National Primate Research Center. All animal procedures in this study were approved by the Institutional Animal Care and Use Committee (IACUC) at Emory University. All documented research in this study was conducted in accordance with the guide for *The Care and Use of Laboratory Animals*.

Generation of transgenic HD rhesus monkey

A transgenic HD monkey (rHD1) was generated by Yang and colleagues (2008). In brief, high titer lentiviruses carried exon1 of human *HTT* with 84 CAG repeats and were co-

injected with lentiviruses carried green fluorescent protein (GFP) gene under the regulation of human polyubiquitin C promoter into the perivitelline space of metaphase II arrested monkey oocytes followed by intracytoplasmic sperm injection (ICSI). The resulting embryos were transferred into surrogate females for the generation of transgenic HD monkeys.⁵⁶

Sperm cryopreservation extender

Each cryopreservation extender (TRIS, TEST, and TTE) was divided into two parts (with and without glycerol). The composition of each extender is listed in Table 1. The primary semen diluting extender (without glycerol) was centrifuged at $7,000 \times g$ at 4°C for one hour to remove yolk granules.⁴⁷ Supernatant was filtered with $0.2 \mu\text{m}$ filter and stored at -80°C . The extender was thawed at 37°C water for 15 min and equilibrated at room temperature (RT) before it was used. The secondary extender is a mixture of the primary extender with 8%, 3%, and 5% glycerol in the TRIS, TEST, and TTE extenders, respectively).

Semen collection procedure

Ejaculates were collected once a week in the morning. The monkey was removed from the cage and situated in the primate chair where the “pole and collar” technique was used. The monkey was sedated with a light dose of Telazol ($0.7\text{--}1.0 \text{ mg/kg}$ body weight), and administered IM. One pre-sized defibrillator gel electrode was wrapped around the base of the penis and connected to the negative lead. The second gel electrode was positioned immediately behind the glans and connected to the positive lead. Then by slow and steady stimulation caused by increasing the output-adjust dial would produce a slight erection, engorgement of the glans, and/or elevation of the testicles into the inguinal region. Typically, engorgement and ejaculation occurred at 10-20 V. When the sample was delivered, the date, time, peak output current, length of time electrical current was delivered, drug administered, behavioral observations, and technician initials were all recorded. The ejaculates were kept at RT for 20 min to liquefy. The liquid portion of semen was transferred into 15 mL conical tube and washed with TALP-HEPES medium supplemented with 4 mg/mL bovine serum albumin (BSA),² and then centrifuged at $112 \times g$ at RT for 7 min. The sperm concentration, motility and viability were determined and recorded.

Sperm freezing procedure

Washed WT rhesus monkey sperm was divided into three groups for three different extenders. The sperm was diluted with primary extender in 15 mL conical tube and kept in 500 mL beaker containing RT water and kept in 4°C refrigerator for 2 h. Equal volume of pre-cooled secondary extender (with glycerol) was added and kept for 30 min at 4°C .⁴⁷ The final concentration of glycerol was 8%, 3% and 5% in TRIS, TEST and TTE, respectively. Final sperm concentration was $30 \times 10^6 \text{ cells mL}^{-1}$. Sperm was then loaded into 0.25 mL straws by using 1 mL syringe and sealed with plug powder. Straws were kept in 4°C water for 30 min. The straws were then laid horizontal on an aluminum rack and then transferred to a styrofoam box containing liquid nitrogen (LN_2) and held for 8 min, positioned 4 cm above the surface of LN_2 . LN_2 was increased by 1 cm every 2 min until the final distance between surface of LN_2 and straw was 1 cm. Straws were then plunged directly into LN_2 and stored for a week before thawing.

For transgenic HD rhesus monkey sperm cryopreservation, the extender that gave the best post-thawed sperm motility and viability in WT was selected for freezing. The cryopreservation process was performed following the protocol described above. Sperm samples were evaluated after a minimum of one-week storage in LN₂.

Sperm thawing

Straws containing frozen spermatozoa were thawed in a 37°C water bath for 1-2 min. Thawed spermatozoa was washed in 15 ml conical tube with 4 mL of TALP-HEPES medium supplemented with 4 mg/mL BSA² by centrifugation at $112 \times g$ under RT for 7 min. The sperm pellet was re-suspended by gently pipetting, and sperm motility and viability were recorded.

Sperm motility evaluation

10 µL thawed semen was placed onto a 37°C pre-warmed microscope slide and covered with a 22 mm square cover-glass. The slides were visualized with 10x positive phase objective on a light microscope. Then sperm motility was evaluated by three different individuals with two of them not knowing about the treatments. The sperm motility recovery was calculated by the following formula: (post-thaw motility % $\times$ 100)/ pre-freeze sperm motility %.

Sperm viability evaluation

The sperm viability was assessed by using a LIVE/DEAD® Sperm Viability Kit (Invitrogen). The pre-freeze or post-thaw sperm concentration was adjusted at 2.5×10^6 mL⁻¹. Sperm samples were incubated with 100 nM of SYBR 14 dye at 37°C water bath for 5-10 min in a dark room, and then they were incubated with 12 µM of Propidium Iodide (PI) for another 5 min at 37°C water bath in a dark room. After that, stained sperm were placed on a microscope slide coated with 0.1% poly-D-lysine (Sigma) and covered with a 22 mm square cover glass. The sperm showing a bright green fluorescence were considered to be alive, but if red fluorescence was viewed, they were rated as dead (Figure 1). Sperm viability was calculated by the following formula: (number of live sperm $\times$ 100)/ number of total sperm count. The sperm viability recovery was calculated by the following formula: (viability of post-thawed sperm % $\times$ 100)/pre-freeze sperm motility %.

Statistical analysis

Sperm motility and viability were analyzed between each treatment group using a general linear model. This test was used to analyze significant differences between freezing extenders in WT frozen sperm and between WT and HD frozen sperm. The analysis was performed using the Statistical Analysis System Software (SAS, version 9.0; SAS, Cary, NC, USA) and a value of $P < 0.05$ was considered as significantly different.

Results

Effect of freezing extender on WT rhesus monkey sperm cryopreservation

The post-thaw motility and motility recovery of WT rhesus monkey sperm are presented in Table 2. The post-thaw sperm motility and motility recovery in TEST group was slightly higher than other groups. However, there is no difference among the three extenders both in terms of motility ($54.78 \pm 3.88\%$, $50.16 \pm 7.52\%$, and $46.99 \pm 4.82\%$, in TEST, TTE, and TRIS respectively) and motility recovery ($60.69 \pm 5.58\%$, $53.99 \pm 6.70\%$, and $52.51 \pm 7.43\%$, in TEST, TTE, and TRIS respectively).

The viability and viability recovery of post-thawed WT rhesus monkey sperm are presented in Table 3. The post-thawed sperm viability in TEST group ($60.75 \pm 2.85\%$) was significant higher than TRIS group ($48.79 \pm 3.46\%$), but no different from TTE group ($54.85 \pm 3.34\%$), while TRIS group was no different from TTE group.

HD rhesus monkey sperm cryopreservation

The comparisons of three extenders using WT rhesus monkey sperm for cryopreservation revealed that TEST extender was slightly better than other extenders. Therefore, TEST was selected for cryopreservation of HD rhesus monkey sperm. The motility and motility recovery of post-thaw HD rhesus monkey sperm is presented in Table 4. The HD sperm motility ($53.67 \pm 8.96\%$) and motility recovery ($56.53 \pm 9.89\%$) showed no significant difference with control group ($58.06 \pm 2.68\%$, in terms of motility and $65.19\% \pm 4.29\%$, in motility recovery).

The sperm viability and viability recovery of HD rhesus monkey sperm is presented in Table 5. The post-thawed HD sperm viability was not different from the control group ($58.18 \pm 1.74\%$ and $61.74 \pm 3.46\%$, respectively). However, viability recovery of HD sperm was significantly lower than the control group ($62.97 \pm 1.57\%$ and $78.48 \pm 3.40\%$, respectively).

Discussion

The composition of the freezing extender plays a key role in sperm cryopreservation. Thus, this study investigated the effect of freezing extenders on rhesus monkey sperm cryopreservation by comparing three different freezing extenders. Due to the limited number of adult WT and HD monkey sperm donors, we were only able to collect sperm from one WT and one HD monkey in this study to develop a cryopreservation protocol that we could further evaluate when more sperm donors are available. The freezing extenders used in our experiments have been previously employed by other research groups in an effort to freeze NHP sperm. Previous studies obtained satisfactory results with TTE in cynomolgus⁴⁷ and rhesus monkey,⁵¹ TEST in rhesus monkey,^{10, 11} and TRIS in rhesus monkey (our laboratory, unpublished results). Post-thaw sperm motility was around 50% for the TEST and TTE in previous studies; however, differences existed among species and freezing methods used. The results of this study showed that three extenders (TRIS, TEST, and TTE) offered good cryoprotection during sperm freezing in rhesus monkey. Although, TEST showed slightly better than other extenders; no significant difference in sperm motility was found among extenders.

Many factors affect successful sperm cryopreservation. These include; freezing and thawing procedures, extenders, types of cryoprotectant, and skill of investigators. The cryoprotectant is one of the major factors that influence sperm survival during sperm cryopreservation. Glycerol is a commonly used permeating cryoprotectant during NHP sperm freezing; however, its toxicity can cause loss in sperm motility and fertility if their concentration is not optimal.²⁸ There are many reports suggesting that 3-5% glycerol offered suitable concentration for cynomolgus⁴⁷ and rhesus monkey^{10, 11, 28, 29, 50} sperm cryopreservation. In the present study, the results showed that 3% (TEST) and 5% (TTE) of glycerol concentration provided better cryoprotection than 8% (TRIS). This confirms that high concentrations of permeating cryoprotectant result in higher toxicity to the sperm cells.

In this study we also conducted transgenic HD rhesus monkey sperm cryopreservation by using TEST extender. The results show that post-thaw sperm motility and viability were not significantly different between HD and non-transgenic sperm. This indicated that mutant huntingtin does not have any adverse effect on the quality of sperm and HD monkey spermatozoa can endure stresses caused during freezing and thawing process.

It is known that, HD can be inherited by the next generation. The gender and age of HD parents influences the transmission of CAG repeats to next generation. In the paternal transmission there is a propensity toward larger CAG repeat expansion, but maternal transmission produces nearly equal numbers of CAG expansion in progeny.^{4, 12, 32} Although, CAG expansion tends to occur more often from the paternal rather than maternal parent; distribution of repeat length in the progeny did not faithfully reflect the distribution of repeats present in the sperm of the founding fathers.²¹ This implies that the molecular basis for the parent of origin effect associated with CAG repeat expansion has not been clarified, and there is a need for further understanding. Therefore, the sperm cryopreservation of HD phenotype in this study may be useful in future HD studies such as germ line transmission, CAG expansion in differing ages of paternal sperm and HD pathology; and it might collectively bring us closer to effective HD treatments.

In conclusion, the present study showed that freezing extender TEST and TTE were excellent extenders which are suitable for rhesus monkey sperm cryopreservation, and no detectable differences of post-thawed sperm motility and viability between HD and WT in TEST extender were found.

Acknowledgments

All protocols involving animal care and handling were approved by Emory University's IACUC. We thank Tim Chi and Adam Neumann for preparation of rhesus monkey semen. We thank Yerkes National Primate Research Center (YNPRC) veterinarian and animal care staff for providing outstanding services. The YNPRC is supported by the base grant P51RR165 awarded by the Animal Resources Program of the NIH. This work was supported by ORIP/NIH (RR018827), and in part by the Royal Golden Jubilee Ph.D program of Thailand Research Fund in cooperation with Suranaree University of Technology. AWSC is supported by the ORIP/NIH (RR018827), NINDS (NS064991) and the American Recovery and Reinvestment Act (ARRA) Fund.

Reference

1. Agca Y. Genome resource banking of biomedically important laboratory animals. *Theriogenology*. 2012; 78:1652–1665.

2. Bavister BD, Leibfried ML, Lieberman G. Development of preimplantation embryos of the golden hamster in a defined culture medium. *Biol Reprod.* 1983; 28:235–247. [PubMed: 6830941]
3. Cho F, Honjo S. A simplified method for collecting and preserving cynomolgus macaque semen. *Jpn J Med Sci Biol.* 1973; 26:261–268. [PubMed: 4546196]
4. Chong SS. Contribution of DNA sequence and CAG size to mutation frequencies of intermediate alleles for Huntington disease: evidence from single sperm analysis. *Hum Mol Genet.* 1997; 6:301–309. [PubMed: 9063751]
5. Clavert A, Cranz C, Brun B, Rumpler YA. Method for sperm cryopreservation of the *lemur fulvus*. *Prim Reprod.* 1986; 14:101.
6. Conradie E, Oettle EE, Seier JV. Assessment of acrosomal integrity of vervet monkey spermatozoa after cryopreservation. *J Med Primatol.* 1994; 23:315–316. [PubMed: 7869362]
7. Courtine G, Bunge MB, Fawcett JW, Grossman RG, Kaas JH, Lemon R, Maier I, Martin J, Nudo RJ, Ramon-Cueto A, Rouiller EM, Schnell L, Wannier T, Schwab ME, Edgerton VR. Can experiments in nonhuman primates expedite the translation of treatments for spinal cord injury in humans? *Nat Med.* 2007; 13:561–566. [PubMed: 17479102]
8. Crook ZR, Housman D. Huntington's disease: can mice lead the way to treatment? *Neuron.* 2011; 69:423–435. [PubMed: 21315254]
9. Denis LT, Poindexter AN, Ritter MB, Seager SWJ, Deter RL. Freeze preservation of squirrel monkey sperm for use in timed fertilization studies. *Fertil Steril.* 1976; 27:723–729. [PubMed: 819313]
10. Dong Q, Correa LM, VandeVoort CA. Rhesus monkey sperm cryopreservation with TEST-yolk extender in the absence of permeable cryoprotectant. *Cryobiology.* 2009; 58:20–27. [PubMed: 18992734]
11. Dong Q, Rodenburg SE, Huang C, VandeVoort CA. Cryopreservation of rhesus monkey (*Macaca mulatta*) epididymal spermatozoa before and after refrigerated storage. *J Androl.* 2008; 29:283–292. [PubMed: 18187397]
12. Duyao M, Ambrose C, Mters R, Noveletto A, Persichetti F, Frontali M, Folstein S, Ross C, Franz M, Abbott M. Trinucleotide repeat length instability and age of onset in Huntington's disease. *Nat Genet.* 1993; 4:387–392. [PubMed: 8401587]
13. Feradis AH, Pawitri D, Suatha IK, Amin MR, Yusuf TL, Sajuthi D. Cryopreservation of epididymal spermatozoa collected by needle biopsy from cynomolgus monkeys (*Macaca fascicularis*). *J Med Primatol.* 2001; 30:100–106. [PubMed: 11491402]
14. Gallo P, Cupic D, Bracco F, Krzalic L, Tavalato B, Battistin L. Experimental allergic encephalomyelitis in the monkey: humoral immunity and blood-brain barrier function. *Ital J Neurol Sci.* 1989; 10:561–565. [PubMed: 2515167]
15. Gould KG, Styperek RP. Improved methods for freeze preservation of chimpanzee sperm. *Am J Primatol.* 1989; 18:275–284.
16. HD Collaborative Research Group. A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington's disease chromosomes. *Cell.* 1993; 72:971–983. [PubMed: 8458085]
17. Hewitson L, Simerly CR, Schatten G. Fate of sperm components during assisted reproduction: implications for infertility. *Hum Fertil (Camb).* 2002; 5:110–116. [PubMed: 12193794]
18. Ho AK, Sahakian BJ, Brown RG, Barker RA, Hodges JR, Ane MN, Snowden J, Thompson J, Esmonde T, Gentry R, Moore JW, Bodner T. Profile of cognitive progression in early Huntington's disease. *Neurology.* 2003; 61:1702–1706. [PubMed: 14694033]
19. Hogstedt S, Lindberg BS, Regardh CG, Mostrom U, Rane A. The rhesus monkey as a model for studies of pregnancy induced changes in metoprolol metabolism. *Pharmacol Toxicol.* 1990; 66:32–36. [PubMed: 2308905]
20. Holt, WV.; Wilton, LJ.; Marshall, VS. Cryopreservation of spermatozoa from the marmoset monkey (*Callithrix jacchus*).. In: Bradley, M.; Cummins, J., editors. *Proc. 7th Symp. Spermatology.* Cairns, Australia 17: 1994.
21. Kovtun IV, Therneau TM, McMurray CT. Gender of embryo contributes to CAG instability in transgenic mice containing a Huntington's disease gene. *Hum Mol Genet.* 2000; 9:2767–2775. [PubMed: 11063736]

22. Kraemer DC, Vera Cruz NC. Collection, gross characteristics and freezing of baboon semen. *J Reprod Fertil.* 1969; 20:345–348. [PubMed: 4981520]
23. Kundu MC, May MC, Chosich J, Bradford AP, Lasley B, Gee N, Santoro N, Appt SE, Polotsky AJ. Assessment of luteal function in the vervet monkey as a means to develop a model for obesity-related reproductive phenotype. *Syst Biol Reprod Med.* 2013; 59:74–81. [PubMed: 23278149]
24. Kusunoki H, Daimaru H, Minam S, Nishimoto S, Yamane K, Fukumoto Y. Birth of a chimpanzee (*pan troglodytes*) after artificial insemination with cryopreserved epididymal spermatozoa collected postmortem. *Zoo Biol.* 2001; 20:135–143.
25. Lambert H, Citino S, Collazo I, Jeyendran RS. Penetration of zona-free hamster oocytes by ejaculated cryopreserved gorilla spermatozoa. *Fertil Steril.* 1991; 56:1201–1203. [PubMed: 1743348]
26. Lanzendorf SE, Holmgren WJ, Schaffer N, Hatasaka H, Wentz AC, Jey-endran RS. In vitro fertilization and gamete micromanipulation in the lowland gorilla. *J Assist Reprod Genet.* 1992; 9:358–364. [PubMed: 1472815]
27. Leverage WE, Valerio DA, Schultz AP, Kingsbury E, Dorey C. Comparative study on the freeze preservation of spermatozoa: primate, bovine and human. *Lab Anim Sci.* 1972; 22:882–889. [PubMed: 4118683]
28. Li Y, Cai K, Kovacs A, Ji W. Effects of various extenders and permeating cryoprotectants on cryopreservation of cynomolgus monkey (*Macaca fascicularis*) spermatozoa. *J Androl.* 2005; 26:387–395. [PubMed: 15867007]
29. Li Y, Cai K, Li J, Dinnyes A, Ji W. Comparative studies with six extenders for sperm cryopreservation in the cynomolgus monkey (*Macaca fascicularis*) and rhesus monkey (*Macaca mulatta*). *Am J Primatol.* 2006; 68:39–49. [PubMed: 16419120]
30. Li Y, Si W, Zhang X, Dinnyes A, Ji W. Effect of amino acids on cryopre-servation of cynomolgus monkey (*Macaca fascicularis*) sperm. *Am J Primatol.* 2003; 59:159–165. [PubMed: 12682924]
31. Mahone JP, Dukelow WR. Semen preservation in *Macaca fascicularis*. *Lab Anim Sci.* 1978; 28:556–561. [PubMed: 31513]
32. Merritt AD, Conneally PM, Rahman NF, Drew AL. Juvenile Huntington's chorea. *Excerpta Medica Int Congr Ser.* 1969; 175:645–673.
33. Morrell JM. Cryopreservation of marmoset sperm (*Callithrix jacchus*). *Cryo-lett.* 1997; 18:45–54.
34. Morrell JM, Hodges JK. Cryopreservation of non-human primate sperm: priorities for future research. *Anim Reprod Sci.* 1998; 53:43–63. [PubMed: 9835366]
35. Ng SC, Martelli P, Liow SL, Herbert S, Oh SH. Intracytoplasmic injection of frozen-thawed epididymal spermatozoa in a nonhuman primate model, the cyno-molgus monkey (*Macaca fascicularis*). *Theriogenology.* 2002; 58:1385–1397. [PubMed: 12387351]
36. Obaldia N, Otero W, Marin C, Aparicio J, Cisneros G. Long-term effect of a simple nest-box on the reproductive efficiency and other life traits of an *Aotus lemurinus* monkey colony: an animal model for malaria research. *J Med Primatol.* 2011; 40:383–391. [PubMed: 21781134]
37. O'Sullivan A, He X, McNiven EM, Hinde K, Haggarty NW, Lonnerdal B, Slupsky CM. Metabolomic phenotyping validates the infant rhesus monkey as a model of human infant metabolism. *J Pediatr Gastroenterol Nutr.* 2012; 56:355–363. [PubMed: 23201704]
38. Paulsen JS, Langbehn DR, Stout JC, Aylward E, Ross CA, Nance M, Guttman M, Johnson S, MacDonald M, Beglinger LJ, Duff K, Kayson E, Biglan K, Shoulson I, Oakes D, Hayden M. Detection of Huntington's disease decades before diagnosis: the Predict-HD study. *J Neurol Neurosurg Psychiatry.* 2008; 79:874–880. [PubMed: 18096682]
39. Polge C, Smith AU, Parkes AS. Revival of spermatozoa after vitrification and dehydration at low temperatures. *Nature.* 1949; 164:666. [PubMed: 18143360]
40. Pope CE, Dresser BL, Chin NW, Liu JH, Loskutoff NM, Behnke J, Brown C, MCrae MA, Sinoway CE, Campbell MK, Cameron KN, Owens O' DM, Johnson CA, Evans RR, Cedars MI. Birth of a lowland gorilla (*Gorilla gorilla gorilla*) following in vitro fertilization and embryo transfer. *Am J Primatol.* 1997; 41:247–260. [PubMed: 9057968]
41. Rice J. Animal models: Not close enough. *Nature.* 2012; 484:S9–S9. [PubMed: 22509510]

42. Ross CA, Shoulson I. Huntington disease: pathogenesis, biomarkers, and approaches to experimental therapeutics. *Parkinsonism Relat Disord*. 2009; 15(Suppl 3):S135–138. [PubMed: 20082975]
43. Roos RA. Huntington's disease: a clinical review. *Orphanet J Rare Dis*. 2010; 5:40. [PubMed: 21171977]
44. Roussel JD, Austin CR. Preservation of primate spermatozoa by freezing. *J. Reprod. Fertil*. 1967; 13:333–335. [PubMed: 4960618]
45. Sanchez-Partida LG, Maginnis G, Dominko T, Martinovich C, McVay B, Fanton J, Schatten G. Live rhesus offspring by artificial insemination using fresh sperm and cryopreserved sperm. *Biol Reprod*. 2000; 63:1092–1097. [PubMed: 10993831]
46. Sankai T, Shimizu K, Cho F, Yoshikawa Y. In vitro fertilization of follicular oocytes by frozen-thawed spermatozoa in Japanese monkeys (*Macaca fuscata*). *Lab Anim Sci*. 1997; 47:58–62. [PubMed: 9051648]
47. Sankai T, Terao K, Yanagimachi R, Cho F, Yoshikawa Y. Cryopreservation of spermatozoa from cynomolgus monkeys (*Macaca fascicularis*). *J Reprod Fertil*. 1994; 101:273–278. [PubMed: 7932359]
48. Seier JV, Conradie E, Oettle EE, Fincham JE. Cryopreservation of vervet monkey semen and recovery of progressively motile spermatozoa. *J Med Primatol*. 1993; 22:355–359. [PubMed: 8138985]
49. Si W, Wang H, Reid C, Hildebrandt TB, Ji W. Effect of sugar type on the survival of frozen-thawed rhesus monkey (*Macaca mulatta*) sperm. *Am J Primatol*. 2006; 68:103–108. [PubMed: 16419104]
50. Si W, Zheng P, Li Y, Dinnyes A, Ji W. Effect of glycerol and dimethyl sulfoxide on cryopreservation of rhesus monkey (*Macaca mulatta*) sperm. *Am J Primatol*. 2004; 62:301–306. [PubMed: 15085534]
51. Si W, Zheng P, Tang X, He X, Wang H, Bavister BD, Ji W. Cryopreservation of rhesus macaque (*Macaca mulatta*) spermatozoa and their functional assessment by in vitro fertilization. *Cryobiology*. 2000; 41:232–240. [PubMed: 11161555]
52. Smith K, Dillingham L, Giddens WE Jr. Metabolic bone disease resembling osteosarcoma in a woolly monkey (*Lagothrix lagotricha*). *Lab Anim Sci*. 1978; 28:451–456. [PubMed: 212639]
53. Thomas J, Carver M, Haisch C, Thomas F, Welch J, Carchman R. Differential effects of intravenous anaesthetic agents on cell-mediated immunity in the rhesus monkey. *Clin Exp Immunol*. 1982; 47:457–466. [PubMed: 6978785]
54. Tollner TL, VandeVoort CA, Overstreet JW, Drobnis EZ. Cryopreservation of spermatozoa from cynomolgus monkeys (*Macaca fascicularis*). *J Reprod Fertil*. 1990; 90:347–352. [PubMed: 2250234]
55. Wolf DP. Artificial insemination and the assisted reproductive technologies in non-human primates. *Theriogenology*. 2009; 71:123–129. [PubMed: 18849064]
56. Yang SH, Cheng PH, Banta H, Piotrowska-Nitsche K, Yang JJ, Cheng EC, Snyder B, Larkin K, Liu J, Orkin J, Fang ZH, Smith Y, Bachevalier J, Zola SM, Li SH, Li XJ, Chan AW. Towards a transgenic model of Huntington's disease in a non-human primate. *Nature*. 2008; 453:921–924. [PubMed: 18488016]
57. Younis AI, Rooks B, Khan S, Gould KG. The effects of antifreeze peptide III (AFP) and insulin transferrin selenium (ITS) on cryopreservation of chimpanzee (*Pan troglodytes*) spermatozoa. *J Androl*. 1998; 19:207–214. [PubMed: 9570745]

Figure 1. Sperm viability evaluation. Live sperm stained with SYBR 14 dye (green fluorescence) and dead sperm stained with PI (red fluorescence).

Table 1

The composition of freezing extenders.

Chemicals	TRIS	TEST	TTE
TES	-	172 mM	47.8 mM
Tris	250 mM	84.7 mM	-
Tris-HCl	-	-	12.7 mM
Glucose	-	55.5 mM	111 mM
Citric acid	87.2 mM	-	-
D-fructose	69.4 mM	-	-
Lactose	-	-	55.4 mM
Raffinose	-	-	3.36 mM
Distill water	50 ml	50 ml	50 ml
pH	7.2-7.4	7.2-7.4	7.2-7.4
Egg yolk	20%	20%	20%
Glycerol	8%	3%	5%
Ref.	Our laboratory	Doing et al., 2008	Sankai et al., 1994

Table 2

The post-thawed motility analysis of WT rhesus monkey spermatozoa in TRIS, TEST, and TTE

Extender	Fresh sperm motility, %	Post-thawed sperm motility, % ^a	Motility recovery rate, % ^a
TRIS		46.99 ± 4.82	52.51 ± 7.43
TEST	87.12 ± 4.56	54.78 ± 3.88	60.69 ± 5.58
TTE		50.16 ± 7.52	53.99 ± 6.70

^aValues within column are not significantly different (P>0.05).

Table 3

Post-thawed viability analysis of WT rhesus monkey spermatozoa in TRIS, TEST, and TTE extenders

Extender	Post-thawed sperm viability, %	Viability recovery rate, %
TRIS	48.79 ± 3.46 ^b	60.19 ± 3.62 ^b
TEST	60.75 ± 2.85 ^a	75.23 ± 4.17 ^a
TTE	54.85 ± 3.34 ^{ab}	67.81 ± 4.01 ^{ab}

^aValues within column are significantly different (P<0.05).^bValues within column are significantly different (P<0.05).

Table 4

Post-thaw sperm motility comparison of HD and WT rhesus monkey

Sperm types	Fresh sperm motility, % ^a	Post-thawed sperm motility, % ^a	Motility recovery rate, % ^a
HD	95.37 ± 1.01	53.67 ± 8.96	56.53 ± 9.89
WT	86.60 ± 2.48	58.06 ± 2.68	65.19 ± 4.29

^aValues within column are not significantly different (P>0.05).

Table 5

Post-thaw sperm viability comparison of HD and WT rhesus monkey

Sperm types	Post-thawed sperm viability, %	Viability recovery rate, %
HD	58.18 ± 1.74	62.97 ± 1.57 ^b
WT	61.74 ± 3.46	78.48 ± 3.40 ^a

^aValues within column are significantly different (P<0.05).^bValues within column are significantly different (P<0.05).