CASE REPORT

Coronary steal or large collateral? Three cases of graft failure in sequential and composite grafts

Jwalit Morakhia,¹ Padmakumar Ramachandran,² Naveen Chandra Ganiga Sanjeeva,¹ Ashok Thakkar³

¹Department of Cardiology, Kasturba Medical College, Manipal University, Manipal, Karnataka, India

²Kasturba Medical College, Manipal University, Manipal, Karnataka, India

³Department of Clinical Trials, Sahajanand Medical Tech Pvt Ltd, Surat, Gujarat, India

Correspondence to

Dr Jwalit Morakhia, jwalitm@gmail.com

Accepted 21 October 2014

SUMMARY

Coronary bypass grafting using conduits with multiple distal anastomoses continues to demand scrutiny. While on one hand these techniques allow the surgeon to avoid or minimise aortic manipulation, the unique flow and pressure characteristics lead to complex forms of graft failure if the anatomy of the target vascular bed is not carefully taken into consideration. We report three cases of graft failure in patients with coronary bypass grafting performed using multiple distal anastomoses, and percutaneous revascularisation in one patient.

BACKGROUND

Long-term outcomes of coronary artery bypass grafts (CABG) with multiple distal anastomoses remain equivocal. Graft failure in these patients can occur in various forms. The three cases of graft failure in composite grafts presented here highlight the need for meticulous planning of the graft anatomy based on the haemodynamic profile of the target coronary bed.

CASE PRESENTATION

Case 1

A 60-year-old man presented with a 2-month history of Canadian Cardiovascular Society (CCS) class III angina. He had undergone CABG 8 years prior for a near-total left anterior descending coronary artery (LAD) occlusion, a total occlusion in the major obtuse marginal (OM) and a 70% stenosis in the second diagonal branch (D2). Left internal mammary artery (LIMA) was anastomosed to LAD and left radial artery to the OM. A saphenous vein graft (SVG) was anastomosed to the D2 with proximal take off from the radial graft. Left ventricle (LV) function was well preserved. An angiogram revealed a new discrete 90% stenosis in the proximal right coronary artery (RCA). LIMA to LAD was patent. However, aortic anastomosis of the radial artery graft revealed a near-total occlusion with retrograde flow up the SVG to D2 supplying the OM territory (figure 1 and videos 1–3). We proceeded with revascularising the proximal RCA with a 3.5×18 mm Endeavor Sprint drug-eluting stents (DES). Our options for OM revascularisation included attempting the native left circumflex (LCX) chronic total occlusion (CTO) or proximal radial artery graft anastomosis. We initially considered the CTO anatomy to be favourable and attempted the native LCX CTO percutaneous coronary intervention (PCI). However, while we wired the lesion successfully

with a Fielder XT guidewire, we could not track a 1 mm CTO balloon across it after trying several manoeuvres for additional backup support. We then proceeded with PCI to the radial artery graft. The graft ostium was engaged with a Judkins right 3.5 6 Fr guide catheter and the radial artery graft wired with a Whisper guidewire and the SVG to D2 with a Balance Middle Weight (BMW) guidewire. The lesion was predilated with a 2.5×12 mm Voyager balloon. A 3.5×24 mm Yukon Choice DES was deployed at 16 atm. Ostium was flared with the stent balloon. A satisfactory angiographic result was noted (video 4). The retrograde flow in the D2 graft was abolished (video 5).

Case 2

A 57-year-old man presented with mild dyspnoea on exertion – New York Heart Association (NYHA) class II. His medical history included an inferior wall myocardial infarction 10 years ago. An angiogram performed at that time was reported as showing a LCX proximal total occlusion, 90% discrete stenosis in proximal RCA with mild distal disease, and 50–60% stenosis in proximal LAD. On the basis of this report, he had then undergone CABG with a SVG to RCA and SVGs to LAD and OM in a ‘Y’ anastomosis. He had mild LV dysfunction with ejection fraction (EF) of 48% and treadmill test (TMT) was positive at 7 metabolic equivalents (METs). We performed an angiogram

Figure 1 Graft anatomy—patient 1. LIMA, left internal mammary artery; D2, second diagonal branch; SVG, saphenous vein graft; LCX, left circumflex; LAD, left anterior descending coronary artery; RCA, right coronary artery.

To cite: Morakhia J, Ramachandran P, Ganiga Sanjeeva NC, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2014-206932

Video 1 Retrograde flow in saphenous vein graft to D2 with subsequent filling of radial artery graft to obtuse marginal (RAO Caudal).

Video 3 Y graft angiogram showing slow flow in obtuse marginal graft and no opacification of D2 graft.

Video 2 Retrograde flow in saphenous vein graft to D2 with subsequent filling of radial artery graft to obtuse marginal (LAO Cranial).

Video 4 Post stenting graft angiogram showing brisk flow in obtuse marginal graft and slow flow in D2 graft.

that showed total occlusion of proximal RCA and Proximal LCX. A patent SVG to RCA was noted. The stem of the 'Y' anastomosis was occluded with retrograde flow in the graft to LAD supplying the LCX territory (figure 2 and videos 6 and 7). The patient opted to continue medical therapy and there has been no symptomatic deterioration at 8 months follow-up.

Case 3

A frail 79-year-old man presented with an episode of unwitnessed syncope associated with diaphoresis. He had undergone CABG 14 years earlier for stable angina and an angiogram revealing RCA total occlusion, ramus 60% stenosis and LAD proximal 80% stenosis. A LIMA graft had been anastomosed to mid-LAD and a radial artery with proximal anastomosis to the LIMA was sequentially anastomosed side-to-side to the ramus

Video 5 Post stenting left coronary artery angiogram showing retrograde flow in D2 graft, but no steal into the obtuse marginal graft.

Figure 2 Graft anatomy—patient 2. PDA, posterior descending artery; SVG, saphenous vein graft; LCX, left circumflex; RCA, right coronary artery; LAD, left anterior descending coronary artery.

Video 8 Patent left internal mammary artery to left anterior descending coronary artery.

Video 6 Saphenous vein graft (SVG) to obtuse marginal filling via retrograde flowing SVG to left anterior descending coronary artery (AP Cranial).

Video 7 Saphenous vein graft (SVG) to obtuse marginal filling via retrograde flowing SVG to left anterior descending coronary artery (LAO Caudal).

Figure 3 Graft anatomy—patient 3. LCX, left circumflex; LAD, left anterior descending coronary artery; RCA, right coronary artery.

Video 9 Radial artery graft segment from ramus to posterior descending artery (AP Caudal).

OUTCOME AND FOLLOW-UP

A satisfactory angiographic result was noted in the patient discussed under case 1. Patients discussed under case 2 and 3 remain clinically stable at 8 months follow-up.

DISCUSSION

A wide variety of grafting techniques are practised today. Composite grafts have been studied since the early 1970s,¹ and have promised several advantages. These include the option of avoiding aortic manipulation entirely with off-pump total arterial revascularisation, shorter procedure times, a potentially better run off when the distal vascular bed is diseased or supplying only partially viable myocardium, and in situations where saphenous veins are limited and of poor quality. Preservation of conduits for a potential re-do bypass surgery in the future is an added benefit.

Video 10 Radial artery graft segment from ramus to posterior descending artery (LAO Cranial).

Planning the anatomy of a composite graft needs deliberation. In an angiographic analysis of arterial grafts with multiple distal anastomoses, 4.5% grafts showed reverse flow.² The authors outlined several issues that needed to be addressed when planning a composite graft. A composite graft trades off pressure capacity for a higher flow capacity. The cut-off stenosis in the target vessel, which could lead to graft obstruction or reversal of flow, is higher for a composite graft compared to a non-composite graft. The authors argued against the use of composite grafts with multiple distal end-to-side anastomoses, and advised maintaining a favourable pressure slope from proximal to distal anastomoses such that the graft terminates in the target vessel with the highest degree of stenosis. Similar findings have been echoed in a systematic review of vein grafts where the authors noted better patency rates with side-to-side compared to end-to-side distal anastomoses.³ On the other hand, the PREVENT IV study found higher graft failure rates and a trend towards worse clinical outcomes with composite SVGs.⁴

The often cited disadvantage of a sequential graft is that a large myocardial territory would be jeopardised in case of occlusion of the proximal segment. However, as seen in our three patients, the composite graft can function as large collateral in case of a proximal occlusion. Similar observations have been made by Christenson *et al* and Meeter *et al*.^{5 6}

Learning points

- ▶ Careful assessment of coronary artery disease severity and meticulous planning of the graft anatomy is cardinal to the long-term success of bypass grafts with multiple distal anastomoses.
- ▶ Graft failure in patients with grafts with multiple distal anastomoses can be complex and render further treatment challenging.
- ▶ While grafts with multiple distal anastomoses are a useful tool in certain subsets, their long-term outcomes compared to grafts with single distal anastomosis merit further study.

Acknowledgements The authors thank Miss Shivani Kothari for detailed assistance in manuscript writing.

Contributors JM, PR, NCGS were involved in the surgical and procedural work. AT was involved in the technical writing and manuscript preparation.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Bartley TD, Bigelow JC, Page US. Aortocoronary bypass grafting with multiple sequential anastomoses to a single vein. *Arch Surg* 1972;105:915–17.
- 2 Nakajima H, Kobayashi J, Tagusari O, *et al*. Functional angiographic evaluation of individual, sequential, and composite arterial grafts. *Ann Thorac Surg* 2006;81:807–14.
- 3 Li J, Liu Y, Zheng J, *et al*. The patency of sequential and individual vein coronary bypass grafts: a systematic review. *Ann Thorac Surg* 2011;92:1292–8.
- 4 Mehta RH, Ferguson TB, Lopes RD, *et al*. Saphenous vein grafts with multiple versus single distal targets in patients undergoing coronary artery bypass surgery one-year graft failure and five-year outcomes from the Project of Ex-Vivo Vein Graft Engineering via Transfection (PREVENT) IV Trial. *Circulation* 2011;124:280–8.
- 5 Meeter K, Veldkamp R, Tijssen J, *et al*. Clinical outcome of single versus sequential grafts in coronary bypass operations at ten years' follow-up. *J Thorac Cardiovasc Surg* 1991;101:1076–81.
- 6 Christenson J, Simonet F, Schmuziger M. Sequential vein bypass grafting: tactics and long-term results. *Vascular* 1998;6:389–97.

Copyright 2014 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow