CASE REPORT

Life-threatening haemoperitoneum secondary to rupture of simple ovarian cyst

Christiane Nyhsen, Syed Umair Mahmood

CHS, Sunderland, UK

Correspondence toDr Syed Umair Mahmood,
umair.mahmood@gmail.com

Accepted 4 November 2014

SUMMARY

A 30-year-old woman with no significant medical or family history presented with epigastric pain radiating to the right shoulder tip. She had an acute drop of haemoglobin within 6 h of admission. She was found to be actively bleeding from a ruptured simple ovarian cyst with no other pathology found. Bleeding was stopped by diathermy.

BACKGROUND

Women of childbearing age with abdominal/pelvic pain are a common presentation. Usually perceived pain does not reflect an acute life-threatening pathology. However, rarely, as demonstrated in this case, prompt imaging and intervention are necessary. It is therefore important to remain vigilant and closely observe all patients, in particular if early discharge is considered.

CASE PRESENTATION

A 30-year-old woman presented with sudden onset of abdominal pain at midnight. Initially described as right flank pain, on admission it was noted mainly in the epigastric region radiating to the right shoulder tip. Associated symptoms included non-bilious vomiting.

Normal regular periods were noted with the last menstrual period being 5–7 days prior to admission. Urine pregnancy test was negative on two separate occasions. There was no significant medical or family history. The patient did not report any pelvic pain.

Examination revealed a tender epigastric region with generalised guarding. Initially, the patient was haemodynamically stable with no evidence of gastrointestinal bleeding.

The initial impression was that the symptoms were owing to a perforated gastric/duodenal ulcer or of gallbladder/biliary pathology.

INVESTIGATIONS

The patient's haemoglobin was normal (138 g/L) on admission. Six hours later, her haemoglobin level dropped to 111 g/L and a marginal rise in urea was noted. Lactate and C reactive protein were normal. White cell count was raised at $24.77 \times 10^9/L$.

CT of the abdomen and pelvis showed a large volume of free turbid fluid (figure 1) and what was thought to be acute extravasation of contrast in the pelvis (figure 2). In addition, a possible left ovarian complex cystic lesion was identified (figure 3).

Subsequent transabdominal ultrasound of the abdomen and pelvis (to clarify pelvic appearances) confirmed a large amount of free fluid containing

Figure 1 Axial contrast-enhanced CT scan with free fluid.

some low-level echoes (fluid with raised cellular count, see figure 4). There was a highly unusual appearance of pelvic organs with layering of echogenic material around the uterus (figure 5). A left ovarian cystic lesion was demonstrated, no definite cyst was seen on the right (figure 6). Unfortunately, the patient was very unwell and trans-vaginal ultrasound was not performed. Decision for immediate surgery was taken.

DIFFERENTIAL DIAGNOSIS

- ▶ Haemorrhage from ruptured ectopic pregnancy
- ▶ Haemorrhage from complex left ovarian cyst
- ▶ Perforated gastric/duodenal ulcer
- ▶ Bleeding from other unknown cause

Figure 2 Axial contrast-enhanced CT scan showing possible extravasation of contrast.

CrossMark

To cite: Nyhsen C, Mahmood SU. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2014-205061

Figure 3 Coronal contrast-enhanced CT scan showing free fluid and apparent left ovarian lesion.

TREATMENT

At laparoscopy, 1.5 L of blood and clots were drained. A bleeding simple right ovarian cyst was identified and the haemorrhage was successfully stopped by diathermy. A simple left ovarian cyst with no other focal pathology was visualised. There was no evidence of endometriosis as underlying pathology. No pathological tissue was demonstrated, therefore histology was not taken. There were no other significant findings.

Figure 4 Transabdominal ultrasound of the abdomen showing a large amount of free fluid.

Figure 5 Transabdominal ultrasound of the pelvis with haematoma surrounding the uterus.

The patient's haemoglobin had dropped to 94 g/L following surgery and as she was very symptomatic she was transfused 2 units of blood, after which she remained stable. She was prescribed antibiotics (co-amoxiclav and metronidazole) due to raised white cell count on admission and spiking temperatures. The white cell count was almost normal at discharge 2 days later (10 vs 24 on admission).

OUTCOME AND FOLLOW-UP

The patient made a quick and full recovery. She remains well several months after the initial presentation with no further complications.

DISCUSSION

Idiopathic spontaneous haemoperitoneum can be life-threatening. Early recognition of this pathology is vital so that appropriate investigations and management may be initiated. Presenting symptoms include acute abdominal pain radiating to the back or shoulder tip, nausea/vomiting, bloating/distension, pain intensely exacerbated by movement, decreased urine output, cold peripheries and decreased consciousness.

Figure 6 Transabdominal ultrasound of the pelvis showing left ovarian lesion.

Massive haemoperitoneums result in signs and symptoms of hypovolaemic shock. They may be due to a variety of aetiologies, such as¹:

- ▶ Gynaecological pathologies: Rupture of *ovarian cysts* is the most common cause, often presenting with pain (may be severe), but significant haemorrhaging requiring intervention is uncommon.² Most incidental ovarian cysts resolve within 60 days without intervention; women can be followed up with serial ultrasonography if needed.^{3,4} *Ectopic pregnancies* are less common but have a much higher risk of significant haemorrhage and should be considered in every female patient of reproductive age with a haemoperitoneum. Performing a β human chorionic gonadotropin (β -HCG) test is very helpful, if results are negative, to exclude this differential diagnosis.⁵ Patients with *endometriosis* can occasionally present with significant bleeding. Other gynaecological causes are rare.
- ▶ Hepatic pathologies: Spontaneous rupture of liver lesions may occur, most commonly due to *hepatocellular carcinoma*, particularly in patients with underlying cirrhosis and/or viral hepatitis. Bleeding from other malignant liver changes such as *metastasis* is unusual. Benign liver lesions such as *larger adenomas* can present with massive bleeding (especially if patients are on oral contraceptives/anabolic steroids or during pregnancy). Other causes of liver rupture are quite rare.
- ▶ Splenic pathologies: The commonest underlying causes for non-traumatic splenic rupture are underlying *viral or other infections* (cytomegalovirus, Epstein-Barr virus or malaria) and splenic infiltration in cases of *lymphoma or leukaemia*. Rarely, other focal splenic lesions or diffuse infiltration of the spleen (*amyloidosis or Gaucher's disease*) are the cause.
- ▶ Vascular pathologies: Ruptured *aneurysms* may lead to spontaneous massive haemoperitoneums (including mycotic and pseudoaneurysms) and a variety of abdominal arteries may be involved. Massive haemorrhage from venous origins is usually related with *varices* (in particular in patients with cirrhosis and known portal hypertension), with very poor outcomes reported. Utero-ovarian vessels may rarely rupture in the later stages of *pregnancy* and haemorrhage as a result of increased intra-abdominal pressures during *labour*; this has also been described.
- ▶ Coagulopathies: Patients who are anticoagulated have a significantly increased risk of spontaneous haemorrhages or haemorrhage due to minor, non-identified trauma. Other patients at significantly increased risk of haemorrhage are those under haemodialysis treatment, patients with abnormal clotting due to liver damage and patients with congenital coagulopathies (such as haemophilia or congenital factor X deficiency).

Imaging

Contrast-enhanced CT scan is often the modality of choice if significant haemorrhage is suspected and if the cause is unclear. Nowadays CT scan is available 24/7 and can be performed relatively quickly. It has the advantage of demonstrating acute

extravasation of contrast if present, therefore indicating the possible location of bleeding and urgency of emergency surgery.

Ultrasound is helpful in the acute setting as it can be performed as a portable examination in the resuscitation room (eg, to identify large amounts of free fluid in the abdomen in unstable patients). Ultrasound is also the superior imaging modality when pelvic pathology is suspected.⁵ It demonstrates female pelvic organs more clearly than CT scan (keeping in mind that endovaginal ultrasound may be needed for clarification as a transabdominal ultrasound may not fully visualise the complexity of ovarian lesions).

MRI is not usually used as first line investigation, but it may be helpful in cases of unclear bleeding sources and in known pregnancy when ultrasound is unable to clarify appearances and where CT scan should be avoided.¹

Learning points

- ▶ Gynaecological causes should be considered in female patients with abdominal pain. Pelvic pain may not always be present.
- ▶ The majority of patients presenting with abdominal or pelvic pain of gynaecological origin do not have life-threatening pathologies.
- ▶ Patients can present with life-threatening bleeding from ovarian pathologies, most often ruptured ectopic pregnancies. However, other causes of serious bleeding can occur, for example, from a simple cyst, as in this case. A pregnancy test is very useful to exclude ectopic pregnancies.
- ▶ Imaging can help guide the surgeon, but a complete diagnosis with accurate localisation of the origin of bleeding may not always be possible.

Acknowledgements The authors would like to thank the library staff at Sunderland Royal Hospital for their great support.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Lucey BC, Varghese JC, Anderson SW, *et al.* Spontaneous hemoperitoneum: a bloody mess. *Emerg Radiol* 2007;14:65–75.
- 2 Bottomley C, Bourne T. Diagnosis and management of ovarian cyst accidents. *Best Pract Res Clin Obstet Gynaecol* 2009;23:711–24.
- 3 Pavlik EJ, Ueland FR, Miller RW, *et al.* Frequency and disposition of ovarian abnormalities followed with serial transvaginal ultrasonography. *Obstet Gynecol* 2013;122(2 Pt 1):210–17.
- 4 Okai T, Kobayashi K, Ryo E, *et al.* Transvaginal sonographic appearance of hemorrhagic functional ovarian cysts and their spontaneous regression. *Int J Gynecol Obstet* 1994;44:47–52.
- 5 Hertzberg BS, Kliewer MA, Paulson EK. Ovarian cyst rupture causing hemoperitoneum: imaging features and the potential for misdiagnosis. *Abdom Imaging* 1999;24:304–8.

Copyright 2014 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow