Published in final edited form as:

Curr Opin Neurol. 2014 October ; 27(5): 591–598. doi:10.1097/WCO.0000000000000129.

Sporadic Inclusion Body Myositis: new insights and potential therapy

Pedro Machado¹, Mazen M. Dimachkie², and Richard J. Barohn²

¹MRC Centre for Neuromuscular Diseases, University College London, London, UK

²Department of Neurology, University of Kansas Medical Center, Kansas City, KS, USA

Abstract

Purpose of the review—To describe new insights and developments in the pathogenesis, diagnosis and treatment of sporadic inclusion body myositis (IBM).

Recent findings—Various hypothesis about the pathogenesis of IBM continue to be investigated, including autoimmune factors, mitochondrial dysfunction, protein dyshomeostasis, altered nucleic acid metabolism, myonuclear degeneration and the role of the myostatin pathway. Serum autoantibodies against cytosolic 5' nucleotidase 1A have been identified in IBM showing moderate diagnostic performance. The differential diagnostic value of histopathological features, including different protein aggregates, continues to be evaluated. MRI may also be of monitoring value in IBM. New therapeutic strategies are being tested in IBM patients, namely the up-regulation of the heat shock response and the antagonism of myostatin.

Summary—Recent important advances have occurred in IBM. These advances, including recent and ongoing clinical trials, may lead to earlier diagnosis and improved understanding and treatment of the disease. Despite improved knowledge, IBM continues to be a puzzling disease and the pathogenesis remains to be clarified. An interdisciplinary, bench to bedside translational research approach is crucial for the successful identification of novel treatments for this debilitating, currently untreatable disorder.

Keywords

Inclusion body myositis; myopathies; pathogenesis; treatment

Introduction

Sporadic inclusion body myositis (IBM) is an acquired muscle disease that predominantly affects individuals older than 45 years of age. The exact prevalence of the disease is uncertain and varies between geographic regions, with prevalence estimates in Caucasian populations ranging between 1–71 people per million, reaching 139 per million above the age of 50 (prevalence figures recently reviewed by Gang et al [1]). However, the prevalence

Correspondence to: Dr Pedro Machado, Box 102, MRC Centre for Neuromuscular Diseases, 8–11 Queen Square, WC1N 3BG London, United Kingdom, p.machado@ucl.ac.uk, Phone number: +442034488256.

Competing interests: None.

of IBM is probably higher as the condition is often undiagnosed or misdiagnosed, which contributes to a diagnostic delay on average of 5 or more years [2, 3].

The pathogenesis of IBM is unknown and despite being classically classified alongside polymyositis (PM), dermatomyositis (DM) and immune-mediated necrotising myopathies as an idiopathic inflammatory myopathy, IBM is a very distinct condition characterized by a slowly progressive course, often with asymmetric weakness, early weakness of the finger flexors and quadriceps muscles, the coexistence of inflammatory and degenerative changes and resistance to immunosuppressive treatment [3–7].

This review focuses on new insights and developments in the pathogenesis, diagnosis and treatment of IBM.

IBM pathogenesis – new insights

Multiple hypothesis about the pathogenesis of IBM have been proposed (figure 1). Environmental factors (eg. viral infection), ageing, genetic susceptibility, autoimmunity, accumulation of toxic proteins, myonuclear degeneration, endoplasmic reticulum stress, impairment of autophagy, disruption of the ubiquitin-proteasome system, myostatin signaling, mitochondrial dysfunction and alterations of nucleic metabolism have all been proposed to play a role in IBM pathogenesis [8–11]. However the interplay between these processes and the primary event that leads to the coexistence of autoimmune and degenerative changes remains uncertain.

Regarding the immune mechanisms, the discovery of the first serum autoantibody marker for IBM, targeting cytosolic 5'-nucleotidase 1A (cN1A), represents an important advance [12–14]. Using a commercial anti-cN1A antibody to stain muscle tissue of IBM patients, it has been shown that cN1A immunoreactivity is predominantly located in rimmed vacuoles and areas of myonuclear degeneration, suggesting a mechanistic link between the inflammatory and degenerative components of IBM [13]. In another study, the analysis of serum cytokines and the immunophenotyping in peripheral blood and muscle tissue of IBM patients, showed that IBM patients had increased levels of Th-1 cytokines and chemokines, increased levels of CD8+CD28⁻ T-cells (IFN- γ producers) and decreased frequency of circulating regulatory T-cells (Tregs, CD4⁺CD25⁺CD127^{low}FOXP3⁺), compared to healthy controls. The specificity of these findings and the therapeutic potential of modulating the Th1 and Treg response in IBM remains to be determined [15].

Recently, a reduced expression of micro-RNA-1 (miRNA-1), miRNA-133a, and miRNA-133b has been reported in IBM (and also in PM and DM) muscle tissue versus non-myositis controls. These miRNAs are critical for muscle differentiation and their reduction correlated with augmented expression of several inflammatory cytokines, namely tumor necrosis factor- α (TNF- α) and interleukin-1 β [16]. In mechanistic experiments, TNF- α inhibited the expression of the above miRNAs and blocked the differentiation of human and mouse myoblasts into myocytes/myotubes in an NF κ B-dependent manner. This inhibition of differentiation was overcome by miRNA-1/133a/133b overexpression. The authors suggest that these results may provide a mechanistic link between the inflammatory and

degenerative components of IBM but this may not be unique to IBM since similar micro-RNA changes were observed in PM and DM [16].

Regarding mitochondrial dysfunction, a correlation between the degree of inflammation, degree of mitochondrial changes and atrophy was recently described in IBM muscle biopsies, suggesting a link between these findings [17]. The authors hypothesise that in IBM, the ongoing inflammation and cytokine environment, the associated production of reactive oxygen and nitrogen species and the associated endoplasmic reticulum stress, have a role in the initiation of mitochondrial DNA damage, leading to the accumulation of clonally-expanded mtDNA deletions and respiratory deficiency, a phenomenon that is not compensated by the malfunctioning cell repair mechanisms. Accumulated mitochondrial mutations could have deleterious effects in the muscle tissue (eg. atrophy, splitting and breakage), explaining the correlation between respiratory chain deficiency and muscle atrophy [17–19].

Protein dyshomeostasis continues to prompt great interest as a potential pathogenic mechanism and therapeutic target in IBM. Impaired protein degradation by autophagy and disruption of the ubiquitin-proteasome system (UPS) may lead to protein accumulation and aggregation and subsequent myofibre injury. Proteins which target ubiquitinated proteins for degradation and autophagy associated proteins can be found in the sarcoplasm of IBM muscle tissue, for example valosin containing protein (VCP), p62, LC3 and neighbour of BRCA 1 gene 1 (NBR1) [20–25]. This suggests either an increased need for protein degradation or impaired UPS and/or autophagy [26]. The colocalization of protein aggregates, immunoproteasomes and MHC class I antigens in IBM muscle fibres suggest a possible link between protein degradation and inflammation [27]. One potential therapeutic approach in IBM could therefore be targeting protein dyshomeostasis, for example with drugs that are capable of inducing the production of molecular chaperones, such as heat shock proteins (HSP), a group of highly conserved molecules involved in preventing the misfolding of proteins and therefore responsible for preventing protein aggregation.

Some of the proteins that accumulate in the sarcoplasm and rimmed vacuoles of IBM muscle tissue are typical of nuclei and lysosomes. These proteins include heterogeneous nuclear ribonucleoproteins (hnRNPs). TDP-43 is one of the most abundant of these proteins [28]. Along with TDP-43, other hnRNPs such as hnRNPA1, hnRNPA2/B1 and hnRNPC1/C2, have also been shown to accumulate in the sarcoplasm of IBM muscle tissue [29, 30]. These results support the hypothesis that myonuclei degeneration and RNA metabolism impairment may be also important phenomena in the pathogenesis of IBM [29–34].

IBM diagnosis – new insights

The muscle biopsy of patients with IBM can show a wide range of changes, including infiltration of the endomysium by mononuclear cells which may surround and invade non-necrotic muscle fibres (partial invasion), rimmed vacuoles, eosinophilic inclusions, amyloid inclusions, ragged red fibres, cytochrome oxidase negative (COX–) and succinate dehydrogenase positive (SDH+) fibres, sarcolemmal expression of MHC Class I,

cytoplasmic and/or nuclear tubulofilaments visualised with electron microscopy (EM) and various morphological features such as fibre hypertrophy/atrophy, increased numbers of internal nuclei, split fibres, endomysial fibrosis and necrotic and regenerating fibres. Several immunoreactive protein aggregates have been described, such as p62, phosphorylated tau, TDP-43 and VCP. However, none of the protein aggregates are IBM-specific and there is controversy regarding its presence, relative abundance and significance [9, 26, 35].

The combination of partial invasion, rimmed vacuoles and either amyloid or 15–18nm tubulofilamentous inclusions visualised with EM is considered very specific of IBM [36]. However it is now widely recognized that patients with clinically typical IBM may not present this combination of features, which lacks sensitivity, especially in early disease stages [3, 37]. However, there is still a paucity of data regarding the differential diagnostic performance of pathological findings, both individually and combined.

Over the last few years there has been a shift from a heavily pathologically-focused diagnosis of IBM into a more clinically-focused diagnosis supported by some but not necessarily all the typical IBM histopathological features. However, it should be noted that despite the existence of an IBM typical early disease pattern characterised by finger flexor and knee extensor weakness, not all patients have this pattern, and the muscle biopsy remains a critical investigation in the differential diagnosis of IBM.

This shift has been reflected in the latest European Neuromuscular Centre (ENMC) diagnostic criteria for IBM (table 1), which include 3 categories: “clinico-pathologically defined IBM”, “clinically defined IBM” and “probable IBM” [38]. The last two categories are more flexible in terms of the required pathological features, contributing to an increase of the overall sensitivity of the ENMC criteria and allowing the selection of patients at earlier disease stages, when they may be more amenable to treatment [38].

Two recent studies assessed the differential diagnostic performance of various pathological features in IBM. Brady et al [39] investigated markers of protein aggregates, inflammation and mitochondrial changes in IBM. In the presence of rimmed vacuoles, the combination of increased MHC class I (or endomysial T-cells) and a characteristic p62 staining pattern (figure 2) distinguished IBM from other myopathies with rimmed vacuoles (93% sensitivity, 100% specificity). In the absence of rimmed vacuoles, and compared to steroid responsive inflammatory myopathies (DM and PM), the presence of COX–/SDH+ fibres had excellent sensitivity (100%) and moderate specificity (73%) for IBM, while the above characteristic p62 staining pattern had excellent specificity (100%) but low sensitivity (44%). Therefore the absence of COX–/SDH+ fibres raises doubt on the diagnosis of IBM while the presence of the characteristic p62 pattern may help to exclude the diagnosis of PM/DM when rimmed vacuoles are absent. In another study, Hiniker et al [40] evaluated the diagnostic utility of the markers LC3, p62 and TDP-43 in differentiating IBM from possible IBM, PM and PM with COX-negative fibres. After receiver operating characteristic (ROC)-curve analysis and cut-off determination, the authors suggested that the following thresholds could be of diagnostic value: >14% LC3 positive fibres (100% specificity, 83% sensitivity), >20% p62 positive fibres (100% specificity, 50% sensitivity) and >7% TDP-43 positive fibres (100% specificity, 67% sensitivity) [40].

The above mentioned identification of anti-cN1A autoantibodies in IBM represents another important diagnostic advance. High antibody reactivity achieved 96–98% specificity (33–34% sensitivity), while moderate reactivity showed 60–70% sensitivity and 89–92% specificity for the diagnosis of IBM in a cohort of patients with neuromuscular diseases, providing a good balance between sensitivity and specificity. When it becomes commercially available, anti-cN1A testing could represent an additional helpful tool to aid in the diagnosis of IBM in clinical practice, particularly at early disease stages [12–14]. Anti-cN1A antibodies are also good candidates to be incorporated in future IBM diagnostic criteria. In addition to the IgG isotype, IgM and IgA antibodies have also been described [41]. The isotype pattern varies between patients and it has been proposed that the levels of all 3 isotypes should be considered to improve diagnostic accuracy. Individually, IgG>0.9 absorbance units (AU) had 51% sensitivity and 94% specificity, IgA>1.2 AU had 49% sensitivity and 95% specificity and IgM>1.9 AU had 53% sensitivity and 96% specificity. The combination of isotypes (IgM>1.9, IgA>1.1, or IgG>1.3) increased sensitivity to 76%, retaining a good level of specificity (91%) [41].

Finally, the role of MRI in the diagnosis and monitoring of IBM and muscle diseases in general is increasing [42]. Fatty replacement can be seen on T1-weighted sequences and muscle oedema can be seen on T2-weighted sequences with fat suppression such as the short tau inversion recovery (STIR) sequence (figure 3). Mimicking the typical clinical presentation, it has been reported that fatty replacement occurs preferentially in the deep finger flexors [43–46] and in the quadriceps femoris muscles [44, 45, 47]. A relative sparing of the rectus femoris [44, 45] and preferential fat accumulation within the medial head of gastrocnemius (in comparison with soleus and lateral gastrocnemius) have also been described [44, 45, 47]. Bone marrow oedema (interpreted as inflammation) on STIR images can also be seen, but inflammation is usually less prominent than fatty replacement [43, 44, 46]. Despite being well described, the differential diagnostic performance of the above MRI pattern has not been established and MRI features are not currently included in IBM diagnostic criteria.

IBM treatment – new insights

There is no effective treatment for IBM [3, 6]. New pathogenesis insights [4, 9] have led to the search of alternative therapeutic approaches [6, 7].

One of these new approaches is the inhibition of the myostatin pathway. Myostatin belongs to the transforming growth factor β (TGF- β) superfamily and is a negative regulator of skeletal muscle mass. It acts by binding to the activin receptor IIB (ActRIIB)/Alk4/5 receptor complex resulting in activation of the SMAD 2/3 and mitogen-activated protein kinase intracellular signalling pathways and in inhibition of the phosphoinositide 3-kinase (PI3K) intracellular signalling pathway, leading to changes in gene transcription and effects on protein synthesis that culminate in muscle atrophy [48]. Myostatin accumulation has been shown in IBM muscle fibres [49] and TGF- β superfamily signalling through the ActRIIB receptor has also been shown to be elevated in IBM muscle tissue (using a SMAD2/3 phosphorylation assay) [50]. Myostatin antagonists can act extracellularly by either binding directly to myostatin (myostatin antibodies, the soluble myostatin receptor ActRIIB-Fc,

follistatin, FSTL3, myostatin propeptide, decorin peptides and GASPI) or by binding to its receptor complex (ActRIIB antibodies). Some of the myostatin antagonists are naturally occurring molecules (follistatin, FSTL3, myostatin propeptide and GASPI) [48].

Bimagrumab (BYM338), a fully human monoclonal antibody that binds competitively to ActRIIB, was recently tested in a small study enrolling 14 IBM patients (11 Bimagrumab, 3 Placebo) [50]. A single high dose (30 mg/kg IV) of Bimagrumab was administered. After 8 weeks, both MRI thigh muscle volume (right leg +6.5%, $p=0.024$; left leg +7.6%, $p=0.009$) and lean body mass (+5.7%, $p=0.014$) increased in the Bimagrumab group versus Placebo. Improvement in the 6-minute walking distance (6MWD) test was also observed, peaking at 16 weeks (+14.6%, $p=0.008$, 12/14 patients with follow-up data) [50].

A phase IIb/III randomized, double-blind, placebo-controlled trial (RDBPCT) is ongoing. [51] The study aims to enrol 240 patients worldwide and will contain 4 treatment groups in equal proportions: Bimagrumab at 10, 3 and 1mg/Kg and Placebo given IV every 4 weeks. Physical function, muscle strength (including knee extension strength), mobility, quality of life and swallowing function, among other outcomes, will be assessed at 52 weeks, the primary outcome being change from baseline in the 6MWD test [51]. Myostatin inhibition using Bimagrumab seems a promising approach. However, its long-term safety profile remains to be determined as well as whether the potential to increase muscle volume will actually translate into improved muscle strength and performance. The relationship between changes in muscle mass, muscle strength and physical function is not linear and it is important to analyse all these outcomes, as they provide complementary information [52].

A gene therapy approach using follistatin to inhibit the myostatin pathway is also being tested. The follistatin gene (FS344) is intramuscularly delivered by adeno-associated virus (AAV). This open-label trial aims to enrol 9 IBM patients, distributed in 3 groups each with 3 patients: low dose (2×10^{11} vg/kg, single leg injection), medium dose (3×10^{11} vg/kg, bilateral quadriceps injection) and high dose (6×10^{11} vg/kg, bilateral quadriceps injection). The primary outcome of the trial is safety; secondary outcomes include changes in muscle function and strength, bilateral MRI and biopsy of quadriceps muscles and thigh circumference measurement [53]. One potential advantage of this approach is that a single administration of AAV-FS344 could conceivably lead to long-term expression of the follistatin gene.

Targeting the heat shock response (HSR) represents another potential therapeutic approach in IBM. This approach is supported by the above described evidence for the role of protein dyshomeostasis in the pathogenesis of IBM. A recent RDBPCT was conducted with Arimoclomol, an orally administered drug that can up-regulate the HSR by amplifying HSP expression [54]. The primary outcome of the study was safety and tolerability. Arimoclomol 100mg TID ($n=16$) and Placebo ($n=8$) were administered for 4 months, with a subsequent 8-month blinded follow-up phase. The drug was safe and well tolerated by patients. Furthermore, at 8 months, we detected a trend of slower decline (change) in muscle strength (manual muscle testing and right hand grip maximum voluntary isometric contraction testing) and physical function (IBM functional rating scale) with Arimoclomol as compared with placebo-treated controls [54]. Targeting protein homeostasis may be a valid therapeutic

option in IBM and these data support further research of Arimoclomol in IBM. Since the relationship between physical function and strength is not linear, capturing both provides complementary information [52].

Conclusion

The last few years have witnessed important progress in the understanding of IBM. These advances may lead to improved diagnosis and the discovery of effective drug treatments in the future. Nevertheless, the disease triggering events and exact way how the various disease pathways interact remain to be elucidated. The growing interest of the scientific community and new therapeutic approaches being tested in clinical trials allows us to be hopeful and confident about the future and in the ultimate goal of translating research findings into improved patient care.

Acknowledgments

This publication was supported by researchers at the National Institute for Health Research University College London Hospitals Biomedical Research Centre. This publication was supported by an Institutional Clinical and Translational Science Award, NIH/NCATS Grant Number UL1TR000001. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the NIH. We would like to thank Dr. Jasper Morrow for providing the MR images and Dr. Stefen Brady for providing the histopathological image.

References

1. Gang Q, Bettencourt C, Machado P, Hanna MG, Houlden H. Sporadic inclusion body myositis: the genetic contributions to the pathogenesis. *Orphanet journal of rare diseases*. 2014 Jun 19;9(1):88. Review of known and candidate genetic susceptibility risk factors for IBM. [PubMed: 24948216]
2. Needham M, James I, Corbett A, Day T, Christiansen F, Phillips B, et al. Sporadic inclusion body myositis: phenotypic variability and influence of HLA-DR3 in a cohort of 57 Australian cases. *Journal of neurology, neurosurgery, and psychiatry*. 2008 Sep; 79(9):1056–1060.
3. Benveniste O, Guiguet M, Freebody J, Dubourg O, Squier W, Maissonobe T, et al. Long-term observational study of sporadic inclusion body myositis. *Brain*. 2011 Nov; 134(Pt 11):3176–3184. [PubMed: 21994327]
4. Machado P, Brady S, Hanna MG. Update in inclusion body myositis. *Current opinion in rheumatology*. 2013 Nov; 25(6):763–771. [PubMed: 24067381]
5. Dimachkie MM, Barohn RJ. Inclusion body myositis. *Current neurology and neuroscience reports*. 2013 Jan;13(1):321. [PubMed: 23250766]
6. Breithaupt M, Schmidt J. Update on treatment of inclusion body myositis. *Current rheumatology reports*. 2013 May;15(5):329. [PubMed: 23529584]
7. Lloyd TE. Novel therapeutic approaches for inclusion body myositis. *Current opinion in rheumatology*. 2010 Nov; 22(6):658–664. [PubMed: 20827206]
8. Needham M, Mastaglia FL. Sporadic inclusion body myositis: a continuing puzzle. *Neuromuscular disorders : NMD*. 2008 Jan; 18(1):6–16. [PubMed: 18160291]
9. Greenberg SA. Theories of the pathogenesis of inclusion body myositis. *Current rheumatology reports*. 2010 Jun; 12(3):221–228. [PubMed: 20425523]
10. Dalakas MC. Sporadic inclusion body myositis--diagnosis, pathogenesis and therapeutic strategies. *Nature clinical practice Neurology*. 2006 Aug; 2(8):437–447.
11. Askanas V, Engel WK, Nogalska A. Inclusion body myositis: a degenerative muscle disease associated with intra-muscle fiber multi-protein aggregates, proteasome inhibition, endoplasmic reticulum stress and decreased lysosomal degradation. *Brain pathology*. 2009 Jul; 19(3):493–506. [PubMed: 19563541]

12. Salajegheh M, Lam T, Greenberg SA. Autoantibodies against a 43 KDa muscle protein in inclusion body myositis. *PLoS One*. 2011; 6(5):e20266. [PubMed: 21629782]
13. Benjamin Larman H, Salajegheh M, Nazareno R, Lam T, Sauld J, Steen H, et al. Cytosolic 5'-nucleotidase 1A autoimmunity in sporadic inclusion body myositis. *Ann Neurol*. 2013 Mar; 73(3): 408–418. Identification and diagnostic performance of autoantibodies against cytosolic 5'-nucleotidase 1A in IBM. [PubMed: 23596012]
14. Pluk H, van Hoeve BJ, van Dooren SH, Stammen-Vogelzangs J, van der Heijden A, Schelhaas HJ, et al. Autoantibodies to cytosolic 5'-nucleotidase 1A in inclusion body myositis. *Ann Neurol*. 2013 Mar; 73(3):397–407. Identification and diagnostic performance of autoantibodies against cytosolic 5'-nucleotidase 1A in IBM. [PubMed: 23460448]
15. Allenbach Y, Chaara W, Rosenzweig M, Six A, Prevel N, Mingozi F, et al. Th1 response and systemic treg deficiency in inclusion body myositis. *PLoS One*. 2014; 9(3):e88788. Study describing the cytokine-profile and the immune cell phenotype in IBM. [PubMed: 24594700]
16. Georgantas RW, Streicher K, Greenberg SA, Greenlees LM, Zhu W, Brohawn PZ, et al. Inhibition of myogenic microRNAs 1, 133, and 206 by inflammatory cytokines links inflammation and muscle degeneration in adult inflammatory myopathies. *Arthritis & rheumatology*. 2014 Apr; 66(4):1022–1033. Study investigating the role of micro RNAs in the interplay between inflammation and degeneration in inflammatory myopathies. [PubMed: 24757153]
17. Rygiel KA, Miller J, Grady JP, Rocha MC, Taylor RW, Turnbull DM. Mitochondrial and inflammatory changes in sporadic Inclusion Body Myositis. *Neuropathology and applied neurobiology*. 2014 Apr 18. Study investigating the prevalence of respiratory deficiency and its relationship to inflammation and atrophy in IBM.
18. Bua EA, McKiernan SH, Wanagat J, McKenzie D, Aiken JM. Mitochondrial abnormalities are more frequent in muscles undergoing sarcopenia. *Journal of applied physiology*. 2002 Jun; 92(6): 2617–2624. [PubMed: 12015381]
19. Krishnan KJ, Reeve AK, Samuels DC, Chinnery PF, Blackwood JK, Taylor RW, et al. What causes mitochondrial DNA deletions in human cells? *Nature genetics*. 2008 Mar; 40(3):275–279. [PubMed: 18305478]
20. Cacciottolo M, Nogalska A, D'Agostino C, Engel WK, Askanas V. Chaperone-mediated autophagy components are upregulated in sporadic inclusion-body myositis muscle fibres. *Neuropathology and applied neurobiology*. 2013 Dec; 39(7):750–761. [PubMed: 23452232]
21. Girolamo F, Lia A, Amati A, Strippoli M, Coppola C, Virgintino D, et al. Overexpression of autophagic proteins in the skeletal muscle of sporadic inclusion body myositis. *Neuropathology and applied neurobiology*. 2013 Dec; 39(7):736–749. [PubMed: 23452291]
22. Temiz P, Weihl CC, Pestronk A. Inflammatory myopathies with mitochondrial pathology and protein aggregates. *J Neurol Sci*. 2009 Mar 15; 278(1–2):25–29. [PubMed: 19101700]
23. D'Agostino C, Nogalska A, Cacciottolo M, Engel WK, Askanas V. Abnormalities of NBR1, a novel autophagy-associated protein, in muscle fibers of sporadic inclusion-body myositis. *Acta neuropathologica*. 2011 Nov; 122(5):627–636. [PubMed: 21935636]
24. D'Agostino C, Nogalska A, Engel WK, Askanas V. In sporadic inclusion body myositis muscle fibres TDP-43-positive inclusions are less frequent and robust than p62 inclusions, and are not associated with paired helical filaments. *Neuropathology and applied neurobiology*. 2011 Apr; 37(3):315–320. [PubMed: 20626631]
25. Ju JS, Weihl CC. p97/VCP at the intersection of the autophagy and the ubiquitin proteasome system. *Autophagy*. 2010 Feb; 6(2):283–285. [PubMed: 20083896]
26. Weihl CC, Pestronk A. Sporadic inclusion body myositis: possible pathogenesis inferred from biomarkers. *Current opinion in neurology*. 2010 Oct; 23(5):482–488. [PubMed: 20664349]
27. Ferrer I, Martin B, Castano JG, Lucas JJ, Moreno D, Olive M. Proteasomal expression, induction of immunoproteasome subunits, and local MHC class I presentation in myofibrillar myopathy and inclusion body myositis. *Journal of neuropathology and experimental neurology*. 2004 May; 63(5): 484–498. [PubMed: 15198127]
28. Salajegheh M, Pinkus JL, Taylor JP, Amato AA, Nazareno R, Baloh RH, et al. Sarcoplasmic redistribution of nuclear TDP-43 in inclusion body myositis. *Muscle Nerve*. 2009 Jul; 40(1):19–31. [PubMed: 19533646]

29. Pinkus JL, Amato AA, Taylor JP, Greenberg SA. Abnormal distribution of heterogeneous nuclear ribonucleoproteins in sporadic inclusion body myositis. *Neuromuscular disorders : NMD*. 2014 Jul; 24(7):611–616. Data suggesting abnormal RNA metabolism in IBM. [PubMed: 24857366]
30. Cortese A, Plagnol V, Brady S, Simone R, Lashley T, Acevedo-Arozena A, et al. Widespread RNA metabolism impairment in sporadic inclusion body myositis TDP43-proteinopathy. *Neurobiology of aging*. 2014 Jun; 35(6):1491–1498. Data suggesting that widespread alterations of RNA metabolism have a pathogenic role in IBM. [PubMed: 24462217]
31. Greenberg SA, Pinkus JL, Amato AA. Nuclear membrane proteins are present within rimmed vacuoles in inclusion-body myositis. *Muscle Nerve*. 2006 Oct; 34(4):406–416. [PubMed: 16823856]
32. Nakano S, Shinde A, Fujita K, Ito H, Kusaka H. Histone H1 is released from myonuclei and present in rimmed vacuoles with DNA in inclusion body myositis. *Neuromuscular disorders : NMD*. 2008 Jan; 18(1):27–33. [PubMed: 17888663]
33. Nalbantoglu J, Karpati G, Carpenter S. Conspicuous accumulation of a single-stranded DNA binding protein in skeletal muscle fibers in inclusion body myositis. *Am J Pathol*. 1994 May; 144(5):874–882. [PubMed: 8178939]
34. Olive M, Janue A, Moreno D, Gamez J, Torrejon-Escribano B, Ferrer I. TAR DNA-Binding protein 43 accumulation in protein aggregate myopathies. *Journal of neuropathology and experimental neurology*. 2009 Mar; 68(3):262–273. [PubMed: 19225410]
35. Greenberg SA. How citation distortions create unfounded authority: analysis of a citation network. *Bmj*. 2009; 339:b2680. [PubMed: 19622839]
36. Griggs RC, Askanas V, DiMauro S, Engel A, Karpati G, Mendell JR, et al. Inclusion body myositis and myopathies. *Annals of neurology*. 1995 Nov; 38(5):705–713. [PubMed: 7486861]
37. Brady S, Squier W, Hilton-Jones D. Clinical assessment determines the diagnosis of inclusion body myositis independently of pathological features. *Journal of neurology, neurosurgery, and psychiatry*. 2013 Nov; 84(11):1240–1246. Study describing and comparing the clinical assessment and histopathological findings at presentation and investigating disease progression in IBM patients.
38. Rose MR, Group EIW. 188th ENMC International Workshop: Inclusion Body Myositis, 2–4 December 2011, Naarden, The Netherlands. *Neuromuscular disorders : NMD*. 2013 Dec; 23(12):1044–1055. [PubMed: 24268584]
39. Brady S, Squier W, Sewry C, Hanna M, Hilton-Jones D, Holton JL. A retrospective cohort study identifying the principal pathological features useful in the diagnosis of inclusion body myositis. *BMJ open*. 2014; 4(4):e004552. Study evaluating the diagnostic utility of a number of markers of protein aggregates together with mitochondrial and inflammatory changes in IBM.
40. Hiniker A, Daniels BH, Lee HS, Margeta M. Comparative utility of LC3, p62 and TDP-43 immunohistochemistry in differentiation of inclusion body myositis from polymyositis and related inflammatory myopathies. *Acta neuropathologica communications*. 2013; 1(1):29. Quantitative study of the diagnostic value of LC3, p62, and TDP-43 immunohistochemistry in IBM. [PubMed: 24252466]
41. Greenberg SA. Cytoplasmic 5'-Nucleotidase Autoantibodies in Inclusion Body Myositis: Isotypes and Diagnostic Utility. *Muscle Nerve*. 2014 Feb 7.
42. Wattjes MP, Kley RA, Fischer D. Neuromuscular imaging in inherited muscle diseases. *European radiology*. 2010 Oct; 20(10):2447–2460. [PubMed: 20422195]
43. Cantwell C, Ryan M, O'Connell M, Cunningham P, Brennan D, Costigan D, et al. A comparison of inflammatory myopathies at whole-body turbo STIR MRI. *Clinical radiology*. 2005 Feb; 60(2):261–267. [PubMed: 15664582]
44. Cox FM, Reijnierse M, van Rijswijk CS, Wintzen AR, Verschuuren JJ, Badrising UA. Magnetic resonance imaging of skeletal muscles in sporadic inclusion body myositis. *Rheumatology*. 2011 Jun; 50(6):1153–1161. [PubMed: 21288962]
45. Phillips BA, Cala LA, Thickbroom GW, Melsom A, Zilko PJ, Mastaglia FL. Patterns of muscle involvement in inclusion body myositis: clinical and magnetic resonance imaging study. *Muscle Nerve*. 2001 Nov; 24(11):1526–1534. [PubMed: 11745956]

46. Sekul EA, Chow C, Dalakas MC. Magnetic resonance imaging of the forearm as a diagnostic aid in patients with sporadic inclusion body myositis. *Neurology*. 1997 Apr; 48(4):863–866. [PubMed: 9109868]
47. Reimers CD, Schedel H, Fleckenstein JL, Nagele M, Witt TN, Pongratz DE, et al. Magnetic resonance imaging of skeletal muscles in idiopathic inflammatory myopathies of adults. *Journal of neurology*. 1994 Mar; 241(5):306–314. [PubMed: 8006684]
48. Smith RC, Lin BK. Myostatin inhibitors as therapies for muscle wasting associated with cancer and other disorders. *Curr Opin Support Palliat Care*. 2013 Dec; 7(4):352–360. [PubMed: 24157714]
49. Wojcik S, Engel WK, McFerrin J, Askanas V. Myostatin is increased and complexes with amyloid-beta within sporadic inclusion-body myositis muscle fibers. *Acta neuropathologica*. 2005 Aug; 110(2):173–177. [PubMed: 15983828]
50. Amato AA, Sivakumar K, Goyal N, David W, Barohn RJ, Salajegheh M, et al. Treatment of Sporadic Inclusion Body Myositis with an Anti-Activin Receptor IIB Antibody. *Annals of Neurology*. 2013 Oct.74:S93–S94.
51. [Access date: 20 June 2014] Efficacy and Safety of Bimagrumab/BYM338 at 52 Weeks on Physical Function, Muscle Strength, Mobility in sIBM Patients (RESILIENT). <http://www.clinicaltrials.gov/ct2/show/NCT01925209>
52. Hogrel JY, Allenbach Y, Canal A, Leroux G, Ollivier G, Mariampillai K, et al. Four-year longitudinal study of clinical and functional endpoints in sporadic inclusion body myositis: Implications for therapeutic trials. *Neuromuscul Disord*. 2014 Jul; 24(7):604–610. [PubMed: 24857365]
53. [Access date: 20 June 2014] Follistatin Gene Transfer to Patients With Becker Muscular Dystrophy and Sporadic Inclusion Body Myositis. <http://www.clinicaltrials.gov/ct2/show/NCT01519349>
54. Machado P, Miller A, Herbelin L, He J, Noel J, Wang Y, et al. Safety and tolerability of arimoclomol in patients with sporadic inclusion body myositis: a randomised, double-blind, placebo-controlled, phase IIa proof-of-concept trial. *Ann Rheum Dis*. 2013 Jun.72(Suppl3):164.

Bullet points

- Evidence for the role of autoimmunity, mitochondrial dysfunction, protein dyshomeostasis, altered nucleic acid metabolism and myonuclear degeneration in IBM pathogenesis continues to accumulate.
- In spite of new insights, the primary disease mechanism and the exact interaction between different pathways involved in IBM pathogenesis is yet to be determined.
- The new European Neuromuscular Centre (ENMC) research diagnostic criteria will allow the enrolment of patients at earlier disease stages in future clinical trials.
- MHC class I up-regulation, mitochondrial abnormalities and positive immunostaining for p62, TDP-43 and LC3 protein aggregates may have diagnostic potential.
- Auto-antibodies against cytosolic 5'-nucleotidase 1A have moderate diagnostic performance in IBM.
- Targeting protein dyshomeostasis and blocking the myostatin pathway may be promising therapeutic approaches in IBM.

Figure 1.
Potential pathogenic mechanisms leading to myofibre injury in sporadic inclusion body myositis

Figure 2. Characteristic p62 staining pattern in inclusion body myositis (as suggested by Brady et al [39])

The following pattern of p62 immunoreactivity is more characteristic of IBM: strongly stained, discrete and clearly delineated, round or angular aggregates, variable in number and size within a muscle fibre but rarely filling it and predominantly located subsarcolemmal, but also perinuclear and adjacent to vacuoles.

Figure 3. Muscle MRI of a patient suffering from sporadic inclusion body myositis (IBM) and from a healthy control

Axial MRI images of a patient with IBM (A–D) and a healthy control (E–H) with MRI T1-weighted images on the left and the short tau inversion recovery (STIR) images on the right side. Increased signal within muscles on T1-weighted sequences indicate fatty infiltration whilst on STIR sequences indicate muscle oedema. The patient shows abnormalities typical of IBM with fatty atrophy of the quadriceps muscles in the thigh (A, arrows) and most marked involvement of medial gastrocnemius in the calf (C, arrowhead). Muscle inflammation is evident within many thigh muscles (B) and within gastrocnemius in the calf (D). Corresponding images for a healthy volunteer show homogenous low signal intensity within muscles.

Table 1

European Neuromuscular Centre Inclusion Body Myositis research diagnostic criteria 2011 (adapted from [38])

Diagnostic sub-group	Clinico-pathologically defined IBM	Clinically defined IBM	Probable IBM
Clinical features			
Duration of weakness > 12 months	X	X	X
Age at onset > 45 years	X	X	X
Creatine kinase ≥ 15x ULN	X	X	X
FF weakness > SA weakness AND/OR KE weakness HF weakness	X	-	-
FF weakness > SA weakness AND KE weakness HF weakness	-	X	-
FF weakness > SA weakness OR KE weakness HF weakness	-	-	X
Pathological features			
Endomysial inflammatory infiltrate	X	1, but not all of the 4 pathological features	1, but not all of the 4 pathological features
Rimmed vacuoles	X		
Protein accumulation* or 15–18nm filaments	X		
Up-regulation of MHC Class I	-		

* Demonstration of amyloid or other protein accumulation by established methods (e.g. for amyloid Congo red, crystal violet, thioflavin T/S, for other proteins p62, SMI-31, TDP-43).

FF, Finger flexion; HF, Hip flexion; KE, Knee extension; SA, Shoulder abduction; MHC Class I, Major histocompatibility complex class I; ULN, upper limit of normal.