secretion? Would transgenic correction of the HPS-2 epithelial defect in HPS1/2 double-mutant mice prevent the development of spontaneous pulmonary fibrosis? Is epithelial-specific correction of the underlying genetic defect or inhibition of AECII apoptosis a potential therapy for HPS pulmonary fibrosis? Further research is indicated to answer these questions and improve the understanding of HPS pulmonary fibrosis and fibrotic lung disease in general.

Author disclosures are available with the text of this article at www.atsjournals.org.

POORNIMA MAHAVADI, PH.D.

*Universities of Giessen and Marburg Lung Center
Justus-Liebig University
Giessen, Germany*

ANDREAS GUENTHER, M.D.

*Universities of Giessen and Marburg Lung Center
Justus-Liebig University
Giessen, Germany
Agaplesion Lung Clinic Waldhof-Elgershausen
Greifensee, Germany
and
German Center for Lung Research
Giessen, Germany*

BERNADETTE R. GOCHUICO, M.D.

*National Human Genome Research Institute
National Institutes of Health
Bethesda, Maryland*

References

- Di Pietro SM, Dell'Angelica EC. The cell biology of Hermansky-Pudlak syndrome: recent advances. *Traffic* 2005;6:525–533.
- Nakatani Y, Nakamura N, Sano J, Inayama Y, Kawano N, Yamanaka S, Miyagi Y, Nagashima Y, Ohbayashi C, Mizushima M. Interstitial pneumonia in Hermansky-Pudlak syndrome: significance of florid foamy swelling/degeneration (giant lamellar body degeneration) of type II pneumocytes. *Virchows Arch* 2002;437:304–313.
- Raghu G, Collard HR, Egan JJ, Martinez FJ, Behr J, Brown KK, Colby TV, Cordier JF, Flaherty KR, Lasky JA, *et al.*; ATS/ERS/JRS/ALAT Committee on Idiopathic Pulmonary Fibrosis. An official ATS/ERS/JRS/ALAT statement: idiopathic pulmonary fibrosis: evidence-based guidelines for diagnosis and management. *Am J Respir Crit Care Med* 2011;183:788–824.
- Brantly M, Avila NA, Shotelersuk V, Lucero C, Huizing M, Gahl WA. Pulmonary function and high-resolution CT findings in patients with an inherited form of pulmonary fibrosis, Hermansky-Pudlak syndrome, due to mutations in HPS-1. *Chest* 2000;117:129–136.
- Anderson PD, Huizing M, Claassen DA, White J, Gahl WA. Hermansky-Pudlak syndrome type 4 (HPS-4): clinical and molecular characteristics. *Hum Genet* 2003;113:10–17.
- Gochuico BR, Huizing M, Golas GA, Scher CD, Tsokos M, Denver SD, Frei-Jones MJ, Gahl WA. Interstitial lung disease and pulmonary fibrosis in Hermansky-Pudlak syndrome type-2, an adaptor protein-3 complex disease. *Mol Med* 2012;18:56–64.
- Young LR, Pasula R, Gulleman PM, Deutsch GH, McCormack FX. Susceptibility of Hermansky-Pudlak mice to bleomycin-induced type II cell apoptosis and fibrosis. *Am J Respir Cell Mol Biol* 2007;37:67–74.
- Mahavadi P, Korfei M, Henneke I, Liebisch G, Schmitz G, Gochuico BR, Markart P, Bellusci S, Seeger W, Ruppert C, *et al.* Epithelial stress and apoptosis underlie Hermansky-Pudlak syndrome-associated interstitial pneumonia. *Am J Respir Crit Care Med* 2010;182:207–219.
- Wang L, Lyerla T. Histochemical and cellular changes accompanying the appearance of lung fibrosis in an experimental mouse model for Hermansky-Pudlak syndrome. *Histochem Cell Biol* 2010;134:205–213.
- Guttentag SH, Akhtar A, Tao JQ, Atochina E, Rusiniak ME, Swank RT, Bates SR. Defective surfactant secretion in a mouse model of Hermansky-Pudlak syndrome. *Am J Respir Cell Mol Biol* 2005;33:14–21.
- Rouhani FN, Brantly ML, Markello TC, Helip-Wooley A, O'Brien K, Hess R, Huizing M, Gahl WA, Gochuico BR. Alveolar macrophage dysregulation in Hermansky-Pudlak syndrome type 1. *Am J Respir Crit Care Med* 2009;180:1114–1121.
- Young LR, Borchers MT, Allen HL, Gibbons RS, McCormack FX. Lung-restricted macrophage activation in the pearl mouse model of Hermansky-Pudlak syndrome. *J Immunol* 2006;176:4361–4368.
- Lyerla TA, Rusiniak ME, Borchers M, Jahreis G, Tan J, Ohtake P, Novak EK, Swank RT. Aberrant lung structure, composition, and function in a murine model of Hermansky-Pudlak syndrome. *Am J Physiol Lung Cell Mol Physiol* 2003;285:L643–L653.
- Atochina-Vasserman EN, Bates SR, Zhang P, Abramova H, Zhang Z, Gonzales L, Tao JQ, Gochuico BR, Gahl W, Guo CJ, *et al.* Early alveolar epithelial dysfunction promotes lung inflammation in a mouse model of Hermansky-Pudlak syndrome. *Am J Respir Crit Care Med* 2011;184:449–458.
- Young LR, Gulleman PM, Bridges JP, Weaver TE, Deutsch GH, Blackwell TS, McCormack FX. The alveolar epithelium determines susceptibility to lung fibrosis in Hermansky-Pudlak syndrome. *Am J Respir Crit Care Med* 2012;186:1014–1024.
- Korfei M, Ruppert C, Mahavadi P, Henneke I, Markart P, Koch M, Lang G, Fink L, Bohle RM, Seeger W, *et al.* Epithelial endoplasmic reticulum stress and apoptosis in sporadic idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med* 2008;178:838–846.
- King TE Jr, Pardo A, Selman M. Idiopathic pulmonary fibrosis. *Lancet* 2011;378:1949–1961.
- Tanjore H, Blackwell TS, Lawson WE. Emerging evidence for endoplasmic reticulum stress in the pathogenesis of idiopathic pulmonary fibrosis. *Am J Physiol Lung Cell Mol Physiol* 2012;302:L721–L729.

Copyright © 2012 by the American Thoracic Society

DOI: 10.1164/rccm.201210-1771ED

CD14⁺S100A9⁺ Myeloid-derived Suppressor Cells Portend Decreased Survival in Patients with Advanced Lung Cancer

It has been clearly shown from multiple preclinical studies that myeloid-derived suppressor cells (MDSC) serve as a target for inhibiting tumor growth (1–3). Although studies in humans have shown the presence of MDSC in different pathological conditions, understanding their clinical significance in cancer requires the full characterization of these cells. The characteristics of MDSC have been described for several human malignancies, including melanoma, colon cancer, and renal cell carcinoma, but the identification of a unique set of markers for human MDSC has

been challenging because MDSC gene expression varies in different tumor types.

Feng and colleagues in this issue of the *Journal* (pp. 1025–1036) describe the characteristics of the CD11b⁺CD14⁺S100A9⁺ monocytic MDSC and their clinical relevance in patients with advanced non-small cell lung cancer (NSCLC) (4). The identification of markers for the characterization of MDSC in lung cancer is useful because only limited data are available on specific markers. The authors evaluated the characteristics of MDSC in the peripheral

blood of patients with NSCLC [adenocarcinoma, 40; squamous cell carcinoma, 6; and nonclassified NSCLC, 6] in comparison to 17 healthy control subjects. In addition, 40 patients who received cisplatin-based chemotherapy in a randomized prospective trial were evaluated for the predictive role of MDSC. The authors demonstrated that there was a higher proportion of CD11b⁺CD14⁺ than CD11b⁺CD14⁻ MDSC in the peripheral blood mononuclear cells of patients with NSCLC, and the frequencies of both subsets were higher in patients with NSCLC in comparison to healthy control subjects. Both subsets of MDSC had similar suppressive activity on CD8 T-cell proliferation and IFN- γ production. In contrast, CD11b⁺CD14⁺ cells from healthy donors showed no suppressive ability on CD8⁺ T-cell proliferation or IFN- γ production. Through inhibitor and antibody-mediated blocking experiments, the authors showed that the CD11b⁺CD14⁺ MDSC-mediated suppression of CD8 T cells *in vitro* was via arginase, inducible nitric oxide synthase, the IL-13/IL-4R α axis, and IL-10. The levels of S100A9⁺ expression by CD11b⁺CD14⁺ correlated with their suppressive ability. Furthermore, high levels of CD11b⁺CD14⁺S100A9⁺ MDSC were associated with poor response to cisplatin-based chemotherapy and predicted shorter progression-free survival. Although the CD11b⁺CD14⁻ subpopulation of MDSC had an inhibitory ability that was comparable to the CD11b⁺CD14⁺ cells, there was no association of these cells with performance status or treatment response of patients with NSCLC to chemotherapy. The findings of this study are significant as they show that S100A9 can serve as an important marker for the further characterization of the CD11b⁺CD14⁺ MDSC in patients with advanced NSCLC. The results of this study corroborate the findings from a previous study that demonstrated an increase in the population of CD14⁺S100A9⁺ MDSC in the peripheral blood from patients with colon cancer in comparison with healthy control subjects (5). Since solid tumors contain a significant population of tumor-infiltrating myeloid cells that promote tumor growth, the lack of evaluation of the characteristics of MDSC from the NSCLC samples is a limitation of this study. It is not known if the CD11b⁺CD14⁺S100A9⁺ MDSC are present in the lung tumor microenvironment of the patients with advanced NSCLC. Nevertheless, the results of this study demonstrate that levels of CD11b⁺CD14⁺S100A9⁺ MDSC in peripheral blood can serve as a predictor of response to chemotherapy in patients with advanced lung cancer. Further investigations are required to determine whether modulating the levels and activities of CD11b⁺CD14⁺S100A9⁺ MDSC would improve patient outcome

in advanced lung cancer. MDSC play a central role in immune suppressive pathways and tumor progression, and modulating the activities of this population could prove beneficial as an innovative therapeutic strategy against lung cancer.

Author disclosures are available with the text of this article at www.atsjournals.org.

SHERVEN SHARMA, Ph.D.

STEVEN DUBINETT, M.D.

UCLA Lung Cancer Research Program

David Geffen School of Medicine at UCLA

Los Angeles, California

Molecular Medicine Laboratory

Veterans Affairs Greater Los Angeles Healthcare System

Los Angeles, California

and

Jonsson Comprehensive Cancer Center

David Geffen School of Medicine at UCLA

Los Angeles, California

RAVI SALGIA, M.D., Ph.D.

Department of Medicine

University of Chicago

Chicago, Illinois

References

1. Fujita M, Kohanbash G, Fellows-Mayle W, Hamilton RL, Komohara Y, Decker SA, Ohlfest JR, Okada H. COX-2 blockade suppresses gliomagenesis by inhibiting myeloid-derived suppressor cells. *Cancer Res* 2011;71:2664–2674.
2. Mundy-Bosse BL, Lesinski GB, Jaime-Ramirez AC, Benninger K, Khan M, Kuppusamy P, Guentherberg K, Kondadasula SV, Chaudhury AR, La Perle KM, *et al*. Myeloid-derived suppressor cell inhibition of the IFN response in tumor-bearing mice. *Cancer Res* 2011;71:5101–5110.
3. Srivastava MK, Zhu L, Harris-White M, *et al*. Myeloid suppressor cell depletion augments antitumor activity in lung cancer. *PLoS* 2012 7:e40677.
4. Feng P-H, Lee K-Y, Chang Y-L, Chan Y-F, Kuo L-W, Lin T-Y, Chung F-T, Kuo C-S, Yu C-T, Lin S-M, *et al*. CD14⁺S100A9⁺ monocytic myeloid-derived suppressor cells and their clinical relevance in non-small cell lung cancer. *Am J Respir Crit Care Med* 2012;186:1025–1036.
5. Zhao F, Hoechst B, Duffy A, Gamrekelashvili J, Fioravanti S, Manns MP, Greten TF, Korangy F. S100A9 a new marker for monocytic human myeloid-derived suppressor cells. *Immunology* 2012;136:176–183.

Copyright © 2012 by the American Thoracic Society

DOI: 10.1164/rccm.201209-1597ED

Maternal Smoking during Pregnancy and Its Effect on Childhood Asthma

Understanding the Puzzle

Maternal smoking during pregnancy is a well-established and modifiable environmental risk factor for infant respiratory viral infection and childhood asthma (1, 2). *In utero* exposure to maternal smoking influences the development of the fetal respiratory system by impairing an infant's immune system and lung development (3–5). However, as most women who smoke during pregnancy continue to smoke after delivery (6), previous findings of the effect of maternal smoking during pregnancy on childhood asthma often represent a combined effect of maternal smoking during pregnancy and secondhand smoke (SHS) exposure after

pregnancy. The independent effect of *in utero* exposure to maternal smoking has not been well differentiated from the SHS exposure after birth. This is an interesting question, as the relative contribution and mechanisms through which SHS contribute to asthma development are likely to be different with *in utero* exposure to maternal smoking compared with SHS inhalational exposure after birth.

Neuman and colleagues, in this issue of the *Journal* (pp. 1037–1043), conducted a pooled analysis of eight European birth cohorts to assess the independent effect of *in utero* exposure to maternal