Published in final edited form as:

Neuropharmacology. 2015 February ; 89: 255–264. doi:10.1016/j.neuropharm.2014.09.029.

Galanin Mediates Features of Neural and Behavioral Stress Resilience Afforded by Exercise

N. R. Sciolino^{a,1}, J.M. Smith^a, A.M. Stranahan^b, K.G. Freeman^c, G. L. Edwards^{a,c}, D. Weinshenker^d, and P.V. Holmes^{a,e,*}

N. R. Sciolino: natale.sciolino@nih.gov; J.M. Smith: jmsmith9@uga.edu; A.M. Stranahan: astranahan@gru.edu; K.G. Freeman: kgf@uga.edu; G. L. Edwards: gedwards@uga.edu; D. Weinshenker: dweinsh@emory.edu; P.V. Holmes: pvholmes@uga.edu

^aNeuroscience, Biomedical and Health Sciences Institute, University of Georgia, Athens, GA 30602 USA

^bPhysiology Department, Georgia Regents University, Augusta, GA 30912 USA

^cPhysiology and Pharmacology, University of Georgia, Athens, GA 30602 USA

^dHuman Genetics, Emory, Atlanta, GA 30322 USA

^ePsychology, University of Georgia, Athens, GA 30602 USA

Abstract

Exercise promotes resilience to stress and increases galanin in the locus coeruleus (LC), but the question of whether changes in galanin signaling mediate the stress-buffering effects of exercise has never been addressed. To test the contributions of galanin to stress resilience, male Sprague Dawley rats received intracerebroventricular (ICV) cannulation for drug delivery and frontocortical cannulation for microdialysis, and were housed with or without a running wheel for 21d. Rats were acutely injected with vehicle or the galanin receptor antagonist M40 and exposed to a single session of either footshock or no stress. Other groups received galanin, the galanin receptor antagonist M40, or vehicle chronically for 21d prior to the stress session. Microdialysis sampling occurred during stress exposure and anxiety-related behavior was measured on the following day in the elevated plus maze. Dendritic spines were visualized by Golgi impregnation in medial prefrontal cortex (mPFC) pyramidal neurons and quantified. Exercise increased galanin levels in the LC. Under non-stressed conditions, anxiety-related behavior and dopamine levels were comparable between exercised and sedentary rats. In contrast, exposure to stress reduced open arm exploration in sedentary rats but not in exercise rats or those treated chronically with

© 2014 Elsevier Ltd. All rights reserved.

*Corresponding Author: Philip Holmes, Ph.D., Chair of the Neuroscience Program, Professor, University of Georgia, 125 Baldwin Street, Athens, GA 30602 USA Phone: 1-706-542-3104, Fax: 1-706-542-3275, pvholmes@uga.edu.

¹Present Address: Laboratory of Neurobiology, National Institute of Environmental Health Science, Research Triangle Park, NC 27709, USA

6. Funding and Disclosure

No real or potential conflicts of interest are inherent in this work.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ICV galanin, indicating improved resilience. Both exercise and chronic, ICV galanin prevented the increased dopamine overflow and loss of dendritic spines observed after stress in sedentary rats. Chronic, but not acute M40 administration blocked the resilience-promoting effects of exercise. The results indicate that increased galanin levels promote features of resilience at both behavioral and neural levels.

Keywords

Galanin; stress resilience; frontal cortex; locus coeruleus; plasticity; dopamine

1. Introduction ²

Stress and adversity are risk factors for mental illness, but the neural mechanisms underlying stress susceptibility and resilience are not well understood (Franklin et al., 2012). Resilience involves adaptive recovery following adverse experiences, and resilient individuals experience fewer deleterious outcomes following stress. The medial prefrontal cortex (mPFC) is particularly sensitive to stress, both in terms of its function and structure (Arnsten, 2009; McEwen and Morrison, 2013), and also contributes to stress resilience (Maier and Watkins, 2010). mPFC activity is sufficient and necessary for many aspects of behavioral resilience. For example, activation of the mPFC induces resilience in behavioral tests of anxiety (Amat et al., 2005; Covington et al., 2010). Inactivation of the mPFC also reverses the anxiolytic effects of environmental enrichment (Lehmann and Herkenham, 2011) and protective effect of behavioral control over stress (Amat et al., 2005), although not all forms of resilience are PFC-dependent (Greenwood et al., 2013). Dendritic spines, the morphological hallmark of excitatory synapses, are reduced in the mPFC after stress (Leuner and Shors, 2012; Radley and Morrison, 2005), whereas antidepressants (Bessa et al., 2009) or intracranial self-stimulation (Ramkumar et al., 2011) prevent stress-induced alterations in dendritic structure. Consistent with these data, clinical literature shows that resilient individuals exhibit greater mPFC activation (New et al., 2009; Peres et al., 2011), while susceptible individuals exhibit impaired mPFC-to-amygdala functional connectivity (Bremner et al., 1999; Tromp et al., 2012).

Exercise promotes behavioral resilience (Sciolino and Holmes, 2012) and reliably increases the neuropeptide galanin in noradrenergic neurons of the locus coeruleus (LC) (Holmes et al., 2006b; Sciolino et al., 2012; Van Hooissen et al., 2004). Galanin receptors (GalR1-3) are expressed throughout stress-responsive circuits, including the LC, dorsal raphe, ventral tegmental area (VTA), hypothalamus, hippocampus, amygdala, and mPFC (Hawes and Picciotto, 2004). Galanin contributes to stress resilience in behavioral tests of anxiety (Karlsson and Holmes, 2006), and galanin signaling is necessary for behavioral effects of antidepressants in animal models (Lu et al., 2005). Microarray profiling also shows that galanin expression distinguishes resilient and susceptible phenotypes (Krishnan et al., 2007). Although widely recognized for its neuromodulatory actions, galanin also has potent

²Analysis of variance (ANOVA), enzyme-linked immunosorbent assay (ELISA), high performance liquid chromatography (HPLC), in situ hybridization (ISH), intracerebroventricular (ICV), locus coeruleus (LC), medial prefrontal cortex (mPFC), ventral tegmental area (VTA)

neurotrophic activity (Abbosh et al., 2011; Cordero-Llana et al., 2014; Hobson et al., 2008) and may alter the plasticity of neurons in regions that mediate resilience. Galanin may modulate catecholamine transmission in stress-responsive neural pathways (Holmes and Picciotto, 2006), either locally within the LC or in distal targets like the VTA or mPFC. Phasic stimulation of VTA dopamine neurons induces susceptible behavior and rapidly reverses resilience (Chaudhury et al., 2013). In contrast, galanin inhibits midbrain dopamine activity (Counts et al., 2002; Weiss et al., 1998) and generally reduces dopamine release (Jansson et al., 1989; Nordstrom et al., 1987; Tsuda et al., 1998). Thus, galanin signaling is poised to regulate resilience through multiple mechanisms throughout stress-responsive circuitry.

In the current studies, we predicted that increasing galanin levels by wheel running or repeated intracranial galanin administration would produce behavioral resilience in tests of anxiety, while chronic galanin receptor blockade would attenuate the stress-buffering effects of exercise. Based on the recognized role of mPFC dopamine as a synaptic marker of stress exposure, we also hypothesized that exercise or galanin administration would prevent stress-induced increases in dopamine overflow. Finally, we predicted that stress will reduce dendritic spine densities in the mPFC in sedentary rats treated with vehicle but not in exercising or galanin-treated rats. Collectively, the results indicate that both exercise and galanin prevent anxiety-like behaviors induced by stress and provide a potential mechanism through which galanin contributes to resilience.

2. Materials and Methods

2.1. Subjects

Male Sprague-Dawley rats ($N=81$; Harlan, Prattville AL, USA) obtained at 175–200 g and given *ad libitum* food and water were housed at $23\pm3^{\circ}\text{C}$ with a 12:12 reverse light:dark cycle. Following 1-week habituation, rats underwent surgery and were housed individually in clear polycarbonate cages (50×30×30 cm). Procedures were conducted in accordance with the National Research Council Guide for the Care and Use of Laboratory Animals (Publication No. 85–23, revised 2013) and approved by University of Georgia IACUC. All efforts were made to minimize animal suffering, to reduce the numbers of animals used, and to utilize alternatives to in vivo techniques, if available.

2.2. General Experimental Methods

All testing and drug administration occurred in the dark phase of the circadian cycle. The general protocol for experiments was: cannulation (day –8), surgical recovery (day –7 to 0), individual housing into exercise or sedentary conditions (day 1 to 21), footshock during microdialysis sampling (day 20), and elevated plus maze testing followed by brain extraction (day 21).

2.3. Stereotaxic Cannulation Surgery

Rats were anesthetized with isoflurane (1–5%) and guide cannulae were positioned and implanted stereotactically. In Experiment 1, unilateral 22G cannulae targeted the lateral ventricle for acute drug delivery (1mm posterior, –1.5mm lateral, 3mm ventral) relative to

bregma using the atlas of Paxinos and Watson (1998). In Experiment 2, unilateral cannulae targeted the lateral ventricle for chronic drug delivery and another targeted the frontal cortex for microdialysis (3.2 mm anterior, 2.2 mm lateral, 1.5 mm ventral; MAB6.6.IC; SciPro, Sanborn NY, USA). Rats received flunixin meglumine (2.5 mg/kg s.c.) immediately and 24 hours post-surgery.

2.4. Voluntary exercise

Rats were assigned to exercise or sedentary groups on day 1 and individually received continuous, free access to a wheel or no wheel in the homecage. Rats were housed in these conditions until the experiment ended (day 21 or 23), as this exercise duration effectively increases galanin (Holmes et al., 2006b; Reiss et al., 2009; Sciolino et al., 2012; Van Hoomissen et al., 2004). No-wheel controls were selected to provide the best distinction between independent variables (e.g., exercise/environmental enrichment vs. no exercise/no environmental enrichment). An electromagnetic counter (Mini Mitter, Bend OR, USA) detected wheel rotations and distance ran was calculated for each subject.

2.5. Drugs

In experiment 1, rats were injected with vehicle (aCSF or dH₂O) or the galanin receptor antagonist M40 (6nmoles; Tocris, Minneapolis MN, USA) twenty minutes before exposure to footshock or no shock. In experiment 2, rats were injected daily with vehicle, galanin (3 nmoles, Tocris), or M40 (6 nmoles) for 21 d during assignment to exercise or sedentary conditions. Thus, unlike experiment 1, rats in experiment 2 were not microinjected before footshock in the open field. Repeated injections were used to study the trophic effects of galanin that presumably result after several weeks of exercise (Holmes et al., 2006b; Sciolino et al., 2012). Microinjections were administered in a volume of 5 μ L using 27G needles extending 1 mm beyond the guide cannulae. Doses of galanin and M40 were chosen based on behavioral effectiveness in rodents (Lewis et al., 2004; McDonald and Crawley, 1996; Reiss et al., 2009).

2.6. Footshock stress in the open field

Rats were exposed to footshock or no shock during *in vivo* microdialysis in an open field (ENV-520, -4145, SOF-810; Med Associates, St. Albans VT, USA). Footshock consisted of 20 shocks across 20 min (1 mA, 0.5 s long, 60 s ITI) and was delivered through the grid floor in the open field. Rats were placed in the open field and ambulatory distance was recorded before (60 min), during (20 min) and after footshock (40 min). A dialysis sample was collected every 20 min. In Experiment 1, an ICV microinjection was also administered 20 min before exposure to footshock or no shock.

2.7. Elevated plus maze

The '+' shaped maze was 50 cm from the floor and consisted of a pair of open (45×9 cm) and closed arms ($45 \times 9 \times 38$ cm) that met in a central platform (9×9 cm). Rats were placed on the center platform facing an open arm, as described previously (Sciolino et al., 2012). The time spent and entries on the arms was recorded for 5 min and analyzed from video by an experimenter blind to treatment. Open arm time is reported as a standard measure,

although we verified that all changes in open arm time also resulted in similar changes in open arm entries and compensatory changes in closed arm time.

2.8. Microdialysis and high performance liquid chromatography (HPLC)

During the rodent's dark cycle (12–14 hours before lights on), stylets were removed from cannulae in the frontal cortex and replaced with a dialysis probe (MAB 9.6.2, PES membrane of 2 mm, .6 mm OD, 6 kDa cutoff; SciPro, Sanborn NY, USA), as published previously (Ogbonmwan et al.). Rats were returned to their homecages and tested in microdialysis experiments the next day. Rats were connected to PE50 tubing and freely explored an open field while aCSF (Sigma Chemicals, St. Louis MO, USA) was delivered through the in-line at 1.5 µL/min. Dialysate was transferred from the outline every 20 min into sterile microcentrifuge vials filled with DHBA (10% sample volume, 0.08 ng/µL) and 0.1% phosphoric acid. Vials were placed in a dark container on ice, and stored at –80°C until use.

2.9. High Performance Liquid Chromatography

Samples were thawed and injected into a system of ESA 584 pumps (ESA, Chelmsford, MA) with a pre-column filter (Synergi Max-RP4u Security Guard, 150×4.6mm, Phenomenex Inc., Torrance CA, USA) and Max-RP cartridges (Phenomenex), as we performed previously (Masini et al., 2004). Mobile phase was delivered at 1 mL/min and contained 100mM sodium phosphate monobasic (Fisher, Pittsburgh PA, USA), 0.1mM EDTA (Sigma), 0.25mM octanesulfonic acid (Sigma), and 5% acetonitrile (JT Baker). Samples and standards were injected at 20 µL using ESA 542 at 4°C. Peaks were detected over 30 min using ESA CoulArray (–150 and 200 mV on the initial and final electrodes). Position and height of dopamine peaks were compared to standards (Sigma; diluted in aCSF). Standards were run in duplicate/12 samples. Dopamine detection limit was 13.7 nmol/mL. Peak chromatogram area was integrated and analyzed by CoulArray 3.05.

2.10. Brain harvesting and tissue sectioning for enzyme-linked immunosorbent assay (ELISA), in situ hybridization (ISH)

Rats were microinjected with Fast Green dye (2 µL/mL; 1–4 µL) and decapitated directly after behavioral testing (Day 21 or 23). Brains were sectioned coronally (12 µm; Microm, Waldorf, Germany) to verify cannulae placement, which was considered on target by dye in the ventricles and correct cannulae track placement (Paxinos and Watson, 1998). Example cannulae tracks are shown in Fig. S1. In Experiment 1, brains were sectioned into hemispheres. One hemisphere was used for ELISA and the other for in ISH in a counterbalanced manner. Tissue was frozen using dry ice and stored at –80 °C. The hemisphere for ISH was sectioned sagittally (12 µm) and mounted on gelatin-coated slides. In Experiment 2, the frontal cortex was dissected (~5×13×9 mm of tissue), processed for Golgi, sectioned coronally (150µm), and slide-mounted.

2.10.1. Enzyme-Linked Immunosorbent Assay (ELISA)—Tissue from the dorsal pons and ventral midbrain was weighed after freezing in isopentane chilled with dry ice, and protein was extracted as we previously described (Primeaux and Holmes, 2000). Tissue was put in test tubes containing buffer (250 µL of 2.5% aprotinin in .5M acetic acid),

homogenized (15s; PowerGen 125, Fisher), heated (100°C for 10 min), and centrifuged (30 min at 4°C, 3000 rpm at 1500× g; Beckman Model TJ-6). Supernatant was poured off into separate tubes, evaporated in a vacuum-sealed concentrator (18 hr at 60°C and 20,000mm Hg; Labconco Centrivap, Kansas City MO, USA), and reconstituted in 250µL buffer. Tissue was processed according to the manufacturer's instructions for the Galanin Rat ELISA kit (S1208, Peninsula Laboratories, San Carlos CA, USA). Wells were read at 450 nm (MiniReader MR590, Dynatech Instruments Inc, Santa Monica CA, USA), averaged across duplicates, and a curve of best fit was used to calibrate to standards. Data are reported as galanin protein in ng/mL per mg tissue.

2.10.2. *In Situ* Hybridization (ISH) and Densitometry—Tissue was fixed (4% formaldehyde in .12M PBS), dehydrated (0.25% acetic anhydride in 0.1M triethanolamine HCl and 0.9% NaCl, followed by 70–100% EtOH washes), and delipidated in chloroform (Murray et al., 2010; Sciolino et al., 2012). An oligonucleotide probe (Human prepro-galanin: 5'-G AAG GTA GCC AGC GCT GTT CAG GGT CCA GCC TCT CTT CTC CTT T - 3'; Oligos etc, Wilsonville OR, USA) was labeled at the 3' end using ³⁵S-dATP (1 mCi; Perkin Elmer, Boston MA, USA), tailing buffer, CoCl₂, and terminal deoxynucleotransferase (Roche, Indianapolis IN, USA). Unbound radionucleotide was removed using column separation (Micro Bio-Spin P30 in Tris, Bio-Rad, Hercules CA, USA) by centrifugation (4000 rpm at 1000×g for 4min; Micro-12, Separation Tech Inc, Sanford FL, USA) and bound radionucleotide was stabilized using 1M dithiothreitol. Sections were covered with radiolabeled probe in hybridization buffer (25% formamide, 72 mMNaCl, 3.2 mM Tris HCl, 0.0032 mM EDTA, 0.001% sodium pyrophosphate, .004% sodium dodecylsulfate, .002mg/mL heparan sulfate, and 2% dextran sulfate) and incubated for 24hr at 37°C. Sections were washed in 1%SSC and 2%SSC-formamide (50:50) series at 40°C and room temperature, distilled H₂O, and EtOH. Sections were dried and opposed to ³⁵S-sensitive film (Kodak BioMax MR, Rochester NY, USA) for 14 d. Films were developed in Kodak GBX fixer and developer. Images were captured on a light table (Northern Light D95, Imaging Research Inc., Piscataway NJ, USA) and digital camera (Nikon D5000, Micro-NIKKOR 55mmf/2.8 lens, Melville NY, USA) under optimized conditions. The LC was traced in NIH ImageJ (<http://rsb.info.nih.gov/ij/>). Mean grayscale was measured in 2–4 sections/rat.

2.11. Golgi impregnation

Tissue from a subset of rats was impregnated according to Rapid GolgiStain Kit instructions (FD NeuroTechnologies, Columbia MD, USA), sectioned and mounted as described above, counterstained in 1% Neutral Red, and coverslipped. Neurons were selected and imaged if they met the following criteria: layer V pyramidal neuron of the prelimbic/infralimbic region between 4.7 to 2.2 mm from bregma (Paxinos and Watson, 1998), completely impregnated, not truncated, and with projections distinct and separate from other neurons as described previously (Stranahan et al., 2007). Images were obtained on a Zeiss AxioImager MTB microscope in z-stacks for later measurement in Reconstruct (<http://synapses.clm.utexas.edu/tools/reconstruct/reconstruct.stm>). Spines were imaged at 100x in secondary and tertiary dendrites in 5–7 neurons per rat, with sampling of five 10µm segments from the apical and basal arbors. Five neurons per rat were also imaged at 10x to

measure dendritic length and complexity. Data were averaged across cells from each animal for subsequent statistical analysis.

2.12. Statistics

Repeated measures analysis of variance (ANOVA) was used to examine weight, wheel running, and open field activity across time. Univariate ANOVA followed by Fisher's LSD post hoc tests was used to examine behavior, dopamine, and dendritic structure. Dopamine is reported in nmol/mL for descriptive purposes, and analyzed as percent change after stress (average analyte at post-stress time points) / (lowest analyte at baseline) X 100) to reduce within-subject variability and enhance statistical power. Significance was $p < 0.05$ for all analyses.

3. Results

3.1. Experiment 1. Exercise increases galanin and promotes resilience to stress in behavioral tests of anxiety

3.1.1. Exercise increases galanin in the locus coeruleus—Exercise increased galanin protein in the dorsal pons ($t_6 = -4.51$, $p < 0.01$), with no effect in the ventral midbrain compared to sedentary counterparts under non-stress conditions ($p > .05$; Fig. 1A). Exercise also increased galanin mRNA in the LC compared to sedentary counterparts exposed to footshock stress ($t_{18} = -2.37$, $p < 0.05$; Fig. 1B–C).

3.1.2. Voluntary running increased across time, and reduced body weight—Running and body weight was recorded as follows from rats assessed for galanin (Fig. 1A) and behavior (Fig. 2A). Wheel running distance increased over time in rats microinjected with vehicle before exposure to no shock ($F_{1,3} = 48.97$, $p < 0.01$; Fig. S2A). Average running distance peaked on day 20 at 4.24 ± 1.58 km. Body weight increased linearly across time in rats exposed to no shock ($F_{1,8} = 316.90$, $p < 0.01$; Fig. S3A). The interaction of time by exercise was significant ($F_{1,8} = 15.39$, $p < 0.01$), and there was a trend for an exercise effect ($F_{1,8} = 4.59$, $p = 0.08$; Fig. S3A). Post hoc tests showed that exercise and sedentary groups were not different in body weight on day 1 ($p = 0.80$) or 14 ($p = 0.06$), but exercise rats exhibited reduced body weight on experimental day 7 ($p < 0.05$) and 21 ($p < 0.05$; Fig. S3A).

Running and body weight was recorded as follows from rats assessed for galanin mRNA (see Fig. 1B) and behavior (see Fig. 2). Wheel running also increased linearly over time in rats that were microinjected before footshock exposure ($F_{1,12} = 20.56$, $p < 0.01$; Fig. S2B). The interaction of time by group ($F_{1,12} = 1.28$, $p = 0.28$) and drug effect ($F_{1,12} = 0.00$, $p = 0.96$; Fig. S2B) were not significant. Running peaked on day 20 at 2.07 ± 0.25 km. Body weight increased linearly across time in rats exposed to footshock ($F_{1,19} = 214.89$, $p = 0.00$; Fig. S3B). There was an interaction of group by time ($F_{2,19} = 5.45$, $p < 0.01$) and group ($F_{2,19} = 5.00$, $p < .05$; Fig. S3B). Groups were not different in weight on day 1 ($F_{2,19} = 1.17$, $p = 0.33$), but were different on day 7 ($F_{2,19} = 4.07$, $p < 0.05$), 14 ($F_{2,19} = 7.26$, $p < 0.01$), and 21 ($F_{2,19} = 5.06$, $p < 0.05$; Fig. S3B). Exercise reduced body weight on days 7 ($p < 0.01$), 14 ($p < 0.01$), and 21 ($p < 0.05$; Fig. S3B) compared to sedentary control. Vehicle- and M40-treated exercise rats were no different in weight at any time ($p > 0.05$; Fig. S3B).

3.1.3. Sedentary rats are susceptible to anxiety-like behavior induced by stress

Behavior in anxiety-related tests was measured in sedentary and exercised rats given an acute microinjection before exposure to footshock or no shock, as shown in the timeline (Fig. 2A). Percent open arm time ($F_{4,27}=3.14$, $p<0.05$; Fig. 2B) and open arm entries ($F_{4,27}=3.79$, $p<0.01$; Fig. 2C) were significantly different across these groups. Exercise did not impact performance in the elevated plus maze under non-stress conditions, as there were no differences between non-stressed exercised and sedentary rats ($p>0.05$; Fig. 2B–C). However, after exposure to stress sedentary rats exhibited anxiety-like behavior in the elevated plus maze compared to all other groups, including exercise rats treated with acute vehicle- ($p<0.01$) or M40 ($p<0.05$), as well as non-stressed sedentary ($p<0.01$) and exercise controls ($p<0.05$; Fig. 2B–C). The dose of M40 used in the present study was selected because previous research shows that M40 (1–8 nmol) effectively blocks a wide range of behaviors induced by galanin or exercise (Crawley et al., 1993; Lewis et al., 2004; McDonald and Crawley, 1996; Reiss et al., 2009). However, we found that acute M40 and vehicle-treated exercise rats were no different in open arm exploration ($p=0.62$; Fig. 2B–C), indicating that acute blockade of galanin signaling during stress did not impact elevated plus maze behavior in exercise rats on the following day. Pilot studies revealed that acute administration of M40 (6 nmoles ICV) tended to reverse the stress-induced impairment in sedentary rats (mean $\pm$ SEM for % open time is 28.7 ± 11.9 and open entries is 5 ± 1 ; unpublished observation), which may indicate that galanin signaling is differently altered in sedentary versus exercise rats. No differences in general locomotion (e.g., closed arm entries, falls off the maze) was observed across groups ($p>0.05$).

Ambulatory distance in the open field was not generally different in sedentary and exercised rats given an acute microinjection before exposure to footshock or no shock ($F_{4,29}=0.94$, $p=0.45$; Fig. 2D), and generally increased across time in a quadratic manner ($F_{1,29}=49.57$, $p<0.01$, Quadratic). The time by exercise interaction ($F_{4,29}=3.82$, $p<0.01$, Quadratic; Fig. 2D) was significant and follow up tests revealed that ambulation was no different at baseline (Pre: $F_{4,29}=1.13$, $p=0.36$) or after injection (Inject: $F_{4,29}=1.34$, $p=0.28$; Fig. 2D), indicating there were no differences in general locomotor activity or response to the injection between exercised and sedentary rats. However, ambulatory distance was reduced after shock exposure (Post: $F_{4,29}=5.84$, $p<0.01$; Fig. 2D), as shock exposed sedentary-vehicle ($p<0.05$), exercise-vehicle ($p<0.05$), and exercise-M40 treated rats ($p<0.05$) exhibited less ambulation compared to non-shock counterparts.

3.2. Experiment 2. Galanin is necessary and sufficient for features of behavioral resilience and prevents neural markers of stress in the frontal cortex

3.2.1. Running increased across time, but body weight was not influenced by exercise

Running and body weight was recorded from rats that received repeated intracranial injections and were assessed for behavior (see Fig. 3) and tissue analysis (see Fig. 4). Wheel running distance increased linearly over time in rats receiving repeated microinjections and shock exposure ($F_{1,21}=11.30$, $p<0.01$; Fig. S2C). The time by drug interaction ($F_{1,21}=0.74$, $p=0.41$) and drug effect ($F_{1,17}=0.73$, $p=0.41$; Fig. S2C) were not significant. Average running on experimental day 20 was 0.84 ± 0.12 km, which was less compared to Experiment 1 likely due to the stress of repeated microinjections that briefly

interrupted (e.g., for 5 min) wheel access during the active-running cycle. Body weight increased linearly over time ($F_{1,41}=67.93$, $p<0.01$), but there was no significant effect of group assignment ($F_{4,41}=0.62$, $p=0.65$; Fig. S3C). Although the time by group interaction was significant ($F_{4,41}=6.69$, $p<0.01$), follow-up ANOVAs revealed no difference in body weight at each time point ($p>0.05$; Fig. S3C).

3.2.2. Exercise-induced resilience behavior is dependent on galanin receptor activation and recapitulated by chronic ICV galanin administration—

Stress-induced changes in behavior were assessed in sedentary rats treated chronically with vehicle or galanin, and in exercised rats given vehicle or M40, as shown in the timeline (Fig. 3A). The stress-buffering effects of exercise on open arm time in the elevated plus maze were mimicked in sedentary rats treated with galanin ($F_{3,37}=11.21$, $p<0.01$; Fig. 3B), while blockade of galanin receptors induced stress susceptibility in exercised rats ($p<0.05$; Fig. 3B). Neither drug nor exercise altered measures that reflect general locomotion (e.g., closed arm entries, falls off the maze) ($p>0.05$).

During the administration of footshocks in the open field on the previous day, pre-shock ambulatory distances were comparable across groups ($F_{3,36}=0.43$, $p=0.73$; Data Not Shown), thereby indicating no effect of exercise or drug treatment on open field habituation or general locomotor activity. After footshock, ambulatory distance increased in a manner dependent on exercise or galanin treatments ($F_{3,36}=8.00$, $p<0.01$). Exercise reduced stress-induced ambulation compared to sedentary controls ($p<0.05$; one-tailed), and M40 reversed this effect in exercised rats ($p<0.05$; Data Not Shown). Galanin increased stress-induced ambulation compared to all other groups ($p<0.05$; Data Not Shown), which suggests that galanin promotes the use of an active defensive, coping strategy during stress.

3.2.3. Chronic ICV galanin administration and exercise each block the stress-induced increase in dopamine overflow in the frontal cortex—

Stress-induced dopamine overflow in the frontal cortex was assessed in sedentary rats treated chronically with vehicle or galanin, and in exercised rats given vehicle or M40 (Fig. 3A). Baseline dopamine levels were no different between exercise- and drug-treated groups (sedentary-vehicle 14.18 ± 23.68 nmol/mL, sedentary-galanin 23.31 ± 25.34 nmol/mL, exercise-vehicle 24.28 ± 23.09 nmol/mL, exercise-M40 19.96 ± 13.10 nmol/mL; $p>0.05$). Footshock increased dopamine levels compared to baseline (Pre-shock; $F_{1,29}=2.76$, $p=.05$, one-tailed; Fig. 3C). There was a significant group by time interaction ($F_{3,29}=4.39$, $p<0.01$) and main effect of group ($F_{3,29}=6.51$, $p<0.01$; Fig. 3C). Groups were no different at the pre-shock timepoint ($F_{3,29}=0.33$, $p=.81$), but differed post-shock ($F_{3,29}=6.65$, $p<0.01$; Fig. 3C). Galanin ($p<0.05$) or exercise ($p<0.01$; Fig. 3C) prevented the stress-induced increase in dopamine overflow compared to vehicle-treated sedentary controls. There was no effect of M40 in exercise rats compared to vehicle ($p=0.52$; Fig. 3C), indicating that the influence of exercise in dampening stress-evoked dopamine release is independent of galanin. Probe placement is shown in Fig. 3D.

3.2.4. Chronic ICV galanin mimics the protective effects of exercise against stress-induced dendritic spine loss in mPFC neurons—

Dendritic structural plasticity was assessed in Golgi-impregnated pyramidal neurons in layer V of the prefrontal

cortex, which were obtained from sedentary and exercise rats treated chronically with vehicle or galanin. Neurons were sampled from the prelimbic and infralimbic regions of the mPFC (Fig. 4A), and dendritic length and complexity, as well as dendritic spine density, were quantified in the apical and basal arbors (Fig. 4B). Apical spine density differed across groups ($F_{3,21}=7.68$, $p<0.01$; Fig. 4C–D). Footshock stress reduced the number of apical spines in sedentary rats ($p<0.05$; Fig. 4C–D). Galanin treatment ($p<0.05$) or exercise ($p<0.01$) increased the density of spines along the apical dendrites of stressed rats relative to the sedentary control, with exercise also increasing spine density compared to the no-shock sedentary group ($p<0.05$; Fig. 4C–D).

Basal spine density also differed across groups ($F_{3,21}=4.77$, $p<0.01$; Fig. 4E–F). While stress did not reduce dendritic spine density in the basal dendritic arbor of sedentary rats ($p=0.23$), galanin treatment ($p<0.05$) or exercise ($p<0.05$; Fig. 4E–F) increased the density of spines along the basal dendrites of stressed rats relative to sedentary controls. No differences were observed in total dendritic length or complexity after exercise, drug, or shock manipulations ($p>0.05$; Fig. S4).

4. Discussion

The present study employed galanin pharmacology in conjunction with an exercise model known to increase galanin expression in the LC (Holmes et al., 2006b; Sciolino et al., 2012; Van Hooymissen et al., 2004). We found that exercise prevented anxiety-like behavior induced by stress. The resilience afforded by exercise was recapitulated in our behavioral tests by repeated intracranial administration of galanin and reversed by similar administration of the galanin receptor antagonist M40. However, a single dose of M40 (given prior to footshock stress) failed to reverse the stress-protective effects of exercise in anxiety tests, which demonstrates that some features of resilience do not depend on acute modulatory actions of galanin. In the chronic drug administration study, the final microinjection occurred 24 hours prior to behavioral testing, diminishing the possibility that galanin reduced anxiety-related behavior through acute actions. Exercise and galanin treatment also each protected the cortex against stress-induced perturbation of dendritic structure. Combined, our data indicate that galanin is necessary and sufficient for key features of resilience/resistance displayed in a well-established rodent test of anxiety.

Exercise and galanin treatment each prevented anxiety-like behavior produced after exposure to stress. Both exercise and galanin-treated rats reacted normally to footshock (e.g., vocalized, defecated, and urinated during shock), and responded to stress initially by altered locomotor behavior in the open field. However, these groups of rats also exhibited reduced anxiety-like behavior the day following stress in the elevated plus maze. These results are likely attributable to resilience/resistance because exercised and galanin-treated rats were behaviorally protected in tests of anxiety after, but not before the initial exposure to stress. Moreover, neither exercise nor galanin groups differed in general locomotor activity (e.g., closed arm entries in plus maze, baseline ambulation in open field) or shock reactivity (Falls et al., 2010; Sciolino et al., 2012), suggesting that differences in locomotor activity or nociception are not a factor. Our data are in agreement with clinical literature that shows exercise attenuates anxiety and depression (Herring et al., 2010; U.S. Department of

Health and Human Services, 2008), and literature from rodent models that show exercise promotes stress resilience in tests of anxiety (De Chiara et al., 2010; Dishman et al., 1997; Fox et al., 2008; Greenwood and Fleshner, In Press; Greenwood et al., 2008; Greenwood et al., 2007a; Maniam and Morris, 2010; Sasse et al., 2008; Sciolino et al., 2012; Zheng et al., 2006). Our data expand upon this evidence and suggest that galanin is necessary for the stress protection afforded by exercise in a standard behavioral test of anxiety.

Several lines of evidence indicate that galanin promotes behavioral resilience. First, intracranial administration of galanin does not reliably influence anxiety or depression-related behavior in the absence of stress (Holmes et al., 2005; Holmes et al., 1994; Karlsson et al., 2005; Weiss et al., 1998), but provides protection specifically under stressful conditions (Bing et al., 1993). Second, transgenic overexpression of galanin in norepinephrine neurons does not alter spontaneous behavior in tests of anxiety, but provides protection in these tests after stress exposure (Holmes et al., 2002). Third, M40 administration (8 nmoles, ICV) blocks fluoxetine-induced activity in the forced swim test (Lu et al., 2005), which is a potent stressor (Connor et al., 1997; Cryan et al., 2005). The present report is in agreement with the galanin literature and suggests that increased galanin in the LC promotes the stress resilience afforded by exercise.

We found that stress reduced dendritic spines in the mPFC in sedentary animals, whereas galanin treatment protected against this loss. This effect was observed in the apical segment of dendritic arbors but not in the basal, which is consistent with literature reporting apical segments are particularly responsive to stress (Leuner and Shors, 2012). Similarly, exercised rats exhibited increased spines in the mPFC after stress, suggesting that exercise also protects against stress-induced dendritic spine loss. The molecular mechanisms that underlie the change in dendritic morphology caused by stress, and the protection provided by manipulations affording resilience, are beginning to be elucidated. Stress-related depletion of dendritic spines in the hippocampus involves activation of RhoA (Chen et al., 2013). Recent evidence links the neurotrophic and neuroprotective actions of galanin to inhibition of RhoA (Hobson et al., 2013), but additional studies will be required to fully characterize the dynamic alterations in dendritic spines following galanin treatment, with the goal to identify how galanin contributes to structural plasticity in the presence and absence of stress.

We also found that manipulations that promote resilience, including repeated exercise and galanin treatment, attenuated shock-evoked dopamine in the frontal cortex. The galanin agonist galnon also attenuates cocaine-evoked dopamine in the frontal cortex (Ogbonmwan et al., 2014), which provides further support that galanin plays a key role in regulating cortical activity. Our data are consistent with the body of literature that shows increased cortical dopamine is a reliable neurochemical signature of acute stress (Abercrombie et al., 1989; Finlay et al., 1995; Pascucci et al., 2007). Galanin administration may blunt stress-induced dopamine release through direct influences on VTA electrophysiological activity (Counts et al., 2002; Weiss et al., 1998). However, we failed to observe a difference in evoked dopamine release between M40 and vehicle-treated exercise rats. Our data therefore suggest that the resilience afforded by exercise in tests measured herein is not dependent on dopaminergic signaling in the mPFC. Although many manipulations that protect against stress do so in a manner that depends on the mPFC (Amat et al., 2005; Amat et al., 2006;

Christianson et al., 2009; Lehmann and Herkenham, 2011), the resilience afforded by these manipulations may require distinct adaptations in other stress-responsive circuitry. For example, prior evidence shows that exercise protects against uncontrollable stress and enhances fear learning regardless of mPFC lesion (Greenwood et al., 2013). Indeed exercise influences the pattern of stress-induced neuronal activation in regions like the mPFC, PVN, amygdala, hippocampus, raphe, sensory/motor integration sites (Campeau et al., 2010; Collins et al., 2009; Greenwood et al., 2005; Greenwood et al., 2003), suggesting that exercise produces resilience by tuning neuronal activity in several corticolimbic circuits.

A variety of factors increase galanin gene expression in the LC, including acute and chronic stressors (see also Austin et al., 1990; Holmes et al., 1995; Kuteeva et al., 2008), voluntary or forced exercise (Holmes et al., 2006a; Murray et al., 2010; O'Neal et al., 2001; Reiss et al., 2009; Sciolino et al., 2012; Van Hooymissen et al., 2004), and antidepressant treatment with fluoxetine or clomipramine (Holmes et al., 2006a; Lu et al., 2005). Future research is needed to systematically identify how exercise and stress impact galanin signaling. Exercise has notably different effects in the presence and absence of stress on measures like HPA reactivity and neurogenesis (Stranahan et al., 2006; Stranahan et al., 2008). It is therefore important to consider the effect of non-manipulated stressors in the context of our data. For example, behavioral testing was likely stressful as running tended to decrease the day following behavioral testing. All of our animals were also singly housed as typically performed in exercise studies to obtain individualized information about running. Notwithstanding, we found that exercise consistently increased LC galanin levels despite that mRNA was measured after footshock and protein measured after no shock. Exercise likely dose-dependently increases galanin as prior research shows that galanin mRNA in the LC is positively associated with running distance (Eisenstein and Holmes, 2007; Holmes et al., 2006b; Sciolino et al., 2012). Although running distance was generally moderate-to-low in the present studies, probably due to surgical implantation of indwelling cannulae, we reliably observed that exercise protected against anxiety-like behavior resulting after stress. These data collectively suggest that the increases in galanin and stress resilience observed in the present study are due to exercise.

The present study is the first to show that galanin contributes to behavioral resilience that results from exercise. Exercise regulates multiple molecular and structural measures of neuroplasticity, including hippocampal neurogenesis and neurotrophic factor expression (Duman and Monteggia, 2006). These phenomena may contribute to the affective consequences of exercise, though the evidence for their specific roles is mixed (Duman et al., 2008; Duman et al., 2009; Fuss et al., 2010; Greenwood et al., 2007b; Li et al., 2008; Llorens-Martin et al., 2010; Trejo et al., 2008). The lack of research regarding the interplay between these structural and molecular signatures represents a major gap in understanding how gene-environment interactions cooperatively induce resilience or susceptibility, and this area deserves further study. Broadly, our data highlight a neurotrophic role for galanin in stress resilience, as chronic, but not acute, galanin receptor antagonism emerged as a determinant of anxious behavior. Our finding that galanin promoted stress resilience at both the behavioral and neural levels suggests a novel interpretation of galanin's role in anxiety that hinges upon its trophic function rather than its well-known neuromodulatory actions. The efficacy of chronic galanin administration alone in sedentary rats is informative in this

regard. Disruption in the galanin system may therefore tip the balance towards susceptibility, as evidenced from the current report and human studies demonstrating that genetic variants in galanin and its receptors (GalR1-3) correlate with risk for anxiety and depression after stressful experiences. Polymorphisms in galanin and its receptor subtypes were found to be more predictive of neuropsychiatric disease than other well-studied polymorphisms, such as the 5-HTTLPR polymorphism (Juhász et al., 2014). Thus, restoring the dysregulation in galanin, endogenously by exercise or exogenously by galanin-based pharmacotherapies, may offer novel approaches to the prevention or treatment of stress-related mental disorders.

5. Conclusions

The neurotrophic actions of galanin represent a potential mechanism for its impact on stress-related behaviors. Galanin may thus produce stress resilience by influencing neural plasticity in the mPFC. The present experiments revealed that manipulations that chronically enhance galanin levels buffer the impact of stress on anxiety-related behaviors and dendritic spines in the mPFC.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by grants DA027535A to Weinshenker and Holmes, and a Diversity Supplement awarded to the parent grant for Sciolino. Some of these data were published in poster form at the Society for Neuroscience (Poster 351.14; Sciolino et al., 2013, San Diego CA, USA). We are grateful for the technical assistance (creation of brain diagrams) provided by Stephanie Herrlinger.

References

- Abbosh C, Lawkowski A, Zaben M, Gray W. GalR2/3 mediates proliferative and trophic effects of galanin on postnatal hippocampal precursors. *J Neurochem*. 2011; 117:425–436. [PubMed: 21281311]
- Abercrombie ED, Keefe KA, DiFrischia DS, Zigmond MJ. Differential effect of stress on in vivo dopamine release in striatum, nucleus accumbens, and medial frontal cortex. *J Neurochem*. 1989; 52:1655–1658. [PubMed: 2709017]
- Amat J, Baratta MV, Paul E, Bland ST, Watkins LR, Maier SF. Medial prefrontal cortex determines how stressor controllability affects behavior and dorsal raphe nucleus. *Nat Neurosci*. 2005; 8:365–371. [PubMed: 15696163]
- Amat J, Paul E, Zarza C, Watkins LR, Maier SF. Previous experience with behavioral control over stress blocks the behavioral and dorsal raphe nucleus activating effects of later uncontrollable stress: role of the ventral medial prefrontal cortex. *J Neurosci*. 2006; 26:13264–13272. [PubMed: 17182776]
- Arnsten AF. Stress signalling pathways that impair prefrontal cortex structure and function. *Nat Rev Neurosci*. 2009; 10:410–422. [PubMed: 19455173]
- Austin MC, Cottingham SL, Paul SM, Crawley JN. Tyrosine hydroxylase and galanin mRNA levels in locus coeruleus neurons are increased following reserpine administration. *Synapse*. 1990; 6:351–357. [PubMed: 1705056]
- Bessa JM, Ferreira D, Melo I, Marques F, Cerqueira JJ, Palha JA, Almeida OF, Sousa N. The mood-improving actions of antidepressants do not depend on neurogenesis but are associated with neuronal remodeling. *Mol Psychiatry*. 2009; 14:764–773. 739. [PubMed: 18982002]

- Bing O, Moller C, Engel JA, Soderpalm B, Heilig M. Anxiolytic-like action of centrally administered galanin. *Neurosci Lett*. 1993; 164:17–20. [PubMed: 7512244]
- Bremner JD, Staib LH, Kaloupek D, Southwick SM, Soufer R, Charney DS. Neural correlates of exposure to traumatic pictures and sound in Vietnam combat veterans with and without posttraumatic stress disorder: a positron emission tomography study. *Biol Psychiatry*. 1999; 45:806–816. [PubMed: 10202567]
- Campeau S, Nyhuis TJ, Sasse SK, Kryskow EM, Herlihy L, Masini CV, Babb JA, Greenwood BN, Fleshner M, Day HE. Hypothalamic pituitary adrenal axis responses to low-intensity stressors are reduced after voluntary wheel running in rats. *Journal of Neuroendocrinology*. 2010; 22:872–888. [PubMed: 20406350]
- Chaudhury D, Walsh JJ, Friedman AK, Juarez B, Ku SM, Koo JW, Ferguson D, Tsai HC, Pomeranz L, Christoffel DJ, Nectow AR, Ekstrand M, Domingos A, Mazei-Robison MS, Mouzon E, Lobo MK, Neve RL, Friedman JM, Russo SJ, Deisseroth K, Nestler EJ, Han MH. Rapid regulation of depression-related behaviours by control of midbrain dopamine neurons. *Nature*. 2013; 493:532–536. [PubMed: 23235832]
- Chen Y, Kramar EA, Chen LY, Babayan AH, Andres AL, Gall CM, Lynch G, Baram TZ. Impairment of synaptic plasticity by the stress mediator CRH involves selective destruction of thin dendritic spines via RhoA signaling. *Mol Psychiatry*. 2013; 18:485–496. [PubMed: 22411227]
- Christianson JP, Thompson BM, Watkins LR, Maier SF. Medial prefrontal cortical activation modulates the impact of controllable and uncontrollable stressor exposure on a social exploration test of anxiety in the rat. *Stress*. 2009; 12:445–450. [PubMed: 19051125]
- Collins A, Hill LE, Chandramohan Y, Whitcomb D, Droste SK, Reul JM. Exercise improves cognitive responses to psychological stress through enhancement of epigenetic mechanisms and gene expression in the dentate gyrus. *PLoS One*. 2009; 4:e4330. [PubMed: 19180197]
- Connor TJ, Kelly JP, Leonard BE. Forced swim test-induced neurochemical endocrine, and immune changes in the rat. *Pharmacol Biochem Behav*. 1997; 58:961–967. [PubMed: 9408201]
- Cordero-Llana O, Rinaldi F, Brennan PA, Wynick D, Caldwell MA. Galanin promotes neuronal differentiation from neural progenitor cells in vitro and contributes to the generation of new olfactory neurons in the adult mouse brain. *Exp Neurol*. 2014
- Counts SE, McGuire SO, Sortwell CE, Crawley JN, Collier TJ, Mufson EJ. Galanin inhibits tyrosine hydroxylase expression in midbrain dopaminergic neurons. *J Neurochem*. 2002; 83:442–451. [PubMed: 12423254]
- Covington HE 3rd, Lobo MK, Maze I, Vialou V, Hyman JM, Zaman S, LaPlant Q, Mouzon E, Ghose S, Tamminga CA, Neve RL, Deisseroth K, Nestler EJ. Antidepressant effect of optogenetic stimulation of the medial prefrontal cortex. *J Neurosci*. 2010; 30:16082–16090. [PubMed: 21123555]
- Crawley JN, Robinson JK, Langel U, Bartfai T. Galanin receptor antagonists M40 and C7 block galanin-induced feeding. *Brain Res*. 1993; 600:268–272. [PubMed: 7679604]
- Cryan JF, Valentino RJ, Lucki I. Assessing substrates underlying the behavioral effects of antidepressants using the modified rat forced swimming test. *Neurosci Biobehav Rev*. 2005; 29:547–569. [PubMed: 15893822]
- De Chiara V, Errico F, Musella A, Rossi S, Mataluni G, Sacchetti L, Siracusano A, Castelli M, Cavasinni F, Bernardi G, Usiello A, Centonze D. Voluntary exercise and sucrose consumption enhance cannabinoid CB1 receptor sensitivity in the striatum. *Neuropsychopharmacology*. 2010; 35:374–387. [PubMed: 19776732]
- Dishman RK, Renner KJ, Youngstedt SD, Reigle TG, Bunnell BN, Burke KA, Yoo HS, Mougey EH, Meyerhoff JL. Activity wheel running reduces escape latency and alters brain monoamine levels after footshock. *Brain Research Bulletin*. 1997; 42:399–406. [PubMed: 9092882]
- Duman CH, Schlesinger L, Russell DS, Duman RS. Voluntary exercise produces antidepressant and anxiolytic behavioral effects in mice. *Brain Res*. 2008; 1199:148–158. [PubMed: 18267317]
- Duman CH, Schlesinger L, Terwilliger R, Russell DS, Newton SS, Duman RS. Peripheral insulin-like growth factor-I produces antidepressant-like behavior and contributes to the effect of exercise. *Behav Brain Res*. 2009; 198:366–371. [PubMed: 19056428]

- Duman RS, Monteggia LM. A neurotrophic model for stress-related mood disorders. *Biol Psychiatry*. 2006; 59:1116–1127. [PubMed: 16631126]
- Eisenstein SA, Holmes PV. Chronic and voluntary exercise enhances learning of conditioned place preference to morphine in rats. *Pharmacol Biochem Behav*. 2007; 86:607–615. [PubMed: 17368736]
- Falls WA, Fox JH, MacAulay CM. Voluntary exercise improves both learning and consolidation of cued conditioned fear in C57 mice. *Behavioral Brain Research*. 2010; 207:321–331.
- Finlay JM, Zigmond MJ, Abercrombie ED. Increased dopamine and norepinephrine release in medial prefrontal cortex induced by acute and chronic stress: effects of diazepam. *Neuroscience*. 1995; 64:619–628. [PubMed: 7715775]
- Fox JH, Hammack SE, Falls WA. Exercise is associated with reduction in the anxiogenic effect of mCPP on acoustic startle. *Behav Neurosci*. 2008; 122:943–948. [PubMed: 18729648]
- Franklin TB, Saab BJ, Mansuy IM. Neural mechanisms of stress resilience and vulnerability. *Neuron*. 2012; 75:747–761. [PubMed: 22958817]
- Fuss J, Ben Abdallah NM, Hensley FW, Weber KJ, Hellweg R, Gass P. Deletion of running-induced hippocampal neurogenesis by irradiation prevents development of an anxious phenotype in mice. *PLoS One*. 2010;5.
- Greenwood, BN.; Fleshner, M. Mechanisms underlying the relationship between physical activity and anxiety: Animal data. Routledge; New York: In Press
- Greenwood BN, Foley TE, Burhans D, Maier SF, Fleshner M. The consequences of uncontrollable stress are sensitive to duration of prior wheel running. *Brain Research*. 2005; 1033:164–178. [PubMed: 15694921]
- Greenwood BN, Kennedy S, Smith TP, Campeau S, Day HE, Fleshner M. Voluntary freewheel running selectively modulates catecholamine content in peripheral tissue and c-Fos expression in the central sympathetic circuit following exposure to uncontrollable stress in rats. *Neuroscience*. 2003; 120:269–281. [PubMed: 12849759]
- Greenwood BN, Spence KG, Crevling DM, Clark PJ, Craig WC, Fleshner M. Exercise-induced stress resistance is independent of exercise controllability and the medial prefrontal cortex. *Eur J Neurosci*. 2013; 37:469–478. [PubMed: 23121339]
- Greenwood BN, Strong PV, Brooks L, Fleshner M. Anxiety-like behaviors produced by acute fluoxetine administration in male Fischer 344 rats are prevented by prior exercise. *Psychopharmacology (Berl)*. 2008; 199:209–222. [PubMed: 18454279]
- Greenwood BN, Strong PV, Dorey AA, Fleshner M. Therapeutic effects of exercise: wheel running reverses stress-induced interference with shuttle box escape. *Behavioral Neuroscience*. 2007a; 121:992–1000. [PubMed: 17907830]
- Greenwood B, Strong P, Foley T, Thompson R, Fleshner M. Learned helplessness is independent of levels of brain-derived neurotrophic factor in the hippocampus. *Neuroscience*. 2007b; 144:1193–1208. [PubMed: 17161541]
- Hawes JJ, Picciotto MR. Characterization of GalR1, GalR2, and GalR3 immunoreactivity in catecholaminergic nuclei of the mouse brain. *J Comp Neurol*. 2004; 479:410–423. [PubMed: 15514977]
- Herring MP, O'Connor PJ, Dishman RK. The effect of exercise training on anxiety symptoms among patients: a systematic review. *Archives of Internal Medicine*. 2010; 170:321–331. [PubMed: 20177034]
- Hobson SA, Bacon A, Elliot-Hunt CR, Holmes FE, Kerr NC, Pope R, Vanderplank P, Wynick D. Galanin acts as a trophic factor to the central and peripheral nervous systems. *Cell Mol Life Sci*. 2008; 65:1806–1812. [PubMed: 18500646]
- Hobson SA, Vanderplank PA, Pope RJ, Kerr NC, Wynick D. Galanin stimulates neurite outgrowth from sensory neurons by inhibition of Cdc42 and Rho GTPases and activation of cofilin. *J Neurochem*. 2013; 127:199–208. [PubMed: 23895321]
- Holmes A, Li Q, Koenig EA, Gold E, Stephenson D, Yang RJ, Dreiling J, Sullivan T, Crawley JN. Phenotypic assessment of galanin overexpressing and galanin receptor R1 knockout mice in the tail suspension test for depression-related behavior. *Psychopharmacology (Berl)*. 2005; 178:276–285. [PubMed: 15365683]

- Holmes A, Picciotto MR. Galanin: a novel therapeutic target for depression, anxiety disorders and drug addiction? *CNS Neurol Disord Drug Targets*. 2006; 5:225–232. [PubMed: 16611095]
- Holmes A, Yang RJ, Crawley JN. Evaluation of an anxiety-related phenotype in galanin overexpressing transgenic mice. *J Mol Neurosci*. 2002; 18:151–165. [PubMed: 11931346]
- Holmes PV, Blanchard DC, Blanchard RJ, Brady LS, Crawley JN. Chronic social stress increases levels of preprogalanin mRNA in the rat locus coeruleus. *Pharmacology Biochemistry and Behavior*. 1995; 50:655–660.
- Holmes PV, Koprivica V, Chough E, Crawley JN. Intraventricular administration of galanin does not affect behaviors associated with locus coeruleus activation in rats. *Peptides*. 1994; 15:1303–1308. [PubMed: 7531843]
- Holmes PV, Yoo HS, Dishman RK. Voluntary exercise and clomipramine treatment elevate preprogalanin mRNA levels in the locus coeruleus in rats. *Neurosci Lett*. 2006a; 408:1–4. [PubMed: 16996684]
- Holmes PV, Yoo HS, Dishman RK. Voluntary exercise and clomipramine treatment elevate preprogalanin mRNA levels in the locus coeruleus in rats. *Neuroscience Letters*. 2006b; 408:1–4. [PubMed: 16996684]
- Jansson A, Fuxe K, Eneroth P, Agnati LF. Centrally administered galanin reduces dopamine utilization in the median eminence and increases dopamine utilization in the medial neostriatum of the male rat. *Acta Physiol Scand*. 1989; 135:199–200. [PubMed: 2466392]
- Juhasz G, Hullam G, Eszlari N, Gonda X, Antal P, Anderson IM, Hokfelt TG, Deakin JF, Bagdy G. Brain galanin system genes interact with life stresses in depression-related phenotypes. *Proc Natl Acad Sci U S A*. 2014
- Karlsson RM, Holmes A. Galanin as a modulator of anxiety and depression and a therapeutic target for affective disease. *Amino Acids*. 2006; 31:231–239. [PubMed: 16733616]
- Karlsson RM, Holmes A, Heilig M, Crawley JN. Anxiolytic-like actions of centrally-administered neuropeptide Y, but not galanin, in C57BL/6J mice. *Pharmacol Biochem Behav*. 2005; 80:427–436. [PubMed: 15740785]
- Krishnan V, Han MH, Graham DL, Berton O, Renthal W, Russo SJ, Laplant Q, Graham A, Lutter M, Lagace DC, Ghose S, Reister R, Tannous P, Green TA, Neve RL, Chakravarty S, Kumar A, Eisch AJ, Self DW, Lee FS, Tamminga CA, Cooper DC, Gershenfeld HK, Nestler EJ. Molecular adaptations underlying susceptibility and resistance to social defeat in brain reward regions. *Cell*. 2007; 131:391–404. [PubMed: 17956738]
- Kuteeva E, Wardi T, Lundstrom L, Sollenberg U, Langel U, Hokfelt T, Ogren SO. Differential role of galanin receptors in the regulation of depression-like behavior and monoamine/stress-related genes at the cell body level. *Neuropsychopharmacology*. 2008; 33:2573–2585. [PubMed: 18172432]
- Lehmann ML, Herkenham M. Environmental enrichment confers stress resiliency to social defeat through an infralimbic cortex-dependent neuroanatomical pathway. *J Neurosci*. 2011; 31:6159–6173. [PubMed: 21508240]
- Leuner B, Shors TJ. Stress, anxiety, and dendritic spines: What are the connections? *Neuroscience*. 2012
- Lewis MJ, Johnson DF, Waldman D, Leibowitz SF, Hoebel BG. Galanin microinjection in the third ventricle increases voluntary ethanol intake. *Alcohol Clin Exp Res*. 2004; 28:1822–1828. [PubMed: 15608598]
- Li Y, Luikart BW, Birnbaum S, Chen J, Kwon CH, Kernie SG, Bassel-Duby R, Parada LF. TrkB regulates hippocampal neurogenesis and governs sensitivity to antidepressive treatment. *Neuron*. 2008; 59:399–412. [PubMed: 18701066]
- Llorens-Martin M, Torres-Aleman I, Trejo JL. Exercise modulates insulin-like growth factor 1-dependent and -independent effects on adult hippocampal neurogenesis and behaviour. *Mol Cell Neurosci*. 2010; 44:109–117. [PubMed: 20206269]
- Lu X, Barr AM, Kinney JW, Sanna P, Conti B, Behrens MM, Bartfai T. A role for galanin in antidepressant actions with a focus on the dorsal raphe nucleus. *Proc Natl Acad Sci U S A*. 2005; 102:874–879. [PubMed: 15647369]
- Maier SF, Watkins LR. Role of the medial prefrontal cortex in coping and resilience. *Brain Res*. 2010; 1355:52–60. [PubMed: 20727864]

- Maniam J, Morris MJ. Voluntary exercise and palatable high-fat diet both improve behavioural profile and stress responses in male rats exposed to early life stress: Role of hippocampus. *Psychoneuroendocrinology*. 2010
- Masini CV, Holmes PV, Freeman KG, Maki AC, Edwards GL. Dopamine overflow is increased in olfactory bulbectomized rats: an in vivo microdialysis study. *Physiol Behav*. 2004; 81:111–119. [PubMed: 15059690]
- McDonald MP, Crawley JN. Galanin receptor antagonist M40 blocks galanin-induced choice accuracy deficits on a delayed-nonmatching-to-position task. *Behav Neurosci*. 1996; 110:1025–1032. [PubMed: 8919005]
- McEwen BS, Morrison JH. The brain on stress: vulnerability and plasticity of the prefrontal cortex over the life course. *Neuron*. 2013; 79:16–29. [PubMed: 23849196]
- Murray PS, Groves JL, Pettett BJ, Britton SL, Koch LG, Dishman RK, Holmes PV. Locus coeruleus galanin expression is enhanced after exercise in rats selectively bred for high capacity for aerobic activity. *Peptides*. 2010; 31:2264–2268. [PubMed: 20850488]
- New AS, Fan J, Murrough JW, Liu X, Liebman RE, Guise KG, Tang CY, Charney DS. A functional magnetic resonance imaging study of deliberate emotion regulation in resilience and posttraumatic stress disorder. *Biol Psychiatry*. 2009; 66:656–664. [PubMed: 19589502]
- Nordstrom O, Melander T, Hokfelt T, Bartfai T, Goldstein M. Evidence for an inhibitory effect of the peptide galanin on dopamine release from the rat median eminence. *Neurosci Lett*. 1987; 73:21–26. [PubMed: 2436097]
- O'Neal HA, Van Hoomissen JD, Holmes PV, Dishman RK. Prepro-galanin messenger RNA levels are increased in rat locus coeruleus after treadmill exercise training. *Neuroscience Letters*. 2001; 299:69–72. [PubMed: 11166940]
- Ogbonmwan YE, Sciolino NR, Groves-Chapman JL, Freeman KG, Edwards GL, Holmes PV, Weinshenker D. The galanin receptor agonist galanin attenuates cocaine-induced reinstatement and dopamine overflow in the frontal cortex. *Addiction Biology*. 2014
- Pascucci T, Ventura R, Latagliata EC, Cabib S, Puglisi-Allegra S. The medial prefrontal cortex determines the accumbens dopamine response to stress through the opposing influences of norepinephrine and dopamine. *Cereb Cortex*. 2007; 17:2796–2804. [PubMed: 17322559]
- Paxinos, G.; Watson, C. The rat brain: in stereotaxic coordinates. Academic Press; San Diego: 1998.
- Peres JF, Foerster B, Santana LG, Ferreira MD, Nasello AG, Savoia M, Moreira-Almeida A, Lederman H. Police officers under attack: resilience implications of an fMRI study. *J Psychiatr Res*. 2011; 45:727–734. [PubMed: 21159352]
- Primeaux SD, Holmes PV. Olfactory bulbectomy increases met-enkephalin- and neuropeptide-Y-like immunoreactivity in rat limbic structures. *Pharmacol Biochem Behav*. 2000; 67:331–337. [PubMed: 11124398]
- Radley JJ, Morrison JH. Repeated stress and structural plasticity in the brain. *Ageing Res Rev*. 2005; 4:271–287. [PubMed: 15993654]
- Ramkumar K, Srikumar BN, Venkatasubramanian D, Siva R, Shankaranarayana Rao BS, Raju TR. Reversal of stress-induced dendritic atrophy in the prefrontal cortex by intracranial self-stimulation. *J Neural Transm*. 2011
- Reiss JI, Dishman RK, Boyd HE, Robinson JK, Holmes PV. Chronic activity wheel running reduces the severity of kainic acid-induced seizures in the rat: possible role of galanin. *Brain Res*. 2009; 1266:54–63. [PubMed: 19248773]
- Sasse SK, Greenwood BN, Masini CV, Nyhuis TJ, Fleshner M, Day HE, Campeau S. Chronic voluntary wheel running facilitates corticosterone response habituation to repeated audiogenic stress exposure in male rats. *Stress*. 2008; 11:425–437. [PubMed: 19065456]
- Sciolino NR, Dishman RK, Holmes PV. Voluntary exercise offers anxiolytic potential and amplifies galanin gene expression in the locus coeruleus of the rat. *Behavioural Brain Research*. 2012; 233:191–200. [PubMed: 22580167]
- Sciolino NR, Holmes PV. Exercise offers anxiolytic potential: A role for stress and brain noradrenergic-galaninergic mechanisms. *Neuroscience & Biobehavioral Reviews*. 2012; 36:1965–1984. [PubMed: 22771334]

- Stranahan AM, Khalil D, Gould E. Social isolation delays the positive effects of running on adult neurogenesis. *Nat Neurosci.* 2006; 9:526–533. [PubMed: 16531997]
- Stranahan AM, Khalil D, Gould E. Running induces widespread structural alterations in the hippocampus and entorhinal cortex. *Hippocampus.* 2007; 17:1017–1022. [PubMed: 17636549]
- Stranahan AM, Lee K, Mattson MP. Central mechanisms of HPA axis regulation by voluntary exercise. *Neuromolecular Med.* 2008; 10:118–127. [PubMed: 18273712]
- Trejo JL, Llorens-Martin MV, Torres-Aleman I. The effects of exercise on spatial learning and anxiety-like behavior are mediated by an IGF-I-dependent mechanism related to hippocampal neurogenesis. *Mol Cell Neurosci.* 2008; 37:402–411. [PubMed: 18086533]
- Tromp DP, Grupe DW, Oathes DJ, McFarlin DR, Hernandez PJ, Kral TR, Lee JE, Adams M, Alexander AL, Nitschke JB. Reduced structural connectivity of a major frontolimbic pathway in generalized anxiety disorder. *Arch Gen Psychiatry.* 2012; 69:925–934. [PubMed: 22945621]
- Tsuda K, Tsuda S, Nishio I, Masuyama Y, Goldstein M. Effects of galanin on dopamine release in the central nervous system of normotensive and spontaneously hypertensive rats. *Am J Hypertens.* 1998; 11:1475–1479. [PubMed: 9880130]
- U.S. Department of Health and Human Services. Physical activity guidelines advisory committee report. 2008.
- Van Hooymissen JD, Holmes PV, Zellner AS, Poudevigne A, Dishman RK. Effects of beta-adrenoreceptor blockade during chronic exercise on contextual fear conditioning and mRNA for galanin and brain-derived neurotrophic factor. *Behavioral Neuroscience.* 2004; 118:1378–1390. [PubMed: 15598146]
- Weiss JM, Bonsall RW, Demetrikopoulos MK, Emery MS, West CH. Galanin: a significant role in depression? *Ann N Y Acad Sci.* 1998; 863:364–382. [PubMed: 9928183]
- Zheng H, Liu Y, Li W, Yang B, Chen D, Wang X, Jiang Z, Wang H, Wang Z, Cornelisson G, Halberg F. Beneficial effects of exercise and its molecular mechanisms on depression in rats. *Behavioral Brain Research.* 2006; 168:47–55.

Highlights

- Exercise promotes resilience by blocking stress-induced behavior in anxiety tests
- Exercise also increases galanin in the locus coeruleus
- Repeated intracranial galanin mimics exercise-induced resilience
- Exercise or galanin treatment prevent dendritic spine loss in the mPFC after stress
- Repeated galanin receptor antagonism blocks resilience behavior induced by exercise

Figure 1. Exercise increases galanin in the locus coeruleus

Galanin levels were assessed in the brain and measured in rats with access to a running wheel or no wheel for 3 weeks (see Fig. 2). Shown are means $\pm$ SEM for (A) galanin protein obtained by ELISA in tissue containing the dorsal pons or ventral midbrain from exercise and sedentary rats ($n=4-5$) exposed to no shock. Also shown are (B) means $\pm$ SEM for galanin mRNA measured by ISH in tissue containing the locus coeruleus (LC) from exercise and sedentary rats ($n=10$) exposed to footshock, and (C) images of galanin mRNA in representative sagittal brain sections from exercise (top) and sedentary rats (bottom). Spectrum scale shows the intensity of (C) galanin mRNA expression. Insets show (C) galanin mRNA in the LC. * $p<0.05$ compared to sedentary.

Figure 2. Exercise blocks stress-induced behavior in an anxiety test, but this resilience behavior is not reversed by acute intra-cranial administration of M40

Rats with access to a running wheel or no wheel were administered ICV vehicle (dH₂O) or the galanin receptor antagonist M40 (6 nmol) immediately before exposure to footshock or no-shock during open field activity. Shown are (A) experimental timeline, means $\pm$ SEM for (B) open arm time and (C) open arm entries in the elevated plus maze, and (D) ambulatory distance during open field activity. N = 6 (sed-veh), 4 (ex-veh), 10 (sed-veh), 9 (ex-veh), and 5 (ex-M40). * $p < 0.05$ compared sed-veh rats exposed to footshock. Ex, exercise. Sed, sedentary. Veh, vehicle.

Figure 3. Exercise blocks stress-induced behavior and -evoked dopamine, effects that are mimicked by repeated intracranial galanin and reversed partially by M40
 Rats received daily administration of ICV vehicle (aCSF), galanin (3 nmol), or galanin receptor antagonist M40 (6 nmol) during exercise or sedentary conditions, and were exposed to footshock during microdialysis in an open field. Shown are (A) experimental timeline, means±SEM for (B) open arm time in the elevated plus maze and (C) cortical dopamine levels, and (D) probe placements. N = 9–16 (Sed Veh), 9–10 (Sed Gal), 8–10 (veh ex), and 5–6 (Ex M40). ** $p < 0.01$, * $p < 0.05$ compared to vehicle-treated sedentary. ++ $p < 0.01$, + $p < 0.05$ compared to galanin-treated sedentary. X $p < 0.05$ compared to vehicle-treated exercise. Ex, exercise. Gal, galanin. Sed, sedentary. Veh, vehicle.

Figure 4. Repeated intracranial galanin administration mimics exercise and prevents dendritic spine loss in mPFC pyramidal neurons after stress

Dendritic spine density was measured in mPFC pyramidal neurons in Golgi-processed tissue (imaged at 100x) obtained from a subset of rats exposed to footshock or no-shock (see Fig. 3). Shown are (A) regions of the mPFC where neurons were selected, and (B) pyramidal neuron cartoon showing the apical and basal segments assessed for spine density. Shown also are (C) mean $\pm$ SEM for apical spine number, (D) representative apical segments, (E) means $\pm$ SEM for basal spine number, and (F) representative basal segments. $N = 6$ (sed-veh no shock), 7 (sed-veh shock), 3 (sed-gal shock), and 6 (ex-veh shock). # $p < 0.05$ compared to no-shock exposed vehicle-treated sedentary. ** $p < 0.01$, * $p < 0.05$ compared to shock exposed vehicle-treated sedentary. Ex, exercise. Gal, galanin. IL, infralimbic. NS, no-shock. PrL, prelimbic. Sed, sedentary. Veh, vehicle.