Coronary Artery Aneurysms in a 15-Year-Old Boy

Stephen J. D'Auria, MD
Meghan Trojan Borden, DO
Sandeep M. Patel, MD
Conrad Smith, MD

A 15-year-old boy with a history of Kawasaki disease presented after an exercise nuclear stress test (performed before a contemplated weight-loss program) revealed a reversible inferolateral and apical defect (left ventricular ejection fraction, 0.53). A coronary angiogram showed the right coronary artery with an $11.5 \times 6 \times 5$ -mm proximal aneurysm, proximal 50% stenosis, and mid 70% stenosis; the left anterior descending coronary artery (LAD) with a proximal 60% stenosis followed by an aneurysm; and the left circumflex coronary artery with a small aneurysm of the proximal segment (Fig. 1). We placed drug-eluting stents in the proximal and mid right coronary artery and in the proximal LAD lesion (Fig. 2). The patient did well and tolerated cardiac rehabilitation. A stress test one year after the procedure revealed no inducible ischemia.

Comment

Kawasaki disease is an acute vasculitis that afflicts children; coronary artery aneurysms (CAAs) are a known sequela.^{1,2} Our patient's largest aneurysm (>8 mm) qualifies as a "giant" CAA. In an analysis of 76 similar patients,³ the survival rates were 88% at both 20 and 30 years; deaths were caused by ischemic events. Patients who develop substantial stenotic lesions need either percutaneous coronary intervention or coronary artery bypass grafting, depending on their coronary anatomy.^{2,3} Surgery is typically recommended for these conditions: obstruction of the left main or more than one major coronary artery, proximal LAD obstruction, compromised collateral flow, or recurrent myocardial infarction.⁴

To reduce the risk of CAAs, patients with Kawasaki disease should initially be treated with aspirin and intravenous immunoglobulin G.⁴ After the diagnosis of Kawasaki disease has been established, the risk level and long-term medical therapy should be determined. Noninvasive coronary imaging can be performed first, followed (if neces-

Section Editor:

Raymond F. Stainback, MD,
Department of Adult
Cardiology, Texas Heart
Institute, 6624 Fannin St.,
Suite 2480, Houston, TX
77030

From: Department of Internal Medicine (Drs. Borden and D'Auria) and the Heart and Vascular Institute (Drs. Patel and Smith), University of Pittsburgh Medical Center, Pittsburgh, Pennsylvania 15213

Address for reprints:

Stephen J. D'Auria, MD,
Department of Internal
Medicine, University of
Pittsburgh Medical Center,
200 Lothrop St., Pittsburgh,
PA 15213

E-mail: daurias@upmc.edu

© 2014 by the Texas Heart®
Institute, Houston

Fig. 1 Diagnostic coronary angiograms. **A)** The right coronary artery has a large proximal aneurysm and hemodynamically significant mid-vessel stenoses. **B)** The left anterior descending coronary artery has a proximal stenosis and aneurysm; the left circumflex coronary artery has a small proximal aneurysm.

Fig. 2 Coronary angiograms after drug-eluting stent placement. **A)** The right coronary artery shows improvement of the stenoses (arrows). **B)** The left anterior descending artery shows no residual stenosis (arrow).

sary) by coronary angiography 6 to 12 months after the inflammatory period.⁴

References

1. Burns JC, Shike H, Gordon JB, Malhotra A, Schoenwetter M, Kawasaki T. Sequelae of Kawasaki disease in adolescents and young adults. *J Am Coll Cardiol* 1996;28(1):253-7.
2. Daniels LB, Gordon JB, Burns JC. Kawasaki disease: late cardiovascular sequelae. *Curr Opin Cardiol* 2012;27(6):572-7.
3. Suda K, Iemura M, Nishiono H, Teramachi Y, Koteda Y, Kishimoto S, et al. Long-term prognosis of patients with Kawasaki disease complicated by giant coronary aneurysms: a single-institution experience. *Circulation* 2011;123(17):1836-42.
4. Newburger JW, Takahashi M, Gerber MA, Gewitz MH, Tani LY, Burns JC, et al. Diagnosis, treatment, and long-term management of Kawasaki disease: a statement for health professionals from the Committee on Rheumatic Fever, Endocarditis and Kawasaki Disease, Council on Cardiovascular Disease in the Young, American Heart Association. *Circulation* 2004;110(17):2747-71.