Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/jobcr

Case Report

Prosthetic rehabilitation of a child with velopharyngeal dysfunction – A case report

Neha Sikka ^{a,*}, Reeta Jain ^b, Ashutosh Kaushik ^c, Reena Rani ^d

^a Senior Lecturer, Department of Prosthodontics and Crown and Bridge, Eklavya Dental College, Kotputli, Jaipur, Rajasthan, India

^b Professor and Head, Department of Prosthodontics and Crown and Bridge, Genesis Institute of Dental Sciences and Research, Ferozepur, Punjab, India

^c PG Student, Department of Orthodontics, SGT Dental College, Budhera, Gurgaon, Haryana, India

^d PG Student, Department of Pedodontics, Post Graduate Institute of Dental Sciences, Rohtak, Haryana, India

ARTICLE INFO

Article history:

Received 5 March 2014

Accepted 9 July 2014

Available online 8 August 2014

Keywords:

Velopharyngeal dysfunction

Velopharyngeal inadequacy

Palatal lift appliance

Speech therapy

ABSTRACT

The speech of person is a unique mode of expression for an individual. Alterations or deviation from normal speech leading to nasal and unintelligible speech have enormous effects on the psychological well being of a person as it hampers social interactions. The anatomic or functional anomalies occurring as a result of congenital or traumatic defects leading to velopharyngeal dysfunctions influence the speech by making it completely atypical. The cleft palate patients may have residual velopharyngeal inadequacy even after surgical repair. A removable palatal lift appliance in conjunction with speech therapy is a viable treatment option for such patients as it is economical, easy to fabricate and easy to use.

Copyright © 2014, Craniofacial Research Foundation. All rights reserved.

1. Introduction

Speech being the oldest media of communication and expression of thoughts has a great effect on the on the psychological status and social interactions. Faulty unintelligible speech is never desirable even for short period. Velopharynx, a muscular sphincter located between the nasopharynx and the oropharynx regulates passage of air during speech.¹ The complete closure of this muscular valve is required for normal physiologic activities like speaking, swallowing, whistling, blowing and sucking. The closure of the sphincter usually involves the movement of soft palate,

lateral and posterior pharyngeal wall.² Velopharyngeal dysfunction due to non closure of sphincter can either be the result of (i) anatomic deficiency, known as velopharyngeal insufficiency, or (ii) inadequate movement called as velopharyngeal incompetence.³ Velopharyngeal dysfunctions make the speech completely atypical. The constant failure to separate the nasal and oral cavities produces hypernasality in speech, compensatory articulation and nasal air leakage.¹

The major causes of velopharyngeal dysfunction include patients with Cleft palate, Cerebrovascular accidents, brain stem tumors and traumatic injuries to neuromotor system. It can be treated by surgery or prosthetic rehabilitation.

* Corresponding author. Tel.: +91 9416330633.

E-mail address: neha.nehasikka@gmail.com (N. Sikka).

<http://dx.doi.org/10.1016/j.jobcr.2014.07.002>

2212-4268/Copyright © 2014, Craniofacial Research Foundation. All rights reserved.

Pharyngeal flap surgery has been a standard treatment approach for years. After surgical correction 20–34% of patients do not acquire desirable amount of velopharyngeal competence due to limited mobility of the pharyngeal flap. An adjunctive prosthodontic therapy by palatal lift appliance is necessary in such cases.⁴ The mechanism of action of the Palatal lift appliances is through elevation of the soft palate in upward and backward direction thereby creating a pressure on the lateral portion of the elevated soft palate, producing a proper velopharyngeal closure.^{2–6}

The present case describes the fabrication of a palatal lift appliance for the correcting the nasality of speech in a child with cleft palate.

2. Case report

A 5-year-old female patient referred to us by a plastic surgeon, reported with the chief complaint of nasality in voice. The history of the patient revealed that she has been operated for cleft palate two weeks back. The sutures in the soft palatal area could be seen, there was complete closure of the defect and the wound was healing uneventfully [Fig. 1a]. The fabrication of a removable tooth supported Palatal lift appliance was chosen to alleviate her problems in speech. The treatment plan with its advantages and disadvantages was explained to the patient's parents and consent was taken.

Step by step procedure for fabrication of palatal lift appliance:

After thorough clinical examination, impression was made in a modified stock tray [Fig. 1b] using irreversible hydrocolloid (Zelgan Plus, Dentsply, USA) and cast prepared.

Stainless steel wire clasps and loop were constructed to be embedded in appliance fabricated [Fig. 1c] using self cure clear acrylic (RR Cold Cure, DPI, India).

The appliance with wire loops was tried in [Fig. 1d]. Necessary adjustments were made to ensure complete seating. For the generation of palatal lift, low fusing impression compound (Green Sticks, DPI, India) was applied to the loop in the posterior part of the appliance in the desired shape and cooled immediately [Fig. 2a].

The hardened green stick displaces the soft palate upwards to produce the lift. Following this green stick wax was added incrementally and patient was asked to perform some physiologic movements including certain phonetic syllables like B and P, blowing through nose against resistance and bending head from side to side. The addition of increments of green stick was terminated at the point where there was improvement in nasality of voice and nasal air leakage [Fig. 2b].

After the generation of sufficient soft palatal lift, the area of appliance consisting of green stick was replaced by clear acrylic resin (RR Cold Cure, DPI, India). The insertion of appliance was done after finishing and polishing [Fig. 2c]. The appliance was checked for nasal air leak by placing a finger near the nose when the patient blows out through mouth after pursing her lips. A lateral cephalogram was taken with appliance in position for determination of generation of lift. The technique of insertion and removal and maintenance was explained to patient and her parents. As the child may not wear the appliance, the parents were advised to ensure the usage during waking hours. Patient was referred to a speech therapist. The speech therapy involved 2 phases, each for a duration of 3 weeks. In the first phase, elimination of the compensatory articulations/inadequate sounds was done by modulating oral and nasal air pressure.^{4,7} The second phase

Fig. 1 – a. Intraoral view showing recent surgical closure for cleft palate. **b.** Preliminary impression made using irreversible hydrocolloid. **c.** Acrylic framework with wire extension fabricated. **d.** Acrylic framework with wire extension placed intraorally.

Fig. 2 – a. Fabrication of palatal lift by addition of green stick on distal portion of appliance. b. Final impression for palatal lift generation. c. Processed palatal lift appliance.

involved the usage of audio-visual and tactile feedback to rectify the anatomical location for production of adequate/target sounds and to monitor their reproduction during normal conversational speech.^{4,7} Constant recalls were maintained to assess the outcome of the therapy and the patient was motivated to do the exercises at home. A marked reduction in nasality of speech and compensatory articulations after 2 months was observed.

3. Discussion

The case report narrates the management of a five year old patient with unintelligible (hypernasal) speech recently operated for cleft palate. Adequate velopharyngeal closure is mandatory for production of normal speech. Children suffering from velopharyngeal dysfunction develop undesirable compensatory mechanisms (in order to deal with the reduced intraoral pressure) at the learning stage.^{4,6,7} It is difficult to get rid of such habits later in life. Thus, if treatment (involving surgical correction, appliance and speech therapy) is done at an early age, establishment of such mechanisms can be avoided and prognosis is improved.^{4,6,7} Premkumar S reported the use of removable palatal lift appliance in a 9 year old patient with chief complaint of nasal speech and history of surgically repaired cleft palate.⁸ Raju H et al⁴ and Aboloyoun AI et al⁹ conducted studies to assess the effectiveness of palatal lift appliances in patients who had previously undergone surgery for cleft palate in different age groups (16–26 years and 8–10 years respectively). The results of both the studies showed significant improvement in speech after treatment with a palatal lift appliance.

The fabrication of a tooth supported removable palatal lift appliance was planned in our case. The children at the age of 5 are mature enough to adapt and care for a removable appliance (Kupietzky A).¹⁰ The basic requirement for palatal lift appliance is retention.² Maximum number of posterior abutments should be strategically used to avoid the disadvantages of cantilever.³ The outcome of the treatment mainly depends on the efficacy of displacement of the soft palate,⁸ degree of

removal of compensatory articulations and the consistent production of target sound in normal conversational speech.^{4,7} A well fabricated palatal lift appliance altogether prevents the air leakage from nostrils which imparts a beneficial effect on the speech therapy. However, this appliance has certain limitations. Like any other removable appliance it is dependent on the compliance of the patient. A lot of effort was done to make the insertion, removal and wearing of the appliance very comfortable. As our patient was very young, the parents were asked to teach the child the importance of wearing the appliance. The parents and the school teachers checked the presence of appliance in the oral cavity. The size of the appliance was not small. So, the chances of swallowing the appliance were feeble. But still it was made sure that the patient was not wearing the appliance during any physical sports activities.

4. Conclusion

A multidisciplinary approach involving surgical repair, palatal lift appliance with adjuvant speech therapy at the earliest possible age is the most accepted mode of treatment for patients with velopharyngeal dysfunction.

Conflicts of interest

All authors have none to declare.

REFERENCES

1. Woo AS. Velopharyngeal dysfunction. *Semin Plast Surg.* 2012;26:170–177.
2. Pinto JHN, Krook MIP. Evaluation of palatal prosthesis for the treatment of velopharyngeal dysfunction. *J Appl Oral Sci.* 2003;3:192–197.

3. Schaefer KS, Taylor TD. Clinical application of palatal lift. In: Thomas D, ed. *Maxillofacial Prosthetics*. Chicago: Quintessence Publishing Co, Inc; 2000:133–142.
4. Raju H, Padmanabhan TV, Narayan A. Effect of palatal lift prosthesis in individuals with velopharyngeal incompetence. *Int J Prosthodont*. 2009;22:579–585.
5. Kipfmüller L, Lang B. Treating velopharyngeal inadequacies with a palatal lift prosthesis. *J Prosthet Dent*. 1972;27:63.
6. Aggarwal KK, Singh BP, Chand P, Patel C. Impact of delayed prosthetic treatment of velopharyngeal insufficiency on quality of life. *Indian J Dent Res*. 2011;22:356–358.
7. Bispo NHM, Whitaker ME, Aferri HC, Neves JDA, Dutka JCR, Pegoraro-Krook MI. Speech therapy for compensatory articulations and velopharyngeal function: a case report. *J Appl Oral Sci*. 2011;19:679–684.
8. Premkumar S. Clinical application of palatal lift appliance in velopharyngeal incompetence. *J Indian Soc Pedod Prev Dent*. 2011;29:70–73.
9. Aboloyoun AI, Ghorab S, Farooq MU. Palatal lifting prosthesis and velopharyngeal insufficiency: preliminary report. *Acta Med Acad*. 2013;42:55–60.
10. Kupietzky A. The treatment and long term management of severe multiple avulsions of primary teeth in a 19-month-old child. *Pediatr Dent*. 2001;23:517–521.

Sub - Mucous fibrosis Oral Leukoplakia Lichen planus

Introducing
1st Time Clinical cure

TurMix

TABLETS
Curcumin 300mg (95% Standardized)+ Piperine 5mg

USFDA Approved

DNA Protecting Fibrinolytic Anti-inflammatory Anti-oxidant

Safe up to 8g/day

Also available

Heal plus

Antiseptic Gel

Turmeric
Tea Tree Oil
Cedar Wood Oil
Aloe Vera

Heal Wounds Naturally

sanat
PRODUCTS LTD.

3rd Floor, Sagar Plaza, District Centre, Laxmi Nagar, Vikas Marg, Delhi 110092

Tel. No. : 011-22518794/95/96, 2205 0535/37

For Availability or Trade Enquires : 93507-98611 / 8009-770777

E-mail : nkhan@sanat.co.in, Website : www.sanat.co.in

For Information of Registered medical practitioners, Hospitals or Laboratories: