Published in final edited form as:

Neuropharmacology. 2015 January ; 88: 171–179. doi:10.1016/j.neuropharm.2014.07.019.

Defects in dendrite and spine maturation and synaptogenesis associated with an anxious-depressive-like phenotype of GABA_A receptor-deficient mice

Zhen Ren^{a,c,*}, Nadia Sahir^{a,1,*}, Shoko Murakami^a, Beth A. Luellen^a, John C. Earnheart^b, Rachnanjali Lal^{a,2}, Ju Young Kim^{d,e}, Hongjun Song^{d,e,f}, and Bernhard Luscher^{a,b,c}

^aDepartment of Biology, The Pennsylvania State University, University Park, PA 16802

^bDepartment of Biochemistry and Molecular Biology, The Pennsylvania State University, University Park, PA 16802

^cCenter for Molecular Investigation of Neurological Disorders, The Huck Institutes of the Life Sciences, The Pennsylvania State University, University Park, PA 16802

^dInstitute for Cell Engineering, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA

^eDepartment of Neurology, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA

^fThe Solomon Snyder Department of Neuroscience, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA

Abstract

Mice that were rendered heterozygous for the $\gamma 2$ subunit of GABA_A receptors ($\gamma 2^{+/-}$ mice) have been characterized extensively as a model for major depressive disorder. The phenotype of these mice includes behavior indicative of heightened anxiety, despair, and anhedonia, as well as defects in hippocampus-dependent pattern separation, HPA axis hyperactivity and increased responsiveness to antidepressant drugs. The $\gamma 2^{+/-}$ model thereby provides strong support for the GABAergic deficit hypothesis of major depressive disorder. Here we show that $\gamma 2^{+/-}$ mice additionally exhibit specific defects in late stage survival of adult-born hippocampal granule cells, including reduced complexity of dendritic arbors and impaired maturation of synaptic spines. Moreover, cortical $\gamma 2^{+/-}$ neurons cultured *in vitro* show marked deficits in GABAergic

© 2014 Elsevier Ltd. All rights reserved.

Corresponding Author: Bernhard Luscher, Ph. D., Professor, Dept. of Biology and Dept. of Biochemistry and Molecular Biology, The Pennsylvania State University, 301 Life Sciences Building, University Park, PA 16801, BXL25@psu.edu, Phone: +1-814-865 5549.

¹Current address: Sanford Children's Health, Research Center, 2301 East 60th Street North, Sioux Falls, SD 57104

²Current address: Johns Hopkins Hospital, 1800 Orleans St., Baltimore, MD 21287

*These authors contributed equally to the work presented

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The authors declare no competing interests.

innervation selectively when grown under competitive conditions that may mimic the environment of adult-born hippocampal granule cells. Finally, brain extracts of $\gamma 2^{+/-}$ mice show a numerical but insignificant trend ($p = 0.06$) for transiently reduced expression of brain derived neurotrophic factor (BDNF) at three weeks of age, which might contribute to the previously reported developmental origin of the behavioral phenotype of $\gamma 2^{+/-}$ mice. The data indicate increasing congruence of the GABAergic, glutamatergic, stress-based and neurotrophic deficit hypotheses of major depressive disorder.

Keywords

Neurogenesis; hippocampal synaptogenesis; synaptic spines; animal model of anxious depression; postnatal development

1. INTRODUCTION

Molecular mechanisms that underlie the pathoetiology of major depressive disorder (MDD) remain poorly understood. However, significant evidence suggests that deficits in GABAergic transmission may play a key role in MDD. The evidence from patients pointing to compromised GABAergic transmission in MDD includes reduced brain concentrations of GABA (Sanacora *et al.*, 1999; Hasler *et al.*, 2007; Gabbay *et al.*, 2012), reduced expression of the GABA-synthesizing enzyme glutamic acid decarboxylase (GAD67) (Karolewicz *et al.*, 2010; Guilloux *et al.*, 2012), altered expression of GABA_A receptors (GABA_ARs) (Merali *et al.*, 2004; Choudary *et al.*, 2005; Sequeira *et al.*, 2007; Klempan *et al.*, 2009; Klumpers *et al.*, 2010), compromised function or loss of GABAergic interneurons (Rajkowska *et al.*, 2007; Maciag *et al.*, 2010; Sibille *et al.*, 2011) and marked functional deficits in cortical inhibition (Levinson *et al.*, 2010). Conversely, antidepressant drugs (Sanacora *et al.*, 2002; Kucukbrahimoglu *et al.*, 2009) and electroconvulsive therapy (Sanacora *et al.*, 2003) can normalize the reduced GABA concentrations in brain and plasma of MDD patients [for review see (Croarkin *et al.*, 2011; Luscher *et al.*, 2011)].

In developing neurons, GABA and GABA_ARs act in concert with brain-derived neurotrophic factor (BDNF) and its receptor tropomyosin receptor kinase (TrkB) to form an interdependent positive feedback loop that is particularly important for maturation of neuronal dendrites and GABAergic synapses (Rico *et al.*, 2002; Chen *et al.*, 2011; Porcher *et al.*, 2011; Waterhouse *et al.*, 2012). Consistent with these studies, reduced expression of BDNF and TrkB are implicated in MDD based on analyses of postmortem brain of suicide victims (Dwivedi *et al.*, 2003; Dwivedi *et al.*, 2005; Karege *et al.*, 2005) and depressed subjects (Tripp *et al.*, 2012).

Mice that were rendered heterozygous for the $\gamma 2$ subunit of GABA_ARs ($\gamma 2^{+/-}$ mice) exhibit behavioral, cognitive, neuroendocrine, and pharmacological aberrations expected of a mouse model of major depressive disorder [for review see (Luscher *et al.*, 2011; Smith & Rudolph, 2012)]. This phenotype has a developmental origin in-between the third and fifth postnatal week of mice (Earnheart *et al.*, 2007; Shen *et al.*, 2012). Moreover, the phenotype of $\gamma 2^{+/-}$ mice includes reduced survival of adult-born hippocampal neurons, which are thought to serve as cellular substrates of antidepressant drug action (Earnheart *et al.*, 2007; Samuels &

Hen, 2011). Adult-born neurons are also critically important for pattern separation and completion (Clelland *et al.*, 2009; Sahay *et al.*, 2011), a cognitive measure that is compromised in MDD (Mogg *et al.*, 2006; Eley *et al.*, 2008; Leal *et al.*, 2014). In the $\gamma 2^{+/-}$ model a defect in pattern separation is illustrated by a significant negative bias in an ambiguous cue discrimination task (Crestani *et al.*, 1999).

Functional interactions between GABA_ARs and BDNF in neural maturation are consistent with the neurotrophic deficit hypothesis of MDD (Duman *et al.*, 1997; Duman & Monteggia, 2006). However, whether the phenotype of $\gamma 2^{+/-}$ mice involves defects in neural maturation and synaptogenesis is not yet known. To address this issue we here have extended our analyses of adult-born hippocampal neurons in the $\gamma 2^{+/-}$ model mouse. We confirm that granule cell progenitors of $\gamma 2^{+/-}$ mice proliferate at normal rates. We also show that they migrate normally but then fail to survive between two and three weeks after exit from the cell cycle. Failure to survive is reflected in significant defects in dendrite and spine maturation of adult-born granule cells. Moreover, experiments in cultured neurons indicate that $\gamma 2$ subunit-deficient neurons exhibit marked deficits in synaptogenesis when these neurons are grown in competition with WT neurons, i. e. conditions that are reminiscent of the competitive environment of adult-born hippocampal neurons. Lastly, we provide evidence that GABA_AR deficits of $\gamma 2^{+/-}$ mice may cause developmental reductions of BDNF expression that could contribute to the developmental depression-related phenotype of $\gamma 2^{+/-}$ mice.

2. EXPERIMENTAL PROCEDURES

2.1. Production and husbandry of mice

GABA_AR $\gamma 2$ subunit heterozygous ($\gamma 2^{+/-}$) mice used for this study were backcrossed onto the 129 $\times$ 1/SvJ genetic background for >40 generations, (Gunther *et al.*, 1995; Crestani *et al.*, 1999). They were produced in our own breeding colony as littermates by crossing WT and $\gamma 2^{+/-}$ mice, genotyped at the time of weaning using PCR of tail biopsies (Alldred *et al.*, 2005) and kept on a standard 12h:12h light-dark cycle with food and water available ad libitum. GFP-transgenic mice (Hadjantonakis *et al.*, 1998) carrying a transgene encoding Enhanced Green Fluorescent Protein (Clontech) driven by the ubiquitously active chicken β -actin promoter and CMV intermediate early enhancer were obtained from JAX mice (Stock # 003116, Jackson Laboratory, Bar Harbor, ME). All animal experiments were approved by the Institutional Animal Care and Use Committee (IACUC) of the Pennsylvania State University and performed in accordance with relevant guidelines and regulations of the National Institutes of Health. All efforts were made to minimize animal suffering, to reduce the number of animals used, and to utilize *in vitro* alternatives to *in vivo* techniques, when available.

2.2. Bromodeoxyuridine labeling and quantitation

Two different bromodeoxyuridine (BrdU) labeling protocols were used to quantify proliferation and neuronal survival of granule cells (Earnheart *et al.*, 2007). To quantify replicating cells, three-week-old mice were injected with a single dose of BrdU of 200 mg/kg (20mg/ml), and the brains were perfused and harvested 24h or 48h later. For

quantification of more mature neurons, three-week-old mice were administered BrdU (4×80 mg/kg i.p. at 2 h intervals, in saline at 8 mg/ml, pH 7.4) and the brains were harvested either 14d or 28d later. The mice were anesthetized with ketamine/xylazine/acepromazine (110, 20, and 3 mg/kg, i.p.) (Schein, Melville, NY), transcardially perfused with ice-cold phosphate buffered saline (PBS), followed by 4% paraformaldehyde in PBS, postfixed for 12 h in the same solution, and cryoprotected by incubation overnight in 30% sucrose. Serial coronal sections (35 μ m) through the hippocampus were cut from frozen brains with a sliding microtome. For quantitation of BrdU-labeled cells the sections were pretreated with 2N HCl for 30 min at 37°C and washed with 0.1M sodium borate and PBS and immunostained with monoclonal rat anti-BrdU antibody (1:500; Accurate Chemical, Westbury, NY). For double labeling with DCX or NeuN and BrdU the sections were first stained with goat anti-DCX (1:1000; Santa Cruz Biotechnology, Dallas, TX) or mouse anti-NeuN (1:1000; Chemicon, Temecula, CA), then fixed in 4% paraformaldehyde (20 min at room temperature), treated with 2N HCl, and stained with anti-BrdU as above. The sections were developed with Cy3-conjugated secondary anti rat (1:500; Molecular probes, Carlsbad, CA) and FITC-conjugated secondary anti guinea pig or mouse antibodies (Jackson ImmunoResearch, West Grove, PA). The number of BrdU or BrdU plus DCX/NeuN positive neurons in the subgranule and granule cell layer of confocal images were counted across sections of the entire bilateral hippocampus as described (Earnheart *et al.*, 2007).

2.3. Analyses of migration and reconstruction and analyses of dendritic arbors

Twelve-week-old female WT and $\gamma 2^{+/-}$ littermate mice were anesthetized with an overdose of Avertin [1.25% (w/v) 2,2,2-tribromoethanol in 5% 2-methyl-2-butanol] (both from Sigma-Aldrich) (375 mg/kg, 30 ml/kg i.p.) and perfused first briefly with PBS and then with 4% paraformaldehyde in 0.1 M phosphate buffer (pH 7.4). The brains were postfixed in the same solution for 24h, rinsed in PBS and sectioned coronally (50 μ m) using a vibratome (Vibratome; St Louis). Floating sections were stained with goat anti DCX (1:1000, Santa Cruz Biotechnology) in 2% goat serum in PBS for 48 h at 4°C and developed with Alexa488-conjugated secondary anti-goat antibody (1:500, Molecular Probes, Carlsbad, CA) for 1 h at room temperature. Stained sections were mounted cover-slipped on glass slides and imaged using a Olympus FV1000 laser scanning confocal microscope equipped with a 40X oil objective (N.A. = 1.3). The radial position of the DCX marked cell body within the granule cell layer was recorded and assigned a position within the inner third (inner granule cell layer), center third or outer third of the granule cell layer. Optical sections covering the full-length of the dendritic tree were collected using 1- μ m z-axis steps. For visualization of complete granule cells dendritic trees individual image stacks were superimposed digitally. Dendrites in 3D image stacks of complete dendritic arbors were traced by means of Neurolucida explorer software (MBF Bioscience, Williston, VT) and subjected to Sholl analyses to determine changes in dendritic complexity.

2.4. Analyses of synaptic spines

To visualize spines, murine Moloney leukemia virus-based CAG-GFP viral vector (Zhao *et al.*, 2006) essentially as described (Ge *et al.*, 2006). Briefly, GP2-293 cells (Clontech) were grown to 80% confluence in 15-cm Petri dishes using DMEM supplemented with 15% fetal calf serum and Pen/Strep. They were co-transfected with pCAG-GFP and pCMV-VSV-G

(both from Addgene, Cambridge, MA) (30 µg each/plate) using the Ca₂PO₄ co-precipitation method. The media were changed 8h later and the culture supernatant harvested and replaced 24, 36 and 48h after the last media change. Culture supernatants were pooled and stored in 50 ml conical tubes at -80°C. On the day of use a 40 ml aliquot of culture supernatant was thawed and centrifuged to remove cell debris (2000 rpm, 4°C, 5 min), filtered through a 0.45 µm filter cartridge and the virus concentrated by ultracentrifugation (25,000 rpm, 4°C, 90 min, SW 32 rotor). The virus pellets were resuspended in 10 µl sterile phosphate buffered saline to a concentration of approximately 10⁸ pfu/ml. Eight-week-old female WT and $\gamma 2^{+/-}$ mice were bilaterally injected with virus (0.5 µl per site) as previously described using the following coordinates relative to bregma: anteroposterior, -0.5 x d mm; lateral, -1.6 mm; ventral, -1.9 mm, with (d) being the distance between bregma and lambda. Two months after injection, the mice were anesthetized with an overdose of avertin (30 ml/kg), perfused and the brains post-fixed and sectioned as above. Stacks of confocal optical sections of GFP fluorescence in dendritic processes were acquired at 0.5 µm intervals with an Olympus FV1000 laser scanning confocal microscope equipped with a 60 × 1.42 N.A. objective and a digital zoom of 2.5. Maximum-density projections of z-stacks were used to classify spines as thin-, stubby-, or mushroom-shaped as described by Gonzalez-Burgos et al. (2000). The spine density of each dendritic fragment was calculated by dividing the manually counted number of each spine category by the length of the recorded dendritic segment (20 µm).

2.5. Analyses of cultured cortical neurons

Cultures of cortical neurons were generated as described (Alldred *et al.*, 2005) using mouse embryonic day 14–15 embryos produced by mating of GFP-transgenic WT and $\gamma 2^{+/-}$ mice. For mixed cultures, GFP-WT and $\gamma 2^{+/-}$ dissociated neurons were mixed 9:1 at the time of plating and then processed for immunostaining at DIV21 under permeabilized conditions as described (Alldred *et al.*, 2005). Guinea pig anti- $\gamma 2$ subunit (gift of J.M. Fritschy, University of Zurich, Switzerland, 1:1500) mAb GAD-6 (0.5 µg/ml; Developmental Studies Hybridoma Bank, University of Iowa, IA) and chicken anti GFP (Abcam, Cambridge, MA) were used as primary antibodies, and the stainings were developed using AlexaFluor 647 donkey anti-mouse (Molecular Probes, Eugene, OR), Cy3 donkey anti-guinea pig (Jackson ImmunoResearch) and Alexa 488 goat anti-chicken antibodies (Molecular Probes, Eugene, OR). Fluorescent images were captured and digitized with a Zeiss Axiophot 2 microscope equipped with a 40 × 1.3 N.A. objective and an ORCA-100 video camera linked to an OpenLab imaging system (PerkinElmer). The density and size of immunoreactive puncta and the degree of colocalization of pre- and postsynaptic markers were quantified as described (Alldred *et al.*, 2005; Fang *et al.*, 2006).

2.6. Quantitation of BDNF

Extraction of BDNF protein from mouse brain homogenates was performed using a high-recovery extraction method (Szapacs *et al.*, 2004) and used in combination with a BDNF enzyme-linked immunosorbent assay (ELISA) (Promega Co., Madison, WI) as recommended by the manufacturer. This protocol was validated by reproducible detection of a 50% reduction of BDNF in BDNF^{+/-} mice (Szapacs *et al.*, 2004). Absorbance of the colored reaction product was measured at 450 nm.

2.7. Data analysis

Simple two group comparisons were done by Mann Whitney tests or t-tests using GraphPad InStat version 3.00 for Macintosh, San Diego California USA. Sholl analyses of granule cell dendrites and spine density data were analyzed by repeated measures ANOVA and followed up by t-tests, using SPSS software (IBM). Probabilities < 0.05 were considered statistically significant.

3. RESULTS

3.1. Characterization of defects in neurogenesis of $\gamma 2^{+/-}$ mice

Previous characterization of 8–12 week old $\gamma 2^{+/-}$ mice revealed significantly reduced four-week-survival of adult-born hippocampal neurons (Earnheart *et al.*, 2007). In addition, cell-specific gene knockout of the $\gamma 2$ subunit in hippocampal radial glia-like astrocytes/stem cells (RGLs) revealed that tonic inhibition of these cells by $\gamma 2$ -containing GABA_ARs inhibits their mitotic activation and dedifferentiation into neuroblast precursor cells, as well as neural cell fate choice (Song *et al.*, 2012). To more precisely characterize the neurogenic consequences of modest GABA_AR reductions in the $\gamma 2^{+/-}$ -model we first analyzed the proliferation and survival of hippocampal neurons in greater detail. We previously had mapped the critical period for development of the anxious-depressive-like phenotype of $\gamma 2^{+/-}$ mice to the third and fourth postnatal week (Shen *et al.*, 2012). Therefore, we chose to conduct these studies in three-week old mice, a time during development when hippocampal neurogenesis is also more prolific than in adulthood. Three-week-old $\gamma 2^{+/-}$ mice and WT littermate controls were metabolically labeled with BrdU, and the number of BrdU-positive cells in the subgranule cell layer were analyzed at various time intervals thereafter. The number of BrdU-positive replicating cells in the SGZ of the DG of $\gamma 2^{+/-}$ mice was indistinguishable from that of WT, independent of whether the cells were quantitated 24 h ($\gamma 2^{+/-}$, 1704 ± 94.9 , WT, 1848 ± 125.0 , $U = 9.5$, $p > 0.05$, $n = 5$ per genotype, Mann Whitney test) (Fig 1D) or 48 h after BrdU labeling ($\gamma 2^{+/-}$, 3381.6 ± 331.5 ; WT, 3885.6 ± 260.9 , $U = 7.0$, $p > 0.05$, $n = 5$) (Fig. 1E). Thus, in contrast to the known role of $\gamma 2$ -containing GABA_ARs in activation of RGLs that is evident upon homozygous cell-specific knock out of the $\gamma 2$ gene (Song *et al.*, 2012), the initial activation and initial proliferation of these cells is not measurably affected by a global 50% reduction of the $\gamma 2$ subunit gene dosage.

To determine the developmental time point during which newborn hippocampal neurons fail to survive we analyzed the brains of BrdU-injected mice either 14 or 28 days post labeling and quantitated the number of BrdU-positive cells in the subgranular and granule cell layers that colocalized with different neuronal markers. At 14 days post labeling the number of cells that were positive for both BrdU and the immature neuron marker DCX was unaffected by genotype ($\gamma 2^{+/-}$, 8190 ± 1333.88 ; WT, 8691 ± 730.97 , $U = 8.0$, $p > 0.05$, $n = 4$, Fig 1F), thereby indicating normal initial differentiation of granule cell precursors. However, when the mice were harvested 28 days post labeling the number of BrdU-labeled cells that were positive for NeuN was markedly reduced compared to WT littermate controls (Fig. 1G, $\gamma 2^{+/-}$, 1248.0 ± 123.8 ; WT = 1779 ± 134.2 , $U = 0.0$, $p < 0.05$, $n = 4$, Mann-Whitney, Fig 1G). The

data indicate that granule cells of $\gamma 2^{+/-}$ mice fail to survive selectively during the late stage of differentiation.

3.2. $\gamma 2$ subunit-containing GABA_ARs regulate dendritic maturation

The reduction in the number of newborn $\gamma 2^{+/-}$ granule cells four weeks after BrdU labeling pointed to possible defects in late stage maturation of these neurons. To test this idea we labeled immature neurons of hippocampal brain sections of 12-week-old $\gamma 2^{+/-}$ and WT mice with DCX and then traced the structure of labeled dendrites in 3-D confocal image stacks using Neurolucida software. In these older mice the rate of neurogenesis is significantly lower than at three or eight weeks of age, a feature that facilitates morphological analyses of isolated dendritic trees of adult-born cells. Sholl analyses of DCX-positive granule cells revealed a significant reduction in the number of concentric circle crossings of granule cell dendrites of $\gamma 2^{+/-}$ vs. WT mice [two-way ANOVA with radial distance as within subject factor, $F(16,15) = 2.43$, $p < 0.05$, $n = 16$]. Posthoc t-tests revealed selective reductions in dendritic complexity in $\gamma 2^{+/-}$ vs. WT mice at a distance of 70 and 90 μm from the soma ($p < 0.05$ and $p < 0.001$, respectively) and a change in the opposite direction at 290 μm from the soma ($p < 0.01$, $n = 16$) (Fig. 2A, B). In dendritic segments between 70–90 μm from the soma, a reduced dendritic complexity was further reflected in a reduced number of branch points of $\gamma 2^{+/-}$ vs. WT neurons ($p < 0.05$, $n = 16$, t-test) (Fig. 2C). By contrast, the total length of dendritic trees was unaltered ($\gamma 2^{+/-}$, $823.8 \pm 57.6 \mu\text{m}$, WT $829 \pm 81.3 \mu\text{m}$).

GABAergic inputs have the potential to affect the radial migration of granule cell progenitors (Ge *et al.*, 2007a). To determine whether the phenotype of $\gamma 2^{+/-}$ mice includes altered migration of granule cell progenitors we divided the granule cell layer radially into equal inner, center, and outer thirds and counted the number of DCX-positive cell bodies in each compartment/brain section and averaged these numbers across sections. DCX-positive cells were rarely found in the outer section of the granule cell layer. Moreover, the average numbers of cells in the inner and center compartments of the granule cell layer were not measurably different from WT controls ($\gamma 2^{+/-}$ mice inner compartment: $88.7 \pm 9.1\%$ of WT, center compartment: $112.7 \pm 57.8\%$ of WT, $n = 3$ mice, $p > 0.5$ for both comparisons, t-tests).

To assess whether the maturational defect of granule cells included changes in spine density or morphology we injected the hippocampus of eight-week old $\gamma 2^{+/-}$ and WT mice with a GFP-encoding retrovirus (CAG-GFP) that specifically infects replicating cells. GFP-labeled spines of adult-born neurons along dendritic segments were manually sorted into thin-, stubby- and mushroom-shaped forms (Gonzalez-Burgos *et al.*, 2000)(Fig. 2D). Quantitation of the density of spines along dendrites revealed a significant genotype effect of GABA_AR deficits on spine density (two-way ANOVA with spine type as within subject factor, $F(2, 29) = 4.3$, $p < 0.05$). Posthoc analyses indicated a selective reduction in the density of mushroom-shaped spines in $\gamma 2^{+/-}$ vs. WT mice ($p = 0.014$, t-test) (Fig. 2E). Given the well-established correlation between density of spines and density of glutamatergic synapses the reduced spine density of $\gamma 2^{+/-}$ granule cell dendrites suggests that these neurons suffer from defects in glutamatergic innervation.

3.3. GABA_ARs regulate GABAergic innervation selectively under competitive conditions

Previous analyses of embryonic neurogenesis in $\gamma 2^{+/-}$ and $\gamma 2^{-/-}$ neurons revealed normal production, migration and survival of cortical neurons (Shen *et al.*, 2012). Moreover, $\gamma 2^{+/-}$ cortical cultures showed normal clustering of postsynaptic GABA_ARs and gephyrin (Essrich *et al.*, 1998). Similarly, conditional knock out of the $\gamma 2^{+/-}$ subunit in glutamatergic neurons of the mouse cortex did not measurably affect GABAergic innervation (Schweizer *et al.*, 2003). By contrast, sparse or mosaic homozygous knock out of the $\gamma 2$ subunit *in vivo* indicated that under some conditions GABA_ARs are important for normal GABAergic innervation (Li *et al.*, 2005; Frola *et al.*, 2013). We hypothesized that the maturational defect in hippocampal granule cells of $\gamma 2^{+/-}$ mice might reflect competition of immature neurons with more mature neurons for proper GABAergic innervation. To examine this idea, we co-cultured GFP-transgenic $\gamma 2^{+/-}$ subunit heterozygous neurons (GFP, $\gamma 2^{+/-}$) with an excess of WT neurons and analyzed the density of pre- and postsynaptic markers at 21 DIV. As predicted, dendrites of GFP-tagged $\gamma 2^{+/-}$ neurons showed dramatic reductions in GABAergic innervation compared to WT neurons analyzed in parallel cultures (density of GAD immunoreactive puncta on dendrites of $\gamma 2^{+/-}$ neurons: $56.2 \pm 4.8\%$ of WT, $n = 13-14$ neurons, $p < 0.001$, t-test), along with deficits in the dendritic density of punctate immunoreactivity for postsynaptic GABA_ARs ($54.4 \pm 2.7\%$ of WT, $p < 0.001$) (Fig. 3A, B). Interestingly, the size of $\gamma 2$ immunoreactive puncta was unchanged ($101.2 \pm 7.3\%$ of WT, $p > 0.05$), whereas the size of GAD puncta was significantly increased in $\gamma 2^{+/-}$ neurons of co-cultures ($130.5 \pm 9.3\%$ of WT, $p < 0.001$) (Fig. 3C), pointing to a presynaptic homeostatic compensatory mechanism. There was also a small decrease in the degree of colocalization in the punctate stainings for GAD and GABA_AR in $\gamma 2^{+/-}$ vs. WT neurons (Fig 3D). Similar defects in GABAergic innervation were seen when untagged $\gamma 2^{+/-}$ neurons were co-cultured with an excess of GFP-tagged WT neurons (not shown). We conclude that $\gamma 2$ -containing GABA_ARs contribute to normal GABAergic innervation, a property that is only evident under conditions where GABAergic axons are forced to choose among target neurons that differ in the level of expression of $\gamma 2$ -containing GABA_ARs.

3.4. Developmental effects of GABA_AR deficits on BDNF expression

In developing neurons, BDNF functions as part of a positive feedback loop that promotes the endocytic stability of GABA_ARs in the plasma membrane (Porcher *et al.*, 2011). Moreover, this mechanism was recently reported to involve BDNF/TrkB-mediated Tyr phosphorylation of the $\gamma 2$ subunit (Vithlani *et al.*, 2013). In turn, GABA_AR-mediated neural depolarization of immature neurons activates a Ca^{2+} -dependent signaling cascade that promotes BDNF gene expression and BDNF release (Baldelli *et al.*, 2002; Obrietan *et al.*, 2002; Porcher *et al.*, 2011). BDNF/TrkB signaling also promotes the postsynaptic accumulation of PSD95 and thereby contributes to maturation of dendritic synaptic spines (Yoshii *et al.*, 2011). These data predict that the dendritic maturational defects seen in the dentate gyrus of $\gamma 2^{+/-}$ mice might involve secondary reductions in BDNF expression that may amplify detrimental consequences of GABA_AR functional deficits. To test this idea we measured BDNF protein expression in frontal cortex, hippocampus and brain stem of one-, three- and nine-week-old female $\gamma 2^{+/+}$ and $\gamma 2^{+/-}$ mice. We found prominent differences in BDNF expression across postnatal development and brain regions, as expected (Fig. 4A–C).

No genotype effects on BDNF expression were evident in samples from individual brain regions across all three developmental stages ($p > 0.05$ for all t-tests). However, a genotype $\times$ brain region ANOVA of data from three-week-old mice normalized to WT values revealed a strong trend towards lower BDNF protein levels in $\gamma 2^{+/-}$ vs. WT mice [$F(1, 35) = 3.72$, $p = 0.06$, $n = 18$] (Fig. 4D). We previously showed that the anxious depression-related increase in emotional reactivity of $\gamma 2^{+/-}$ mice has a developmental origin between the third and end of fourth postnatal week (Earnheart *et al.*, 2007; Shen *et al.*, 2012). The data suggest that developmental brain-region-independent reductions in BDNF expression might contribute to the known behavioral abnormalities of $\gamma 2^{+/-}$ mice.

4. DISCUSSION

We previously reported that the $\gamma 2$ subunit of GABA_ARs is critically important for the survival of adult-born neurons of the hippocampus (Earnheart *et al.*, 2007). We here have extended these findings and show that similarly compromised neurogenesis is also present in juvenile mice. Moreover, we show that neurons fail to survive at a relatively late stage of differentiation, at a time when they no longer express the immature marker DCX. Loss of neurons was associated with a defect of dendritic arbor maturation in the molecular layer proximal to the granule cell layer. Impaired differentiation of granule cells overlaps temporally with a period of enhanced synaptic plasticity of adult born granule cells (Ge *et al.*, 2007b) that is thought to be pivotal for pattern separation and completion (Sahay *et al.*, 2011), a hippocampus-dependent form of cognitive disambiguation of similar life situations (Schmidt *et al.*, 2012). Consistent with defects in pattern separation, $\gamma 2^{+/-}$ mice show selective defects in ambiguous cue discrimination learning while performing normally or better than WT mice in other hippocampus-dependent learning and memory tasks (Crestani *et al.*, 1999). Deficits in the resolution of ambiguity reminiscent of those of $\gamma 2^{+/-}$ mice are commonly associated with MDD (Austin *et al.*, 1999; Schatzberg *et al.*, 2000; Rogers *et al.*, 2004; Anda *et al.*, 2006), suggesting that defects in granule cell maturation may contribute to the anxious-depressive phenotype of $\gamma 2^{+/-}$ mice.

In addition to dendritic maturation, the $\gamma 2$ subunit of GABA_ARs is known to be essential for GABA-mediated inhibition of RGL activation, and for the control of neural vs. glial cell fate choice of RGL-derived neural progenitors (Song *et al.*, 2012). This was shown by cell type specific knock out of the $\gamma 2$ subunit in RGLs, which results in their rapid dedifferentiation and mitotic activation. Increased proliferation of neural precursor cells was similarly observed upon deletion of the $\alpha 4$ subunit (but not the δ subunit) of GABA_ARs (Duveau *et al.*, 2011), indicating that the receptors that mediate the tonic GABA input that inhibits activation of RGLs likely have an $\alpha 4\beta\gamma 2$ subunit composition. However, we showed here that heterozygous deletion of the $\gamma 2$ subunit did not measurably affect this mechanism, as evidence by normal numbers of BrdU labeled cells 24 and 48 h after labeling. This is consistent with evidence that the $\gamma 2$ -subunit is largely dispensable for assembly of $\alpha\beta$ receptor complexes, and that $\alpha\beta$ complexes can form GABA-gated channels in non-synaptic membranes, albeit with significantly lower channel conductance than $\alpha\beta\gamma 2$ receptors (Baer *et al.*, 1999; Lorez *et al.*, 2000; Mortensen & Smart, 2006). By contrast, the $\gamma 2$ subunit is absolutely essential for accumulation of GABA_ARs at synapses, the typical function of $\gamma 2$ -containing GABA_ARs in mature neurons (Essrich *et al.*, 1998; Schweizer *et al.*, 2003). In

four-week-old granule cells the $\gamma 2$ subunit is thought to be part of $\alpha 2\beta\gamma 2$ receptors that are invariably enriched at synapses. This view is supported by analyses of knockout mice lacking the $\alpha 2$ subunit, which show selective maturational defects of granule cells comparable to those of $\gamma 2^{+/-}$ mice (Duveau *et al.*, 2011).

Sparse knock down of the $\gamma 2$ subunit by in utero electroporation and analyses of mice with a mosaic $\gamma 2$ loss of function allele have indicated that GABA_ARs play a role in GABAergic synapse formation (Li *et al.*, 2005; Frola *et al.*, 2013). Defects in GABAergic innervation were also observed following sparse shRNA-mediated knockdown of GABA_AR interacting trafficking proteins (Fang *et al.*, 2006; Yuan *et al.*, 2008). This was unexpected given that GABAergic innervation is unaffected in $\gamma 2$ knock-out mice and primary cultured neurons (Essrich *et al.*, 1998). However, we now show that even a partial reduction in the expression of the $\gamma 2$ subunit results in a marked reduction in GABAergic innervation when $\gamma 2^{+/-}$ neurons are grown in competition with WT neurons. This competitive environment explains the apparent discrepancies above between phenotypes observed following sparse vs. global knock out of the $\gamma 2$ subunit. Moreover, this situation is reminiscent of the condition faced by developing adult-born granule cells that have to compete with mature neurons for GABAergic innervation. Consistent with a direct role of GABAergic innervation GABA_ARs overexpressed in heterologous cells can initiate GABAergic innervation when co-cultured with neurons (Fuchs *et al.*, 2013). However, this synaptogenic effect of GABA_ARs is substantially weaker than that of GABA_ARs co-expressed with the synaptic cell adhesion protein neuroligin-2 (Dong *et al.*, 2007).

The deficits in granule cell maturation seen in $\gamma 2^{+/-}$ mice are reminiscent of similar deficits seen in multiple mouse lines with genetically induced defects in BDNF/TrkB signaling (Sairanen *et al.*, 2005; Kaneko *et al.*, 2012; Waterhouse *et al.*, 2012). This is consistent with substantial evidence that neurotrophic mechanisms of BDNF/TrkB are intrinsically intertwined with mechanisms of GABAergic transmission. First, BDNF functions upstream of GABA_ARs by enhancing the relative excitability of GABAergic versus glutamatergic neurons (Wardle & Poo, 2003) and by promoting the release of GABA (Jovanovic *et al.*, 2000) from parvalbumin-positive interneurons (Waterhouse *et al.*, 2012), a mechanism that delimits the mitotic activation of RGLs (Song *et al.*, 2012). Second, BDNF/TrkB signaling serves to stabilize GABA_ARs at the cell surface, both in immature (Porcher *et al.*, 2011) and mature neurons (Vithlani *et al.*, 2013). In immature neurons, BDNF/TrkB facilitates GABA_AR cell surface accumulation (Porcher *et al.*, 2011) and hence GABA/GABA_AR mediated membrane depolarization, which enables Ca²⁺ entry by V-gated Ca²⁺ channels (Maric *et al.*, 2001; Borodinsky *et al.*, 2003; Fiszman & Schousboe, 2004; Schmidt-Hieber *et al.*, 2004; Gascon *et al.*, 2006) and NMDARs (Tashiro *et al.*, 2006), followed by activation of diverse Ca²⁺ dependent Ser/Thr kinases that promote CREB phosphorylation and CREB-dependent gene transcription (Shaywitz & Greenberg, 1999; Nakagawa *et al.*, 2002; Fujioka *et al.*, 2004; Gur *et al.*, 2007; Jagasia *et al.*, 2009). In addition, GABA_AR-mediated membrane depolarization facilitates BDNF release (Porcher *et al.*, 2011). Moreover, one of the most prominent target genes activated by CREB is BDNF (Shieh *et al.*, 1998; Tao *et al.*, 1998; Obrietan *et al.*, 2002). These mechanisms indicate that BDNF functions not only upstream but also downstream of GABAergic transmission. Consistent

with GABAergic control of BDNF expression and release we found a near significant trend ($p = 0.06$) for reduced BDNF protein expression in developing $\gamma 2^{+/-}$ vs. WT mice. Our analyses of genotype dependent changes in individual brain regions were likely underpowered due to dynamic, behavioral state- and brain region-dependent variation in BDNF expression. Nevertheless, our data strongly suggest that BDNF expression is regulated by GABAergic transmission *in vivo*. Therefore, it is conceivable that reduced BDNF expression and function contributes to the postnatal developmental origin of the anxious-depressive-like phenotype of $\gamma 2^{+/-}$ mice (Earnheart *et al.*, 2007; Shen *et al.*, 2010; Shen *et al.*, 2012). Although far from detectable by biochemical means, it is likely that reduced BDNF transcription also applies to immature granule cells of $\gamma 2^{+/-}$ mice. These neurons are likely more vulnerable than embryo-derived neurons as they need to compete for GABAergic innervation with mature surrounding neurons. Consistent with this interpretation, we showed previously that the rate of proliferation, migration and survival of embryonic cortical neurons is unaffected in $\gamma 2^{+/-}$ and $\gamma 2^{-/-}$ mice (Shen *et al.*, 2012).

In addition to a deficit in dendritic complexity and indirect evidence for reduced GABAergic innervation, granule cells of $\gamma 2^{+/-}$ mice showed a reduction in the dendritic density of mushroom type synaptic spines. Mushroom type spines are known to serve as sites of stable and functionally mature glutamatergic innervation (Matsuzaki *et al.*, 2001; Ashby *et al.*, 2006). Thus, the dendritic spine abnormalities of $\gamma 2^{+/-}$ mice suggest association of functional defects in glutamatergic transmission with the established anxious-depressive phenotype of these mice. Atrophy of dendrites and spines are hallmarks also of stress-based, glutamatergic (Popoli *et al.*, 2012) and neurotrophic deficit hypotheses of MDD (Duman & Li, 2012). In sum, our findings reported here suggest increasing congruence of all these hypotheses with the GABAergic deficit hypothesis of MDD (Luscher *et al.*, 2011).

Acknowledgments

We are grateful to Yao Guo for technical assistance. We thank Steven Schiff and Laura Bianco (Center for Neural Engineering, Penn State University) for access to and technical help with the Neurolucida software and imaging system. We thank J.M. Fritschy (University of Zurich) for generous gifts of GABA_AR antibodies and Thomas Fuchs for critical reading of the manuscript. This publication was made possible by grants MH097247, MH089111 and MH099851 to B.L. from the National Institutes of Mental Health (NIMH) and a grant from the Pennsylvania Department of Health using Tobacco Settlement Funds. Its contents are solely the responsibility of the authors and do not necessarily represent the views of the NIMH or the NIH. The Pennsylvania Department of Health specifically disclaims responsibility for any analyses, interpretations or conclusions.

REFERENCES

- Allred MJ, Mulder-Rosi J, Lingenfelter SE, Chen G, Luscher B. Distinct gamma2 subunit domains mediate clustering and synaptic function of postsynaptic GABA_A receptors and gephyrin. *J. Neurosci.* 2005; 25:594–603. [PubMed: 15659595]
- Anda RF, Felitti VJ, Bremner JD, Walker JD, Whitfield C, Perry BD, Dube SR, Giles WH. The enduring effects of abuse and related adverse experiences in childhood. A convergence of evidence from neurobiology and epidemiology. *Eur Arch Psychiatry Clin Neurosci.* 2006; 256:174–186. [PubMed: 16311898]
- Ashby MC, Maier SR, Nishimune A, Henley JM. Lateral diffusion drives constitutive exchange of AMPA receptors at dendritic spines and is regulated by spine morphology. *J. Neurosci.* 2006; 26:7046–7055. [PubMed: 16807334]

- Austin MP, Mitchell P, Wilhelm K, Parker G, Hickie I, Brodaty H, Chan J, Eyers K, Milic M, Hadzi-Pavlovic D. Cognitive function in depression: a distinct pattern of frontal impairment in melancholia? *Psychol Med*. 1999; 29:73–85. [PubMed: 10077295]
- Baer K, Essrich C, Benson JA, Benke D, Bluethmann H, Fritschy J-M, Luscher B. Postsynaptic clustering of GABA(A) receptors by the gamma3 subunit *in vivo*. *Proc. Natl. Acad. Sci. USA*. 1999; 96:12860–12865. [PubMed: 10536013]
- Baldelli P, Novara M, Carabelli V, Hernandez-Guijo JM, Carbone E. BDNF up-regulates evoked GABAergic transmission in developing hippocampus by potentiating presynaptic N- and P/Q-type Ca²⁺ channels signalling. *Eur J Neurosci*. 2002; 16:2297–2310. [PubMed: 12492424]
- Borodinsky LN, O'Leary D, Neale JH, Vicini S, Coso OA, Fisman ML. GABA-induced neurite outgrowth of cerebellar granule cells is mediated by GABA(A) receptor activation, calcium influx and CaMKII and erk1/2 pathways. *J. Neurochem*. 2003; 84:1411–1420. [PubMed: 12614341]
- Chen AI, Nguyen CN, Copenhagen DR, Badurek S, Minichiello L, Ranscht B, Reichardt LF. TrkB (tropomyosin-related kinase B) controls the assembly and maintenance of GABAergic synapses in the cerebellar cortex. *J. Neurosci*. 2011; 31:2769–2780. [PubMed: 21414899]
- Choudary PV, Molnar M, Evans SJ, Tomita H, Li JZ, Vawter MP, Myers RM, Bunney WE Jr, Akil H, Watson SJ, Jones EG. Altered cortical glutamatergic and GABAergic signal transmission with glial involvement in depression. *Proc Natl Acad Sci U S A*. 2005; 102:15653–15658. [PubMed: 16230605]
- Clelland CD, Choi M, Romberg C, Clemenson GD Jr, Fragniere A, Tyers P, Jessberger S, Saksida LM, Barker RA, Gage FH, Bussey TJ. A functional role for adult hippocampal neurogenesis in spatial pattern separation. *Science*. 2009; 325:210–213. [PubMed: 19590004]
- Crestani F, Lorez M, Baer K, Essrich C, Benke D, Laurent JP, Belzung C, Fritschy JM, Luscher B, Mohler H. Decreased GABA(A)-receptor clustering results in enhanced anxiety and a bias for threat cues. *Nat. Neurosci*. 1999; 2:833–839. [PubMed: 10461223]
- Croarkin PE, Levinson AJ, Daskalakis ZJ. Evidence for GABAergic inhibitory deficits in major depressive disorder. *Neurosci Biobehav Rev*. 2011; 35:818–825. [PubMed: 20946914]
- Dong N, Qi J, Chen G. Molecular reconstitution of functional GABAergic synapses with expression of neuroligin-2 and GABA(A) receptors. *Mol Cell Neurosci*. 2007; 35:14–23. [PubMed: 17336090]
- Duman RS, Heninger GR, Nestler EJ. A molecular and cellular theory of depression. *Arch Gen Psychiatry*. 1997; 54:597–606. [PubMed: 9236543]
- Duman RS, Li N. A neurotrophic hypothesis of depression: role of synaptogenesis in the actions of NMDA receptor antagonists. *Philos Trans R Soc Lond B Biol Sci*. 2012; 367:2475–2484. [PubMed: 22826346]
- Duman RS, Monteggia LM. A neurotrophic model for stress-related mood disorders. *Biol Psychiatry*. 2006; 59:1116–1127. [PubMed: 16631126]
- Duveau V, Laustela S, Barth L, Gianolini F, Vogt KE, Keist R, Chandra D, Homanics GE, Rudolph U, Fritschy JM. Spatiotemporal specificity of GABAA receptor-mediated regulation of adult hippocampal neurogenesis. *Eur J Neurosci*. 2011; 34:362–373. [PubMed: 21722213]
- Dwivedi Y, Mondal AC, Rizavi HS, Conley RR. Suicide brain is associated with decreased expression of neurotrophins. *Biol Psychiatry*. 2005; 58:315–324. [PubMed: 15939410]
- Dwivedi Y, Rizavi HS, Conley RR, Roberts RC, Tamminga CA, Pandey GN. Altered gene expression of brain-derived neurotrophic factor and receptor tyrosine kinase B in postmortem brain of suicide subjects. *Arch Gen Psychiatry*. 2003; 60:804–815. [PubMed: 12912764]
- Earnheart JC, Schweizer C, Crestani F, Iwasato T, Itohara S, Mohler H, Luscher B. GABAergic control of adult hippocampal neurogenesis in relation to behavior indicative of trait anxiety and depression states. *J. Neurosci*. 2007; 27:3845–3854. [PubMed: 17409249]
- Eley TC, Gregory AM, Lau JY, McGuffin P, Napolitano M, Rijdsdijk FV, Clark DM. In the face of uncertainty: a twin study of ambiguous information, anxiety and depression in children. *J Abnorm Child Psychol*. 2008; 36:55–65. [PubMed: 17687641]
- Essrich C, Lorez M, Benson J, Fritschy J-M, Luscher B. Postsynaptic clustering of major GABA(A) receptor subtypes requires the gamma2 subunit and gephyrin. *Nat. Neurosci*. 1998; 1:563–571. [PubMed: 10196563]

- Fang C, Deng L, Keller CA, Fukata M, Fukata Y, Chen G, Luscher B. GODZ-mediated palmitoylation of GABA(A) receptors is required for normal assembly and function of GABAergic inhibitory synapses. *J. Neurosci.* 2006; 26:12758–12768. [PubMed: 17151279]
- Fiszman ML, Schousboe A. Role of calcium and kinases on the neurotrophic effect induced by gamma-aminobutyric acid. *J Neurosci Res.* 2004; 76:435–441. [PubMed: 15114615]
- Frola E, Patrizi A, Goetz T, Medrihan L, Petrini EM, Barberis A, Wulff P, Wisden W, Sassoe-Pognetto M. Synaptic competition sculpts the development of GABAergic axo-dendritic but not perisomatic synapses. *PloS one.* 2013; 8:e56311. [PubMed: 23457547]
- Fuchs C, Abitbol K, Burden JJ, Mercer A, Brown L, Iball J, Anne Stephenson F, Thomson AM, Jovanovic JN. GABA(A) receptors can initiate the formation of functional inhibitory GABAergic synapses. *Eur J Neurosci.* 2013; 38:3146–3158. [PubMed: 23909897]
- Fujioka T, Fujioka A, Duman RS. Activation of cAMP signaling facilitates the morphological maturation of newborn neurons in adult hippocampus. *J. Neurosci.* 2004; 24:319–328. [PubMed: 14724230]
- Gabbay V, Mao X, Klein RG, Ely BA, Babb JS, Panzer AM, Alonso CM, Shungu DC. Anterior cingulate cortex gamma-aminobutyric acid in depressed adolescents: relationship to anhedonia. *Arch Gen Psychiatry.* 2012; 69:139–149. [PubMed: 21969419]
- Gascon E, Dayer AG, Sauvain MO, Potter G, Jenny B, De Roo M, Zraggen E, Demareux N, Muller D, Kiss JZ. GABA regulates dendritic growth by stabilizing lamellipodia in newly generated interneurons of the olfactory bulb. *J. Neurosci.* 2006; 26:12956–12966. [PubMed: 17167085]
- Ge S, Goh EL, Sailor KA, Kitabatake Y, Ming GL, Song H. GABA regulates synaptic integration of newly generated neurons in the adult brain. *Nature.* 2006; 439:589–593. [PubMed: 16341203]
- Ge S, Pradhan DA, Ming GL, Song H. GABA sets the tempo for activity-dependent adult neurogenesis. *Trends Neurosci.* 2007a; 30:1–8. [PubMed: 17116335]
- Ge S, Yang CH, Hsu KS, Ming GL, Song H. A critical period for enhanced synaptic plasticity in newly generated neurons of the adult brain. *Neuron.* 2007b; 54:559–566. [PubMed: 17521569]
- Gonzalez-Burgos G, Barrionuevo G, Lewis DA. Horizontal synaptic connections in monkey prefrontal cortex: an in vitro electrophysiological study. *Cereb Cortex.* 2000; 10:82–92. [PubMed: 10639398]
- Guilloux JP, Douillard-Guilloux G, Kota R, Wang X, Gardier AM, Martinowich K, Tseng GC, Lewis DA, Sibille E. Molecular evidence for BDNF- and GABA-related dysfunctions in the amygdala of female subjects with major depression. *Mol Psychiatry.* 2012; 17:1130–1142. [PubMed: 21912391]
- Gunther U, Benson J, Benke D, Fritschy J-M, Reyes G, Knoflach F, Crestani F, Aguzzi A, Arigoni M, Lang Y, Bluethmann H, Mohler H, Luscher B. Benzodiazepine-insensitive mice generated by targeted disruption of the gamma 2 subunit gene of gamma-aminobutyric acid type A receptors. *Proc Natl Acad Sci U S A.* 1995; 92:7749–7753. [PubMed: 7644489]
- Gur TL, Conti AC, Holden J, Bechtholt AJ, Hill TE, Lucki I, Malberg JE, Blendy JA. cAMP response element-binding protein deficiency allows for increased neurogenesis and a rapid onset of antidepressant response. *J. Neurosci.* 2007; 27:7860–7868. [PubMed: 17634380]
- Hadjantonakis AK, Gertsenstein M, Ikawa M, Okabe M, Nagy A. Non-invasive sexing of preimplantation stage mammalian embryos. *Nat. Genet.* 1998; 19:220–222. [PubMed: 9662390]
- Hasler G, van der Veen JW, Tumonis T, Meyers N, Shen J, Drevets WC. Reduced prefrontal glutamate/glutamine and gamma-aminobutyric acid levels in major depression determined using proton magnetic resonance spectroscopy. *Arch Gen Psychiatry.* 2007; 64:193–200. [PubMed: 17283286]
- Jagasia R, Steib K, Englberger E, Herold S, Faus-Kessler T, Saxe M, Gage FH, Song H, Lie DC. GABA-cAMP response element-binding protein signaling regulates maturation and survival of newly generated neurons in the adult hippocampus. *J. Neurosci.* 2009; 29:7966–7977. [PubMed: 19553437]
- Jovanovic JN, Czernik AJ, Fienberg AA, Greengard P, Sihra TS. Synapsins as mediators of BDNF-enhanced neurotransmitter release. *Nat Neurosci.* 2000; 3:323–329. [PubMed: 10725920]
- Kaneko M, Xie Y, An JJ, Stryker MP, Xu B. Dendritic BDNF synthesis is required for late-phase spine maturation and recovery of cortical responses following sensory deprivation. *J. Neurosci.* 2012; 32:4790–4802. [PubMed: 22492034]

- Karege F, Vaudan G, Schwald M, Perroud N, La Harpe R. Neurotrophin levels in postmortem brains of suicide victims and the effects of antemortem diagnosis and psychotropic drugs. *Brain Res Mol Brain Res*. 2005; 136:29–37. [PubMed: 15893584]
- Karolewicz B, Maciag D, O'Dwyer G, Stockmeier CA, Feyissa AM, Rajkowska G. Reduced level of glutamic acid decarboxylase-67 kDa in the prefrontal cortex in major depression. *Int J Neuropsychopharmacol*. 2010; 13:411–420. [PubMed: 20236554]
- Klempan TA, Sequeira A, Canetti L, Lalovic A, Ernst C, French-Mullen J, Turecki G. Altered expression of genes involved in ATP biosynthesis and GABAergic neurotransmission in the ventral prefrontal cortex of suicides with and without major depression. *Mol Psychiatry*. 2009; 14:175–189. [PubMed: 17938633]
- Plumbers UM, Veltman DJ, Drent ML, Boellaard R, Comans EF, Meynen G, Lammertsma AA, Hoogendijk WJ. Reduced parahippocampal and lateral temporal GABA_A-[11C]flumazenil binding in major depression: preliminary results. *Eur J Nucl Med Mol Imaging*. 2010; 37:565–574. [PubMed: 19890631]
- Kucukibrahimoglu E, Saygin MZ, Caliskan M, Kaplan OK, Unsal C, Goren MZ. The change in plasma GABA, glutamine and glutamate levels in fluoxetine- or S-citalopram-treated female patients with major depression. *Eur J Clin Pharmacol*. 2009; 65:571–577. [PubMed: 19373461]
- Leal SL, Tighe SK, Jones CK, Yassa MA. Pattern separation of emotional information in hippocampal dentate and CA3. *Hippocampus*. 2014
- Levinson AJ, Fitzgerald PB, Favalli G, Blumberger DM, Daigle M, Daskalakis ZJ. Evidence of cortical inhibitory deficits in major depressive disorder. *Biol Psychiatry*. 2010; 67:458–464. [PubMed: 19922906]
- Li RW, Yu W, Christie S, Miralles CP, Bai J, Loturco JJ, De Blas AL. Disruption of postsynaptic GABA receptor clusters leads to decreased GABAergic innervation of pyramidal neurons. *J Neurochem*. 2005; 95:756–770. [PubMed: 16248887]
- Lorez M, Benke D, Luscher B, Mohler H, Benson JA. Single-channel properties of neuronal GABA_A receptors from mice lacking the 2 subunit. *J Physiol*. 2000; 527(Pt 1):11–31. [PubMed: 10944167]
- Luscher B, Shen Q, Sahir N. The GABAergic deficit hypothesis of major depressive disorder. *Mol Psychiatry*. 2011; 16:383–406. [PubMed: 21079608]
- Maciag D, Hughes J, O'Dwyer G, Pride Y, Stockmeier CA, Sanacora G, Rajkowska G. Reduced density of calbindin immunoreactive GABAergic neurons in the occipital cortex in major depression: relevance to neuroimaging studies. *Biol Psychiatry*. 2010; 67:465–470. [PubMed: 20004363]
- Maric D, Liu QY, Maric I, Chaudry S, Chang YH, Smith SV, Sieghart W, Fritschy JM, Barker JL. GABA expression dominates neuronal lineage progression in the embryonic rat neocortex and facilitates neurite outgrowth via GABA(A) autoreceptor/Cl⁻ channels. *J Neurosci*. 2001; 21:2343–2360. [PubMed: 11264309]
- Matsuzaki M, Ellis-Davies GC, Nemoto T, Miyashita Y, Iino M, Kasai H. Dendritic spine geometry is critical for AMPA receptor expression in hippocampal CA1 pyramidal neurons. *Nat Neurosci*. 2001; 4:1086–1092. [PubMed: 11687814]
- Merali Z, Du L, Hrdina P, Palkovits M, Faludi G, Poulter MO, Anisman H. Dysregulation in the suicide brain: mRNA expression of corticotropin-releasing hormone receptors and GABA(A) receptor subunits in frontal cortical brain region. *J Neurosci*. 2004; 24:1478–1485. [PubMed: 14960621]
- Mogg K, Bradbury KE, Bradley BP. Interpretation of ambiguous information in clinical depression. *Behav Res Ther*. 2006; 44:1411–1419. [PubMed: 16487479]
- Mortensen M, Smart TG. Extrasynaptic α subunit GABA_A receptors on rat hippocampal pyramidal neurons. *J Physiol*. 2006; 577:841–856. [PubMed: 17023503]
- Nakagawa S, Kim JE, Lee R, Malberg JE, Chen J, Steffen C, Zhang YJ, Nestler EJ, Duman RS. Regulation of neurogenesis in adult mouse hippocampus by cAMP and the cAMP response element-binding protein. *J Neurosci*. 2002; 22:3673–3682. [PubMed: 11978843]
- Obrietan K, Gao XB, Van Den Pol AN. Excitatory actions of GABA increase BDNF expression via a MAPK-CREB-dependent mechanism--a positive feedback circuit in developing neurons. *J Neurophysiol*. 2002; 88:1005–1015. [PubMed: 12163549]

- Popoli M, Yan Z, McEwen BS, Sanacora G. The stressed synapse: the impact of stress and glucocorticoids on glutamate transmission. *Nat. Rev. Neurosci.* 2012; 13:22–37. [PubMed: 22127301]
- Porcher C, Hatchett C, Longbottom RE, McAinch K, Sihra TS, Moss SJ, Thomson AM, Jovanovic JN. Positive feedback regulation between gamma-aminobutyric acid type A (GABA(A)) receptor signaling and brain-derived neurotrophic factor (BDNF) release in developing neurons. *J. Biol. Chem.* 2011; 286:21667–21677. [PubMed: 21474450]
- Rajkowska G, O'Dwyer G, Teleki Z, Stockmeier CA, Miguel-Hidalgo JJ. GABAergic neurons immunoreactive for calcium binding proteins are reduced in the prefrontal cortex in major depression. *Neuropsychopharmacology.* 2007; 32:471–482. [PubMed: 17063153]
- Rico B, Xu B, Reichardt LF. TrkB receptor signaling is required for establishment of GABAergic synapses in the cerebellum. *Nat Neurosci.* 2002; 5:225–233. [PubMed: 11836532]
- Rogers MA, Bellgrove MA, Chiu E, Mileskin C, Bradshaw JL. Response selection deficits in melancholic but not nonmelancholic unipolar major depression. *J Clin Exp Neuropsychol.* 2004; 26:169–179. [PubMed: 15202537]
- Sahay A, Scobie KN, Hill AS, O'Carroll CM, Kheirbek MA, Burghardt NS, Fenton AA, Dranovsky A, Hen R. Increasing adult hippocampal neurogenesis is sufficient to improve pattern separation. *Nature.* 2011; 472:466–470. [PubMed: 21460835]
- Sairanen M, Lucas G, Ernfors P, Castren M, Castren E. Brain-derived neurotrophic factor and antidepressant drugs have different but coordinated effects on neuronal turnover, proliferation, and survival in the adult dentate gyrus. *J. Neurosci.* 2005; 25:1089–1094. [PubMed: 15689544]
- Samuels BA, Hen R. Neurogenesis and affective disorders. *Eur J Neurosci.* 2011; 33:1152–1159. [PubMed: 21395859]
- Sanacora G, Mason GF, Rothman DL, Behar KL, Hyder F, Petroff OA, Berman RM, Charney DS, Krystal JH. Reduced cortical gamma-aminobutyric acid levels in depressed patients determined by proton magnetic resonance spectroscopy. *Arch Gen Psychiatry.* 1999; 56:1043–1047. [PubMed: 10565505]
- Sanacora G, Mason GF, Rothman DL, Hyder F, Ciarcia JJ, Ostroff RB, Berman RM, Krystal JH. Increased cortical GABA concentrations in depressed patients receiving ECT. *Am J Psychiatry.* 2003; 160:577–579. [PubMed: 12611844]
- Sanacora G, Mason GF, Rothman DL, Krystal JH. Increased occipital cortex GABA concentrations in depressed patients after therapy with selective serotonin reuptake inhibitors. *Am J Psychiatry.* 2002; 159:663–665. [PubMed: 11925309]
- Schatzberg AF, Posener JA, DeBattista C, Kalehzan BM, Rothschild AJ, Shear PK. Neuropsychological deficits in psychotic versus nonpsychotic major depression and no mental illness. *Am J Psychiatry.* 2000; 157:1095–1100. [PubMed: 10873917]
- Schmidt B, Marrone DF, Markus EJ. Disambiguating the similar: the dentate gyrus and pattern separation. *Behav Brain Res.* 2012; 226:56–65. [PubMed: 21907247]
- Schmidt-Hieber C, Jonas P, Bischofberger J. Enhanced synaptic plasticity in newly generated granule cells of the adult hippocampus. *Nature.* 2004; 429:184–187. [PubMed: 15107864]
- Schweizer C, Balsiger S, Bluethmann H, Mansuy M, Fritschy JM, Mohler H, Luscher B. The gamma2 subunit of GABA(A) receptors is required for maintenance of receptors at mature synapses. *Mol. Cell. Neurosci.* 2003; 24:442–450. [PubMed: 14572465]
- Sequeira A, Klempan T, Canetti L, French-Mullen J, Benkelfat C, Rouleau GA, Turecki G. Patterns of gene expression in the limbic system of suicides with and without major depression. *Mol Psychiatry.* 2007; 12:640–655. [PubMed: 17353912]
- Shaywitz AJ, Greenberg ME. CREB: a stimulus-induced transcription factor activated by a diverse array of extracellular signals. *Annu Rev Biochem.* 1999; 68:821–861. [PubMed: 10872467]
- Shen Q, Fuchs T, Sahir N, Luscher B. GABAergic control of critical developmental periods for anxiety- and depression-related behavior in mice. *PLoS One.* 2012; 7:e47441. [PubMed: 23071808]
- Shen Q, Lal R, Luellen BA, Earnheart JC, Andrews AM, Luscher B. gamma-Aminobutyric acid-type A receptor deficits cause hypothalamic-pituitary-adrenal axis hyperactivity and antidepressant

- drug sensitivity reminiscent of melancholic forms of depression. *Biol Psychiatry*. 2010; 68:512–520. [PubMed: 20579975]
- Shieh PB, Hu SC, Bobb K, Timmusk T, Ghosh A. Identification of a signaling pathway involved in calcium regulation of BDNF expression. *Neuron*. 1998; 20:727–740. [PubMed: 9581764]
- Sibille E, Morris HM, Kota RS, Lewis DA. GABA-related transcripts in the dorsolateral prefrontal cortex in mood disorders. *Int J Neuropsychopharmacol*. 2011; 14:721–734. [PubMed: 21226980]
- Smith KS, Rudolph U. Anxiety and depression: mouse genetics and pharmacological approaches to the role of GABA(A) receptor subtypes. *Neuropharmacology*. 2012; 62:54–62. [PubMed: 21810433]
- Song J, Zhong C, Bonaguidi MA, Sun GJ, Hsu D, Gu Y, Meletis K, Huang ZJ, Ge S, Enikolopov G, Deisseroth K, Luscher B, Christian KM, Ming GL, Song H. Neuronal circuitry mechanism regulating adult quiescent neural stem-cell fate decision. *Nature*. 2012; 489:150–154. [PubMed: 22842902]
- Szapacs ME, Numis AL, Andrews AM. Late onset loss of hippocampal 5-HT and NE is accompanied by increases in BDNF protein expression in mice co-expressing mutant APP and PS1. *Neurobiol Dis*. 2004; 16:572–580. [PubMed: 15262269]
- Tao X, Finkbeiner S, Arnold DB, Shaywitz AJ, Greenberg ME. Ca²⁺ influx regulates BDNF transcription by a CREB family transcription factor-dependent mechanism. *Neuron*. 1998; 20:709–726. [PubMed: 9581763]
- Tashiro A, Sandler VM, Toni N, Zhao C, Gage FH. NMDA-receptor-mediated, cell-specific integration of new neurons in adult dentate gyrus. *Nature*. 2006; 442:929–933. [PubMed: 16906136]
- Tripp A, Oh H, Guilloux JP, Martinowich K, Lewis DA, Sibille E. Brain-derived neurotrophic factor signaling and subgenual anterior cingulate cortex dysfunction in major depressive disorder. *Am J Psychiatry*. 2012; 169:1194–1202. [PubMed: 23128924]
- Vithlani M, Hines RM, Zhong P, Terunuma M, Hines DJ, Revilla-Sanchez R, Jurd R, Haydon P, Rios M, Brandon N, Yan Z, Moss SJ. The ability of BDNF to modify neurogenesis and depressive-like behaviors is dependent upon phosphorylation of tyrosine residues 365/367 in the GABA(A)-receptor gamma2 subunit. *J. Neurosci*. 2013; 33:15567–15577. [PubMed: 24068823]
- Wardle RA, Poo MM. Brain-derived neurotrophic factor modulation of GABAergic synapses by postsynaptic regulation of chloride transport. *J. Neurosci*. 2003; 23:8722–8732. [PubMed: 14507972]
- Waterhouse EG, An JJ, Orefice LL, Baydyuk M, Liao GY, Zheng K, Lu B, Xu B. BDNF promotes differentiation and maturation of adult-born neurons through GABAergic transmission. *J. Neurosci*. 2012; 32:14318–14330. [PubMed: 23055503]
- Yoshii A, Murata Y, Kim J, Zhang C, Shokat KM, Constantine-Paton M. TrkB and protein kinase Mzeta regulate synaptic localization of PSD-95 in developing cortex. *J. Neurosci*. 2011; 31:11894–11904. [PubMed: 21849550]
- Yuan X, Yao J, Norris D, Tran DD, Bram RJ, Chen G, Luscher B. Calcium-modulating cyclophilin ligand regulates membrane trafficking of postsynaptic GABA(A) receptors. *Mol Cell Neurosci*. 2008; 38:277–289. [PubMed: 18424167]
- Zhao C, Teng EM, Summers RG Jr, Ming GL, Gage FH. Distinct morphological stages of dentate granule neuron maturation in the adult mouse hippocampus. *J. Neurosci*. 2006; 26:3–11. [PubMed: 16399667]

- GABA_A receptor deficient mice represent a model for major depressive disorder
- This phenotype includes deficits in dendritic maturation of hippocampal granule cells
- Deficits in spine maturation suggest additional deficits at glutamatergic synapses
- GABA_A receptors are also required for normal formation of GABAergic synapses
- GABA_A receptors may regulate expression of BDNF

Figure 1. Heterozygosity of the $\gamma 2^{+/-}$ subunit results in a deficit in the number of NeuN positive dentate gyrus granule cell neurons born at three weeks of age

A-C. Representative images showing BrdU labeled brain sections of mice labeled at three weeks of age and euthanized 24h (A), 48h (B) or 28d (C) after BrdU injection. Arrows indicate neurons that are doubly positive for BrdU and the immature neuronal marker DCX (B) or the mature neuronal marker NeuN (C). **D. E.** P21 mice were injected once with BrdU (200 mg/kg) and brains harvested 24 h (D) or 48h (E) later. There was no difference in the number of BrdU positive cells in the subgranule cell layer of $\gamma 2^{+/-}$ mice vs. WT controls ($p > 0.05$ for both time points). **F. G.** To label immature neurons, P21 mice were metabolically labeled (4 times 80 mg/kg every 2h) with BrdU and the brains harvested 14 days (F) or 28 days later (G). There was no difference in BrdU/DCX double positive cells in $\gamma 2^{+/-}$ mice compared to WT controls for brains harvested 14 days after BrdU injection (F) ($p > 0.05$). However, there was a significant deficit in BrdU/NeuN double positive mature neurons in $\gamma 2^{+/-}$ mice analyzed 28d post BrdU injection (G) ($70.15 \pm 7.0\%$ of WT; $n = 4$ per genotype, $p < 0.05$, $U = 0$, Mann Whitney). Data indicate means $\pm$ SEM. * $p < 0.05$, Scale bar, 20 μ m.

Figure 2. Heterozygosity of the $\gamma 2$ subunit results in reduced complexity of dendrites and defects in spine maturation

A. Example of concentric rings placed on the soma of a DCX-labeled granule cell (green) used for Sholl analysis (rings are not drawn to scale). The section was counterstained with DRAQ5 (blue) to visualize cell nuclei and demarcate the granule cell and molecular layers. ML: molecular layer. GCL: granule cell layer. **B.** Quantification of the number of circle crossings of GFP labeled granule cell dendrites revealed a significant genotype effect [$F(16,15) = 2.4$, $p < 0.05$, $n = 16$]. Posthoc t-tests revealed significant reductions in dendritic complexity in $\gamma 2^{+/-}$ vs. WT mice at a distance of 70 and 90 μm from the soma ($p < 0.05$ and $p < 0.001$) and a change in the opposite direction at 290 μm ($p < 0.01$, $n = 16$). **C.** There was a significant reduction in the number of branch points in $\gamma 2^{+/-}$ vs. WT mice in the radial interval of 70-90 μm from the soma of granule cells ($p < 0.05$, $n = 16$, t-test). **D.** Representative image of a retrovirus-labeled dendritic segment of an eight-week-old granule cell. The boxed segment in the left micrograph is shown magnified on the right with thin-, stubby- and mushroom-type spines indicated. **E.** Quantification of the density of different types of spines. A two-way genotype x spine density ANOVA with spine types as within subject factors revealed a significant interaction between genotype and total spine density [$F(2, 29) = 4.3$, $p < 0.05$]. Posthoc analyses revealed a significant reduction selectively in mushroom type spines in granule cells of $\gamma 2^{+/-}$ vs. WT mice ($p < 0.05$, $n = 16$, t-test with Bonferroni correction). Data represent means $\pm$ SEM, * $p < 0.05$, ** $p < 0.01$.

Figure 3. Subtle deficits in $\gamma 2$ -containing GABA_ARs result in deficits of GABAergic innervation

A, Cortical neurons (DIV21) derived from WT neurons were either cultured alone (top row) or mixed together with cortical neurons from GFP-transgenic $\gamma 2^{+/-}$ embryos (bottom row, 9:1 excess of WT neurons). The neurons were immunostained for the $\gamma 2$ subunit (red, **a2**, **b2**) and GAD (blue, **a3**, **b3**). Merged red and blue channels are shown in panels (**a4**) and (**b4**). Boxed dendritic segments are enlarged in separate panels on the right. Arrows in enlarged panels point to a WT axon(s) that faithfully grows along a WT dendrite. By contrast, arrowheads show an axon of presynaptic WT neurons that grows across the dendrite of a $\gamma 2^{+/-}$ neuron, and failing to adhere to the dendrite (**a**). Mutant neurons ($\gamma 2^{+/-}$) were identified by the transgene-encoded GFP fluorescence (green, **b1**). **B**. When $\gamma 2^{+/-}$ neurons were co-cultured in the presence of an excess of WT neurons the density of immunoreactive puncta for $\gamma 2$ ($54.4 \pm 2.7\%$ of WT, $n = 13-14$ neurons, $p < 0.001$, t-test) and GAD ($56.2 \pm 4.8\%$ of WT, $n = 13-14$, $p < 0.001$, t-test) along their dendrites was significantly reduced compared to WT neurons grown in pure cultures. **C**. The size of $\gamma 2$ puncta was unchanged ($101.2 \pm 7.3\%$ of WT, $p > 0.05$, t-test), whereas the size of GAD puncta (of WT presynaptic interneurons) was significantly increased when $\gamma 2^{+/-}$ neurons were co-cultured with an excess of WT neurons ($130.5 \pm 9.3\%$ of WT, $p < 0.001$), suggesting presynaptic compensation for limiting amounts of postsynaptic GABA_ARs. **D**. The number of $\gamma 2$ puncta colocalized with GAD in $\gamma 2^{+/-}$ neurons co-cultured with WT was

reduced ($87.7 \pm 3.62\%$ of WT; $n = 13-14$, $p < 0.01$, t-test). Similarly, the number of GAD puncta colocalized with $\gamma 2$ was reduced in $\gamma 2^{+/-}$ neurons co-cultured with WT ($91.4 \pm 2.8\%$ of WT; $n = 13-14$, $p < 0.05$). Data represent means $\pm$ SEM; * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Figure 4. BDNF protein levels in different brain regions across postnatal development

A-C. BDNF protein levels were measured in extracts from frontal cortex (FC), brain stem (BS) and hippocampus (Hippo) of one-, three-, and nine-week-old $\gamma 2^{+/-}$ and WT littermate mice. No differences in BDNF expression were observed in any of the three brain regions analyzed (t-tests, $p > 0.05$ for all comparisons). **D.** However, a genotype x brain region ANOVA of data from three-week-old mice normalized to WT revealed a strong trend

towards lower BDNF protein expression in $\gamma 2^{+/-}$ vs. WT mice [$F(1, 34) = 3.72, p = 0.06, n = 18$]. BDNF levels are reported as ng BDNF/g wet weight tissue $\pm$ SEM.