Published in final edited form as:

Mol Biosyst. 2015 January 2; 11(1): 262–274. doi:10.1039/c4mb00604f.

Decomposition of RNA methylome reveals co-methylation patterns induced by latent enzymatic regulators of epitranscriptome

Lian Liu¹, Shao-Wu Zhang^{1,*}, Yu-Chen Zhang¹, Hui Liu², Lin Zhang², Runsheng Chen^{1,3}, Yufei Huang⁴, and Jia Meng^{5,*}

¹Key Laboratory of Information Fusion Technology of Ministry of Education, School of Automation, Northwestern Polytechnical University, Xi'an, Shaanxi, 710027, China

²School of Information and Electrical Engineering, China University of Mining and Technology, Xuzhou, 221116, China

³Institute of Biophysics, Chinese Academy of Sciences, Beijing, 100101, China

⁴Department of Electrical and Computer Engineering, University of Texas at San Antonio, TX, 78249, USA

⁵Department of Biological Sciences, Xi'an Jiaotong-Liverpool University, Suzhou, Jiangsu, 215123, China

Abstract

Biochemical modifications to mRNA, especially N6-methyladenosine (m⁶A) and 5-methylcytosine (m⁵C), are recently shown to be associated with crucial biological functions. Despite the intriguing advancements, little is known so far about the dynamic landscape of RNA methylome across different cell types and how the epitranscriptome is regulated at system level by enzymes, i.e., RNA methyltransferases and demethylases. To investigate this issue, a meta-analysis of m⁶A MeRIP-Seq datasets from 10 different experimental conditions (cell type/tissue or treatment) are collected, and the combinatorial epitranscriptome, which consists of 42758 m⁶A sites, is extracted and divided into 3 clusters, in which the methylation sites are likely to be hyper- or hypo-methylated simultaneously (or co-methylated), indicating the sharing of a common methylation regulator. Four different clustering approaches are used, including, K-means, hierarchical clustering (HC), Bayesian factor regression model (BFRM) and nonnegative matrix factorization (NMF) to unveil the co-methylation patterns.

To validate whether the patterns are corresponding to enzymatic regulators, i.e., RNA methyltransferases or demethylases, the target sites of a known m⁶A regulator, fat mass and obesity-associated protein (FTO), are identified from an independent mouse MeRIP-Seq dataset and lifted to human. Our study shows that, 3 out of the 4 clustering approaches used can successfully identify a group of methylation sites overlapping with FTO target sites at significance

*Correspondence: zhangsw@nwpu.edu.cn (SWZ) and jia.meng@xjtlu.edu.cn (JM).

Software Availability: The open source R package "RMT" for extracting combinatorial epitranscriptome from multiple MeRIP-Seq experiments is provided in Supplement Materials.

level 0.05 (after multiple hypothesis adjustment), among which, the result of NMF is the most significant (p-value 2.81e-06). We defined a new approach evaluating the consistency between two clustering results and shows that clustering results of different methods are highly correlated indicating strongly the existence of co-methylation patterns. Consistent with recent studies, a number of cancer and neuronal disease-related bimolecular functions are enriched in the identified clusters, which are biological functions can be regulated at epitranscriptional level, indicating the pharmaceutical prospect of RNA N6-methyladenosine-related studies.

This result successfully reveals the linkage between the global RNA co-methylation patterns embedded in the epitranscriptomic data under multiple experimental conditions and the latent enzymatic regulators, suggesting a promising direction towards more comprehensive understanding of the epitranscriptome.

Introduction

The dynamic chemical modifications of DNA together with their functions have been well established through intensive studies ranging from simple model organisms to human in the past decade [1–3]. Compared with DNA modifications, RNA modifications are largely neglected, and have yet drawn extensive attention until very recently [4–7]. The presence of chemical modifications to RNA was established as early as 1970s [8–10]. Until today, people have identified over 100 post-transcriptional RNA modifications in all 3 kingdoms of life [11]; however, not much is known so far about their biological, physiological and pathological functions due to technical limitations. Recently, a new technique (differently named as “MeRIP-Seq” [12] or “m⁶A-Seq” [13]) was proposed in 2012 by two independent groups, for the first time, enabled the global unbiased profiling of mRNA N6-methyladenosine methylome, one component of the epitranscriptome layer of gene regulation [14]. Currently, the only practically feasible unbiased approach for measuring mRNA m⁶A is MeRIP-Seq technique [12, 13, 15]. MeRIP-Seq pulls down and sequences the methylated mRNA fragments with anti-m⁶A antibody, and an input control sample is also generated to measure the basal abundance of all genes (see Figure 1).

Similar to ChIP-Seq data analysis [16–18], since MeRIP-Seq cannot provide base-resolution, the detection of RNA methylation sites from MeRIP-Seq has been mainly formulated as the “peak detection” problem [19, 20]; however, as point out previously, a single RNA methylation site may be split into 2 sections due to the existence of introns, the peak calling of MeRIP-Seq data should ideally be splicing-aware [14, 19]. Nevertheless, MeRIP-Seq technique successfully combines the essence of methylated DNA immunoprecipitation sequencing (MeDIP-Seq) and RNA sequencing (RNA-Seq) for high-resolution detection of transcriptome-wide RNA methylation modifications. Within 2 years’ time, the technique has been applied to a number of important studies in human, mouse, yeast, etc. [21–25]. Meanwhile, some RNA (de)methylation enzymes are identified [24, 26–28]. These studies together greatly enhanced our understanding of the reversible modifications to mRNA [5, 29].

However, one question remains to be answered is how the epitranscriptome, which consists of tens of thousands of RNA methylation sites, is regulated at system level across multiple

conditions by RNA methyltransferases and demethylases. While it is important to cumulate additional knowledge for the function of a specific RNA methylation enzyme under a particular condition to obtain all pieces of a Jigsaw puzzle, it is also necessary to integrate what we have so far for a big picture and untangle the high dimensional RNA methylome of tens of thousands of RNA methylation sites to shape an interpretable picture. The RNA methylome embraces a number of features that make a system level computational analysis necessary and feasible:

- **The RNA m⁶A methylome consists of a large number of RNA methylation sites** (ranging from 9124 to 46293 m⁶A sites predicted in different conditions [14]). It has been even speculated that, every RNA molecule may be methylated under a specific condition. Conceivably, dimension reduction technique is necessary to make sense of high-dimensional information. The RNA residuals are methylated or demethylated by a relatively small number of regulators (RNA methyltransferases or demethylases), each of which regulates thousands of RNA methylation sites. In this sense, high dimensional RNA methylation data can be conveniently projected into lower dimensional space of RNA methylation regulators to reveal directly the biochemical causes of the observed phenomenon.
- **RNA methylation is non-stoichiometric**, i.e., a specific residual can be methylated only on a fraction of transcripts, not necessarily all or none. This process is influenced by the “methylation potential” [30], which is the ratio of S-adenosylmethionine (SAM, the universal methyl donor cosubstrate) and S-adenosylhomocysteine (SAH, the by-product of SAM that acts as competitive inhibitor). With the simplest approximation of an equilibrium condition, the ratio between methylated and unmodified residuals is directly proportional to the SAM/SAH ratio, and is independent from the absolute RNA abundance. The fact that the same nucleic acids are not methylated at the same level indicates the specificity of the enzymes, which is more complicatedly determined by the methylation complex.
- **RNA co-methylation pattern exists due to enzymatic regulators.** Consistent with the “SAM/SAH” ratio, the group of RNA methylation sites controlled by the same RNA methylation factor will show hyper- or hypo-methylation simultaneously. This is analogous to transcription factor (TF) network or microRNA (miRNA) network, where the regulated target genes show co-expression pattern consistent with their regulator. Conceivably, the hyper-methylation of a large number of RNA methylation sites may indicates the increase of methyltransferases or the decreases of demethylase. Co-regulation pattern is the key for the identification of latent regulatory factors, which may function at protein level and cannot be directly observable from the high-throughput RNA methylation sequencing data. Due to the activities of RNA methylation enzymes and their specificity, the methylation level of the large number of RNA methylation sites are not independent of each other but show some clustering effect. On one hand, it is likely that a single methylation factor may regulate a large number of RNA methylation sites simultaneously; on the other hand, if we consider a single RNA methylation enzyme can be a protein complex consists of the protein products of

several genes, then it is possible that the methylation status of a single site is determined by multiple proteins. Although the real regulatory relationship between RNA methylation sites and enzymes can be more complex, it is practically more convenient to start with simpler computation methods, such as, K-means, and gradually increases the complexity of tools.

- **RNA co-methylation patterns are likely to have specific biological functions.**
From evolutionary point of view, to maximize the functionality of RNA methylation as a means of regulation, rather than aimlessly targeting a number of unrelated genes by random, natural selection should favor that an RNA methylation regulator targets functional-related genes to control specific functions so as to add the adaptability of the liver organism. It should be reasonable to assume that the targets of the same RNA methylation enzyme are likely to share functions in common; and on the other hand, for the purpose of validation, if some functions are statistically related to an RNA methylation enzyme, it is likely that identification of the enzyme and its targets is successful.
- **Transcriptional regulation may indirectly affect the epitranscriptome.**
Although existing studies mostly focus on the changes of the absolute amount of methylation with the basal RNA expression levels ignored, it is important to notice whether the increase is triggered by transcriptional regulation or by the enzymatic regulation (RNA methyltransferases or demethylases). While the RNA “methylation potential” moderates the ratio of methylated and unmodified molecules, transcriptional regulation directly control the absolute amount of RNA transcripts with the ratio unchanged. In practice, the changes in the absolute amount of RNA methylation can be due to a joint effect of the two (See Figure 2).

In this study, the combinatorial RNA m⁶A methylome from 10 experimental conditions (different tissues, cell lines or treatments) is firstly extracted using an R package we developed to study its dynamics. Four different clustering approaches, representing different rationales, are applied to dissect the RNA methylation sites. The results confirm the existence of co-methylation patterns and their relationship with RNA methylation enzymes.

Material and Method

Multiple N6-methyladenosine MeRIP-Seq datasets from different conditions are collected for this analysis. Samples include HEK293, HepG2, U2OS cell lines and brain tissue with different treatments, some with more than 1 biological replicates. The raw data in FASTQ format (SRA) was obtained directly from Gene Expression Omnibus (GEO), and then aligned to the human reference genome assembly (hg19) with spliced aligner Tophat2 [31] for the following analysis. The information of these datasets is summarized in Table 1.

To dissect the RNA methylome with clustering approach, the joint epitranscriptome must be firstly extracted. Currently, there is no convenient tool provided with this function. For this purpose, we developed an open source R package **RNA Methylation Tool (RMT)** for the processing of multiple MeRIP-Seq datasets and extracting the combinatorial

epitranscriptome, i.e., all the RNA methylation sites detected under one or more conditions. The general work flow of RMT is shown in Figure 3.

We will in the next detail each step of RMT work flow.

- Extract individual RNA methylome using exomePeak R/Bioconductor package:** The very first step of the analysis is to extract all the RNA methylation sites (or the “epitranscriptome”) from each individual MeRIP-Seq experiment. As aforementioned, there are two types of samples in MeRIP-Seq, i.e., IP and control samples. As essentially an enrichment-based approach, since the pull-down reads in the IP sample are enriched with methylated fragments, there is likely a higher number of reads (or a “peak”) appearing near the methylation sites in the IP sample compared with the input control sample, thus the methylation sites may be detected with “peak calling” method. We previously developed the exomePeak R/Bioconductor package [19] for this purpose. The exomePeak package is based on the C-test for comparison of two Poisson means [32] to detect the methylation sites on RNA molecule. As a splicing-aware peak caller focus on the exons only, its effectiveness on MeRIP-Seq peak calling has been demonstrated previously [19].
- Merge all detected methylation sites for a combinatorial methylome:** RNA methylation is reversible and dynamic under different conditions. The RNA methylome with tens of thousands of RNA methylation sites identified under different conditions in previous step are further merged into a combined set. It is worth mentioning that, the difference between RNA methylome under different conditions is not only due to the context-specificity but also related to noise and the detectability of MeRIP-Seq, i.e., when the expression level of a gene is low, its methylation site can be difficult to detect. The combination of multiple RNA methylomes is conducted in the following way: (1) RNA methylation sites that do not overlap with those detected under a different condition are context-specific and unique, thus all are kept. (2) RNA methylation sites that overlap with those detected under a different condition are not context specific and may appear multiple times. Under this scenario, only the widest methylation sites are kept.
- Quantification of RNA methylation level:** A natural way to quantify the RNA methylation level (percentage of methylated RNA molecules) based on MeRIP-Seq data is the “IP/Input ratio”, which is defined as ratio of the number of reads in IP and input control sample after compensating for the sequencing depth (or total number of reads). However, infinite “IP/Input ratio” might be generated when there are no reads detected in the input sample, which is not rare. Here, we adopt the way of computing gene expression in RNA-Seq with “RPKM” for a specific methylation site. The RNA methylation level is then quantified with:

$$x_{m,j} = \log_2 \left(\frac{t_{m,j} + 0.01}{c_{m,j} + 0.01} \right) \quad (1)$$

Where, $t_{m,j}$ indicates the RPKM value of m -th methylation site in the IP sample from j -th biological replicate, and $c_{m,j}$ input control sample. The introduction of

RPKM and its fold change into MeRIP-Seq is intuitive and conceptually sound. Please note that, under an ideal scenario of reversible chemical reaction, the methylation level is independent of RNA abundance (transcriptional regulation) and determined by only the “methylation potential”. For the aforementioned reason, the co-methylation patterns embedded is irrelevant to the gene expression. when calculating RPKM value, “R” refers to reads mapped to a specific methylation site, and “K” refers to 1000bp of methylation site. In all the following analysis, multi-reads (reads can be mapped to multiple genomic locations) are excluded to eliminate mapping ambiguity. The redefined methylation level in (1) is then more robust and will not generate infinite value. For most methylation sites, where short reads are more enriched in IP sample compared with input control sample, it should be a positive number.

After extracting the combinatorial RNA methylome and the methylation levels, feature selection was conducted to select the most informatics features (RNA methylation sites) for clustering purpose. For each feature, the methylation level defined in (1) can be determined on every single biological replicate provided the corresponding paired IP and Input control MeRIP-Seq sample is available. For the purpose of detecting co-methylated RNA methylation sites, of interests are those varies significantly across different conditions/replicates. For best clustering effect, it is important to exclude those with small variance in methylation level. We will select features that have larger variance in methylation level across different conditions, then the methylation levels are standardize prior to clustering analysis with:

$$\hat{x}_{m,j} = \frac{x_{m,j} - \mu_m}{s_m} \quad (2)$$

Where μ_m and s_m are the mean and sample standard deviation of m -th methylation site, respectively. Based on the standardized matrix of $\hat{x}_{m,j}$, co-methylation pattern embedded may be extracted. Each co-methylation pattern consists of a number of methylation sites are hyper- and hypo-methylated simultaneously, indicating a common latent regulatory factor at epitranscriptomic level. For this purpose, 4 widely used clustering approaches are adopted, including, K-means, hierarchical clustering, Bayesian factor model [33] and nonnegative matrix factorization [34], each reflects a different underlying assumption. The clustering results are then passed for gene ontology analysis with topGO R/Bioconductor package [35] for functional analysis.

Result and Discussion

RNA methylome of individual sample

The newly developed RMT package internally adapts exomePeak package for peak calling on every individual IP sample. Specifically, the RNA methylation sites under each condition are shown in Table 2, the number of peaks (RNA methylation sites) under a specific condition ranges from 1200 (brain) to 21127 (HepG2). The strand-specific consensus sequences for each condition are then obtained at MEME-ChIP webserver [36] by extracting all the peak regions from hg19 whole genome FASTA file provided from Illumina iGenome.

Despite the large variation in the numbers of detected RNA methylation sites, the enriched consensus sequences are similar to the known RRACH motif of m⁶A [12, 13], indicating the determined RNA methylation sites are accurate and consistent with previous studies.

Combinatorial RNA methylome

The combinatorial RNA methylome of 10 different datasets (Table 1) is then extracted by the newly developed RMT package. Thanks to the GenomicFeatures R/Bioconductor package [37], peak merging stage takes only seconds to complete, the most time consuming step is peak calling by exomePeak package. The 258465 RNA methylation sites detected on 18 replicates from 10 different conditions are further merged into 42758 unique RNA methylation sites using the rules previously specified. (See Figure 4 and Supplementary Material S2)

Under individual conditions, different number of merged methylation sites can be observed (see Figure 5A), and consistent with previous studies, the RNA methylation sites are more enriched near the stop codon of mRNA and on 3'UTR region (see Figure 5B). One interesting new observation is that, the RNA methylation sites on lncRNA are consistently more enriched on the 5'end compared with 3'end, whose cause is yet clear (see Figure 5B).

For the merged 42758 unique RNA methylation sites, each on average appears on 1/3 (6 of 18) of biological replicates obtained under 10 different conditions. Specifically, 35.4% (15126) appears only once, and only 33.1% (14415) appears on more than 5 biological replicates, indicating a highly dynamic landscape of RNA N⁶-methyladenosine. Please note that, this highly specific behavior may partially attribute to detection noise and the dynamics of transcription, i.e., given the current protocol of MeRIP-Seq, it is impossible to detect an RNA methylation site on lowly expressed genes, so dynamics of transcription will also be cumulated in MeRIP-Seq peak calling. We then compared the genomic distribution of the most frequent (appear more than 5 times) and most rare methylation sites (appear only once). As shown in Figure 6, compared with the most frequent methylation sites, the highly specific sites are highly enriched in 5'UTR region of mRNA. On lncRNA, however, they do not show distinct differences.

After extract the combinatorial epitranscriptome of more than 40k RNA m⁶A sites, their RPKM values are determined for each individual sample, and the methylation levels are calculated and standardized based on equation (1) and (2). Feature selection was conducted to select 3274 methylation sites with sample variance larger than 30. The selected methylation sites having methylation levels changing significantly across different conditions are then passed to clustering algorithm for discovery of the co-methylation patterns. Specifically, we applied four clustering algorithms to cluster the 3274 RNA methylation sites, including K-means, hierarchical clustering (HC), non-negative matrix factorization (NMF) and Bayesian Factor Regression Model (BFRM). An important predefined parameter for clustering analysis is the optimal number of clusters, which is pre-determined with cluster silhouettes of K-means method in a model selection procedure (see Figure 7). In our analysis, the other 3 clustering approaches adapt the same number of clusters for easy comparison purpose. It is worth mentioning that, the optimal number of clusters 3 is determined with a small subset of the data available (10 different conditions),

and the actual number of RNA methylation regulatory factors can be much larger with a lot more different cell types.

After applying all 4 clustering approaches to the selected 3274 unique RNA methylation sites, as it can be seen from Figure 8, the methylation sites were divided into 3 clusters. Considering the 4 clustering methods each embraces a different rationale, it is not surprising to see that there exist distinct differences among them. While the BFRM clusters are of approximately similar size, the other 3 algorithms, especially for K-means and HC, generated clusters with quite uneven sizes.

Given the aforementioned clustering results, it is important to check whether different clustering approaches capture relatively consistent co-methylation pattern. For this purpose, we define a new method for evaluating the consistency between two clustering approaches. Given the clustering results of N elements by two methods $\mathbf{c} = [c_1, c_2, \dots, c_N]$ and $\mathbf{r} = [r_1, r_2, \dots, r_N]$, where $c_i, r_i \in \{1, 2, \dots, K\}$ represents the cluster ID of the i -th element, and K represents the total number of clusters. We first convert the clustering IDs into a pair-wise resembling matrix (PRM) $\mathbf{C}$ and $\mathbf{R}$, specifically, for an element in pair-wise resembling matrix $c_{i,j}$ generated from $\mathbf{c} = [c_1, c_2, \dots, c_N]$, we define,

$$c_{i,j} = \begin{cases} -1 & i=j \\ 1 & i \neq j, c_i = c_j \\ 0 & i \neq j, c_i \neq c_j \end{cases} \quad (3)$$

Intuitively, $c_{i,j}$ represents whether the i -th element and j -th element are from the same cluster. In this way, the clustering results can be converted into a matrix recording whether two elements belong to the same cluster or not, and the conversion is unique and reversible as long as the cluster IDs have no order. The sensitivity and specificity of a clustering result $\mathbf{c} = [c_1, c_2, \dots, c_N]$ compared with a reference clustering result $\mathbf{r} = [r_1, r_2, \dots, r_N]$ can be conveniently defined as:

$$\begin{cases} \rho_{sen}(\mathbf{c}|\mathbf{r}) = p(r_{i,j}=1|c_{i,j}=1) \\ \rho_{spe}(\mathbf{c}|\mathbf{r}) = p(r_{i,j}=0|c_{i,j}=0) \end{cases} \quad (4)$$

Please note that, by random, the expected value of sensitivity and specificity are $1/K$ and $(K-1)/K$, respectively. And clustering IDs and reference clustering IDs are not switchable, i.e., $\rho_{sen}(\mathbf{c}|\mathbf{r}) \neq \rho_{sen}(\mathbf{r}|\mathbf{c})$. We may further define a consistency score ρ when a true reference clustering ID set is not available using,

$$\begin{aligned} \rho(\mathbf{c}, \mathbf{r}) &= \rho(\mathbf{r}, \mathbf{c}) \\ &= \frac{1}{4} [\rho_{sen}(\mathbf{c}|\mathbf{r}) + \rho_{spe}(\mathbf{c}|\mathbf{r})] + \frac{1}{4} [\rho_{sen}(\mathbf{r}|\mathbf{c}) + \rho_{spe}(\mathbf{r}|\mathbf{c})] \end{aligned} \quad (5)$$

In this way, the consistency between two clustering results $\mathbf{c} = [c_1, c_2, \dots, c_N]$ and $\mathbf{r} = [r_1, r_2, \dots, r_N]$ can be evaluated. It is worth mentioning that, the defined consistent score ρ in (5) have the following good properties: (1) The definition gives a score of 1 when the two clustering results are identical, and gives a score of 0.5 when the two clustering ID sets are

totally independent. (2) The definition of ρ is parameter-free. The definition of pair-wise resembling matrix (PRM) in (5) doesn't require the numbers of clusters are the same for the two clustering results. The consistency of the 4 clustering approaches is then compared pair-wisely, and the result is summarized in Table 3. Not surprisingly, the clustering results are highly correlated and well above the random 0.5 threshold. Specifically, the pair-wise consistency score between K-means, HC and NMF are always larger than 0.7; the only exception is BFRM with much lower consistency with the others. This is not surprising because BFRM represents a rationale fundamentally different from K-means, HC and NMF, i.e., the regulators (or factors) of BFRM model can up- or down-regulate some targets simultaneously, so 2 methylation sites belonging to the same BFRM cluster can be reversely correlated in methylation level. This phenomenon is not possible for the other 3 clusters.

The highly consistency between different clustering approaches indicate that co-methylation patterns are successfully captured. To find out whether the identified co-methylation patterns are corresponding to actual RNA methylation regulators, we compared them with the target genes of a known RNA m⁶A demethylase FTO [26]. Specifically, the FTO target sites in mouse midbrain are firstly identified using exomePeak package based on the m⁶A MeRIP-Seq data obtained from wild type mouse midbrain and FTO knockout condition [22]. Since it has been shown previously that m⁶A are conserved between human and mouse, these FTO target sites are lifted to human genome hg19 assembly with UCSC LiftOver tool [38]. Not surprisingly, more than 90% of the lifted FTO target sites overlap with our combinatorial RNA methylome. We then use Fisher's exact test (FET) to compare with the identified co-methylation patterns are significantly overlapping with the FTO target genes. Multiple hypothesis correction was also conducted to calculate the false discovery rate (FDR) using BH method [39]. As it is shown in Table 4, 3 out of 4 clustering approaches can successfully identify a co-methylation pattern significantly overlapping with FTO targets at significance level 0.05 after multiple hypothesis correction, showing that clustering methods can indeed identify biologically meaningful co-methylation pattern corresponding to the latent regulator of epitranscriptome. The most significant co-methylation pattern overlapping with FTO targets is from NMF with p -value 2.81e-06 and FDR 3.372e-05, suggesting the superior performance of NMF clustering approach over the other 3 methods at the current setting in finding the co-methylation pattern in the epitranscriptome. Interestingly, for all 4 clustering approaches, FTO targets sites are enriched in the largest of the 3 clusters. Indeed, FTO is the first mRNA demethylase discovered and plays a crucial role in human metabolism [5, 6, 29]. It is reasonable to assume FTO is one of the most influential regulators at epitranscriptomic layer. Please note that, the captured RNA co-methylation pattern is not due to any of the transcriptional regulations. Under the ideal case of a reversible chemical reaction, the methylation potential is independent from RNA abundance and determines directly the ratio of methylated and unmodified RNA molecules.

Among the 4 clustering approaches, NMF gives the most biologically significant result (see Figure 9). NMF decomposes the RNA methylome into the product of two non-negative matrices, which essentially assume a regression-like model. Under the framework of NMF, a single methylation site may be regulated by multiple RNA methylation regulators, which is

substantially different from Kmeans and HC. The recent identification of METTL3-WTAP-METTL14 RNA methylation complex [24, 27] provides a good example of complex dependence relationship. All 3 genes are required to form an RNA methylation complex. It is worth mentioning that, although the actual regulatory relationship at epitranscriptomic layer is more complex, it doesn't prohibit using a simpler and more straightforward approach.

Conceivably, the 3 clusters of RNA methylation sites identified from NMF are roughly corresponding to 3 regulators of the epitranscriptome, through which different biomolecular functions can be regulated at epitranscriptomic level. To reveal these functions, gene ontology enrichment analysis was conducted with topGO R/Bioconductor package [35] against the Biological processes category of gene ontology database. As it is shown in Figure 10, relative distinct biological functions are enriched in different clusters. There is essentially no overlap in function between different clusters when keeping the top 20 mostly enriched functions for each cluster, indicating the regulatory mechanism at epitranscriptomic level is potentially highly specific.

Specifically, the functions enriched in cluster 2, which is corresponding to FTO target sites, are shown in Figure 11. There are a number of important biological functions enriched in this cluster, including, neuron differentiation (p-value 0.0014), neurogenesis (0.0044), etc. Consistent previous studies of FTO, these functions are highly related to neuronal diseases [22] and cancer, indicating the prospect of RNA m⁶A and FTO studies.

Conclusion and Discussion

In summary, with a newly developed R package RMT, we extracted the combinatorial epitranscriptome of more than 40000 RNA methylation sites from 18 biological replicates obtained under 10 different conditions, and then partially clustered them into 3 groups using 4 different clustering approaches after feature and model selection. The methylation sites belonging to the same cluster are likely to be hyper- and hypo-methylated simultaneously (or co-methylated), indicating that they are regulated by a common RNA methylation factor (methyltransferase or demethylase). We defined a new approach evaluating the consistency between two clustering results and show that, despite the discrepancy between different clustering approaches, the 4 clustering results are still highly correlated, capturing consistent patterns embedded in the epitranscriptome. To examine whether the co-methylation patterns are biologically meaningful, the target RNA methylation sites of a known RNA demethylase (FTO) is identified from mouse and lifted to human, and we show that FTO target sites are significantly overlapping with identified co-methylation pattern. Many important biological functions are significantly enriched and thus may be regulated by different RNA methylation factors at the epitranscriptomic level, a layer of gene regulation that has been missing for decades [14].

Computational reconstruction of the entire RNA methylation network is especially difficult due to the following reasons: Firstly, till this day, only a small number of genes related to RNA methylation are identified [11]. Although it is possible to predict enzyme targets from enzyme knockout dataset, to our best knowledge, no effort has been made so far to

computationally predict the target sites of a specific RNA methyltransferase or demethylase. Secondly, so far there are only very limited amount of MeRIP-Seq dataset available. And the methylation levels for lowly expressed genes are difficult to estimate. It is clearly that precision achieved on lowly and highly expressed transcripts are different. From computational perspective, existing methods are not yet optimized for the intrinsic features of MeRIP-Seq, and advanced computational approaches that can take care of this discrepancy are still needed. With more data cumulated and revised mathematical models, it is promising that, future work may more explicitly and specifically associate methylation sites with RNA methylation factors.

Limited by the available data and knowledge, this study provides only the RNA methylation sites that are likely to share a common regulator, but fails to specify what actually those regulators (gene or protein) are. Nevertheless, it demonstrated, for the first time, the feasibility of dissecting the RNA methylome based on the RNA co-methylation patterns induced by RNA methyltransferases and demethylases, implying a promising direction in untangling the RNA methylome based on its regulators, through which, the biological meaning and underlying mechanism can be revealed in a deeper and more concise manner. Conceivably, with the eliminated regulatory relationship, the manipulating of the entire RNA methylome can be achieved through much less number of enzymatic regulators. This work can also inspire RNA methylation is an open question, many methods are not implanted. This work may also potentially be integrated with other related studies, e.g., a joint analysis of RNA methylation sites and transcription starting sites [40–42], or suggests the combinatorial patterns of different post-transcriptional RNA modifications, like in chromatin modifications and transcription factor binding [43, 44].

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the support from National Natural Science Foundation of China (61473232, 61401370, 91430111 and 61170134) to SWZ and JM; National Science Foundation of USA (CCF-1246073) to YH; The graduate starting seed fund of Northwestern Polytechnical University (Z2014145) to YCZ and LL; Fundamental Research Funds for the Central Universities (2014QNB47, 2014QNA84) to LZ and HL. We also thank computational support from the UTSA Computational Systems Biology Core, funded by the National Institute on Minority Health and Health Disparities (G12MD007591) from the National Institutes of Health.

References

1. Bernstein BE, Meissner A, Lander ES. The mammalian epigenome. *Cell*. Feb 23.2007 128:669–81. [PubMed: 17320505]
2. Bock C. Analysing and interpreting DNA methylation data. *Nat Rev Genet*. Oct.2012 13:705–19. [PubMed: 22986265]
3. Laird PW. Principles and challenges of genomewide DNA methylation analysis. *Nat Rev Genet*. Mar.2010 11:191–203. [PubMed: 20125086]
4. He C. Grand challenge commentary: RNA epigenetics? *Nat Chem Biol*. Dec.2010 6:863–5. [PubMed: 21079590]
5. Fu Y, Dominissini D, Rechavi G, He C. Gene expression regulation mediated through reversible m6A RNA methylation. *Nat Rev Genet*. 2014 vol. advance online publication.

6. Meyer KD, Jaffrey SR. The dynamic epitranscriptome: N6-methyladenosine and gene expression control. *Nat Rev Mol Cell Biol.* 2014 vol. advance online publication.
7. Liebers R, Rassoulzadegan M, Lyko F. Epigenetic Regulation by Heritable RNA. *PLoS Genet.* 2014; 10:e1004296. [PubMed: 24743450]
8. Desrosiers R, Friderici K, Rottman F. Identification of methylated nucleosides in messenger RNA from Novikoff hepatoma cells. *Proceedings of the National Academy of Sciences.* 1974; 71:3971–3975.
9. Schibler U, Kelley DE, Perry RP. Comparison of methylated sequences in messenger RNA and heterogeneous nuclear RNA from mouse L cells. *Journal of molecular biology.* 1977; 115:695–714. [PubMed: 592376]
10. Dubin DT, Taylor RH. The methylation state of poly A-containing-messenger RNA from cultured hamster cells. *Nucleic acids research.* 1975; 2:1653–1668. [PubMed: 1187339]
11. Machnicka MA, Milanowska K, Osman Oglou O, Purta E, Kurkowska M, Olchowik A, Januszewski W, Kalinowski S, Dunin-Horkawicz S, Rother KM, Helm M, Bujnicki JM, Grosjean H. MODOMICS: a database of RNA modification pathways--2013 update. *Nucleic Acids Res.* Jan.2013 41:D262–7. [PubMed: 23118484]
12. Meyer KD, Saletore Y, Zumbo P, Elemento O, Mason CE, Jaffrey SR. Comprehensive analysis of mRNA methylation reveals enrichment in 3' UTRs and near stop codons. *Cell.* Jun 22.2012 149:1635–46. [PubMed: 22608085]
13. Dominissini D, Moshitch-Moshkovitz S, Schwartz S, Salmon-Divon M, Ungar L, Osenberg S, Cesarkas K, Jacob-Hirsch J, Amariglio N, Kupiec M, Sorek R, Rechavi G. Topology of the human and mouse m6A RNA methylomes revealed by m6A-seq. *Nature.* May 10.2012 485:201–6. [PubMed: 22575960]
14. Saletore Y, Meyer K, Korlach J, Vilfan ID, Jaffrey S, Mason CE. The birth of the Epitranscriptome: deciphering the function of RNA modifications. *Genome Biol.* Oct 31.2012 13:175. [PubMed: 23113984]
15. Dominissini D, Moshitch-Moshkovitz S, Salmon-Divon M, Amariglio N, Rechavi G. Transcriptome-wide mapping of N(6)-methyladenosine by m(6)A-seq based on immunocapturing and massively parallel sequencing. *Nat Protoc.* Jan.2013 8:176–89. [PubMed: 23288318]
16. Zhang Y, Liu T, Meyer CA, Eeckhoutte J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM, Brown M, Li W. Model-based analysis of ChIP-Seq (MACS). *Genome Biol.* 2008; 9:R137. [PubMed: 18798982]
17. Park PJ. ChIP-seq: advantages and challenges of a maturing technology. *Nature reviews genetics.* 2009; 10:669–680.
18. Harris RA, Wang T, Coarfa C, Nagarajan RP, Hong C, Downey SL, Johnson BE, Fouse SD, Delaney A, Zhao Y. Comparison of sequencing-based methods to profile DNA methylation and identification of monoallelic epigenetic modifications. *Nature biotechnology.* 2010; 28:1097–1105.
19. Meng J, Cui X, Rao MK, Chen Y, Huang Y. Exome-based analysis for RNA epigenome sequencing data. *Bioinformatics.* Jun 15.2013 29:1565–1567. [PubMed: 23589649]
20. Li Y, Song S, Li C, Yu J. MeRIP-PF: an easy-to-use pipeline for high-resolution peak-finding in MeRIP-Seq data. *Genomics Proteomics Bioinformatics.* Feb.2013 11:72–5. [PubMed: 23434047]
21. Fustin J-M, Doi M, Yamaguchi Y, Hida H, Nishimura S, Yoshida M, Isagawa T, Masaki Morioka S, Kakeya H, Manabe I, Okamura H. RNA-Methylation-Dependent RNA Processing Controls the Speed of the Circadian Clock. *Cell.* 2013; 155:793–806. [PubMed: 24209618]
22. Hess ME, Hess S, Meyer KD, Verhagen LA, Koch L, Bronneke HS, Dietrich MO, Jordan SD, Saletore Y, Elemento O, Belgardt BF, Franz T, Horvath TL, Ruther U, Jaffrey SR, Kloppenburg P, Bruning JC. The fat mass and obesity associated gene (Fto) regulates activity of the dopaminergic midbrain circuitry. *Nat Neurosci.* Aug.2013 16:1042–8. [PubMed: 23817550]
23. Schwartz S, Agarwala SD, Mumbach MR, Jovanovic M, Mertins P, Shishkin A, Tabach Y, Mikkelsen TS, Satija R, Ruvkun G. High-Resolution Mapping Reveals a Conserved, Widespread, Dynamic mRNA Methylation Program in Yeast Meiosis. *Cell.* 2013

24. Liu J, Yue Y, Han D, Wang X, Fu Y, Zhang L, Jia G, Yu M, Lu Z, Deng X, Dai Q, Chen W, He C. A METTL3-METTL14 complex mediates mammalian nuclear RNA N6-adenosine methylation. *Nat Chem Biol.* 2014; 10:93–95. [PubMed: 24316715]
25. Wang Y, Li Y, Toth JI, Petroski MD, Zhang Z, Zhao JC. N6-methyladenosine modification destabilizes developmental regulators in embryonic stem cells. *Nat Cell Biol.* 2014 vol. advance online publication.
26. Jia G, Fu Y, Zhao X, Dai Q, Zheng G, Yang Y, Yi C, Lindahl T, Pan T, Yang Y-G. N6-methyladenosine in nuclear RNA is a major substrate of the obesity-associated FTO. *Nature chemical biology.* 2011; 7:885–887.
27. Ping X-L, Sun B-F, Wang L, Xiao W, Yang X, Wang W-J, Adhikari S, Shi Y, Lv Y, Chen Y-S, Zhao X, Li A, Yang Y, Dahal U, Lou X-M, Liu X, Huang J, Yuan W-P, Zhu X-F, Cheng T, Zhao Y-L, Wang X, Danielsen JMR, Liu F, Yang Y-G. Mammalian WTAP is a regulatory subunit of the RNA N6-methyladenosine methyltransferase. *Cell Res.* 2014
28. Khoddami V, Cairns BR. Identification of direct targets and modified bases of RNA cytosine methyltransferases. *Nat Biotechnol.* May.2013 31:458–64. [PubMed: 23604283]
29. Lee M, Kim B, Kim VN. Emerging Roles of RNA Modification: m6A and U-Tail. *cell.* 2014; 158:980–987. [PubMed: 25171402]
30. Carmel, R.; Jacobsen, DW. Homocysteine in health and disease. Cambridge University Press; 2001.
31. Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome Biol.* 2013; 14:R36. [PubMed: 23618408]
32. Przyborowski J, Wilenski H. Homogeneity of results in testing samples from Poisson series: With an application to testing clover seed for dodder. *Biometrika.* 1940; 31:313–323.
33. Carvalho CM, Chang J, Lucas JE, Nevins JR, Wang Q, West M. High-Dimensional Sparse Factor Modeling: Applications in Gene Expression Genomics. *J Am Stat Assoc.* Dec 1.2008 103:1438–1456. [PubMed: 21218139]
34. Lee DD, Seung HS. Algorithms for non-negative matrix factorization. *Advances in neural information processing systems.* 2001:556–562.
35. Alexa, A.; Rahnenfuhrer, J. R package version 2.8. 2010. topGO: enrichment analysis for gene ontology.
36. Machanick P, Bailey TL. MEME-ChIP: motif analysis of large DNA datasets. *Bioinformatics.* Jun 15.2011 27:1696–1697. [PubMed: 21486936]
37. Lawrence M, Huber W, Pages H, Aboyoun P, Carlson M, Gentleman R, Morgan MT, Carey VJ. Software for computing and annotating genomic ranges. *PLoS Comput Biol.* Aug.2013 9:e1003118. [PubMed: 23950696]
38. Lawrence, M.; Carey, V.; Gentleman, R.; XML, I.; IRanges, L.; Lawrence, MM. Package 'tracklayer'. 2013.
39. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B (Methodological).* 1995:289–300.
40. Chen W, Lei T-Y, Jin D-C, Lin H, Chou K-C. PseKNC: A flexible web server for generating pseudo K-tuple nucleotide composition. *Analytical biochemistry.* 2014; 456:53–60. [PubMed: 24732113]
41. Guo S-H, Deng E-Z, Xu L-Q, Ding H, Lin H, Chen W, Chou K-C. iNuc-PseKNC: a sequence-based predictor for predicting nucleosome positioning in genomes with pseudo k-tuple nucleotide composition. *Bioinformatics.* Jun 1.2014 30:1522–1529. [PubMed: 24504871]
42. Chen W, Feng P-M, Deng E-Z, Lin H, Chou K-C. iTIS-PseTNC: A sequence-based predictor for identifying translation initiation site in human genes using pseudo trinucleotide composition. *Analytical Biochemistry.* 2014; 462:76–83. [PubMed: 25016190]
43. Ferraris L, Stewart AP, Kang J, DeSimone AM, Gemberling M, Tantin D, Fairbrother WG. Combinatorial binding of transcription factors in the pluripotency control regions of the genome. *Genome research.* 2011; 21:1055–1064. [PubMed: 21527551]

44. Kato M, Hata N, Banerjee N, Futcher B, Zhang MQ. Identifying combinatorial regulation of transcription factors and binding motifs. *Genome biology*. 2004; 5:R56. [PubMed: 15287978]

Figure 1. Illustration of MeRIP-Seq Protocol

In MeRIP-Seq, two types of samples (IP and control samples) can be generated. In the beginning of the protocol, RNA molecules are firstly sheared into fragments of around 100nt. Through anti-m⁶A antibody, the IP sample provides unbiased measurement of the methylated RNA fragments; the control sample reflects the basal RNA abundance, including both the methylated and unmodified RNA molecules.

Figure 2. Regulation of RNA methylome

The absolute amount of RNA methylation is affected by transcriptional regulation and enzymatic regulation. On one hand, transcriptional regulation changes the absolute amount of RNA molecules, keeping the relative amount unchanged; on the other hand, enzymatic regulation changes the percentage of methylated RNA with the total amount of RNA molecules unchanged. In this figure, the absolute amount of RNA methylation increases 4 times in cancer compared with the control condition, which is due to a joint effect of transcriptional up-regulation (fold change of 2) and enzymatic hyper-methylation (fold change of 2).

Figure 3. Work Flow of RMT package

The RMT package is developed for convenient extracting the combinatorial epitranscriptome from multiple MeRIP-Seq datasets obtained from different conditions. Specifically, it requires the input of multiple MeRIP-Seq datasets in the form of aligned BAM files, and outputs all the RNA methylation sites together with their estimated methylation levels. The RMT package is available in Supplementary Material S1.

Figure 4. An RNA methylation site spanning a splicing site

The figure shows an RNA methylation site spanning a splicing site on the CDS of ASXL2 (Entrez Gene ID: 55252). The reads are consistently more enriched in the IP sample compared with the input control sample across 10 different experimental conditions. As shown in the HEK293 and HepG2 tracks, this methylation site consists of several sub-peaks and thus may represent more than 1 methylated residual. It is possible that this peak is detected as several non-joint peaks under some conditions, but this merged unique RNA methylation site is chosen with our aforementioned procedure.

Figure 5. Distribution of RNA methylation sites

A. The number of unique RNA methylation sites on each biological replicate. Please note that, the sites here represents the merged peaks, which can be different from the number of peaks called by ExomePeak on this individual sample. **B.** This subfigure shows the distribution of unique RNA methylation sites on mRNA and lncRNA with each curve representing one IP sample (total 18). Consistent with previous studies, on mRNA, the RNA methylation sites are highly enriched near stop codon and on 3'UTR region. Interestingly, the RNA methylation sites on lncRNA are slightly more enriched on the 5'end compared with 3'end.

Figure 6. Distribution of unique RNA methylation sites with different occurrence frequency
 Compared with the most frequent methylation sites (blue curve), the highly specific sites (red curve) are highly enriched in 5'UTR region of mRNA. On lncRNA, the two sets do not show distinct differences.

Figure 7. Silhouette value of K-means clustering method

Silhouette Value compares the intra-group similarity to closest group similarity. Using K-means approach, the selected 3274 RNA methylation sites are clustered into 3 groups with 1780, 595 and 899 sites, respectively. And in our analysis, the other 3 clustering approaches adapt the same number of clusters for easy comparison. It is worth mentioning that, the optimal number of clusters 3 is determined with a small subset of the data available (10 different conditions), and the actual number of RNA methylation regulatory factors can be much larger with a lot more different cell types.

Figure 8. clustering results of 4 approaches

Figure shows the number of peaks in 3 clusters determined by 4 different clustering approaches. The largest cluster was generated by HC with 2272 RNA methylation site (69.4% of 3274). Compared with the other 3 clustering approaches, BFRM generates a well-balanced result with clusters of almost equal size. It is worth mentioning that, the cluster IDs are generated with no specific order, and the clusters generated by different numbers with the same ID are not necessarily corresponding to the same physical RNA co-methylation pattern.

Figure 9. Decompose RNA methylome by NMF

NMF decomposes the RNA methylome into the product of two nonnegative matrices, i.e., the basis components, which represents the loading or regulatory relationship, and the mixture components, which represents the activities of the regulators. Interestingly, although the biological replicates under the same condition are modeled independently by NMF, the estimated regulator activities (mixture coefficients) are highly consistent on the biological replicates from the same or similar conditions. E.g., the activities of regulars are highly consistent on U2OS, HEK293K and HepG2, respectively. The clustering result of NMF is obtained by assigning an RNA methylation site to the factor with the largest absolute value of regulatory coefficient. Please note that, although NMF in theory allows a single RNA methylation site regulated by multiple factors, we consider only the strongest regulator in the analysis.

Figure 10. Gene ontology enrichment result of K-means
Relative distinct biological functions are enriched in different clusters. There is essentially no overlap in function between different clusters when keeping the top 20 mostly enriched functions for each cluster. This analysis was conducted by topGO R/Bioconductor package [35].

Figure 11. Biological processes in FTO-related co-methylation pattern

Cluster 3 of K-means result is sharing significantly with FTO targeted genes. A number of important biological functions are enriched in this cluster, including, regulation of neuron apoptotic process (p-value 0.0052), programmed cell death (p-value 0.0451), neuron death (0.0265), neuron apoptotic process, etc. These functions are highly related to neuronal diseases and cancer.

Table 1

Human MeRIP-Seq Datasets

ID	Tissue/Cell	Treatment	Other Info	# of Replicates IP & Input	# of Reads (Million) IP & Input	Source
h1	HEK293T		SYSY Ab	2 & 3	145 & 217	[12]
h2			NEB Ab	1 & 3	33 & 217	[12]
h3	Brain			1 & 1	22 & 17	[13]
h4				4 & 3	68 & 85	[13]
h5	HepG2	UV		1 & 1	21 & 7	[13]
h6		HS		1 & 1	34 & 52	[13]
h7		HGF		1 & 1	33 & 23	[13]
h8		IFN		1 & 1	47 & 27	[13]
h9	U2OS			3 & 3	86 & 83	[21]
h10		DAA		3 & 3	80 & 87	[21]

Table 2

RNA methylome of a specific biological replicate

ID	Tissue	Treatment	Other Info	Sample	# of peaks	Motif	E-value
h1	HEK293T		SYSY Ab	S1	21114		2.8e-147
				S2	17492		2.2e-146
h2	HEK293T		NEB Ab	S3	12118		1.8e-134
h3				S4	1200		1.5e-019
h4	Brain			S5	18166		1.3e-191
				S6	21127		1.9e-168
				S7	16441		2.6e-196
				S8	15126		8.3e-102
h5	HepG2	UV		S9	7545		4.1e-124
h6		HS		S10	16301		2.2e-140

ID	Tissue	Treatment	Other Info	Sample	# of peaks	Motif	E-value
h7		HGF		S11	9477		2.9e-106
		IFN		S12	8629		6.3e-011
h9	U2OS			S13	20330		2.1e-085
				S14	15931		4.7e-076
				S15	15698		1.8e-109
h10		DAA		S16	13342		1.2e-054
				S17	15985		8.7e-060
				S18	12442		2.7e-055

Table 3

Consistency of the 4 clustering approaches

	K-means	HC	NMF	BFRM
K-means	NA	0.772	0.805	0.594
HC	0.772	NA	0.701	0.574
NMF	0.805	0.701	NA	0.584
BFRM	0.594	0.574	0.584	NA

Table 4

Result of FTO target enrichment analysis

Algorithms	Cluster ID	# of sites	p-value	FDR	Odds ratio
K-means	Cluster 1	1780	0.012	0.048	1.651
	Cluster 2	595	0.977	0.999	0.568
	Cluster 3	899	0.884	0.999	0.769
HC	Cluster 1	316	0.734	0.999	0.837
	Cluster 2	2272	0.051	0.153	1.523
	Cluster 3	686	0.963	0.999	0.629
NMF	Cluster 1	998	0.999	0.999	0.408
	Cluster 2	1597	2.81e-06	3.372e-05	2.793
	Cluster 3	679	0.995	0.999	0.481
BFRM	Cluster 1	1249	0.003	0.018	1.791
	Cluster 2	990	0.655	0.999	0.934
	Cluster 3	1035	0.999	0.999	0.484

Note: Three out of 4 clustering approaches can successfully identify a co-methylation pattern corresponding to FTO target sites at significance level 0.05 after multiple hypothesis correction. Interestingly, for all 4 clustering approaches, FTO targets sites are enriched in the largest of the 3 clusters. Indeed, FTO is the first mRNA demethylase discovered and plays a crucial role in human metabolism [5, 6, 29]. It is reasonable to assume FTO is one of the most influential regulators at epitranscriptomic layer.