Published in final edited form as:

Cytokine. 2014 December ; 70(2): 134–140. doi:10.1016/j.cyto.2014.06.018.

Longitudinal Changes in Serum Proinflammatory Markers across Pregnancy and Postpartum: Effects of Maternal Body Mass Index

Lisa M. Christian, PhD^{1,2,3,4} and Kyle Porter, MAS⁵

¹Department of Psychiatry, The Ohio State University, Medical Center

²The Institute for Behavioral Medicine Research, The Ohio State University, Medical Center

³Department of Obstetrics and Gynecology, The Ohio State University, Medical Center

⁴Department of Psychology, The Ohio State University, Medical Center

⁵Center for Biostatistics, The Ohio State University, Medical Center

Abstract

Background—The maternal immune system undergoes substantial changes to support healthy pregnancy. Although obesity is a primary driver of inflammation and predictive of perinatal complications, additive effects of pregnancy and obesity on changes in inflammatory processes are not well delineated.

Methods—This study examined serum proinflammatory markers interleukin(IL)-6, IL-8, tumor necrosis factor(TNF)- α , IL-1 β , and C-reactive protein(CRP) during each trimester of pregnancy and 4-6 weeks postpartum among 57 women.

Results—Overall, IL-6 showed an increasing trend across pregnancy and significant increase at postpartum. Similarly, TNF- α increased significantly across gestation, with a further increase at postpartum. Both IL-8 and IL-1 β showed a U-shaped curve, decreasing from early to later pregnancy, and increasing at postpartum. Finally, serum CRP decreased significantly across pregnancy, with further decreases at postpartum. Maternal obesity predicted higher IL-6 at each study visit. Obese women showed a trend toward elevated serum CRP during pregnancy, and significantly higher levels at postpartum.

Discussion—The course of pregnancy and postpartum is characterized by significant changes in serum proinflammatory mediators. Obese women show elevations in serum proinflammatory

© 2014 Elsevier Ltd. All rights reserved.

Address correspondence and reprint requests to Lisa M. Christian, PhD, Institute for Behavioral Medicine Research Room 112, 460 Medical Center Drive, The Ohio State University Medical Center, Columbus, Ohio 43210. Lisa.Christian@osumc.edu Phone: 614-293-0936 Fax: 614-293-4200.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflicts of Interest

The authors report no conflicts of interest.

markers relative to normal weight women during pregnancy and postpartum. Further research is needed to determine the extent to which obesity-induced inflammation affects maternal and fetal health.

Keywords

inflammation; pregnancy; postpartum; cytokines; obesity; body mass index; inflammatory markers; longitudinal

The maternal immune system undergoes substantial changes to support healthy pregnancy. Compared to nonpregnancy, normal pregnancy is characterized by mild elevations in both serum pro- and antiinflammatory cytokine levels [1-8]. It has been proposed that at the local uterine/placental level, early pregnancy (i.e., implantation) and late pregnancy (i.e., approaching delivery) are inflammatory states while mid-pregnancy is an anti-inflammatory state [9]. However, few studies have examined longitudinal changes in circulating serum cytokines as pregnancy progresses or during the postpartum transition. Moreover, the majority of existing studies on this topic are small, cross-sectional in design, and/or do not include assessment at every trimester or at postpartum. Thus, the typical course of change in serum inflammatory markers in normal pregnancy and postpartum is not fully delineated.

While available evidence suggests that healthy pregnancy is typified by an enhanced inflammatory state, studies also show that *excessive* inflammation is incompatible with healthy pregnancy. Elevations in proinflammatory cytokines in maternal serum and amniotic fluid are causally implicated in risk of preterm delivery in the context of infection as well as idiopathic cases [10-14]. Proinflammatory cytokines can promote preterm labor by triggering preterm contractions, encouraging cervical ripening, and causing rupture of the membranes [15, 16]. Inflammatory pathways are also implicated in the development of gestational hypertension [17-23] and gestational diabetes [24]. Moreover, maternal inflammation has been associated with effects on fetal development including risk for neurobehavioral disorders and adverse metabolic changes [25-27].

Because adipocytes secrete proinflammatory cytokines, obesity is a primary promoter of inflammation. Obesity in pregnancy is linked to risk of gestational hypertension and gestational diabetes with inflammatory mechanisms serving as clear drivers [24, 28]. Moreover, maternal obesity has been linked to adverse perinatal outcomes including fetal death and premature birth as well as increased risk of metabolic syndrome, diabetes, obesity, and neurodevelopmental disorders in offspring [29, 30]. Despite clear health relevance, limited data are available regarding the potential synergistic relationship between pregnancy and obesity in affecting inflammatory processes.

The current study examined longitudinal changes in the serum proinflammatory markers interleukin(IL)-6, IL-8, tumor necrosis factor (TNF)- α , IL-1 β , and C-reactive protein (CRP) during each trimester of pregnancy and at 4-6 weeks postpartum in a racially diverse sample of predominately lower income women. These markers were selected based on their relevance to pregnancy-related health outcomes including preterm birth, gestational hypertension/preeclampsia, and fetal brain injury [31-34] as well as obesity [35-37]. It was hypothesized that, in the sample overall, there would be significant changes in inflammatory

markers across the course of pregnancy and postpartum. It was also hypothesized that greater maternal body mass index (BMI) would be associated with elevated inflammatory markers across the study period.

Methods

Study Design

Sixty pregnant women were recruited from the Ohio State University Medical Center (OSUMC) Prenatal Clinic. Study visits were conducted during the 1st trimester (Mean= 11.1 $\pm$ 2.3 weeks gestation), 2nd trimester (Mean= 22.9 $\pm$ 2.3 weeks gestation), 3rd trimester (Mean= 31.4 $\pm$ 1.8 weeks gestation) and at 4-6 weeks postpartum (Mean= 4.1 $\pm$ 2.1 weeks). At each visit, women provided a blood sample. Underweight women (BMI < 18.5) were excluded from the current analyses due to low representation (n=3), resulting in a final sample of 57.

Participants

All women were born and raised in the United States. Women were not eligible if they had current hypertension, diabetes, chronic conditions with implications for immune function (e.g., rheumatoid arthritis, multiple sclerosis, or human immunodeficiency virus), fetal anomaly, illicit drug use or more than two alcoholic drinks per week during pregnancy (per self-report or medical record). Women reporting acute illness (e.g., cold or flu-like symptoms) or antibiotic use within 10 days of a study visit were rescheduled. Each completed informed consent and received modest compensation. The study was approved by the OSU Biomedical Institutional Review Board.

Demographics

Age, race/ethnicity, marital status, education, annual family income, and number of prior pregnancies (parity) were collected by self-report.

Body Mass Index

Pre-pregnancy body mass index (BMI; kg/m²) was calculated using self-reported pre-pregnancy weight and height measured at the first visit.

Health Behaviors

At the initial study visit, information on current and past smoking as well as frequency of participation in vigorous physical activity was collected.

Statistical Analysis

Women were categorized based on BMI categories using the following standard ranges: 18.5 - 24.9 = normal weight, 25 - 29.9 = overweight, and ≥ 30 = obese. To evaluate demographic similarity between groups, participant characteristics were compared between groups by ANOVA for continuous variables and chi-square tests for categorical variables. The primary endpoints were serum levels of IL-6, TNF- α , IL-1 β , IL-8, and CRP at each trimester of pregnancy and postpartum. Each endpoint was log transformed for analysis to

meet normality assumptions. For descriptive purposes, Pearson's correlations were conducted to determine associations between cytokines at each study visit. A linear mixed model was fit to each endpoint across the four visits. The model included main effects for BMI group and study visit, the interaction between BMI group and study visit, and a random effect for subject with a compound symmetry covariance structure to account for correlation among measures from the same subject. Parameter estimate contrasts were constructed to test for differences over pregnancy for all subjects (comparing each pair of visits) and differences between BMI groups at each study visit. Each comparison was performed at the $\alpha = 0.05$ significance level. With this testing approach, the expected number of false positives for each endpoint was 0.9. All analyses were performed using SAS® software, version 9.2.

Results

Demographics and Health Behaviors

Participant characteristics for each BMI group are reported in Table 1. As shown, the BMI groups did not differ significantly in age, race, parity, marital status, income, education, smoking history, or exercise ($p \leq 0.07$).

Correlations Between Biomarkers

To determine associations between biomarkers at each study visit, Pearson's correlations were conducted. As shown in Tables 2-5, inflammatory markers were highly correlated with each other at each study visit. These patterns of association were highly similar at each stage of assessment.

Longitudinal Changes in Biomarkers Overall

Trajectories of change for each inflammatory marker are presented in Figures 1-5. In the sample overall, a non-significant trend was observed for increases in IL-6 across pregnancy (1st versus 3rd trimester $p = 0.12$). In addition, IL-6 increased significantly from the 3rd trimester to postpartum ($p = 0.01$). Similarly, TNF- α increased significantly from early ($p = 0.001$) and middle to later pregnancy ($p = 0.01$), with a further increase at postpartum relative to the 3rd trimester ($p = 0.03$). In contrast, IL-8 showed a U-shaped curve, decreasing in the 2nd trimester relative to the 1st trimester ($p = 0.03$) and increasing significantly at postpartum follow-up ($p < 0.001$). Similarly, IL-1 β showed a U-shaped curve, decreasing from early to late pregnancy and increasing significantly at postpartum ($p = 0.01$). CRP decreased from the 1st to 3rd trimester ($p = 0.02$), and decreased further at postpartum ($p = 0.01$).

Longitudinal Changes in Biomarkers by BMI

Next, analyses examined the moderating effect of BMI on changes in biomarkers across pregnancy and postpartum. Both obese and overweight women had significantly higher serum IL-6 than normal weight women at all three pregnancy study visits ($p < 0.04$). At postpartum IL-6 was higher in obese women than in normal weight women ($p = 0.004$). For TNF- α , overweight women showed elevations at each pregnancy study visit relative to normal weight women ($p < 0.04$).

During pregnancy, there was a trend toward higher CRP in obese women relative to normal weight women at each pregnancy study visit ($p = 0.052$). Obese women had significantly elevated CRP at postpartum as compared to overweight ($p = 0.04$) or normal weight women ($p = 0.045$). No significant differences were observed in IL-1 β or IL-8 based on BMI category.

Discussion

The role of inflammation in pregnancy has been a topic of debate; the allograft paradigm forwards that in order to prevent rejection of the fetus, the maternal immune system must acquire an anti-inflammatory state during pregnancy. More recently, it has been suggested that, at least at the local level, each stage of pregnancy is characterized by a unique inflammatory environment with the first and third trimesters typified by a proinflammatory state and the second trimester an antiinflammatory state [38]. However, the typical course of changes in inflammatory markers at the peripheral level has not been well-delineated. Moreover, the extent to which obesity, a known promoter of chronic low-grade inflammation, may alter such trajectories is not known. Thus, delineating the course of immune change across the perinatal period in relation to maternal weight is of clinical importance.

IL-6 is a key cytokine associated with adverse perinatal health outcomes [39-44]. Notably, IL-6 is secreted by adipose tissue [35]. In this study, IL-6 showed a non-significant trend toward increases across the course of pregnancy with significant increases at postpartum relative to the 3rd trimester. As hypothesized, both overweight and obese women showed elevated IL-6 relative to normal weight women at each pregnancy visit. A similar pattern at postpartum was statistically significant for obese women only. These findings are consistent with prior evidence that IL-6 is elevated during pregnancy as compared to non-pregnancy, with greater elevations in women with higher body mass [1, 45]. The current data replicate and extend prior findings to the postpartum period.

TNF- α is also secreted by adipose tissue. In this study, serum TNF- α increased significantly from early to late pregnancy, with further increases at postpartum. This is consistent with prior studies showing increasing TNF- α across gestation [46, 47]. Overweight women showed elevations in TNF- α at each pregnancy visit. However, serum TNF- α values in obese women were between normal weight and overweight women; thus, an incremental increase in TNF- α based on BMI was not observed.

Both IL-8 and IL-1 β showed a U-shaped curve, with decreases in mid or late pregnancy compared to early pregnancy, and subsequent increases at postpartum. This pattern of response could indicate an initial inflammatory response in early pregnancy that is down-regulated over time. Contrary to prediction, no differences in IL-1 β or IL-8 were observed based on maternal body mass. As adipokines, both IL-1 β and IL-8 tend to be increased in in obesity [36, 48, 49], however central adiposity may be a more sensitive predictor of inflammatory status than BMI [50]. Thus, although difficult to accurately quantify in pregnant women, future studies should include other measures of adiposity in addition to

BMI; in particular, waist-hip ratio measured in early pregnancy should be reasonably valid as limited weight gain occurs during the first trimester.

C-reactive protein (CRP) is an acute-phase protein synthesized by the liver in response to inflammatory mediators, particularly IL-6. In the current study, serum CRP decreased significantly from early to late pregnancy in the overall sample, with further decreases at postpartum. This is consistent with prior longitudinal studies showing decreases in CRP across gestation [51, 52]. However, other studies have found increasing levels [53] or no significant changes in serum CRP across pregnancy [8]. Thus, findings in relation to CRP are inconsistent. In the hypothesized direction, obese women showed a trend toward higher CRP during pregnancy as well as significantly higher CRP at postpartum than normal weight women.

IL-6 is a potent inducer of the acute-phase response which is characterized by hepatic release of CRP [54]. Thus, as expected, IL-6 and CRP levels were highly correlated at each study visit. Despite these correlations, different patterns of change in IL-6 versus CRP across time were observed. IL-6 release does not result in concomitant CRP production in certain circumstances. For example, in response to acute exercise, substantial increases in serum IL-6 do not result in corresponding increases in CRP [55]. In addition to instigating the acute phase response, IL-6 serves multiple functions including effects on metabolism, regulation of bone homeostasis, and pain regulation [56]. The functional roles for inflammatory markers and the relative importance of given roles at different stages of gestation remain to be fully explicated.

Notably, with the exception of CRP, all inflammatory mediators assessed (IL-6, TNF- α , IL-8, and IL-1 β) increased at 4-6 weeks postpartum relative to the 3rd trimester. This may be related to the substantial neuroendocrine changes occurring at postpartum, particularly the considerable drop in cortisol. In addition, regardless of mode of delivery, healing processes occur following delivery which may instigate inflammatory responses [57]. In addition to affecting the healing process, functioning of the immune system following childbirth may have clinical implications for risk of postpartum mood disorders [58, 59]. Moreover, certain autoimmune diseases including rheumatoid arthritis and multiple sclerosis tend to remit during pregnancy, but become exacerbated at postpartum [60, 61]. Thus, propensity toward inflammatory responses following childbirth also has clinical relevance in this context.

A strength of this study is utilization of a longitudinal rather than cross-sectional design. In addition, inclusion of a postpartum timepoint extends prior findings. However, this study did not include a pre-pregnancy assessment or a non-pregnant comparison group. Thus, we were unable to determine the extent to which first trimester values may be altered relative to the non-pregnant state. Although we assessed women at postpartum, this does not represent a typical non-pregnant state since substantial neuroendocrine changes and healing processes occur following childbirth, as described. However, based on prior studies which have included a non-pregnant control group or women assessed several months postpartum [e.g., 6, 7, 8], we expect that women in this study had elevations in inflammatory markers during each trimester as compared to a non-pregnant state, including markers such as IL-8 and CRP which showed a decreasing course from earlier to later pregnancy.

This study was observational and did not address possible mechanisms behind the observed changes. Changes in cytokine levels during pregnancy likely reflect a combination of 1) responses to the changing neuroendocrine environment, 2) functional changes to support pregnancy (which may be mechanistically supported by neuroendocrine changes) and 3) a reflection of the physiological stress of pregnancy. This study did not examine the extent to which concomitant changes in neuroendocrine function (e.g., cortisol, progesterone) correspond to the observed changes. However, even with such data, delineating causal relationships between neuroendocrine and immune parameters in human pregnancy within an observational design is not possible because numerous complex changes occur relatively simultaneously within short time periods. With regard to functional changes, many cytokines, including IL-6 as described above, are pleiotropic – having multiple effects [62]. Moreover, the ultimate effects of cytokines are affected by cytokine receptor function and related sensitivity of immune cells. Future studies should aim to identify the functional effects of specific cytokines during each stage of gestation, both at the peripheral and local level, as these functions should provide insight into mechanisms underlying observed changes. Finally, pregnancy is a considerable physical stressor on the body; it is marked by nearly 50% increase in blood volume, changes in lung volume and kidney function, as well as weight increases that reflect the developing baby as well as placental growth, blood volume, and fat stores. These considerable physical stressors of pregnancy likely affect inflammatory processes.

This study examined serum proinflammatory proteins, but did not examine inflammatory responses to a stimulus. Such data would be highly informative, as prior studies show that conditions with an inflammatory component including depression and coronary artery disease as well as obesity are associated with dysregulation of inflammatory responses [63-66]. Moreover, despite elevations in serum proinflammatory markers during pregnancy versus non-pregnancy, available evidence from both human and animal studies indicates that inflammatory *responses* are attenuated [67-71]. Because responses to *in vivo* as well as *in vitro* stimuli provide greater variability across individuals than serum levels, examination of responses to challenges may ultimately provide greater ability to differentiate women showing typical versus atypical (i.e., dysregulated) immune function during pregnancy. This should be considered in future studies.

In conclusion, these data demonstrate that pregnancy and postpartum are characterized by different trajectories of change for specific inflammatory markers. In addition, women with higher body mass show elevations in inflammatory markers across pregnancy and postpartum, but similar trajectories of change in these markers as do normal weight women. As reviewed, excessive inflammation has been associated with a variety of adverse perinatal health outcomes, including preterm delivery, effects on fetal development, and risk of serious maternal health conditions including gestational diabetes and gestational hypertension [72, 73]. Given such associations, there is growing interest in describing typical changes in immune parameters across pregnancy, identifying deviations with health relevance, and determining risk factors for such deviations. The current investigation advances our knowledge in this regard. Further research is needed to determine the extent to

which obesity-induced inflammation during pregnancy and postpartum may mediate the association between maternal obesity and adverse perinatal health outcomes.

Acknowledgments

We appreciate the contributions of Clinical Research Assistants Colleen Sagrilla, Kelly Marceau, and Rebecca Long to data collection and Research Associate Hui Xu for conducting the cytokine assays. We thank Jay Iams, MD, for his support in participant recruitment. We would like to thank our study participants and the staff at the OSU Clinical Research Center and Wexner Medical Center Prenatal Clinic.

Role of the Funding Sources

This study was supported by NINR (R01 NR013661, LMC) and NICHD (R21 HD067670, LMC). The project described was supported by Award Number Grant UL1TR000090 from the National Center For Advancing Translational Sciences. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Center For Advancing Translational Sciences or the National Institutes of Health. NIH had no further role in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the paper for publication.

References

- [1]. Curry AE, Vogel I, Skogstrand K, Drews C, Schendel DE, Flanders WD, et al. Maternal plasma cytokines in early- and mid-gestation of normal human pregnancy and their association with maternal factors. *J Reprod Immunol*. 2008; 77:152–60. [PubMed: 17692390]
- [2]. Vassiliadis S, Ranella A, Papadimitriou L, Makrygiannakis A, Athanassakis I. Serum levels of pro- and anti-inflammatory cytokines in non-pregnant women, during pregnancy, labour and abortion. *Mediators Inflamm*. 1998; 7:69–72. [PubMed: 9836491]
- [3]. Austgulen R, Lien E, Liabakk NB, Jacobsen G, Arntzen KJ. Increased Levels of Cytokines and Cytokine Activity Modifiers in Normal-Pregnancy. *Eur J Obstet Gynecol Reprod Biol*. 1994; 57:149–55. [PubMed: 7713287]
- [4]. Opsjon SL, Wathen NC, Tingulstad S, Wiedswang G, Sundan A, Waage A, et al. Tumor-Necrosis-Factor, Interleukin-1, and Interleukin-6 in Normal Human-Pregnancy. *Am J Obstet Gynecol*. 1993; 169:397–404. [PubMed: 8362955]
- [5]. Makhseed M, Raghupathy R, Azizieh F, Farhat R, Hassan N, Bandar A. Circulating cytokines and CD30 in normal human pregnancy and recurrent spontaneous abortions. *Hum Reprod*. 2000; 15:2011–7. [PubMed: 10967006]
- [6]. Sharma A, Satyam A, Sharma JB. Leptin, IL-10 and inflammatory markers (TNF-alpha, IL-6 and IL-8) in pre-eclamptic, normotensive pregnant and healthy non-pregnant women. *Am J Reprod Immunol*. 2007; 58:21–30. [PubMed: 17565544]
- [7]. Kraus TA, Sperling RS, Engel SM, Lo YT, Kellerman L, Singh T, et al. Peripheral Blood Cytokine Profiling During Pregnancy and Post-partum Periods. *Am J Reprod Immunol*. 2010; 64:411–26. [PubMed: 20712812]
- [8]. Belo L, Santos-Silva A, Rocha S, Caslake M, Cooney J, Pereira-Leite L, et al. Fluctuations in C-reactive protein concentration and neutrophil activation during normal human pregnancy. *Eur J Obstet Gynecol Reprod Biol*. 2005; 123:46–51. [PubMed: 16260340]
- [9]. Mor G, Cardenas I, Abrahams V, Guller S. Inflammation and pregnancy: the role of the immune system at the implantation site. *Ann N Y Acad Sci*. 2011; 1221:80–7. [PubMed: 21401634]
- [10]. Dizon-Townson DS. Preterm labour and delivery: a genetic predisposition. *Paediatr Perinat Epidemiol*. 2001; 15:57–62. [PubMed: 11520400]
- [11]. Romero R, Avila C, Santhanam U, Sehgal PB. Amniotic fluid interleukin-6 in pre-term labor: association with infection. *J Clin Invest*. 1990; 85:1392–400. [PubMed: 2332497]
- [12]. Romero R, Yoon BH, Mazor M, Gomez R, Diamond MP, Kenney JS, et al. The diagnostic and prognostic value of amniotic fluid white blood cell count, glucose, interleukin-6, and Gram stain in patients with preterm labor and intact membranes. *Am J Obstet Gynecol*. 1993; 169:805–16. [PubMed: 7694461]

- [13]. Romero R, Yoon BH, Mazor M, Gomez R, Gonzales R, Diamond MP. A comparative study of the diagnostic performance of amniotic fluid glucose, white blood cell count, interleukin-6, and Gram stain in the detection of microbial invasion in patients with preterm premature rupture of membranes. *Am J Obstet Gynecol.* 1993; 169:839–51. [PubMed: 7694463]
- [14]. Murtha AP, Sinclair T, Hauser ER, Swamy GK, Herbert WN, Heine RP. Maternal serum cytokines in preterm premature rupture of membranes. *Obstet Gynecol.* 2007; 109:121–7. [PubMed: 17197597]
- [15]. Hagberg H, Mallard C, Jacobsson B. Role of cytokines in preterm labour and brain injury. *Br J Obstet Gynaecol.* 2005; 112:16–8.
- [16]. Romero R, Espinoza J, Goncalves LF, Kusanovic JP, Friel LA, Nien JK. Inflammation in preterm and term labour and delivery. *Sem Fetal Neonatal Med.* 2006; 11:317–26.
- [17]. Kurki T, Hiilesmaa V, Raitasalo R, Mattila H, Ylikorkala O. Depression and anxiety in early pregnancy and risk for preeclampsia. *Obstet Gynecol.* 2000; 95:487–90. [PubMed: 10725477]
- [18]. Landsbergis PA, Hatch MC. Psychosocial work stress and pregnancy-induced hypertension. *Epidemiology.* 1996; 7:346–51. [PubMed: 8793358]
- [19]. Hetzel BS, Bruer B, Poidevin L. A survey of the relation between certain common antenatal complications in primiparae and stressful life situations during pregnancy. *J Psychosom Res.* 1961; 5:175–82. [PubMed: 13713965]
- [20]. Rofe Y, Goldberg J. Prolonged exposure to a war environment and its effects of the blood pressure of pregnant women. *Br J Med Psychol.* 1983; 56:305–11. [PubMed: 6661400]
- [21]. Kanayama N, Tsujimura R, She L, Maehara K, Terao T. Cold-induced stress stimulates the sympathetic nervous system, causing hypertension and proteinuria in rats. *J Hypertens.* 1997; 15:383–9. [PubMed: 9211173]
- [22]. Khatun S, Kanayama N, Belayet HM, Masui M, Sugimura M, Kobayashi T, et al. Induction of preeclampsia like phenomena by stimulation of sympathetic nerve with cold and fasting stress. *Eur J Obstet Gynecol Reprod Biol.* 1999; 86:89–97. [PubMed: 10471148]
- [23]. Arck PC. Stress and pregnancy loss: Role of immune mediators, hormones, and neurotransmitters. *Am J Reprod Immunol.* 2001; 46:117–23. [PubMed: 11506075]
- [24]. Retnakaran R, Hanley AJ, Raif N, Connelly PW, Sermer M, Zinman B. C-reactive protein and gestational diabetes: the central role of maternal obesity. *J Clin Endocrinol Metab.* 2003; 88:3507–12. [PubMed: 12915627]
- [25]. Du M, Yan X, Tong JF, Zhao J, Zhu MJ. Maternal obesity, inflammation, and fetal skeletal muscle development. *Biol Reprod.* 2010; 82:4–12. [PubMed: 19516021]
- [26]. Entringer S, Wadhwa PD. Developmental Programming of Obesity and Metabolic Dysfunction: Role of Prenatal Stress and Stress Biology. *Nestle Nutr Works Se.* 2013; 74:107–20.
- [27]. Bilbo SD, Schwarz JM. Early-life programming of later-life brain and behavior: a critical role for the immune system. *Front Behav Neurosci.* 2009; 3:14. [PubMed: 19738918]
- [28]. Denison FC, Roberts KA, Barr SM, Norman JE. Obesity, pregnancy, inflammation, and vascular function. *Reproduction.* 2010; 140:373–85. [PubMed: 20215337]
- [29]. Schmatz M, Madan J, Marino T, Davis J. Maternal obesity: the interplay between inflammation, mother and fetus. *J Perinatol.* 2010; 30:441–6. [PubMed: 19907427]
- [30]. Buss C, Entringer S, Davis EP, Hobel CJ, Swanson JM, Wadhwa PD, et al. Impaired Executive Function Mediates the Association between Maternal Pre-Pregnancy Body Mass Index and Child ADHD Symptoms. *Plos One.* 2012; 7
- [31]. Coussons-Read ME, Lobel M, Carey JC, Kreither MO, D'Anna K, Argys L, et al. The occurrence of preterm delivery is linked to pregnancy-specific distress and elevated inflammatory markers across gestation. *Brain, Behav, Immun.* 2012; 26:650–9. [PubMed: 22426431]
- [32]. Freeman DJ, McManus F, Brown EA, Cherry L, Norrie J, Ramsay JE, et al. Short- and long-term changes in plasma inflammatory markers associated with preeclampsia. *Hypertension.* 2004; 43:708–14. [PubMed: 15452036]
- [33]. Brown AS, Hooton J, Schaefer CA, Zhang H, Petkova E, Babulas V, et al. Elevated maternal interleukin-8 levels and risk of schizophrenia in adult offspring. *Am J Psychiatry.* 2004; 161:889–95. [PubMed: 15121655]

- [34]. Buka SL, Tsuang MT, Torrey EF, Klebanoff MA, Wagner RL, Yolken RH. Maternal cytokine levels during pregnancy and adult psychosis. *Brain, Behav, Immun.* 2001; 15:411–20. [PubMed: 11782107]
- [35]. Eder K, Baffy N, Falus A, Fulop AK. The major inflammatory mediator interleukin-6 and obesity. *Inflammation Res.* 2009; 58:727–36.
- [36]. Kim CS, Park HS, Kawada T, Kim JH, Lim D, Hubbard NE, et al. Circulating levels of MCP-1 and IL-8 are elevated in human obese subjects and associated with obesity-related parameters. *Int J Obesity.* 2006; 30:1347–55.
- [37]. Tzanavari T, Giannogonas P, Karalis KP. TNF-alpha and obesity. Current directions in autoimmunity. 2010; 11:145–56. [PubMed: 20173393]
- [38]. Moorman NJ, Cristea IM, Terhune SS, Rout MP, Chait BT, Shenk T. Human cytomegalovirus protein UL38 inhibits host cell stress responses by antagonizing the tuberous sclerosis protein complex. *Cell Host Microbe.* 2008; 3:253–62. [PubMed: 18407068]
- [39]. Coussons-Read ME, Lobel M, Carey JC, Kreither MO, D'Anna K, Argys L, et al. The occurrence of preterm delivery is linked to pregnancy-specific distress and elevated inflammatory markers across gestation. *Brain Behavior and Immunity.* 2012; 26:650–9.
- [40]. Hatzidaki E, Gourgiotis D, Manoura A, Korakaki E, Bossios A, Galanakis E, et al. Interleukin-6 in preterm rupture of membranes as an indicator of neonatal outcome. *Acta Obstetrica et Gynaecologica Scandinavica.* 2005; 84:632–8.
- [41]. Holst RM, Mattsby-Baltzer I, Wennerholm UB, Hagberg H, Jacobsson B. Interleukin-6 and interleukin-8 in cervical fluid in a population of Swedish women in preterm labor: relationship to microbial invasion of the amniotic fluid, intra-amniotic inflammation, and preterm delivery. *Acta Obstet Gynecol Scand.* 2005; 84:551–7. [PubMed: 15901266]
- [42]. Lockwood CJ, Ghidini A, Wein R, Lapinski R, Casal D, Berkowitz RL. Increased Interleukin-6 Concentrations in Cervical Secretions Are Associated with Preterm Delivery. *Am J Obstet Gynecol.* 1994; 171:1097–102. [PubMed: 7943078]
- [43]. Samuelsson A-M, Ohrn I, Dahlgren J, Eriksson E, Angelin B, Folkow B, et al. Prenatal exposure to interleukin-6 results in hypertension and increased hypothalamic-pituitary-adrenal axis activity in adult rats. *Endocrinology.* 2004; 145:4897–911. [PubMed: 15284195]
- [44]. Yoon BH, Romero R, Yang SH, Jun JK, Kim IO, Choi JH, et al. Interleukin-6 concentrations in umbilical cord plasma are elevated in neonates with white matter lesions associated with periventricular leukomalacia. *Am J Obstet Gynecol.* 1996; 174:1433–40. [PubMed: 9065108]
- [45]. Ramsay JE, Ferrell WR, Crawford L, Wallace AM, Greer IA, Sattar N. Maternal obesity is associated with dysregulation of metabolic, vascular, and inflammatory pathways. *J Clin Endocrinol Metab.* 2002; 87:4231–7. [PubMed: 12213876]
- [46]. Coussons-Read ME, Okun ML, Nettles CD. Psychosocial stress increases inflammatory markers and alters cytokine production across pregnancy. *Brain, Behav, Immun.* 2007; 21:343–50. [PubMed: 17029703]
- [47]. Winkler G, Cseh K, Baranyi E, Melczer Z, Speer G, Hajos P, et al. Tumor necrosis factor system in insulin resistance in gestational diabetes. *Diabetes Res Clin Pract.* 2002; 56:93–9. [PubMed: 11891016]
- [48]. Ferrante AW. Obesity-induced inflammation: a metabolic dialogue in the language of inflammation. *J Intern Med.* 2007; 262:408–14. [PubMed: 17875176]
- [49]. Trayhurn P, Wood IS. Adipokines: inflammation and the pleiotropic role of white adipose tissue. *Br J Nutr.* 2004; 92:347–55. [PubMed: 15469638]
- [50]. Hermsdorff HHM, Zulet MA, Puchau B, Martinez JA. Central Adiposity Rather Than Total Adiposity Measurements Are Specifically Involved in the Inflammatory Status from Healthy Young Adults. *Inflammation.* 2011; 34:161–70. [PubMed: 20467889]
- [51]. Friis CM, Paasche Roland MC, Godang K, Ueland T, Tanbo T, Bollerslev J, et al. Adiposity-related inflammation: effects of pregnancy. *Obesity (Silver Spring).* 2013; 21:E124–30. [PubMed: 23505192]
- [52]. Stewart FM, Freeman DJ, Ramsay JE, Greer IA, Caslake M, Ferrell WR. Longitudinal assessment of maternal endothelial function and markers of inflammation and placental function

- throughout pregnancy in lean and obese mothers. *J Clin Endocrinol Metab.* 2007; 92:969–75. [PubMed: 17192290]
- [53]. Larsson A, Palm M, Hansson LO, Basu S, Axelsson O. Reference values for alpha1-acid glycoprotein, alpha1-antitrypsin, albumin, haptoglobin, C-reactive protein, IgA, IgG and IgM during pregnancy. *Acta Obstet Gynecol Scand.* 2008; 87:1084–8. [PubMed: 18792844]
- [54]. Heinrich PC, Castell JV, Andus T. Interleukin-6 and the Acute Phase Response. *Biochem J.* 1990; 265:621–36. [PubMed: 1689567]
- [55]. Czarkowska-Paczek B, Bartłomiejczyk I, Gabrys T, Przybylski J, Nowak M, Paczek L. Lack of relationship between interleukin-6 and CRP levels in healthy male athletes. *Immunol Lett.* 2005; 99:136–40. [PubMed: 15894122]
- [56]. Scheller J, Chalaris A, Schmidt-Arras D, Rose-John S. The pro- and anti-inflammatory properties of the cytokine interleukin-6. *Biochim Biophys Acta.* 2011; 1813:878–88. [PubMed: 21296109]
- [57]. Menger MD, Vollmar B. Surgical trauma: hyperinflammation versus immunosuppression? *Langenbecks Arch Surg.* 2004; 389:475–84. [PubMed: 15173946]
- [58]. Boufidou F, Lambrinoudaki I, Argeitis J, Zervas IM, Pliatsika P, Leonardou AA, et al. CSF and plasma cytokines at delivery and postpartum mood disturbances. *J Affect Disord.* 2009; 115:287–92. [PubMed: 18708264]
- [59]. Maes M, Lin AH, Ombet W, Stevens K, Kenis G, De Jongh R, et al. Immune activation in the early puerperium is related to postpartum anxiety and depressive symptoms. *Psychoneuroendocrinology.* 2000; 25:121–37. [PubMed: 10674277]
- [60]. De Man YA, Dolhain RJ, Van De Geijn FE, Willemsen SP, Hazes JM. Disease activity of rheumatoid arthritis during pregnancy: results from a nationwide prospective study. *Arthritis Care Res.* 2008; 59:1241–8.
- [61]. Finkelsztejn A, Brooks J, Paschoal F, Fragoso Y. What can we really tell women with multiple sclerosis regarding pregnancy? A systematic review and meta-analysis of the literature. *BJOG: An International Journal of Obstetrics & Gynaecology.* 2011; 118:790–7. [PubMed: 21401856]
- [62]. Ozaki K, Leonard WJ. Cytokine and cytokine receptor pleiotropy and redundancy. *J Biol Chem.* 2002; 277:29355–8. [PubMed: 12072446]
- [63]. Gregor MF, Hotamisligil GS. Inflammatory Mechanisms in Obesity. *Annual Review of Immunology.* 2011; 29:415–45. 29.
- [64]. Carty CL, Heagerty P, Nakayama K, McClung EC, Lewis J, Lum D, et al. Inflammatory response after influenza vaccination in men with and without carotid artery disease. *Arterio Thromb Vasc Biol.* 2006; 26:2738–44.
- [65]. Glaser R, Robles TF, Sheridan J, Malarkey WB, Kiecolt-Glaser JK. Mild depressive symptoms are associated with amplified and prolonged inflammatory responses after influenza virus vaccination in older adults. *Arch Gen Psychiatry.* 2003; 60:1009–14. [PubMed: 14557146]
- [66]. Christian LM, Franco A, Iams JD, Sheridan J, Glaser R. Depressive symptoms predict exaggerated inflammatory response to *in vivo* immune challenge during human pregnancy. *Brain, Behav, Immun.* 2010; 24:49–53. [PubMed: 19464358]
- [67]. Elenkov IJ, Wilder RL, Bakalov VK, Link AA, Dimitrov MA, Fisher S, et al. IL-12, TNF-alpha, and hormonal changes during late pregnancy and early postpartum: Implications for autoimmune disease activity during these times. *J Clin Endocrinol Metab.* 2001; 86:4933–8. [PubMed: 11600565]
- [68]. Marzi M, Vigano A, Trabattini D, Villa ML, Salvaggio A, Clerici E, et al. Characterization of type 1 and type 2 cytokine production profile in physiologic and pathologic human pregnancy. *Clin Exp Immunol.* 1996; 106:127–33. [PubMed: 8870710]
- [69]. Fofie AE, Fewell JE, Moore SL. Pregnancy influences the plasma cytokine response to intraperitoneal administration of bacterial endotoxin in rats. *Exp Physiol.* 2004; 90:95–101. [PubMed: 15466457]
- [70]. Aguilar-Valles A, Poole S, Mistry Y, Williams S, Luheshi GN. Attenuated fever in rats during late pregnancy is linked to suppressed interleukin-6 production after localized inflammation with turpentine. *Journal of Physiology-London.* 2007; 583:391–403.

- [71]. Ashdown H, Poole S, Boksa P, Luheshi GN. Interleukin-1 receptor antagonist as a modulator of gender differences in the febrile response to lipopolysaccharide in rats. *Am J Physiol-Reg I*. 2007; 292:R1667–R74.
- [72]. Christian LM. Psychoneuroimmunology in pregnancy: Immune pathways linking stress with maternal health, adverse birth outcomes, and fetal development. *Neurosci Biobehav Rev*. 2012; 36
- [73]. Entringer S, Buss C, Wadhwa PD. Prenatal stress and developmental programming of human health and disease risk: concepts and integration of empirical findings. *Curr Opin Endocrinol Diabetes Obes*. 2010; 17:507–16. [PubMed: 20962631]

Highlights

- Excessive inflammation has been associated with a variety of adverse perinatal health
- Serum cytokines show different trajectories of change across pregnancy and postpartum
- Deviations from typical trajectories may predict adverse perinatal outcomes
- Obese women have elevated serum proinflammatory markers across gestation and postpartum
- Effects of obesity-induced inflammation on maternal and fetal health are not fully explicated

Longitudinal Changes in Serum Proinflammatory Markers by Pre-pregnancy Body Mass Index

Figures 1-5. Key Findings in Relation to Obesity

Compared to normal weight women, obese women had significantly higher serum IL-6 at each assessment timepoint. There was a similar, non-significant trend for CRP during pregnancy. Obese women showed significantly higher CRP than normal weight women at postpartum.

Error bars ± 1 SE.

Table 1
Subject Characteristics by Body Mass Index (BMI)

Characteristic	Normal Weight (n=17)	Overweight (n=16)	Obese (n=24)	p-value
Age				0.99
mean (SD)	24.3 (3.9)	24.1 (3.5)	24.3 (3.7)	
Race				0.16
African-American	11 (65%)	8 (50%)	19 (79%)	
White	6 (35%)	8 (50%)	5 (21%)	
Parity				0.72
0	1 (6%)	1 (6%)	3 (13%)	
1	9 (53%)	7 (44%)	8 (33%)	
2	7 (41%)	8 (50%)	13 (54%)	
Marital Status				0.47
Married	2 (12%)	3 (19%)	2 (8%)	
In a relationship (unmarried)	13 (76%)	10 (63%)	14 (58%)	
Single	2 (12%)	3 (19%)	8 (33%)	
Income				0.26
< \$15,000	12 (71%)	8 (50%)	17 (81%)	
\$15,000 - \$30,000	4 (24%)	4 (25%)	6 (25%)	
\$30,000	1 (6%)	4 (25%)	1 (4%)	
Education				0.88
Some high school or less	5 (29%)	5 (31%)	7 (29%)	
High School Graduate	7 (41%)	4 (25%)	6 (25%)	
Some college or college degree	5 (29%)	7 (44%)	11 (46%)	
Smoking				0.07
Current	5 (29%)	4 (25%)	2 (8%)	
Past	6 (35%)	1 (6%)	10 (42%)	
Never	6 (35%)	11 (69%)	12 (50%)	
Vigorous activity				0.87
1 hour/ week	8 (47%)	8 (50%)	10 (42%)	

Pearson's correlation coefficient for log-transformed cytokines at each study visit
Tables 2-5

1 st trimester					
	IL-6	IL-8	TNF-α	IL-1β	CRP
IL-6	1	0.49***	0.59***	0.48***	0.46***
IL-8		1	0.52***	0.39**	0.25
TNF-α			1	0.49***	0.13
IL-1β				1	0.10
CRP					1

2 nd trimester					
	IL-6	IL-8	TNF-α	IL-1β	CRP
IL-6	1	0.50***	0.62***	0.40**	0.46***
IL-8		1	0.39**	0.16	0.25
TNF-α			1	0.33*	0.15
IL-1β				1	0.21
CRP					1

3 rd trimester					
	IL-6	IL-8	TNF-α	IL-1β	CRP
IL-6	1	0.45***	0.57***	0.34**	0.47***
IL-8		1	0.37**	0.13	0.13
TNF-α			1	0.40**	0.24
IL-1β				1	0.29*
CRP					1

Postpartum					
	IL-6	IL-8	TNF-α	IL-1β	CRP
IL-6	1	0.30*	0.53***	0.25	0.36**

Postpartum					
	IL-6	IL-8	TNF- α	IL-1 β	CRP
IL-8		1	0.49***	0.26*	-0.08
TNF- α			1	0.32*	0.20
IL-1 β				1	0.15
CRP					1

*** p<0.001,
** p<0.01,
* p<0.05