Published in final edited form as:

Perspect Psychol Sci. 2013 January 1; 8(1): 72–78. doi:10.1177/1745691612469031.

Measuring Memory Reactivation with Functional MRI: Implications for Psychological Theory

Benjamin J. Levy^a and Anthony D. Wagner^{a,b}

^aDepartment of Psychology, Stanford University

^bNeurosciences Program, Stanford University

Abstract

Environmental cues often remind us of earlier experiences, with cues triggering the reactivation of memories of events past. Recent evidence suggests that memory reactivation can be observed using functional MRI, and that distributed pattern analyses can even provide evidence of reactivation on individual trials. The ability to measure memory reactivation offers unique and powerful leverage on theoretical issues of long-standing interest in cognitive psychology, providing a means to address questions that have proven difficult to answer with behavioral data alone. Here we consider three instances. First, reactivation measures can indicate whether memory-based inferences (i.e., generalization) arise through the encoding of integrated cross-event representations or through the flexible expression of separable event memories. Second, online measures of memory reactivation may inform theories of forgetting by providing information about when competing memories are reactivated during competitive retrieval situations. Finally, neural reactivation may provide a window onto the role of replay in memory consolidation. The ability to track memory reactivation, including at the individual trial level, provides unique leverage that is not afforded by behavioral measures, and thus promises to shed light on such varied topics as generalization, integration, forgetting, and consolidation.

Cues in the environment frequently remind us of past experiences, which allows the past to inform current behavior. Memory reactivation can occur with or without the intention to remember, and with or without awareness of the connection between the cues and the memories they activate. Reactivated memories frequently inform our current goals, but in some instances they may be irrelevant to, and even distract us from, our goals. As such, memory reactivation plays a critical role in guiding behavior, as it influences the knowledge upon which decisions and actions are based.

Although memory reactivation is central to many psychological theories, it is often difficult to directly observe reactivation in behavior. Researchers can explicitly ask participants whether they are currently reminded of past experiences, but individuals may not always be aware when reactivation occurs. Moreover, asking participants to reflect upon whether they are remembering may influence the frequency and type of reminders they experience. As an alternative approach, researchers may indirectly probe memory and then infer when

reactivation has occurred by observing effects on behavior (e.g., Anderson & McCulloch, 1999). In many instances, however, the link between the observed behavior and memory reactivation is ambiguous.

Neuroimaging methods, such as functional MRI (fMRI), offer a novel approach to measuring memory reactivation that complements behavioral indices (e.g., Kahn, Davachi, & Wagner, 2004; Nyberg, Habib, McIntosh, & Tulving, 2000; Wheeler, Petersen, & Buckner, 2000; for review, see Danker & Anderson, 2010; Mitchell & Johnson, 2009). In some instances, fMRI can even provide powerful evidence of reminding independent of behavior, and fMRI can yield measures of memory reactivation at the individual trial level (e.g., Chadwick et al., 2010; Johnson et al., 2009; Polyn et al., 2005; for review, see Rissman & Wagner, 2012). Here, we review recent fMRI studies that demonstrate how neural measures of memory reactivation can be used to inform psychological theories about the representations and processes underlying generalization, forgetting, and consolidation (e.g., Kuhl, Rissman, Chun, & Wagner, 2011; Kuhl, Shah, DuBrow, & Wagner, 2010; Rasch et al., 2007; Shohamy & Wagner, 2008; Zeithamova & Preston, 2010).

Integrative encoding as a mechanism for generalizing across experiences

One situation where memory influences behavior is when we encounter a novel choice that requires that we generalize knowledge across multiple prior experiences to determine the appropriate decision/action. From one prominent view, such generalization occurs at the time of choice, and is based on the retrieval of multiple discrete episodic memories, from which we logically infer the relationship between the stimuli and the decision that should be made (e.g., Eichenbaum, 2000; Kumaran & McClelland, in press). By contrast, an alternative mechanism for generalization is integrative encoding that occurs during the acquisition of the original experiences, before the novel choice probe is ever encountered (e.g., Eichenbaum, 2000; O'Reilly & Rudy, 2000; Shohamy & Wagner, 2008). By this account, we are reminded of similar prior events during encoding of subsequent related events, and when this reminding occurs an integrated mnemonic representation is formed that spans past and present episodes. This integrated memory directly encodes the relations between stimuli from temporally distinct events, with this knowledge being retrieved later when the novel choice is encountered. A key question is whether memory-based inferences that support generalization can occur because of encoding of integrated cross-event representations or whether generalization exclusively occurs through the flexible expression of separable event memories at test.

Shohamy and Wagner (2008) provided support for integrative encoding, reporting fMRI evidence of reactivation during the learning phase of an acquired equivalence task (e.g., Myers et al., 2003). Participants initially learned associations between stimuli (e.g., A-X, B-X, A-Y) through trial-and-error feedback (see Fig. 1a). Because some of the associations overlapped (i.e., both A and B predict X), it was possible that during the encoding of a specific association participants reactivated other related associations (i.e., while encoding A-Y they may recall the A-X and B-X associations). To the extent that such reactivation occurs during learning, participants can form a representation that integrates information from past experience with that present in the immediate environment. This integrated

representation could then support subsequent decisions when the participant is faced with a novel choice that can be solved through generalization (i.e., generalizing the A-Y association to B-Y, because of A and B's equivalence due to their common association with X).

Consistent with the integrative encoding account, Shohamy and Wagner (2008) observed that activity in the hippocampus increased across the phase where participants learned the original associations (see Fig. 1b). Because prior data indicate that repeated encoding of the same association results in decreasing hippocampal activation (e.g., Kohler, Danckert, Gati, & Menon, 2005; Zeineh, Engel, Thompson, & Bookheimer, 2003), whereas reactivation of past associations results in increased hippocampal activation (e.g., Eldridge et al., 2000), Shohamy and Wagner interpreted this hippocampal increase as a marker of memory reactivation during learning. Critically, the magnitude of this increase across learning predicted participants' later generalization performance, suggesting that reactivation during learning supported subsequent memory-based inferences. These findings suggest that during subsequent events associative mechanisms in the hippocampus reactivate prior experiences allowing for knowledge integration across distinct episodes; the resulting integrative representations can support later generalization.

Of course, evidence of reactivation during encoding does not preclude a role for retrieval-based inference in generalization (e.g., Greene, Gross, Elsinger, & Rao, 2006; Heckers et al., 2004; Preston, Shrager, Dudukovic, & Gabrieli, 2004). Using a single-shot learning paradigm, Zeithamova and Preston (2010) provided (a) further evidence for integrative encoding – hippocampal activity during encoding of an event that overlapped with a prior experience (i.e., during B-C encoding, after having studied A-B) predicted later transfer during a novel choice (i.e., choosing C over D when presented with A), and (b) evidence consistent with a contribution of retrieval-based inference – participants who generalized well demonstrated greater test phase activation during generalization probes (A-C) than during directly learned probes (A-B). Thus, although computational modeling work suggests that reactivating event content from distinct memories at test is sufficient to explain generalization (Kumaran & McClelland, in press), extant neuroimaging data clearly suggest that the ability to generalize from past experiences depends upon both integrative encoding, arising from reactivation of past experiences during subsequent learning, and retrieval-based inference. Moreover, to the extent that learning accompanies retrieval-based inferences, then the mechanisms that give rise to such inferences, when combined with learning, might also provide a computational account of integrative encoding.

The relationship between activation of competing memories and forgetting

Sometimes memory reactivation can be counter-productive, particularly when an association that comes to mind is not relevant to current goals. In these situations, individuals suffer interference, which is thought to be a major cause of forgetting (Anderson, 2003; Crowder, 1976). According to many psychological theories, forgetting due to interference is a retrieval phenomenon where stronger memories tend to come to mind readily and block retrieval of weaker memories (e.g., Anderson, 1983; McGeoch, 1942; Raaijmakers & Shiffrin, 1981). An alternative theory, however, posits that when interference occurs it elicits a secondary

process (or set of processes) that counteract(s) interference by inhibiting competing memories (e.g., Anderson, 2003; Melton & Irwin, 1940; Osgood, 1949). Thus, from this perspective, it is the process of overcoming interference during a prior retrieval attempt that renders competing memories harder to recall later.

Considerable behavioral and neuroimaging evidence has been gathered in favor of the inhibitory account of forgetting (for reviews, see Anderson, 2003; Bäuml, Pastötter, & Hanslmayr, 2010; Levy & Anderson, 2002; Storm, 2011), but inhibitory forgetting effects are often relatively small and are not always replicated (e.g., Bulevich, Roediger, Balota, & Butler, 2006; Perfect et al., 2004; Williams & Zacks, 2001). Although some of this variability may be attributable to variations in cognitive control abilities across individuals (Levy & Anderson, 2008), there are also boundary conditions on when inhibition will occur (for reviews, see Anderson, 2003; Levy & Anderson, 2002). In particular, participants in interference paradigms are thought to sometimes reactivate “competing” memories along with the target memory, potentially forming an integrated representation that spans both events (Anderson & McCulloch, 1999). This integration would protect the “competing” memory, reducing the need for inhibitory control, and therefore reducing forgetting on later memory tests. Neural measures of reactivation may be able to provide evidence about when this type of integration occurs.

Although fMRI data clearly indicate that competitive retrieval is associated with greater engagement of cognitive control processes (e.g., Danker, Gunn, & Anderson, 2008; Kuhl, Dudukovic, Kahn, & Wagner, 2007; Wimber et al., 2008), initial studies did not provide clear evidence about when competing memories were reactivated. Kuhl et al. (2011) recently reported evidence demonstrating this type of reactivation using distributed pattern analyses of fMRI data. In the prior section on generalization, the magnitude of activation within the hippocampus was used as a marker of reactivation, whereas here reactivation was assessed through the reinstatement of distributed patterns of activity across many voxels within the ventral temporal cortex. Specifically, in a retroactive interference paradigm where participants learned overlapping associations (A-B and A-C pairs), pattern analyses revealed evidence that the competing associate was reactivated during attempts to retrieve the target associate (i.e., evidence that B was reactivated during attempts to retrieve the A-C association). Critically, greater evidence for reactivation of the earlier B memory was correlated with better memory for those items on a final post-test, consistent with the hypothesis that reactivation led to the formation of an integrated representation that protected earlier memories from forgetting. Similarly, Kuhl et al. (2010) found that hippocampally-mediated reactivation of cortical patterns during the *encoding* of associations that overlapped with previously learned associations predicted better subsequent memory for the earlier associations (i.e., greater hippocampal activity during A-C encoding predicted better subsequent memory for the A-B association). Both of these findings suggest that reactivation can lead to integrated cross-event representations that protect earlier memories from forgetting.

Prior behavioral evidence suggests that encoding, unlike retrieval, does not induce competition and therefore does not trigger the need for interference-resolution processes (e.g., Bäuml, 1996; Bäuml & Hartinger, 2002; Ciranni & Shimamura, 1999). Therefore, it is

not surprising that reactivation during encoding trials did not induce forgetting. It is less clear, however, whether reactivation during retrieval should always be beneficial for later remembering, as observed by Kuhl et al. (2011). It seems likely that, at least in some retrieval situations, reactivating competing memories should trigger the need for inhibitory control and thus greater forgetting. The relationship between reactivation and inhibition, however, is not necessarily straightforward (for discussions, see Anderson & Levy, 2011; Norman, Newman & Detre, 2007; Bäuml, Pastötter, & Hanslmayr, 2010) and this relationship may be obscured by integrative encoding mechanisms. Nevertheless, neural measures that can provide continuous indices of reactivation on a trial-by-trial basis hold great promise for observing competitive dynamics during retrieval and promise to provide critical leverage on testing theories about the relationship between reactivation during retrieval attempts and subsequent forgetting (see also Öztekin & Badre, 2011).

The role of memory reactivation in consolidation

Memory reactivation is thought to be critical for ‘systems consolidation’, wherein a memory that initially depends on hippocampal representation is gradually encoded in neocortex (e.g., Squire & Alvarez, 1995). Research in rodents suggests that recent memories are reactivated during sleep (e.g., Foster & Wilson, 2006; Lee & Wilson, 2002; Wilson & McNaughton, 1994) and this reactivation is thought to reflect the hippocampus playing back recent experiences to the neocortex, hypothesized to be a critical process in consolidation (e.g., McClelland, McNaughton, & O’Reilly, 1995). Recent neuroimaging work in humans has sought to experimentally manipulate memory reactivation during sleep, examining effects at the neural and behavioral levels. For example, Rasch, Büchel, Gais, and Born (2007) had participants study object-location associations while being exposed to a specific odor and then re-exposed the participants to this odor during slow-wave sleep. This re-exposure led to increased hippocampal activity and improved subsequent memory for the object-location associations (relative to exposure to a control odor), consistent with the odor triggering reactivation of associated events, putatively facilitating their consolidation and rendering them more readily recallable. Using scalp electroencephalography (EEG), Rudoy, Voss, Westerberg, and Paller (2009) expanded on this finding by cueing with unique sounds that were each associated with a particular object-location association. Exposure to a subset of the sounds during sleep led to better subsequent memory for the cued associations, and EEG responses to the cues during sleep significantly predicted subsequent memory performance.

These two studies are consistent with the possibility that memory reactivation was induced by environmental cues during sleep, contributing to memory consolidation and enhanced post-sleep memory. Neither of these studies, however, provides compelling evidence that directly links clear evidence for reactivation with improved subsequent memory. For example, these measures might simply reflect further cue encoding, which then allows the cue to act as a more potent reminder during the post-sleep memory test. Future fMRI studies can borrow from the approaches described above to determine if and when off-line memory reactivation occurs, be it during sleep or during awake, resting states (Tambini, Ketz & Davachi, 2010). Although the evidence in this section may be more speculative and less certain, this is certainly another area where fMRI measures of reactivation hold considerable promise for informing psychological theory.

Conclusion

Here we have described how functional MRI methods, based on either activity from regions of interest or from distributed patterns of activity, can be used to track memory reactivation as it occurs, even on a trial-by-trial basis. In this manner, fMRI provides unique leverage on psychological theory that is not afforded by behavioral measures, which do not allow investigators to observe when a past experience has been covertly reactivated. Moving forward, we anticipate that neuroimaging will be informative for numerous open questions, as many psychological theories invoke covert retrieval as a key mechanism. For example, environmental cues are thought to activate negative stereotype-consistent information in groups vulnerable to social identity threat (e.g., Murphy, Steele, & Gross, 2007). Similarly, cues are thought to trigger retrieval of threat-related information in depression, anxiety, and post-traumatic stress disorder (e.g., Mathews & MacLeod, 2005; McNally, Kaspi, Riemann, & Zeitlin, 1990). Researchers in these and other areas can leverage fMRI measures of reactivation to determine when memories are retrieved, which specific cues trigger the strongest reactivation, and what are the psychological and behavioral consequences of such reactivation. In our view, fMRI measures of covert and overt reactivation hold great promise for testing psychological theories, illustrating the complementary power of functional neuroimaging as a fundamental tool for psychological science.

Acknowledgments

Supported by the National Institute of Mental Health (R01-MH080309; R01-MH076932; F32-MH079648).

References

- Anderson, JR. The architecture of cognition. Harvard University Press; Cambridge, MA: 1983.
- Anderson MC. Rethinking interference theory: Executive control and the mechanisms of forgetting. *Journal of Memory and Language*. 2003; 49:415–445.
- Anderson, MC.; Levy, BJ. On the relationship between interference and inhibition in cognition.. In: Benjamin, A., editor. *Successful Remembering and Successful Forgetting: A Festschrift in Honor of Robert A. Bjork*. Psychology Press; New York, NY: 2011. p. 107-132.
- Anderson MC, McCulloch KC. Integration as a general boundary condition on retrieval-induced forgetting. *Journal of Experimental Psychology: Learning, Memory, and Cognition*. 1999; 25:608–629.
- Bäuml K-H. Revisiting an old issue: Retroactive interference as a function of the degree of original and interpolated learning. *Psychonomic Bulletin & Review*. 1996; 3:380–384. [PubMed: 24213942]
- Bäuml K-H, Hartinger A. On the role of item similarity in forgetting. *Memory*. 2002; 10:215–224. [PubMed: 11958725]
- Bäuml K-H, Pastötter B, Hanslmayr S. Binding and inhibition in episodic memory: Cognitive, emotional, and neural processes. *Neuroscience & Biobehavioral Reviews*. 2010; 34:1047–1054. [PubMed: 19397925]
- Bulevich JB, Roediger HL, Balota DA, Butler AC. Failures to find suppression of episodic memories in the think/no-think paradigm. *Memory & Cognition*. 2006; 34:1569–1577. [PubMed: 17489284]
- Chadwick MJ, Hassabis D, Weiskopf N, Maguire EA. Decoding individual episodic memory traces in the human hippocampus. *Current Biology*. 2010; 20:544–471. [PubMed: 20226665]
- Ciranni MA, Shimamura AP. Retrieval-induced forgetting in episodic memory. *Journal of Experimental Psychology: Learning, Memory, and Cognition*. 1999; 25:1403–1414.
- Crowder, RG. *Principles of learning and memory*. Erlbaum; Hillsdale, NJ: 1976.

- Danker JF, Anderson JR. The ghosts of brain states past: Remembering reactivates the brain regions engaged during encoding. *Psychological Bulletin*. 2010; 136:87–102. [PubMed: 20063927]
- Danker JF, Gunn P, Anderson JR. A rational account of memory predicts left prefrontal activation during controlled retrieval. *Cerebral Cortex*. 2008; 18:2674–2685. [PubMed: 18321871]
- Eichenbaum H. A cortical-hippocampal system for declarative memory. *Nature Reviews Neuroscience*. 2000; 1:41–50.
- Eldridge LL, Knowlton BJ, Furmanski CS, Bookheimer SY, Engel SA. Remembering episodes: A selective role for the hippocampus during retrieval. *Nature Neuroscience*. 2000; 3:1149–1152.
- Foster DJ, Wilson MA. Reverse replay of behavioural sequences in hippocampal place cells during the awake state. *Nature*. 2006; 440:680–683. [PubMed: 16474382]
- Greene AJ, Gross WL, Elsinger CL, Rao SM. An fMRI analysis of the human hippocampus: inference, context, and task awareness. *Journal of Cognitive Neuroscience*. 2006; 18:1156–1173. [PubMed: 16839289]
- Heckers S, Zalesak M, Weiss AP, Ditman T, Titone D. Hippocampal activation during transitive inference in humans. *Hippocampus*. 2004; 14:153–162. [PubMed: 15098721]
- Johnson JD, McDuff SGR, Rugg MD, Norman KA. Recollection, familiarity, and cortical reinstatement: A multivoxel pattern analysis. *Neuron*. 2009; 63:697–708. [PubMed: 19755111]
- Kahn I, Davachi L, Wagner AD. Functional-neuroanatomic correlates of recollection: Implications for models of recognition memory. *Journal of Neuroscience*. 2004; 24:4172–4180. [PubMed: 15115812]
- Kohler S, Danckert S, Gati JS, Menon RS. Novelty responses to relational and non-relational information in the hippocampus and parahippocampal region: a comparison based on event-related fMRI. *Hippocampus*. 2005; 15:763–774. [PubMed: 15999342]
- Kuhl BA, Dudukovic NM, Kahn I, Wagner AD. Decreased demands on cognitive control reveal the neural processing benefits of forgetting. *Nature Neuroscience*. 2007; 10:908–914.
- Kuhl BA, Rissman J, Chun MM, Wagner AD. Fidelity of neural reactivation reveals competition between memories. *Proceedings of the National Academy of Sciences, USA*. 2011; 108:5903–5909.
- Kuhl BA, Shah AT, DuBrow S, Wagner AD. Resistance to forgetting associated with hippocampus-mediated reactivation during new learning. *Nature Neuroscience*. 2010; 13:501–506.
- Kumaran, D.; McClelland, JL. Beyond episodic memory: A complementary learning systems account of the hippocampal contribution to generalization. *in press*
- Lee AK, Wilson MA. Memory of sequential experience in the hippocampus during slow wave sleep. *Neuron*. 2002; 36:1183–1194. [PubMed: 12495631]
- Levy BJ, Anderson MC. Inhibitory processes and the control of memory retrieval. *Trends in Cognitive Sciences*. 2002; 6:299–305. [PubMed: 12110363]
- Levy BJ, Anderson MC. Individual differences in the suppression of unwanted memories: The executive deficit hypothesis. *Acta Psychologica*. 2008; 127:623–635. [PubMed: 18242571]
- Mathews A, MacLeod C. Cognitive vulnerability to emotional disorders. *Annual Review of Clinical Psychology*. 2005; 1:167–195.
- McClelland JL, McNaughton BL, O'Reilly RC. Why there are complementary learning systems in the hippocampus and neocortex: Insights from the successes and failures of connectionist models of learning and memory. *Psychological Review*. 1995; 102:419–457. [PubMed: 7624455]
- McGeoch, JA. *The psychology of human learning: An introduction*. Longmans, Green and Co.; New York: 1942.
- McNally RJ, Kaspi SP, Riemann BC, Zeitlin SB. Selective processing of threat cues in posttraumatic stress disorder. *Journal of Abnormal Psychology*. 1990; 99:398–402. [PubMed: 2266215]
- Melton AW, Irwin JM. The influence of degree of interpolated learning on retroactive inhibition and the overt transfer of specific responses. *American Journal of Psychology*. 1940; 3:173–203.
- Mitchell KJ, Johnson MK. Source monitoring 15 years later: What have we learned from fMRI about the neural mechanisms of source memory? *Psychological Bulletin*. 2009; 135:638–677. [PubMed: 19586165]
- Murphy MC, Steele CM, Gross JJ. Signaling threat: How situational cues affect women in math, science, and engineering settings. *Psychological Science*. 2007; 18:879–885. [PubMed: 17894605]

- Myers CE, Shohamy D, Gluck MA, Grossman S, Kluger A, Ferris S, Golomb J, Schnirman G, Schwartz R. Dissociating hippocampal versus basal ganglia contributions to learning and transfer. *Journal of Cognitive Neuroscience*. 2003; 15:185–193. [PubMed: 12676056]
- Norman KA, Newman EL, Detre G. A neural network model of retrieval-induced forgetting. *Psychological Review*. 2010; 114:887–953. [PubMed: 17907868]
- Nyberg L, Habib R, McIntosh AR, Tulving E. Reactivation of encoding-related brain activity during memory retrieval. *Proceedings of the National Academy of Sciences, USA*. 2000; 97:11120–11124.
- O'Reilly RC, Rudy JW. Conjunctive representations in learning and memory: principles of cortical and hippocampal function. *Psychological Review*. 2000; 108:311–345. [PubMed: 11381832]
- Osgood CE. The similarity paradox in human learning: A resolution. *Psychological Review*. 1949; 56:132–143. [PubMed: 18139121]
- Öztekin I, Badre D. Distributed patterns of brain activity that lead to forgetting. *Frontiers in Human Neuroscience*. 2011; 5:1–8. [PubMed: 21283556]
- Perfect TJ, Stark L-J, Tree JJ, Moulin CJA, Ahmed L, Hutter R. Transfer appropriate forgetting: The cue-dependent nature of retrieval-induced forgetting. *Journal of Memory and Language*. 2004; 51:399–417.
- Polyn SM, Natu VS, Cohen JD, Norman KA. Category-specific cortical activity precedes retrieval during memory search. *Science*. 2005; 310:1963–66. [PubMed: 16373577]
- Preston AR, Shrager Y, Dudukovic NM, Gabrieli JD. Hippocampal contribution to the novel use of relational information in declarative memory. *Hippocampus*. 2004; 14:148–152. [PubMed: 15098720]
- Raaijmakers JGW, Shiffrin RM. Search of associative memory. *Psychological Review*. 1981; 88:93–134.
- Rasch B, Büchel C, Gais S, Born J. Odor cues during slow-wave sleep prompt declarative memory consolidation. *Science*. 2007; 315:1426–1429. [PubMed: 17347444]
- Rissman J, Wagner AD. Distributed representations in memory: Insights from functional brain imaging. *Annual Review of Psychology*. 2012; 63:101–28.
- Rudoy JD, Voss JL, Westerberg CE, Paller KA. Strengthening individual memories by reactivating them during sleep. *Science*. 2009; 326:1079. [PubMed: 19965421]
- Shohamy D, Wagner AD. Integrating memories in the human brain: Hippocampal-midbrain encoding of overlapping events. *Neuron*. 2008; 60:378–389. [PubMed: 18957228]
- Squire LR, Alvarez P. Retrograde amnesia and memory consolidation: a neurobiological perspective. *Current Opinion in Neurobiology*. 1995; 5:169–177. [PubMed: 7620304]
- Storm BC. Retrieval-induced forgetting and the resolution of competition.. In: Benjamin, AS., editor. *Successful Remembering and Successful Forgetting: A Festschrift in honor of Robert A. Bjork*. Psychology Press; New York, NY: 2011. p. 89-105.
- Tambini A, Ketz N, Davachi L. Enhanced brain correlations during rest are related to memory for recent experiences. *Neuron*. 2010; 65:280–290. [PubMed: 20152133]
- Wheeler ME, Petersen SE, Buckner RL. Memory's echo: Vivid remembering reactivates sensory-specific cortex. *Proceedings of the National Academy of Sciences, USA*. 2000; 97:11125–11129.
- Williams CC, Zacks RT. Is retrieval-induced forgetting an inhibitory process? *American Journal of Psychology*. 2001; 114:329–354. [PubMed: 11641884]
- Wilson MA, McNaughton BL. Reactivation of hippocampal ensemble memories during sleep. *Science*. 1994; 265:676–679. [PubMed: 8036517]
- Wimber M, Bäuml K-H, Bergström Z, Markopoulos G, Heinze H-J, Richardson-Klavehn A. Neural markers of inhibition in human memory retrieval. *Journal of Neuroscience*. 2008; 28:13419–13427. [PubMed: 19074015]
- Zeineh MM, Engel SA, Thompson PM, Bookheimer SY. Dynamics of the hippocampus during encoding and retrieval of face-name pairs. *Science*. 2003:299–577-580.
- Zeithamova D, Preston AR. Flexible memories: Differential roles for medial temporal lobe and prefrontal cortex in cross-episode binding. *Journal of Neuroscience*. 2010; 30:14676–14684. [PubMed: 21048124]

Figure 1.

Evidence for integrative encoding from Shohamy and Wagner (2008). (A) Behavioral paradigm for the acquired equivalence task. Participants learned face-scene associations through trial-and-error feedback (upper left). Critically, during the learning phase some cues shared a common associate (lower left). During a subsequent test (upper right), participants were tested on the trained pairs and on generalization trials. Generalization trials could be correctly responded to by way of two different mechanisms (lower right): during test, participants may retrieve multiple distinct trained associations and draw inferences from them about what response should be made; alternatively, during learning, prior associations may be reactivated and then integrated into a single cross-event representation that is then available to be retrieved when the generalization trials are encountered at test. (B) Neuroimaging evidence for integrative encoding. Activity in the right and left hippocampus (top) increased across the learning phase. Across participants, the degree to which activation increased from early to late learning was predictive of performance (% correct) on the generalization trials (middle). A median split of the participants revealed that the “good” generalizers showed increased activity from the early to late learning phase, whereas “poor” generalizers did not (bottom).