An intravascular lymphoma with extravascular tendencies

Cody A. Thomas, MD, Joseph M. Guileyardo, MD, and John R. Krause, MD

Intravascular large B-cell lymphoma (IVL) is rare and characterized by selective growth of neoplastic cells within the lumina of small blood vessels. We present the case of a 69-year-old woman who died of a widespread IVL with extravascular involvement of the lymph nodes, liver, bladder, and adrenal gland. This report discusses the unique features of IVL with concurrent extravascular components.

Intravascular large B-cell lymphoma (IVL), also known as angiotrophic lymphoma, was first described in 1959 as an endothelial neoplasm with vascular propagation (1). Only within the last 15 years has IVL been considered distinctive enough for World Health Organization (WHO) classification (2). The 2008 WHO definition states that IVL is a type of extranodal large B-cell lymphoma that selectively grows within the lumina of vessels, particularly capillaries, with the exception of larger arteries and veins. The WHO classification also states that this lymphoma is often widely disseminated in extranodal sites, and lymph nodes are usually spared (3). It is also important to note that extravascular infiltration by tumor cells is observed in only 11% of IVL cases, and patients presenting with classic features of IVL have later developed a nodal diffuse large B-cell lymphoma with shared immunophenotypic markers (1). There are also reports of nodal B-cell lymphoma recurring as IVL (4, 5) and IVL presenting concurrently with a lymphomatous mass in the central nervous system (6). We report the autopsy findings of a widely disseminated IVL with extensive extravascular involvement by tumor of the lymph nodes, bladder, liver, and adrenal gland.

CASE REPORT

A 69-year-old woman was admitted to Baylor University Medical Center at Dallas with symptoms of fatigue and generalized weakness for the past year, as well as night sweats and fever of 2 weeks' duration. On admission she had a blood pressure of 91/42 mm Hg, a pulse of 140 beats per minute, a temperature of 100.5°F, and oxygen saturation of 95% on room air. The clinical diagnosis was probable sepsis. Laboratory studies revealed the following: white blood cell count of 8200/uL (82% segmented neutrophils, 6% normal lymphocytes, 7% monocytes, 4% bands, and 1% metamyelocytes), hemoglobin

of 11.4 g/dL, hematocrit of 34.5%, platelet count of 128,000/uL, sodium of 125 mEq/L, potassium of 4.8 mEq/L, bicarbonate of 17 mEq/L, creatinine of 0.9 mg/dL, glucose of 98 mg/dL, calcium of 7.4 mg/dL, lactate dehydrogenase of 2070 U/L (normal level: 100–190 U/L), erythrocyte sedimentation rate of 58 mm/h (normal level: 0–30 mm/h), and C-reactive protein of 19.8 mg/dL (normal level: 0–0.3 mg/dL). A chest computed tomography (CT) scan revealed a paratracheal mass, raising clinical suspicion for paraneoplastic syndrome due to lung cancer, and an abdominal CT scan revealed only diverticulosis and abdominopelvic ascites. Despite therapy, she continued with fevers and hypovolemia and developed severe metabolic acidosis (pH 7.12) and acute respiratory failure. She died on the third day after admission.

At necropsy the paratracheal mass seen on imaging proved to be a hematoma contiguous with a central catheter insertion site at the right jugular vein. There was a focal thrombus in the left brachiocephalic vein associated with central catheter placement through the left jugular vein. Grossly within the liver there were numerous tan nodules ranging from 0.2 to 1.8 cm in diameter. Also present was a 1.6 cm tan, firm mucosal nodule in the urinary bladder, nodular expansion of the left adrenal gland, and two enlarged periaortic and perinephric lymph nodes. Both lymph nodes were approximately 1.5 cm and firm.

Microscopic examination demonstrated atypical lymphoid cells within the lumina of small vessels in most organs and tissues throughout the body, most prominent in the lungs (*Figure 1a*). These intravascular lymphoid cells were enlarged with pleomorphic, vesicular nuclei, prominent nucleoli, and multiple mitotic figures. The thrombus from the left brachiocephalic vein also contained these atypical cells but the paratracheal hematoma did not.

From the Department of Pathology, Baylor University Medical Center at Dallas and the Charles A. Sammons Cancer Center, Dallas, Texas.

Corresponding author: John R. Krause, MD, Section of Hematopathology, Department of Pathology, Baylor University Medical Center at Dallas, 3500 Gaston Avenue, Dallas, TX 75246 (e-mail: john.krause@BaylorHealth.edu); Joseph M. Guileyardo, MD, Department of Pathology, Baylor University Medical Center at Dallas, 3500 Gaston Avenue, Dallas, TX 75246 (e-mail: guileyardo@sbcglobal.net).

Figure 1. (a) Intravascular neoplastic cells in the lung (hematoxylin and eosin, ×200). (b) Extravascular lymphomatous involvement of bladder mucosa. The neoplastic cells in the lymph nodes, liver, and adrenal gland had the same morphologic appearance (hematoxylin and eosin, ×400). (c) CD20 immunostain of the periaortic lymph node highlighting both intra- and extravascular lymphoma cells (CD20 immunoperoxidase stain, ×200).

While most of the tumor was intravascular, there was also significant extravascular tumor infiltration. The bladder, liver, and adrenal gland nodules were dense populations of lymphoma cells (*Figure 1b*). The enlarged lymph nodes were sclerotic and diffusely involved by intravascular and extravascular lymphoma. The bone marrow was 60% to 70% cellular with scattered atypical lymphoid cells present within vascular spaces, but no lymphomatous infiltration was identified. Immunohistochemical staining revealed that both the intra- and extravascular lymphoma cells of an involved node were CD20 and MUM-1 positive and CD3, CD30, CD15, CD10, BCL6, and Alk-1 negative (*Figure 1c*). The MUM-1 positivity and the CD10 and BCL6 negativity indicate that this lymphoma was of postgerminal center origin (activated B-cell type). A Ki-67 proliferation index revealed approximately 50% of the tumor cells to be positive.

DISCUSSION

Although 55 years have passed since IVL was first described, our understanding of this entity is still incomplete. Presenting clinical features are usually nonspecific and quite variable due to the widespread nature of the disease. There is no laboratory, radiologic, or cytogenetic study specific for this diagnosis, and despite being intravascular, peripheral blood involvement is

only demonstrated in 5% to 9% of patients (7). All of these factors commonly delay or prevent antemortem diagnosis. The neoplastic cells tend to remain confined to small- and intermediate-sized blood vessels with no particular organ preference. Defects in adhesion molecules CD29 (beta 1 integrin), which is essential for leukocyte extravasation, and CD54 (ICAM-1), which is involved in transvascular lymphocyte migration, have been suggested as an explanation for the intravascular localization (8); however, the mechanism is likely multifactorial and more studies are needed.

The current case of IVL is unusual because this patient had a prominent, diffuse intravascular component of tumor cells associated with a significant extravascular component. The morphologic and immunohistochemical characteristics of both components were identical. We believe this aggressive IVL extravasated into the lymph nodes, liver, bladder, and adrenal parenchyma as a diffuse large B-cell lymphoma. Nodal lymphoma transformation into intravascular lymphoma (4, 5) is less likely due to the overwhelming intravascular component of lymphoma cells. However, it is not possible to prove the actual temporal sequence without previous biopsies exhibiting the pattern of disease progression. Nevertheless, the distinction can be important since most patients diagnosed with IVL are treated with regimens used for advanced diffuse large B-cell lymphoma (7).

IVL continues to present diagnostic challenges due to its elusive and diverse pathologic and clinical findings, and according to Ponzoni and colleagues (9), a comprehensive analysis is needed to find analogies and differences between the intravascular and extravascular components of large B-cell lymphomas to further characterize them and potentially formulate more focused treatment depending on each patient's unique presentation.

1. Ponzoni M, Ferreri AJ. Intravascular lymphoma: a neoplasm of 'homeless' lymphocytes? *Hematol Oncol* 2006;24(3):105–112.
2. Harris NL, Jaffe ES, Diebold J, Flandrin G, Muller-Hermelink HK, Vardiman J, Lister TA, Bloomfield CD. World Health Organization classification of neoplastic diseases of the hematopoietic and lymphoid tissues: report of the Clinical Advisory Committee meeting—Airlie House, Virginia, November 1997. *J Clin Oncol* 1999;17(12):3835–3849.
3. Swerdlow SH, Campo E, Harris NL, Jaffe ES, Pileri SA, Stein H, Thiele J, Vardiman JW, eds. *WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues*. Lyon, France: International Agency for Research on Cancer, 2008:252–253.
4. Zhao XF, Sands AM, Ostrow PT, Halbiger R, Conway JT, Bagg A. Recurrence of nodal diffuse large B-cell lymphoma as intravascular large B-cell lymphoma: is an intravascular component at initial diagnosis predictive? *Arch Pathol Lab Med* 2005;129(3):391–394.
5. Katz DA, Miller IJ, Gregory SA. Intravascular B-cell lymphoma following nodal diffuse large B-cell lymphoma. *Clin Adv Hematol Oncol* 2010;8(9):637–641.
6. Imai H, Kajimoto K, Taniwaki M, Miura I, Hatta Y, Hashizume Y, Watanabe M, Shiraishi T, Nakamura S. Intravascular large B-cell lymphoma presenting with mass lesions in the central nervous system: a report of five cases. *Pathol Int* 2004;54(4):231–236.
7. Ponzoni M, Ferreri AJ, Campo E, Facchetti F, Mazzucchelli L, Yoshino T, Murase T, Pileri SA, Doglioni C, Zucca E, Cavalli F, Nakamura S. Definition, diagnosis, and management of intravascular large B-cell lymphoma: proposals and perspectives from an international consensus meeting. *J Clin Oncol* 2007;25(21):3168–3173.
8. Ponzoni M, Arrigoni G, Gould VE, Del Curto B, Maggioni M, Scapinello A, Paolino S, Cassisa A, Patriarca C. Lack of CD 29 (beta1 integrin) and CD 54 (ICAM-1) adhesion molecules in intravascular lymphomatosis. *Hum Pathol* 2000;31(2):220–226.
9. Ponzoni M, Campo E, Ferreri AJ. Intravascular lymphoma occurring in patients with other non-Hodgkin lymphomas. *Clin Adv Hematol Oncol* 2010;8(9):641–642.