Published in final edited form as:

Can J Aging. 2014 December ; 33(4): 413–425. doi:10.1017/S0714980814000282.

The Role of Partnership Status on Late-Life Physical Function*

Sean Clouston¹, Andrea Lawlor², and Ashton Verdery³

¹ Program in Public Health and Department of Preventive Medicine, Stony Brook University

² Center for the Study of Democratic Citizenship, McGill University, and Institute for Governmental Studies, University of California, Berkeley

³ Department of Sociology and Carolina Population Center, University of North Carolina, Chapel Hill

Abstract

This study examined the socioeconomic pathways linking partnership status to physical functioning, assessed using objective measures of late life physical functioning including peak flow and grip strength. Using Wave 4 of the Survey of Health, Ageing and Retirement in Europe (SHARE), we ran multilevel models to examine the relationship between partnership status and physical function in late life, adjusting for social-network characteristics, socioeconomic factors, and health behaviours. We found a robust relationship between partnership status and physical function. Incorporating social-network characteristics, socioeconomic factors, and health behaviours showed independent robust relationships with physical function. Co-variables attenuated the impact of cohabitation, separation, and widowhood on physical function; robust effects were found for singlehood and divorce. Sex-segregated analyses suggest that associations between cohabitation, singlehood, divorce, and widowhood were larger for men than for women. Results suggest that social ties are important to improved physical function.

Keywords

aging; European health; family; physical function; social networks; multilevel models

*Sean Clouston was supported by funding from the U.S. National Institute on Aging (P01 AG043362). Andrea Lawlor is supported by the Social Sciences and Humanities Research Council (SSHRC). Ashton Verdery would like to thank the Duke Network Analysis Center and the Department of Sociology at the University of North Carolina at Chapel Hill for support during the time this study was conducted. This study used data from SHARE Wave 4 release 1.1.1, as of March 28, 2013. The SHARE data collection has been primarily funded by the European Commission through the 5th Framework Programme (project QLK6-CT-2001-00360 in the thematic programme Quality of Life), through the 6th Framework Programme (projects SHARE-I3, RII-CT-2006-062193, COMPARE, CIT5-CT-2005-028857, and SHARELIFE, CIT4-CT-2006-028812) and through the 7th Framework Programme (SHARE-PREP, No. 211909, SHARE-LEAP, No. 227822, and SHARE M4, No. 261982). Additional funding from the U.S. National Institute on Aging (U01 AG09740-13S2, P01 AG005842, P01 AG08291, P30 AG12815, R21 AG025169, Y1-AG-4553-01, IAG BSR06-11, and OGHA 04-064) and the German Ministry of Education and Research as well as from various national sources is gratefully acknowledged (see www.share-project.org for a full list of funding institutions). The funders played no role in the completion of this study.

Correspondence and requests for offprints should be sent to /La correspondance et les demandes de tirés-à-part doivent être adressées à : Sean Clouston, Ph.D. Program in Public Health Health Sciences Center 3-071 Stony Brook University Stony Brook, NY 11794-8338 (sean.clouston@stonybrookmedicine.edu).

Healthy aging is a life course process that incorporates multiple domains of aging (Kuh, Cooper, Hardy, Richards, & Shlomo, 2014). Marital status has been robustly linked to death in a number of countries and across time (Trovato, 1991; 1998; Trovato & Lauris, 1989; Waite, 1995; Waite & Gallagher, 2000), and has been associated with as much as a 10-year extension in lifespan (Waite). Many researchers have thus focused on the role of partnerships in broadly benefiting health (Clouston & Quesnel-Vallée, 2012), reducing mortality (Gove, 1973; Travato & Lauris; Travato, 1987) and maintaining function (Unger, McAvay, Bruce, Berkman, & Seeman, 1999). However, while partnership has been robustly associated with health, estimates of the partnership “benefit” are due in part to the stressful and detrimental aspects of divorce (Amato, 2000; Gahler, 2006) and to the negative association between singlehood and health (Carr & Springer, 2010). Moreover, the harms and benefits of partnership status tend to be more concentrated among men (Kiecolt-Glaser & Newton, 2001; Waite; Waite & Gallagher).

Although there may be a robust association between partnership status and health, the mechanisms that structure that relationship are influenced by a range of economic, social, and behavioural factors (Umberson, Crosnoe, & Reczek, 2010). For example, partnerships consistently provide a source of engagement that influences healthy living across the life course (Waite, 1995; Waite & Gallagher, 2000; Stolzenberg & Waite, 2005). Partnerships may encourage individuals to “clean up” health behaviours (Duncan, Wilkerson, & England, 2006) and may influence the types of activities in which individuals engage (Albert et al., 1995; Angevaren, Aufdemkampe, Verhaar, Aleman, & Vanhees, 2008; Kuh et al., 2014;). It may also be that those who have unobserved predispositions to good health are preferentially selected into partnerships (Goldman, 1993). Berkman, Glass, Brissette, and Seeman (2000) have extended this argument to social networks, arguing that they broadly influence health and function over the life course by contributing to factors such as social support (Callaghan & Morrissey, 2008), influence of alters (individuals in a person's social network) over health-related activities and exposures (Christakis & Fowler, 2007; 2008), embedding of individuals with others with similar health lifestyles (Cockerham, 2005; 2007; Murray & Stafford, 2013), or through improved access to material goods (Phelan & Link, 2013). As a result, researchers have linked a person's social-connectedness to a number of health-related outcomes including health behaviours (Christakis & Fowler), self-perceived health (Cohen & Janicki-Deverts, 2009), cancer (Pinquart & Duberstein, 2010), and all-cause mortality (Holt-Lunstad, Smith & Layton, 2010). Furthermore, partnership, across the life course, is likely to improve the general “health reserve”: the ability for individuals to resist damage and successfully maintain health including, for instance, cognitive (Stern, 2002; 2009) or physiological (Buchner & Wagner, 1992; Xue, 2011) functioning.

Although partnership has been generally associated with health, a better understanding of how partnership “gets under the skin” or is “embodied” is needed (Krieger, 2005). Embodiment may occur as partnership cumulatively influences physical function over the life course factor that influences health. Although there is a good understanding of the association between partnership and self-rated health, the association between partnership and physical function remains under-examined. Physical function, using objective biomarkers of aging-related decline such as grip strength and peak flow, is a robust predictor of health (Cooper et al., 2011b; Warburton, Gledhill, & Quinney, 2001a), healthy aging

(Warburton, Geldhill, & Quinney, 2001b), brain aging (Clouston et al., 2013), and mortality (Cooper, Kuh, & Hardy, 2010; Rantanen et al., 2003). Physical function is interesting to life course analysts because it is susceptible to a lifetime of exposure to modifiable behavioural and structural factors (Franklin & Tate, 2009). For example, stressful work environments have been shown to negatively influence physical function in later life (Russo et al., 2006). Health behaviours, including physical activity (Paterson & Warburton, 2010) and smoking status (van den Borst et al., 2011), similarly influence physical functioning in late life. Finally, socioeconomic status is a persistent and common cause of health and functioning over the entire life course (Link & Phelan, 2010; Phelan & Link, 2013), and it has been associated with physical function in late life (Brunner et al., 2009; Hurst et al., 2013).

Hypotheses

Partnership may influence physical function; however, this influence is likely to be mediated by its strong association with social, behavioural, and economic factors. We hypothesize that partnership will influence physical function. Furthermore, associations between partnership and physical function will be mediated by adjusting for (1) social network characteristics, (2) household income and education, and (3) health behaviours. Finally, we expect that insofar as partnership is related to physical function, that such a relationship will be reduced among women as compared to men.

Methods

Data Source

The Survey of Health, Ageing, and Retirement in Europe (SHARE) biennially interviews a longitudinal panel of individuals aged 50 and older since it began in 2006 (Börsch-Supan et al., 2013). Although SHARE includes data from 19 countries across Europe, three countries – Ireland, Israel, and Greece – were not included in 2010 when SHARE incorporated the first large-scale social networks data collection. Non-response in this sample was 61.6 per cent on average at the household level and 85.3 per cent at the individual level. Nearly 60,000 individuals provided marital status, gender, and age ($n = 56,900$). We further limited the sample to household members aged 55 and older who were living alone or with a spouse for a final sample size of 31,712.

Measures

Objective measures of *physical function* were collected during in-person interviews. Because physical functioning may indicate different domains of functioning (Clouston et al., 2013), we used two indicators of physical function: grip strength and peak flow. To allow comparability between measures and to ease interpretation, we standardized grip strength and peak flow so that one unit equals one standard deviation. *Grip strength* is measured in kilograms (kg) using a handgrip dynamometer, and indicates the function for individuals to grab and hold onto physical objects. We used the maximum physical function over two trials for the dominant hand; in this sample, the dominant hand is the right hand 93 per cent of the time. *Peak flow* is measured in litres per minute (l/m) using a peak flow meter.

Partnership status was measured using self-reported relationship status. Individuals living in cohabiting relationships were identified either as married or living in a common-law partnership. Individuals further noted if they were not in a partnership, whether they were single (never-married), separated, divorced, or widowed.

Social network characteristics were measured using network size, satisfaction, and closeness. *Size* measures the number of non-spouse alters with whom the respondent reported sharing a close social tie (ranging from 0 to a maximum of 7); the average social network includes 2.4 members. We further subdivided this measure to indicate the number of friends and family members that respondents reported seeing regularly (none, one, two, or three or more). *Network dissatisfaction* was measured with a binary indicator developed from a 0–10 score recording whether individuals reported that they were satisfied with the size and functionality of their social networks. Due to the highly skewed data (the average overall was 8.9, or highly satisfied), we dichotomized this indicator to note those who reported being dissatisfied (less than 8 of 10). *Emotionality* indicates the feeling of closeness with alters and was measured using an average across non-spouse alters. Emotionality ranged from 1 (not close at all) to 4 (very close), and averages 3.25 (close to very close).

Education was measured in years to help ensure comparability between countries. *Household income* was measured in euros earned from all sources before taxes in the year preceding Wave 4. Over 10 per cent of values were missing for household income, so we tested the robustness of our analyses to selection by replicating these analyses using multiply-imputed income data; sensitivity analyses did not alter our results. To mark behavioural factors, we include indicators of *smoking status* (never-smoked, not currently smoking, or currently smoking), and *physical activity* (whether respondents reported having been physically active at least once per week over the past month).

Finally, we also adjusted for demographic and physical factors including age, sex, and height. We measured age, centred at 65, using the difference between birth year and year and month of interview. Cooper et al. (2011a) noted that the rate of decline in physical function depends on gender; we therefore included in all models an interaction between gender and age, and further completed sensitivity analyses separately among men and women. We measured *height* in centimetres (cm).

Statistical Analysis

The data were cluster-sampled within households, which were in turn nested within countries. Contextual factors, such as cultural or economic differences, are likely to modify late-life function, thus ignoring this structural inequality is inadvisable. We used restricted maximum likelihood in mixed-effects multilevel models (MLMs) to examine the relationship between partnership status and late-life function. An MLM allows us to estimate the population average “fixed effects”, while also accounting for “random intercepts”: the household- and country-level variance structures under the assumption that the sum of the expected deviations at any particular level is zero. An MLM is advantageous in this context because it can allow us to implicitly account for within-group shared exposures including, for instance, differences in culture, genetic heritage, or geography that may positively or negatively influence physical functioning.

In our analysis, accounting for shared household variability may account for different micro-climates and local context while accounting for shared country-level variability may help to account for shared genetic, historical, educational, cultural, or geographic influences. We used t -tests to determine whether estimates differed between nested models. We calculated pseudo- R^2 to examine model fit and provide R^2 change to examine the role of additional predictors in improving model fit. We assumed independent error structures and estimate random intercepts at both the country and household levels. We used Huber-White robust standard errors. Here, we provide analysis on the whole sample, and on sex-specific subsets. To ensure comparability, we used sex-standardized measures of physical function. To compare model estimates between these samples, we provided differences, which were tested for significance using two-tailed t -tests and pooled standard.

Results

Table 1 provides sample means, standard deviations, and a preliminary examination of the variation between countries and across households. Examining country-level variability using the range in deviations from the sample means adjusted for age, sex, and height, we find substantial differences. There is substantial between-country variation in grip strength and peak flow: mean grip strength ranges from 4.5 kg below average in Spain to 3.4 kg above average in Denmark. Similarly, the percentage of respondents suggesting that they had three or more friends ranges from 0.5 per cent to 6.2 per cent in Poland and Switzerland respectively. Using the country-level variance ratio, we see that country-level differences explain between 1 and 30 per cent of the between-person variability, depending on the measure. Examining household variability, the variability ratio shows that household-level factors may explain a substantial portion of the variation between spouses, often upwards of 70 per cent.

Table 2 provides MLM estimates of associations between partnership status and grip strength. First, we provide bivariate estimates (adjusting for age, sex, age $\times$ sex, and height) between marital status and partnership. Model fit is significant (pseudo- $R^2 = .15$, $p < .001$). Analyses of model fit (results not shown) suggest that incorporating the household-level significantly improved model fit over the one-level model ($p < .001$), and further, that adding country significantly improved the model ($p < .001$). Random-effects estimates in Model 1 similarly exhibit shared variation at the household and country levels.

Examining results in Model 1, we find that marital status is associated with peak flow: those who were single, divorced, or widowed had lower peak flow than those who were married. The model provided a good fit for the outcome. Results show significant within-country and within-household shared variance.

Incorporating social network characteristics (Model 2), we find significant relationships between friend and family network size and peak flow, such that larger networks are associated with improved physical functioning. We also find that the negative impact of divorce (difference = $-.022$, $p < .05$) and widowhood (difference = $-.025$, $p < .05$) increased significantly in size.

Adjusting for socioeconomic factors (Model 3), we find positive associations linking education and income to peak flow. Each year increase in education is associated with an increase in peak flow of approximately 3.4 l/m. Incorporating socioeconomic co-variables attenuates the relationship between cohabitation and peak flow, and it also significantly mediates the relationship between widowhood and peak flow (difference = .03, $p < .05$).

Finally, adjusting for behavioural factors (Model 4), we find a significant relationship between both current smoking status and physical activity with peak flow such that smoking substantially decreases peak flow, while being physically active increases it. Incorporating behavioural factors significantly improved model fit ($p < .001$). Furthermore, we find that the influence of widowhood was attenuated.

Examining the relationship between partnership status and grip strength (see Table 3), we find first that the baseline models fit the data well (pseudo- $R^2 = .33$, $p < .001$). We further note that there is a significant relationship between being single, divorced, or widowed and lowered grip strength. We also find substantial within-country and within-household shared variance in grip strength.

Incorporating social network characteristics (Model 2), we find only limited results. Grip strength is significantly related to having three or more family members, with having two (but not more) friends, and with being dissatisfied with one's social network. However, incorporating these does not improve model fit (R^2 change = .000, n.s.).

Adjusting for income and education significantly improves model fit. Results suggest that having another year of education is associated with an increase in grip strength. Furthermore, we find that having higher household income was similarly associated with improved grip strength. Incorporating socioeconomic factors reduces the overall impact of social network ties and partnership status, and attenuates the relationship between friendship networks and grip strength. Moreover, adjustment for socioeconomic factors attenuates the relationship between widowhood and grip strength.

Accounting for health behaviours (Model 4), we find that being physically active is associated with improved grip strength while being a current smoker is associated with lower grip strength. Incorporating health behaviours significantly improves model fit; however, it does little to mediate associations between partnership status and grip strength.

Table 4 provides analyses separately for females and males using sex-standardized grip strength and peak flow using fully adjusted models (Model 4 from Tables 2 and 3 above). These results generally show stronger associations between partnership status and both grip strength and peak flow among men. For example, there is a significant difference between genders in the influence of cohabitation, which is associated with both grip strength and peak flow among males but not among females. Similarly, the influence of singlehood is larger for men, although it is significant and negative for women.

Sensitivity Analyses

Missing data can be problematic. We assessed the sensitivity of these results to missing data in two ways. First, we incorporated multiply-imputed income data (using 20 imputations separately calculated by sex), which resulted in similar results. Second, we adjusted for the estimated likelihood of sample selection on the outcome variables, which showed similar results. Insofar as missing data are not missing at random, our results linking partnership to physical function are likely to be unbiased, but our results linking less-permanent factors including social network status or smoking status may be conservative because those who are worst off are most likely to be excluded both from the sample and from engaging in such activities. We present results using unimputed data for two reasons: first, multilevel models provide results that are robust when data are missing at random (Atkins, 2005); and second, incorporating estimated data generally reduces the variability without adding information, an unnecessary trade-off when sample sizes are sufficiently large. We have 15 country-level groups – fewer than 20, which is an often-measured optimal benchmark – which may result in a small bias (5%) in the random effects estimates (Maas & Hox, 2005). We continued using country-level variation because national context plays a significant role in determining physical function, with some countries such as Sweden having particularly high physical function (1 kg above average) while others such as Portugal or Spain have particularly low physical function (3 kg below average). Ignoring this country-level variation results in poorer model fit and yields similar results.

Discussion

We used a large and generalizable cross-national dataset to provide robust estimates of the impact of partnership status on physical function in late life. Using multilevel models, we found partnership status to be a robust predictor of physical function. Specifically, after adjustment for social networks, economic factors, and health behaviours, those who were single or divorced had poorer peak flow and lower grip strength than those who were partnered. Furthermore, being dissatisfied with one's social network and having fewer social ties outside of one's partnership were also associated with poorer peak flow but not with grip strength. Our results suggest that non-spousal social interaction may be sizable, and may even be large enough to replace some of the substantial negative impact of partnership status on peak flow, but only in specific contexts (e.g., having three or more family members in one's network) does this sort of interaction impact grip strength. Moreover, these results further support research suggesting that social factors influence late-life function.

Strengths and Limitations

Our results have a number of limitations. We used cross-sectional data, which are biased by a lack of consideration for change over time (Hofer, Berg, & Era, 2003). This is particularly problematic because researchers have suggested that change in physical functioning can be modified by social networks (Unger et al., 1999) and by other life course factors (Peeters, Dobson, Deeg, & Brown, 2013). Our social networks measures are self-reported, truncated, and do not measure informational spread between alters. Similarly, we generally do not have information about the characteristics of alters in the social network, which substantially limits the capability of our analysis to judge selective versus causal patterns. Non-response

was substantial, with around 38 per cent of households not reporting; however, within households, over 85 per cent of individuals responded. Such a pattern of non-response supports a view that those who did not respond did so at the household level because of reasons, such as lack of time or interest, and are thus unlikely to bias these results; however, insofar as those who did not respond did so because they were both living alone and of poor functional status, then our results are likely to be conservative.

Finally, because we lack longitudinal data on social networks, we cannot adequately guarantee that results are consistent over time. However, this is the first study of its kind to test the effects on peak flow and grip strength of the relationship of partnership status and social network characteristics in a large generalizable cohort. These analyses use objective measures of function, which improve over self-reported measures. Furthermore, our analyses use multilevel modeling to adjust for potential clustering at the household and country levels, both of which were found to be significant in our results, in a way that is robust to missing data.

Social Inequalities and Health

Fundamental cause theory posits that resources, including money, power, prestige, knowledge, and beneficial social connections can be used to influence health, and their unequal distribution may cause inequalities in health (Link & Phelan, 1995; Phelan & Link, 2013). Our results show, alternatively, that higher household income, more years of education, and larger social networks were associated with improved physical function. Future analysis should seek to examine whether these characteristics are due to causal influences, or matching-related selection.

Berkman et al. (2000) noted that networks are used to seek and receive social support, influence social engagement and attachment, and, finally, access material goods. A similar argument is routinely made when linking partnership status to health (Amato, 2000; Cheung, 2000; Gahler, 2006; Waite, 1995; Waite & Gallagher, 2000; Williams & Umberson, 2004). We noted this similarity and posited that insofar as social networks can replicate or replace spousal ties, incorporating network data may result in mediation of the relationship between partnership status and late-life function. Our results suggest that social network characteristics are predictive of physical function, and further, that having more extra-marital social ties – either family or friends – could help to replace the substantial negative effects of exiting or lacking a spousal relationship.

Shared Variability

Social science researchers have long highlighted the role that context plays in influencing health (Diez-Roux, 1998; Murray & Stafford, 2013). Researchers have noted that societies can work to alter their health profiles by shaping the context in which health is experienced (Clouston & Quesnel-Vallée, 2012; Clouston et al., 2012; Link, 2008; Link & Phelan, 2009). In our results, we found that countries predicted shared variation, and that adjusting for country-level random intercepts significantly improved model fit. Insofar as this is due to contextual, and not genetic, factors, we might expect that societies are shaping physical function. Such a conclusion does receive some support in the literature; for example, a

recent systematic review noted that area context influences health and functioning later in life (Yen, Michael, & Perdue, 2009).

Results showed substantial shared variance at the household level. Shared variance could indicate one of two effects: selective or causal. Selection may occur if physical function acts as a marital selector (Goldman, 1993), such that healthier people tend to directly or indirectly attract others who are healthy (Clouston & Quesnel-Vallée, 2012; Hewitt & Turell, 2011; Mare, 2008). However, it is also likely that household dynamics help to shape the behaviours, exposures, and health lifestyles of household members (Cockerham, 2005). These results may reflect contextual factors, such as community-level opportunities for physical activity, stressful environments, or vulnerability at the household or community levels, which may influence both partners' capabilities.

Partnership and Functioning

We found support for research suggesting that there is a substantial health benefit to partnerships (Waite, 1995; Waite & Gallagher, 2000). Specifically, we found that individuals in marital partnerships had higher physical function than those who remained single or were divorced. There are substantial stressors involved in divorce, both financial and emotional that may negatively impact overall health but may not be as substantial during separation (Hewitt & Turell, 2011). These analyses support such a conclusion, with divorce – but not separation – showing a robust negative impact on physical functioning. Indeed, divorce broadly influenced health, with those who were divorced having, on average –6.5 l/m lower peak flow and –0.44 kg lower grip strength than those who were not; effects that roughly correspond to the negative effect of having an extra 1.5–1.7 years of extra aging.

Cohabitation is often shown to be a significant predictor of marital dissolution (Mard, 2011), intimate partner violence and depression (Urquia, O'Campo, & Ray, 2012), and mortality (Liu & Reczek, 2012; Scafato et al., 2008). We did not find that those in marital partnerships had better physical functioning over cohabiting couples, but we did find that men in cohabiting partnerships fared worse than did those in marital partnerships. This provides marginal support for prior results showing that marital relationships are healthier than cohabiting ones among men (Waite, 1995), but also suggests that cohabiting partnerships may simply be different from other types of partnership (Wu, 2000). One reason for such may be that even in socially supportive regions, they are less stable than marital unions (Ménard, 2011). Because partnership structure depends on social context, these results may not be widely generalizable beyond societies, like those in Europe or Canada, where cohabitation is a common form of long-term stable relationships (Beaujot, 2000).

The effect of partnership status was substantially smaller among women. These results extend prior work noting that women are less susceptible to partnership-related benefits or detriments than are men (Gahler, 2006) to domains of physical functioning. Differing relationships between partnership status and physical functioning between genders could reflect differences in partnership benefits, selective factors, or coping mechanisms during divorce or singlehood. Our results do not clarify the mechanisms through which inequalities

may arise. Longitudinal research is needed that examines these different explanations in more detail.

Mechanisms

Individuals who engage in physical and cognitive activities may be better protected from declines in function and are more likely to live longer, healthier lives (Cooper et al., 2011; Dumurgier et al., 2009; Thompson et al., 2003; Valenzuela, Sachdev, Wen, Chen, & Brodaty, 2008). We found that larger and more satisfying social networks were protective of peak flow, even after adjusting for smoking status and socioeconomic status. We also found that engagement in regular physical activity was associated with better physical functioning overall. This extends earlier work showing that social engagement can help to protect health and improve healthy aging (Barnes, De Leon, Wilson, Bienias, & Evans, 2004; Krueger et al., 2009; Zunzunegui, Alvarado, Del Ser, & Otero, 2003).

One reason that men may benefit more from partnership than women is that they engage in more risky behaviours prior to partnership (Duncan et al., 2006). We found that incorporating health behaviours explained some of the influence of divorce and of widowhood on peak flow, but were not explanative of relationships between partnership and grip strength. Nevertheless, such behaviours were commonly associated with physical function as a whole and are believed to be integral to social network dynamics. There are substantively interesting reasons to believe that partnership does not uniformly improve health behaviours (Reczek, 2012); more research is needed to examine the extent to which associations between partnership status and health behaviours may be explained by shared, improved contexts or by selective forces early in life.

Incorporating social network characteristics was a significant addition to the model for peak flow, but provided less robust results for grip strength. For peak flow, we found that incorporating social networks also increased the influence of partnership status. Such an increase is suggestive of a “buffering” mechanism if individuals actively supplement partnerships with non-spousal friends and family. Such friendships, if lasting and stable, may effectively replace the role of spouse during times of high stress or when spouses are unavailable, and may further replicate the social engagement benefits that spouses otherwise provide.

Policy Implications

Marital policies have been previously proposed to incentivize marriage and discourage separation or divorce. However, these policies tend to ignore the substantial difficulties that can face individuals when trying to find partners: for example, that the chances of finding a partner are heavily dependent on the partnership market (Lichter, LeClere, & McLaughlin, 1991). Our results suggest that a substantial part of the partnership benefit in physical functioning is due to factors, such as socioeconomic inequalities and health behaviours, which may be more tractable to intervention. Similarly, efforts to increase social connectivity may lead to an improvement in the overall functioning among older individuals. These factors may further reduce partnership-related inequalities if change is focused on increasing involvement among those who are otherwise socially isolated. This

offers further support for the growing policy interest in “network interventions” (Valente, 2012).

References

- Albert MS, Jones K, Savage CR, Berkman L, Seeman T, Blazer D. Predictors of cognitive change in older persons: MacArthur studies of successful aging. *Psychology and Aging*. 1995; 10(4):578–589. [PubMed: 8749585]
- Amato PR. The consequences of divorce for adults and children. *Journal of Marriage and Family*. 2000; 62(4):1269–1287.
- Angevaren M, Aufdemkampe G, Verhaar H, Aleman A, Vanhees L. Physical activity and enhanced fitness to improve cognitive function in older people without known cognitive impairment. *Cochrane database of systematic reviews (Online)*. 2008; 16(3):1–73.
- Atkins DC. Using multilevel models to analyze couple and family treatment data: Basic and advanced issues. *Journal of Family Psychology*. 2005; 19(1):98–110. [PubMed: 15796656]
- Barnes LL, De Leon CM, Wilson RS, Bienias JL, Evans DA. Social resources and cognitive decline in a population of older African Americans and whites. *Neurology*. 2004; 63(12):2322–2326. [PubMed: 15623694]
- Beaujot, R. Earning and caring in Canadian families. Broadview Press; Peterborough, ON: 2000.
- Berkman LF, Glass T, Brissette I, Seeman TE. From social integration to health: Durkheim in the new millennium. *Social Science & Medicine*. 2000; 51(6):843–857. [PubMed: 10972429]
- Börsch-Supan A, Brandt M, Hunkler C, Kneip T, Korbmacher J, Malter F. Data Resource Profile: The Survey of Health, Ageing and Retirement in Europe (SHARE). *International Journal of Epidemiology*. 2013; 42(4):992–1001. [PubMed: 23778574]
- Brunner E, Shipley M, Spencer V, Kivimaki M, Chandola T, Gimeno D, et al. Social inequality in walking speed in early old age in the Whitehall II study. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 2009; 64(10):1082.
- Buchner DM, Wagner EH. Preventing frail health. *Clinics in Geriatric Medicine*. 1992; 8(1)
- Callaghan P, Morrissey J. Social support and health: A review. *Journal of Advanced Nursing*. 2008; 18(2):203–210. [PubMed: 8436711]
- Carr D, Springer KW. Advances in families and health research in the 21st century. *Journal of Marriage and Family*. 2010; 72(3):743–761.
- Cheung YB. Marital status and mortality in British women: A longitudinal study. *International Journal of Epidemiology*. 2000; 29(1):93–99. [PubMed: 10750609]
- Christakis NA, Fowler JH. The spread of obesity in a large social network over 32 years. *New England Journal of Medicine*. 2007; 357(4):370–379. [PubMed: 17652652]
- Christakis NA, Fowler JH. The collective dynamics of smoking in a large social network. *New England Journal of Medicine*. 2008; 358(21):2249–2258. [PubMed: 18499567]
- Clouston S, Brewster P, Kuh D, Richards M, Cooper R, Hardy R. The dynamic relationship between physical function and cognition in longitudinal aging cohorts. *Epidemiologic Reviews*. 2013; 35(1):33–50.
- Clouston S, Kuh D, Herd P, Elliott J, Richards M, Hofer SM. Benefits of educational attainment on adult fluid cognition: International evidence from three birth cohorts. *International Journal of Epidemiology*. 2012; 41(6):1729–1736. [PubMed: 23108707]
- Clouston S, Quesnel-Vallée A. The role of defamilialization in the relationship between partnership and self-rated health: A cross-national comparison of Canada and the United States. *Social Science & Medicine*. 2012; 75:1342–1350. [PubMed: 22800920]
- Cockerham WC. Health lifestyle theory and the convergence of agency and structure. *Journal of Health and Social Behavior*. 2005; 46(1):51–67. [PubMed: 15869120]
- Cockerham, WC. Social causes of health and disease. Polity Press; Cambridge, UK: 2007.
- Cohen S, Janicki-Deverts D. Can we improve our physical health by altering our social networks? *Perspectives on Psychological Science*. 2009; 4(4):375–378. [PubMed: 20161087]

- Cooper R, Hardy R, Sayer AA, Ben-Shlomo Y, Birnie K, Cooper C. Age and gender differences in physical capability levels from mid-life onwards: The harmonisation and meta-analysis of data from eight UK cohort studies. *PLoS ONE*. 2011a; 6(11):1–14. doi:0.1371/journal.pone.0027899.
- Cooper R, Kuh D, Cooper C, Gale CR, Lawlor DA, Matthews F. Objective measures of physical capability and subsequent health: a systematic review. *Age and Ageing*. 2011b; 40(1):14–23. [PubMed: 20843964]
- Cooper R, Kuh D, Hardy R. Objectively measured physical capability levels and mortality: Systematic review and meta-analysis. *British Medical Journal*. 2010; 341(25):639.
- Diez-Roux AV. Bringing context back into epidemiology: Variables and fallacies in multilevel analysis. *American Journal of Public Health*. 1998; 88(2):216–622. [PubMed: 9491010]
- Dumurgier J, Elbaz A, Ducimetière P, Tavernier B, Alperovitch A, Tzourio C. Slow walking speed and cardiovascular death in well functioning older adults: Prospective cohort study. *British Medical Journal*. 2009; 339(b4460):1–7.
- Duncan G, Wilkerson B, England P. Cleaning up their act: The effects of marriage and cohabitation on licit and illicit drug use. *Demography*. 2006; 43(4):691–710. [PubMed: 17236542]
- Franklin NC, Tate CA. Lifestyle and successful aging: An overview. *American Journal of Lifestyle Medicine*. 2009; 3(1):6–11.
- Gahler M. “To divorce is to die a bit ...”: A longitudinal study of marital disruption and psychological distress among Swedish women and men. *The Family Journal*. 2006; 14(4):372–382.
- Goldman N. Marriage selection and mortality patterns: Inferences and fallacies. *Demography*. 1993; 30(2):189–208. [PubMed: 8500636]
- Gove W. Sex, marital status, and mortality. *American Journal of Sociology*. 1973; 79(1):45–67.
- Hewitt B, Turrell G. Short-term functional health and well-being after marital separation: Does initiator status make a difference? *American Journal of Epidemiology*. 2011; 173(11):1308–1318. [PubMed: 21454825]
- Hofer SM, Berg S, Era P. Evaluating the interdependence of aging-related changes in visual and auditory acuity, balance, and cognitive functioning. *Psychology and Aging*. 2003; 18(2):285–305. [PubMed: 12825777]
- Holt-Lunstad J, Smith TB, Layton JB. Social relationships and mortality risk: A meta-analytic review. *PLoS mMedicine*. 2010; 7(7):e1000316.
- Hurst L, Stafford M, Cooper R, Hardy R, Richards M, Kuh D. Lifetime socioeconomic inequalities in physical and cognitive aging. *American Journal of Public Health*. 2013; 103(9):1641–1648. [PubMed: 23865666]
- Kiecolt-Glaser JK, Newton TL. Marriage and health: His and hers. *Psychological Bulletin*. 2001; 127(4):472–503. [PubMed: 11439708]
- Krieger, N. *Embodying inequality: Epidemiologic perspectives*. Baywood; Amityville, NY: 2005.
- Krueger KR, Wilson RS, Kamenetsky JM, Barnes LL, Bienias JL, Bennett DA. Social engagement and cognitive function in old age. *Experimental Aging Research*. 2009; 35(1):45–60. [PubMed: 19173101]
- Kuh, D.; Cooper, R.; Hardy, R.; Richards, M.; Ben Shlomo, Y. *A life course approach to healthy ageing*. Oxford University Press; London, UK: 2014.
- Lichter D, LeClere FB, McLaughlin DK. Local marriage markets and the marital behavior of Black and White women. *The American Journal of Sociology*. 1991; 96(4):843–867.
- Link BG. Epidemiological sociology and the social shaping of population health. *Journal of Health and Social Behavior*. 2008; 49(4):367–384. [PubMed: 19181044]
- Link BG, Phelan JC. Social conditions as fundamental causes of disease. *Journal of Health and Social Behavior*. 1995; 35(Extra Issue):80–94. [PubMed: 7560851]
- Link BG, Phelan JC. The social shaping of health and smoking. *Drug and Alcohol Dependence*. 2009; 104(Supplement 1):S6–S10. [PubMed: 19477607]
- Link, BG.; Phelan, JC. Social conditions as fundamental causes of health inequalities.. In: Bird, C.; Conrad, P.; Fremont, A.; Timmermans, S., editors. *Handbook of Medical Sociology*. 6th ed.. Vanderbilt University Press; Nashville, TN: 2010. p. 3-17.

- Liu H, Reczek C. Cohabitation and US adult mortality: An examination by gender and race. *Journal of Marriage and Family*. 2012; 74(4):794–811.
- Maas CJ, Hox JJ. Sufficient sample sizes for multilevel modeling. *Methodology: European Journal of Research Methods for the Behavioral and Social Sciences*. 2005; 1(3):86–92.
- Mare, R. [1 August 2013] Educational assortative mating in two generations. 2008. from www.econ.ucla.edu/workshops/papers/History/mare_eam_08.pdf
- Ménard F-P. What makes it fall apart? The determinants of the dissolution of marriages and common-law unions in Canada. *McGill Sociological Review*. 2011; 2(1):59–76.
- Murray, ET.; Stafford, M. Lifetime lifestyles III: where we live, the life course and ageing.. In: Kuh, D.; Cooper, R.; Hardy, R.; Richards, M.; Ben Shlomo, Y., editors. *A life course approach to healthy ageing*. Oxford University Press; London, UK: 2013. p. 246-260.
- Paterson DH, Warburton DE. Review Physical activity and functional limitations in older adults: a systematic review related to Canada's Physical Activity Guidelines. *International Journal of Behavioral Nutrition and Physical Activity*. 2010; 7(38):1–22. [PubMed: 20145731]
- Peeters G, Dobson AJ, Deeg DJ, Brown WJ. A life-course perspective on physical functioning in women. *Bulletin of the World Health Organization*. 2013; 91(9):661–670. [PubMed: 24101782]
- Phelan, JC.; Link, BG. Fundamental cause theory.. In: Cockerham, W., editor. *Medical sociology on the move*. Springer; New York, NY: 2013. p. 105-125.
- Pinquart M, Duberstein PR. Associations of social networks with cancer mortality: A meta-analysis. *Critical Reviews in Oncology/Hematology*. 2010; 75(2):122–137. [PubMed: 19604706]
- Rantanen T, Volpato S, Ferrucci L, Heikkinen E, Fried LP, Guralnik JM. Handgrip strength and cause specific and total mortality in older disabled women: Exploring the mechanism. *Journal of the American Geriatrics Society*. 2003; 51(5):636–641. [PubMed: 12752838]
- Reczek C. The promotion of unhealthy habits in gay, lesbian, and straight intimate partnerships. *Social Science & Medicine*. 2012; 75(6):1114–1121. [PubMed: 22703888]
- Russo A, Onder G, Cesari M, Zamboni V, Barillaro C, Capoluongo E. Lifetime occupation and physical function: A prospective cohort study on persons aged 80 years and older living in a community. *Occupational and Environmental Medicine*. 2006; 63(7):438–442. [PubMed: 16782827]
- Scafato E, Galluzzo L, Gandin C, Ghirini S, Baldereschi M, Capurso A. Marital and cohabitation status as predictors of mortality: A 10-year follow-up of an Italian elderly cohort. *Social Science & Medicine*. 2008; 67(9):1456–1464. [PubMed: 18675500]
- Stern Y. What is cognitive reserve? Theory and research application of the reserve concept. *Journal of the International Neuropsychological Society*. 2002; 8:448–460. [PubMed: 11939702]
- Stern Y. Cognitive reserve. *Neuropsychologia*. 2009; 47(10):2015–2028. [PubMed: 19467352]
- Stolzenberg, RM.; Waite, L. Effects of marriage, divorce, and widowhood on health.. In: Bianchi, SM.; Casper, LM.; King, RB., editors. *Work, family, health, and well-being*. Lawrence Erlbaum; Mayway N.J.: 2005. p. 361-378.
- Thompson PD, Buchner D, Piña IL, Balady GJ, Williams MA, Marcus BH. Exercise and physical activity in the prevention and treatment of atherosclerotic cardiovascular disease. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2003; 23(8):e42–e49.
- Trovato F. A longitudinal analysis of divorce and suicide in Canada. *Journal of Marriage and the Family*. 1987; 49(1):193–203.
- Trovato F. Sex, marital status, and suicide in Canada: 1951-1981. *Sociological Perspectives*. 1991; 34(4):427–445. [PubMed: 12343800]
- Trovato F. Nativity, marital status and mortality in Canada. *Canadian Review of Sociology and Anthropology*. 1998; 35(1):65–91.
- Trovato F, Lauris G. Marital status and mortality in Canada: 1951-1981. *Journal of Marriage and the Family*. 1989; 51(4):907–922.
- Umberson D, Crosnoe R, Reczek C. Social relationships and health behavior across life course. *Annual Review of Sociology*. 2010; 36:139–157.
- Unger JB, McAvay G, Bruce ML, Berkman L, Seeman T. Variation in the impact of social network characteristics on physical functioning in elderly persons: MacArthur Studies of Successful Aging.

- The Journals of Gerontology Series B: Psychological Sciences and Social Sciences. 1999; 54B(5):S245–S251.
- Urquia ML, O'Campo PJ, Ray JG. Marital status, duration of cohabitation, and psychosocial well-being among childbearing women: A Canadian nationwide survey. *American Journal of Public Health*. 2012; 103(2):e8–e15. [PubMed: 23237180]
- Valente TW. Network interventions. *Science*. 2012; 337(6090):49–53. [PubMed: 22767921]
- Valenzuela MJ, Sachdev P, Wen W, Chen X, Brodaty H. Lifespan mental activity predicts diminished rate of hippocampal atrophy. *PLoS ONE*. 2008; 3(7):e2598. [PubMed: 18612379]
- van den Borst B, Koster A, Yu B, Gosker HR, Meibohm B, Bauer DC. Is age-related decline in lean mass and physical function accelerated by obstructive lung disease or smoking? *Thorax*. 2011; 66(11):961–969. [PubMed: 21724748]
- Waite L. Does marriage matter? *Demography*. 1995; 32(4):483–507. [PubMed: 8925942]
- Waite, L.; Gallagher, M. *The case for marriage: Why married people are happier, healthier, and better off financially*. Doubleday; New York, NY: 2000.
- Warburton DE, Gledhill N, Quinney A. Musculoskeletal fitness and health. *Canadian Journal of Applied Physiology*. 2001a; 26(2):217–237. [PubMed: 11312417]
- Warburton DE, Gledhill N, Quinney A. The effects of changes in musculoskeletal fitness on health. *Canadian Journal of Applied Physiology*. 2001b; 26(2):161–216. [PubMed: 11312416]
- Williams K, Umberson D. Marital status, marital transitions, and health: A gendered life course perspective. *Journal of Health and Social Behavior*. 2004; 45(1):81–98. [PubMed: 15179909]
- Wu, Z. *Cohabitation: An alternative form of family living*. Oxford University Press; Toronto, ON: 2000.
- Xue Q-L. The frailty syndrome: definition and natural history. *Clinics in Geriatric Medicine*. 2011; 27(1):1–15. [PubMed: 21093718]
- Yen IH, Michael YL, Perdue L. Neighborhood environment in studies of health of older adults: A systematic review. *American Journal of Preventive Medicine*. 2009; 37(5):455–463. [PubMed: 19840702]
- Zunzunegui M-V, Alvarado BE, Del Ser T, Otero A. Social networks, social integration, and social engagement determine cognitive decline in community-dwelling Spanish older adults. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*. 2003; 58(2):S93–S100.

Table 1

Descriptive characteristics including per cent, means, standard deviations, country-level ranges, and both country and household variance ratios (Survey of Health, Ageing and Retirement in Europe, 2010-12)

Variable name	Categories	%	Country Range (Min, Max)		
Male		44.14	(43.40, 48.87)		
Partnership Status	Married	71.79	(61.35, 80.07)		
	Common-law	1.63	(0.00, 8.69)		
	Separated	1.12	(0.26, 2.31)		
	Single	4.84	(2.63, 8.50)		
	Divorced	8.28	(2.06, 12.22)		
	Widowed	12.34	(9.16, 18.54)		
Family Size	None	53.81	(44.89, 75.29)		
	One	20.56	(13.06, 23.86)		
	Two	14.60	(7.84, 18.15)		
	Three or more	11.03	(3.81, 15.53)		
Friendship Size	None	80.03	(67.23, 94.32)		
	One	12.43	(4.11, 19.12)		
	Two	4.82	(1.04, 8.80)		
	Three or more	2.71	(0.52, 6.17)		
Network Dissatisfaction		10.72	(4.99, 13.57)		

Variable name	Mean	SD	Country Variance Ratio		Household Variance Ratio
Emotionality	1.74	0.50	(1.49, 1.92)	0.95	0.19
Age: 65	0.88	9.97	(-0.43, 4.87)	0.99	0.05
Years of Education	10.75	4.35	(6.14, 13.50)	0.82	0.09
Household Income	9.34	1.28	(8.36, 10.40)	0.70	0.00
Grip Strength	33.19	12.10	(28.69, 36.55)	0.98	0.36
Peak Flow	332.21	160.15	(261.28, 399.6)	0.94	0.19

SD = standard deviation

Table 2

Multilevel models estimating the relationship between partnership status and peak flow adjusting for social network characteristics, health behaviours, and socioeconomic factors (Source: Survey of Health, Ageing and Retirement in Europe, 2010-12)

Variable name	Category	Model 1		Model 2		Model 3		Model 4	
		B	SE	B	SE	B	SE	B	SE
Marital Status	Married	Reference		Reference		Reference		Reference	
	Cohabiting	-0.066	0.039	-0.063	0.038	-0.058	0.040	-0.041	0.038
	Separated	-0.021	0.054	-0.041	0.055	-0.038	0.058	-0.025	0.053
	Single	-0.172	0.023	***	-0.186	0.024	***	-0.169	0.026
Divorced		-0.045	0.012	***	-0.067	0.010	***	-0.056	0.013
								***	-0.034
Widowed		-0.041	0.012	***	-0.065	0.014	***	-0.036	0.011
								***	-0.021
Age		-0.028	0.001	***	-0.027	0.001	***	-0.025	0.001
								***	-0.025
Male		0.779	0.031	***	0.804	0.031	***	0.787	0.030
								***	0.801
Age × Male		-0.012	0.001	***	-0.012	0.001	***	-0.013	0.001
								***	-0.013
Height (cm)		0.002	0.002		0.002	0.002	0.002	0.002	0.002
Family Network	None	Reference		Reference		Reference		Reference	
	One	0.057	0.011	***	0.051	0.011	***	0.043	0.011
	Two	0.085	0.015	***	0.075	0.015	***	0.059	0.015
	Three or more	0.152	0.026	***	0.140	0.024	***	0.121	0.025
Friend Network	None	Reference		Reference		Reference		Reference	
	One	0.059	0.017	**	0.040	0.016	*	0.036	0.015
	Two	0.098	0.029	**	0.069	0.028	*	0.062	0.028
	Three or more	0.111	0.028	***	0.071	0.026	**	0.055	0.025
Network Dissatisfaction		0.004	0.034		0.011	0.035		0.020	0.034
Emotionality		0.000	0.017		0.008	0.017		0.014	0.017
Years of Education					0.022	0.003	***	0.020	0.003
								***	0.020
Ln Household Income					0.044	0.011	***	0.039	0.011
								***	0.039

Variable name	Category	Model 1		Model 2		Model 3		Model 4	
		B	SE	B	SE	B	SE	B	SE
Smoking Status	Non-smoker								
	Smoked in past	0.07	0.02	0.07	0.02	0.05	0.02	-0.003	0.018
	Smoker	0.31	0.11	0.30	0.11	0.29	0.11	-0.226	0.016
Physically Active		0.36	0.01	0.36	0.01	0.36	0.01	0.192	0.028
Random Effects									
Country		0.07	0.02	0.07	0.02	0.05	0.02	0.05	0.02
Household		0.31	0.11	0.30	0.11	0.29	0.11	0.29	0.11
Residual		0.36	0.01	0.36	0.01	0.36	0.01	0.35	0.01
Model Fit									
Pseudo- R^2			0.152	***	0.154	***	0.158	***	0.165
R^2 Change				0.002	***	***	0.005	***	0.011
B = coefficient									
SE = standard error									
*** $p < .001$									
** $p < .01$									
* $p < .05$									

Table 3

Multilevel models estimating the relationship between partnership status and grip strength adjusting for social network characteristics, health behaviours, and socioeconomic factors (Source: Survey of Health, Ageing and Retirement in Europe, 2010-12)

Variable name	Category	Model 1		Model 2		Model 3		Model 4	
		B	SE	B	SE	B	SE	B	SE
Marital Status	Married	Reference		Reference		Reference		Reference	
	Cohabiting	-0.035	0.034	-0.033	0.034	-0.030	0.034	-0.024	0.031
	Separated	0.007	0.035	0.005	0.034	0.010	0.032	0.012	0.034
	Single	-0.145	0.013	***	-0.144	0.015	***	-0.132	0.015
Divorced		-0.053	0.014	**	-0.054	0.013	***	-0.044	0.013
								**	-0.040
Widowed		-0.035	0.007	***	-0.035	0.007	***	-0.017	0.009
Age		-0.029	0.001	***	-0.029	0.001	***	-0.028	0.001
								***	-0.026
Male		1.357	0.024	***	1.361	0.024	***	1.352	0.023
								***	1.346
Age × Male		-0.022	0.001	***	-0.022	0.001	***	-0.022	0.001
								***	-0.023
Height (cm)		0.003	0.002		0.003	0.002		0.003	0.002
Family Network	None	Reference		Reference		Reference		Reference	
	One	0.002	0.014		0.002	0.014		-0.008	0.014
	Two	0.003	0.011		0.003	0.011		-0.013	0.011
	Three or more	0.036	0.014	**	0.031	0.014	**	0.015	0.014
Friend Network	None	Reference		Reference		Reference		Reference	
	One	0.012	0.012		0.012	0.011		-0.005	0.011
	Two	0.037	0.012	**	0.021	0.012		0.011	0.013
	Three or more	0.017	0.022		-0.003	0.018		-0.019	0.019
Network Dissatisfaction		-0.032	0.014	**	-0.028	0.014		-0.019	0.014
Emotionality		-0.010	0.009		-0.005	0.010		-0.004	0.010
Years of Education					0.009	0.004	**	0.008	0.004
Ln Household Income					0.032	0.006	***	0.029	0.005
Smoking Status	Non-smoker	Reference		Reference		Reference		Reference	

Variable name	Category	Model 1		Model 2		Model 3		Model 4	
		B	SE	B	SE	B	SE	B	SE
	Smoked in past							0.009	0.010
	Smoker							−0.026	0.012
									**
Physically Active								0.211	0.023

Random Effects									
Country		0.02	0.01	0.02	0.01	0.01	0.00	0.01	0.00
Household		0.05	0.01	0.05	0.01	0.05	0.01	0.05	0.01
Residual		0.33	0.01	0.33	0.01	0.33	0.01	0.33	0.01
Model Fit									
Pseudo- R^2		0.330	***	0.330	***	0.333	***	0.338	***
R^2 Change				0.000		0.002	***	0.007	***

Table 4Influence of partnership status by sex on both grip strength and peak flow using fully adjusted models^a

Variable name	Category	Female		Male		Comparison			
Grip Strength		B _{Female}	SE		B _{Male}	SE	B _{Female} – B _{Male}		
Marital Status	Married	Reference		Reference					
	Cohabiting	0.014	0.072		–0.063	0.040	**	0.077	**
	Separated	–0.030	0.072		0.047	0.040		–0.077	**
	Single	–0.061	0.020	**	–0.271	0.029	***	0.210	***
	Divorced	–0.039	0.026		–0.073	0.026	**	0.034	
	Widowed	0.019	0.013		–0.075	0.028	**	0.095	***
Peak Flow									
Marital Status	Married	Reference		Reference					
	Cohabiting	0.012	0.076		–0.094	0.035	**	0.105	**
	Separated	–0.062	0.081		–0.011	0.059		–0.051	
	Single	–0.094	0.028	***	–0.257	0.038	***	0.162	***
	Divorced	–0.030	0.017	**	–0.063	0.042		0.033	**
	Widowed	–0.022	0.022		–0.056	0.038		0.034	

B = Coefficient

SE = Standard error

 $p < .001$ **
 $p < .01$ *
 $p < .05$ ^a Results are presented using sex-standardized coefficients. Models adjust for social network characteristics, years of education, household income, smoking status, and physical activity. Results further account for shared country-level variance.