Published in final edited form as:

Organic Synth. 2012 ; 2012: . doi:10.1002/0471264229.os088.13.

PHOSPHINE-CATALYZED [3 + 2] ANNULATION: SYNTHESIS OF ETHYL 5-(*tert*-BUTYL)-2-PHENYL-1-TOSYL-3-PYRROLINE-3-CARBOXYLATE

Ian P. Andrews and Ohyun Kwon¹

¹Department of Chemistry and Biochemistry, University of California, Los Angeles, 607 Charles E. Young Drive East, Los Angeles, California 90095-1569

1. Procedure

A. Ethyl 5,5-dimethylhexa-2,3-dienoate (1)

A flame-dried 2-L 3-necked, round-bottomed flask equipped with an overhead mechanical stirrer (Teflon paddle, 6 × 2 × 0.3 cm), a temperature probe, and a nitrogen inlet is charged with ethyl (triphenylphosphoranylidene)acetate (46.4 g, 133 mmol, 1.00 equiv) (Note 1), dichloromethane (665 mL) (Note 9), and triethylamine (20.4 mL, 146 mmol, 1.10 equiv) (Note 10). The mixture is stirred under nitrogen until a homogeneous solution is achieved. 3,3-Dimethylbutyryl chloride (18.5 mL, 133 mmol, 1.00 equiv) (Note 11) is added over 1 h using an addition funnel. The reaction mixture is stirred at room temperature for an additional 18 h during which time a precipitate forms, resulting in a cloudy heterogeneous mixture. The reaction mixture is added in three portions to a 1-L round-bottomed flask and concentrated to a volume of ca. 230 mL using a rotary evaporator (20 °C, 20–25 mmHg). Hexanes (300 mL) (Note 12) is added, and the flask is manually swirled for 2 min, resulting in a precipitate. The heterogeneous suspension is filtered through silica gel (100 g) (Note 13) in a 600-mL glass-fritted medium porosity Büchner funnel into a 2-L round-bottomed flask. The filter cake is rinsed with hexanes (600 mL). The resulting slightly yellow solution is concentrated under rotary evaporation (20 °C, 20–25 mmHg) until 750 mL of the solvent

¹Ethyl (triphenylphosphoranylidene)acetate is commercially available from Aldrich or can be prepared readily from ethyl bromoacetate (Note 2) and triphenylphosphine (Note 3) using the following procedure provided by the submitters: A 1-L round-bottom flask equipped with a magnetic stir bar is charged with triphenylphosphine (39.4 g, 150 mmol) and toluene (200 mL) (Note 4). The mixture is stirred until a homogeneous solution is achieved. Ethyl bromoacetate (16.6 mL, 150 mmol) is added via syringe. The mixture is capped with a plastic stopper and stirred at room temperature for 24 h, during which time a white precipitate is formed, resulting in a thick slurry. The precipitate is collected under vacuum filtration using a glass-fritted Büchner funnel. The filtrate is washed with 75 mL of toluene (Note 5) and 75 mL of diethyl ether (Note 6). The white solid is transferred to a 1-L round-bottomed flask, placed under vacuum for 2 h, and then dissolved in 450 mL of deionized water (Note 7). A magnetic stir bar is added, followed by phenolphthalein indicator solution (200 µL) (Note 8). While stirring, 2 M potassium hydroxide is added dropwise until a pink color persists. The aqueous layer is extracted with dichloromethane (3 × 200 mL) (Note 9) and the combined organic extracts are washed with brine (400 mL), dried over sodium sulfate, filtered, and concentrated under rotary evaporation (30 °C, 20–25 mmHg). The resulting thick oil is redissolved in 20 mL of diethyl ether (Note 6) and concentrated under rotary evaporation (30 °C, 20–25 mmHg) to give an off-white solid. Drying under high vacuum (0.1–0.2 mmHg) yields 48.1 g (92%) of ethyl (triphenylphosphoranylidene)acetate as an off-white solid, mp 118–121 °C. The checkers used commercially available material from Aldrich (95%).

⁹Dichloromethane (99.9%) as purchased from Fluka and used as received.

¹⁰Triethylamine (99.5%) was purchased from Aldrich and used as received.

¹¹3,3-Dimethylbutyryl chloride (99%) was purchased from Aldrich and used as received.

¹²Hexanes (98.5%) was purchased from Aldrich and used as received.

¹³Aldrich (200–400 mesh) silica gel was used as received. The silica gel is packed in the funnel from a slurry in hexanes. The surface of the packed silica is covered with filter paper to prevent disruption.

has been removed (Note 14). The heterogeneous mixture is filtered through silica gel (100 g) (Note 13) in a 600-mL glass fritted medium porosity Büchner funnel into a 2-L round-bottomed flask. The filter cake is rinsed with 500 mL of hexanes/diethyl ether (19:1) (Note 15). The solution is concentrated (20 °C, 20–25 mmHg) and left on vacuum overnight to give ethyl 5,5-dimethylhexa-2,3-dienoate (**1**) (16.4–16.6 g, 73–74%) (Notes 16 and ¹⁷) as a pale yellow oil.

B. Ethyl 5-(tert-butyl)-2-phenyl-1-tosyl-3-pyrroline-3-carboxylate (**2**)

A flame-dried, 1-L 3-necked, round-bottomed flask, equipped with an overhead mechanical stirrer (Teflon paddle, 6 × 2 × 0.3 cm), temperature probe, and a nitrogen inlet, is charged with (*E*)-*N*-benzylidene-4-methylbenzenesulfonamide (7.80 g, 30.0 mmol, 1.00 equiv) (Note 18) and purged with argon. Benzene (300 mL) (Note 19) is added to the flask via syringe, and the mixture is stirred until a homogeneous solution is achieved. Tributylphosphine (1.5 mL, 6.0 mmol, 0.20 equiv) (Note 20) is added via syringe and the reaction mixture is stirred for 5 min. Ethyl 5,5-dimethylhexa-2,3-dienoate (**1**) (6.06 g, 36.0 mmol, 1.20 equiv) is added over 5 min using a syringe pump (Note 21). The reaction is monitored using TLC (Note 22). Upon completion, the mixture is transferred into a 1-L round-bottomed flask and concentrated under rotary evaporation (30 °C, 20–25 mmHg) to a thick orange/brown oil and placed under high vacuum for 10 min (0.1–0.2 mmHg) to ensure the removal of residual solvent (Note 23). A sample of the crude oil (est. 0.5 g) is purified by column chromatography to afford seed crystals of **2** (Notes 24 and ²⁵). A mixture of hexanes/ethyl acetate (9:1, 75 mL) is added to the crude oil in a 1-L round-bottomed flask and a reflux

¹⁴Upon concentration of the solution, a white precipitate forms.

¹⁵After the rinse is complete, thin layer chromatography (TLC) is used to check the flow-through for the presence of compound **1**. The filter cake is rinsed with 6 × 100 mL portions of hexanes/diethyl ether (19:1) until TLC of the flow-through reveals the absence of **1**.

¹⁶The physical properties are as follows: *R*_f = 0.53 (hexanes/ethyl acetate, 19:1); IR: 2964, 2906, 2870, 1959, 1716, 1463, 1413, 1365, 1286, 1255, 1240, 1152, 1095, 1039, 874, 803 cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ: 1.12 (s, 9 H); 1.27 (t, *J* = 7.1 Hz, 3 H), 4.11–4.25 (m, 2 H), 5.59–5.63 (m, 2 H). ¹³C NMR (100 MHz, CDCl₃) δ: 14.1, 29.9, 32.5, 60.5, 89.6, 106.6, 166.1, 210.4. HRMS [C₁₀H₁₆O₂ + H]⁺: calcd. for 169.1223, found 169.1220. Anal. calcd. for C₁₀H₁₆O₂: C, 71.39; H, 9.59. Found: C, 70.79; H, 9.75.

¹⁷Differential scanning calorimetry of (**1**) was completed (DSC, 30 °C to 300 °C, ramp 10 °C/min). Exothermic decomposition was observed with an onset temperature of 90 °C. Heating of the product beyond 30 °C and purification by distillation is therefore not recommended without further safety testing.

¹⁸(*E*)-*N*-Benzylidene-4-methylbenzenesulfonamide was prepared as previously described.²

¹⁹Benzene (>99.5%) was purchased from Fluka and used as received.

²⁰Tributylphosphine (97%) was purchased from Aldrich and used without purification. ³¹P NMR spectroscopy revealed that the reagent comprised 91% tributylphosphine (δ –30.65 ppm) and less than 5% tributylphosphine oxide (δ 48.49 ppm). Three other resonances (δ –17.35, <3%; δ 34.79, <1%; δ 132.33, <2%) were also observed.

²¹The mixture slowly transforms from clear to yellow after the addition of ethyl 5,5-dimethylhexa-2,3-dienoate (**1**).

²²The reaction is monitored using TLC (hexanes/ethyl acetate, 3:1), observing the disappearance of (*E*)-*N*-benzylidene-4-methylbenzene-sulfonamide (*R*_f = 0.54) and the appearance of the product, ethyl 5-(*tert*-butyl)-2-phenyl-1-tosyl-3-pyrroline-3-carboxylate (**2**) (*R*_f = 0.66). Visualization is performed under UV light and through staining with potassium permanganate (prepared by dissolving 20 g of potassium carbonate in 300 mL of water followed by the sequential addition of 5 mL of 5% NaOH(aq) and 3 g of KMnO₄). Potassium permanganate was purchased from Fisher and used as received. The reaction typically reaches completion within 4–7 h. The reaction was checked for completion by TLC after 18 h prior to work-up.

²³If a vacuum of 0.1 mmHg is not reached, the oil can be dried overnight under vacuum (20–30 mmHg), then concentrated from diethyl ether (2 × 50 mL) and hexanes (50 mL) to provide a viscous brown oil.

²⁴The reproducibility of the crystallization is improved by generating a small amount of seed material through column chromatography.

²⁵Column chromatography was performed using a 40 mm-wide glass column packed with Aldrich (200–400 mesh) silica gel (90 g). The column is packed from a slurry of silica gel in hexanes/ethyl acetate (9:1). The crude residue is loaded onto the column in the oily state obtained immediately after rotary evaporation. The flask is rinsed twice with 2 mL portions of hexanes/ethyl acetate (9:1). The column is eluted with hexanes/ethyl acetate (9:1) and fractions are collected. Each collected fraction is analyzed using TLC, and those containing the product are collected, concentrated under rotary evaporation (30 °C, 20–25 mm Hg), and dried under vacuum to afford an oil that crystallized on standing.

condenser is attached. A Teflon-coated stir bar is added and the crude residue is dissolved by heating in an oil bath with stirring until a homogeneous solution is achieved. After cooling to room temperature, the reflux condenser is removed and the flask is capped with a plastic stopper and left to stand overnight. The seed crystals are added as the reaction mixture is cooled. The following day the mixture is placed in a freezer at $-20\text{ }^{\circ}\text{C}$ for 2 h. The crystals that precipitated are collected under vacuum filtration using a glass-fritted funnel (Notes 26 and ²⁷). The crystals are rinsed with hexanes (200 mL) to give 11.4–11.8 g (88–92% yield) of **2** as a white to pale yellow solid (Note 28).

2. Notes

¹Ethyl (triphenylphosphoranylidene)acetate is commercially available from Aldrich or can be prepared readily from ethyl bromoacetate (Note 2) and triphenylphosphine (Note 3) using the following procedure provided by the submitters: A 1-L round-bottom flask equipped with a magnetic stir bar is charged with triphenylphosphine (39.4 g, 150 mmol) and toluene (200 mL) (Note 4). The mixture is stirred until a homogeneous solution is achieved. Ethyl bromoacetate (16.6 mL, 150 mmol) is added via syringe. The mixture is capped with a plastic stopper and stirred at room temperature for 24 h, during which time a white precipitate is formed, resulting in a thick slurry. The precipitate is collected under vacuum filtration using a glass-fritted Büchner funnel. The filtrate is washed with 75 mL of toluene (Note 5) and 75 mL of diethyl ether (Note 6). The white solid is transferred to a 1-L round-bottomed flask, placed under vacuum for 2 h, and then dissolved in 450 mL of deionized water (Note 7). A magnetic stir bar is added, followed by phenolphthalein indicator solution (200 μL) (Note 8). While stirring, 2 M potassium hydroxide is added dropwise until a pink color persists. The aqueous layer is extracted with dichloromethane ($3 \times 200\text{ mL}$) (Note 9) and the combined organic extracts are washed with brine (400 mL), dried over sodium sulfate, filtered, and concentrated under rotary evaporation ($30\text{ }^{\circ}\text{C}$, 20–25 mmHg). The resulting thick oil is redissolved in 20 mL of diethyl ether (Note 6) and concentrated under rotary evaporation ($30\text{ }^{\circ}\text{C}$, 20–25 mmHg) to give an off-white solid. Drying under high vacuum (0.1–0.2 mmHg) yields 48.1 g (92%) of ethyl (triphenylphosphoranylidene)acetate as an off-white solid, mp $118\text{--}121\text{ }^{\circ}\text{C}$. The checkers used commercially available material from Aldrich (95%).

²Ethyl bromoacetate (98%) was purchased from Alfa Aesar and used as received.

²⁶The submitters obtained additional product from the filtrates using column chromatography (Note 27). The checkers did not isolate additional product from the filtrate, but determined by quantitative HPLC analysis that an additional 960 mg of **2** was present in the combined mother liquors and wash.

²⁷Any small amounts of solid left on the filter funnel are dissolved in ethyl acetate and combined with the filtrate prior to concentrating. The filtrate is concentrated under rotary evaporation ($35\text{ }^{\circ}\text{C}$, 20–25 mmHg) and purified through column chromatography to yield an additional 1.0 g (7.8%) of **2** as a white solid. Column chromatography was performed using a 4.75-cm-wide glass column packed with Aldrich (200–400 mesh) silica gel (65 g). The column is packed from a slurry of silica gel in hexanes/ethyl acetate (9:1). The crude residue is loaded onto the column in the oily state obtained immediately after rotary evaporation. The flask is rinsed twice with 2 mL portions of hexanes/ethyl acetate (9:1). The column is eluted with hexanes/ethyl acetate (9:1) and fractions of ca. 22 mL are collected. Each collected fraction is analyzed using TLC and those containing the product are collected, concentrated under rotary evaporation ($30\text{ }^{\circ}\text{C}$, 20–25 mmHg), and dried under vacuum.

²⁸The physical properties are as follows: mp $93\text{--}98\text{ }^{\circ}\text{C}$; IR: 2957, 1714, 1335, 1232, 1164, 1102, 1089, 980 cm^{-1} . ¹H NMR (400 MHz, CDCl_3) δ : 0.80 (s, 9 H), 1.14 (t, $J = 7.1\text{ Hz}$, 3 H), 2.40 (s, 3 H), 4.12 (q, $J = 7.1\text{ Hz}$, 2 H), 4.37 (d, $J = 2.3\text{ Hz}$, 1 H), 5.89 (s, 1 H), 6.73 (dd, $J = 1.2, 2.5\text{ Hz}$, 1 H), 7.32–7.23 (comp, 5 H), 7.43 (d, $J = 7.4\text{ Hz}$, 2 H), 7.71 (d, $J = 8.2\text{ Hz}$, 2 H). ¹³C NMR (100 MHz, CDCl_3) δ : 14.0, 21.5, 27.9, 35.9, 60.8, 68.4, 77.8, 127.5, 127.9, 128.0, 129.6, 134.1, 134.2, 139.7, 141.2, 143.8, 162.7. HRMS [$\text{C}_{24}\text{H}_{29}\text{NO}_4\text{S} + \text{H}^+$]: calcd. for 428.1890, found 428.1890. Anal. calcd. for $\text{C}_{24}\text{H}_{29}\text{NO}_4\text{S}$: C, 67.42; H, 6.84; N, 3.28. Found: C, 67.51; H, 7.14; N, 3.20.

- ³Triphenylphosphine (99%) was purchased from Alfa Aesar and used as received.
- ⁴Toluene (99.8%) was purchased from Fisher Scientific and used as received.
- ⁵Some of the precipitate may stick to the walls of the flask. If so, toluene can be used to rinse the round-bottomed flask.
- ⁶Diethyl ether (99.9%) was purchased from Fisher Scientific and used without further purification.
- ⁷If the addition of water does not result in a homogeneous solution, heating is applied to aid the solubilization of the phosphonium salt.
- ⁸Phenolphthalein indicator solution is prepared by dissolving phenolphthalein powder (100 mg) (purchased from Mallinckrodt and used as received) in 2-propanol (10 mL) (99.9% purchased from Fisher Scientific and used as received). The addition of this indicator is used to help avoid saponification of the ethyl ester, which occurs upon the addition of excess potassium hydroxide.
- ⁹Dichloromethane (99.9%) as purchased from Fluka and used as received.
- ¹⁰Triethylamine (99.5%) was purchased from Aldrich and used as received.
- ¹¹3,3-Dimethylbutyryl chloride (99%) was purchased from Aldrich and used as received.
- ¹²Hexanes (98.5%) was purchased from Aldrich and used as received.
- ¹³Aldrich (200–400 mesh) silica gel was used as received. The silica gel is packed in the funnel from a slurry in hexanes. The surface of the packed silica is covered with filter paper to prevent disruption.
- ¹⁴Upon concentration of the solution, a white precipitate forms.
- ¹⁵After the rinse is complete, thin layer chromatography (TLC) is used to check the flow-through for the presence of compound **1**. The filter cake is rinsed with 6 × 100 mL portions of hexanes/diethyl ether (19:1) until TLC of the flow-through reveals the absence of **1**.
- ¹⁶The physical properties are as follows: R_f = 0.53 (hexanes/ethyl acetate, 19:1); IR: 2964, 2906, 2870, 1959, 1716, 1463, 1413, 1365, 1286, 1255, 1240, 1152, 1095, 1039, 874, 803 cm^{-1} . ^1H NMR (400 MHz, CDCl_3) δ : 1.12 (s, 9 H); 1.27 (t, J = 7.1 Hz, 3 H), 4.11–4.25 (m, 2 H), 5.59–5.63 (m, 2 H). ^{13}C NMR (100 MHz, CDCl_3) δ : 14.1, 29.9, 32.5, 60.5, 89.6, 106.6, 166.1, 210.4. HRMS [$\text{C}_{10}\text{H}_{16}\text{O}_2 + \text{H}$] $^+$: calcd. for 169.1223, found 169.1220. Anal. calcd. for $\text{C}_{10}\text{H}_{16}\text{O}_2$: C, 71.39; H, 9.59. Found: C, 70.79; H, 9.75.
- ¹⁷Differential scanning calorimetry of (**1**) was completed (DSC, 30 °C to 300 °C, ramp 10 °C/min). Exothermic decomposition was observed with an onset temperature of 90 °C. Heating of the product beyond 30 °C and purification by distillation is therefore not recommended without further safety testing.
- ¹⁸(*E*)-*N*-Benzyldiene-4-methylbenzenesulfonamide was prepared as previously described.²
- ¹⁹Benzene (99.5%) was purchased from Fluka and used as received.
- ²⁰Tributylphosphine (97%) was purchased from Aldrich and used without purification. ^{31}P NMR spectroscopy revealed that the reagent comprised 91% tributylphosphine (δ –30.65 ppm) and less than 5% tributylphosphine oxide (δ 48.49 ppm). Three other resonances (δ –17.35, <3%; δ 34.79, <1%; δ 132.33, <2%) were also observed.
- ²¹The mixture slowly transforms from clear to yellow after the addition of ethyl 5,5-dimethylhexa-2,3-dienoate (**1**).
- ²²The reaction is monitored using TLC (hexanes/ethyl acetate, 3:1), observing the disappearance of (*E*)-*N*-benzyldiene-4-methylbenzene-sulfonamide (R_f = 0.54) and the appearance of the product, ethyl 5-(*tert*-butyl)-2-phenyl-1-tosyl-3-pyrroline-3-carboxylate

(2) ($R_f = 0.66$). Visualization is performed under UV light and through staining with potassium permanganate (prepared by dissolving 20 g of potassium carbonate in 300 mL of water followed by the sequential addition of 5 mL of 5% $\text{NaOH}_{(\text{aq})}$ and 3 g of KMnO_4). Potassium permanganate was purchased from Fisher and used as received. The reaction typically reaches completion within 4–7 h. The reaction was checked for completion by TLC after 18 h prior to work-up.

²³If a vacuum of 0.1 mmHg is not reached, the oil can be dried overnight under vacuum (20–30 mmHg), then concentrated from diethyl ether (2×50 mL) and hexanes (50 mL) to provide a viscous brown oil.

²⁴The reproducibility of the crystallization is improved by generating a small amount of seed material through column chromatography.

²⁵Column chromatography was performed using a 40 mm-wide glass column packed with Aldrich (200–400 mesh) silica gel (90 g). The column is packed from a slurry of silica gel in hexanes/ethyl acetate (9:1). The crude residue is loaded onto the column in the oily state obtained immediately after rotary evaporation. The flask is rinsed twice with 2 mL portions of hexanes/ethyl acetate (9:1). The column is eluted with hexanes/ethyl acetate (9:1) and fractions are collected. Each collected fraction is analyzed using TLC, and those containing the product are collected, concentrated under rotary evaporation (30 °C, 20–25 mm Hg), and dried under vacuum to afford an oil that crystallized on standing.

²⁶The submitters obtained additional product from the filtrates using column chromatography (Note 27). The checkers did not isolate additional product from the filtrate, but determined by quantitative HPLC analysis that an additional 960 mg of **2** was present in the combined mother liquors and wash.

²⁷Any small amounts of solid left on the filter funnel are dissolved in ethyl acetate and combined with the filtrate prior to concentrating. The filtrate is concentrated under rotary evaporation (35 °C, 20–25 mmHg) and purified through column chromatography to yield an additional 1.0 g (7.8%) of **2** as a white solid. Column chromatography was performed using a 4.75-cm-wide glass column packed with Aldrich (200–400 mesh) silica gel (65 g). The column is packed from a slurry of silica gel in hexanes/ethyl acetate (9:1). The crude residue is loaded onto the column in the oily state obtained immediately after rotary evaporation. The flask is rinsed twice with 2 mL portions of hexanes/ethyl acetate (9:1). The column is eluted with hexanes/ethyl acetate (9:1) and fractions of ca. 22 mL are collected. Each collected fraction is analyzed using TLC and those containing the product are collected, concentrated under rotary evaporation (30 °C, 20–25 mmHg), and dried under vacuum.

²⁸The physical properties are as follows: mp 93–98 °C; IR: 2957, 1714, 1335, 1232, 1164, 1102, 1089, 980 cm^{-1} . ^1H NMR (400 MHz, CDCl_3) δ : 0.80 (s, 9 H), 1.14 (t, $J = 7.1$ Hz, 3 H), 2.40 (s, 3 H), 4.12 (q, $J = 7.1$ Hz, 2 H), 4.37 (d, $J = 2.3$ Hz, 1 H), 5.89 (s, 1 H), 6.73 (dd, $J = 1.2, 2.5$ Hz, 1 H), 7.32–7.23 (comp, 5 H), 7.43 (d, $J = 7.4$ Hz, 2 H), 7.71 (d, $J = 8.2$ Hz, 2 H). ^{13}C NMR (100 MHz, CDCl_3) δ : 14.0, 21.5, 27.9, 35.9, 60.8, 68.4, 77.8, 127.5, 127.9, 128.0, 129.6, 134.1, 134.2, 139.7, 141.2, 143.8, 162.7. HRMS [$\text{C}_{24}\text{H}_{29}\text{NO}_4\text{S} + \text{H}]^+$: calcd. for 428.1890, found 428.1890. Anal. calcd. for $\text{C}_{24}\text{H}_{29}\text{NO}_4\text{S}$: C, 67.42; H, 6.84; N, 3.28. Found: C, 67.51; H, 7.14; N, 3.20.

Safety and Waste Disposal Information

All hazardous materials should be handled and disposed of in accordance with “Prudent Practices in the Laboratory”; National Academy Press; Washington, DC, 1995.

3. Discussion

The use of nucleophilic tertiary phosphines as catalysts in organic transformations has become a powerful tool for organic synthesis over the past two decades. The discovery that the zwitterions generated from the addition of a phosphine to an electron-deficient allene or alkyne can engage with various electrophiles to produce annulated products has generated much interest in the development of related reactions.³ This flurry of activity is due, in part, to the complexity that can be generated from relatively simple starting materials, as well as the simplicity of the experimental procedures. For instance, workup typically requires no more than rotary evaporation and column chromatography. Most of the developed reactions have entailed the preparation of carbo- and heterocyclic products from simple starting materials, such as olefins, allenes, or acetylenes in combination with alkenes, imines, aldehydes, or dinucleophiles.⁴ In the first examples of such reactions, reported by Xiyan Lu, activated allenes were reacted with electron-deficient alkenes and imines to produce cyclopentenones and pyrrolines, respectively.⁵ Since Lu's original report of such [3 + 2] annulations, many enantioselective variants of the procedure have been described, as well as several applications in the realm of natural product synthesis.^{6,7} Our laboratory discovered that α -alkyl allenates behave as 1,4-dipole-like species in novel [4 + 2] annulations with imines to form tetrahydropyridines.⁸ This broadly applicable reaction has since been applied to natural product synthesis⁹ and has been developed into an asymmetric process.¹⁰ The utility of the phosphine-catalyzed [4 + 2] annulation was further enhanced upon the successful application of activated alkenes to form functionalized cyclohexenones¹¹ and of trifluoromethyl ketones to form dihydropyrans.¹² Of particular importance to this discussion is the synthesis of 2,5-disubstituted pyrroline-3-carboxylates through the reactions of γ -substituted allenates and imines.¹³ 2,5-Disubstituted pyrroline-3-carboxylates are important scaffolds that are commonly used as intermediates en route to other important nitrogen atom-containing heterocycles.¹⁴ Aside from their use as synthetic intermediates, some 2,5-disubstituted-3-pyrrolines themselves exhibit significant biological activity.¹⁵ Recently, our laboratory synthesized a library of 2,5-disubstituted pyrroline-3-carboxylates through the phosphine-catalyzed [3 + 2] annulations of resin-bound allenates and imines. Several of these compounds were identified as geranylgeranyltransferase type-I (GGTase-1) inhibitors (GGTIs) and one, P61A6, exhibits excellent antitumor effects when tested in mice.

Employing the procedure described above provides access to various 2,5-disubstituted pyrroline-3-carboxylates through the reactions of imines and 4-alkyl-2,3-butadienates in the presence of a catalytic amount of tributylphosphine. Table 1 reveals that the reactions of various 4-alkyl- and 4-aryl-2,3-butadienates all led to the desired 2,5-disubstituted pyrrolines in excellent yields (Table 1, entries 1–6). The process is highly diastereoselective, with preferential formation of the syn isomer ranging from 91:9 for the γ -methyl group (Table 1, entry 1) to exclusive for γ -isopropyl, *tert*-butyl, and phenyl groups (Table 1,

entries 4–6). Note that product **2c** (Table 1, entry 3) was formed in a similar reaction employing the 2-butyrate, but in only 63% yield.^{5b}

The scope of the reaction was further established by reacting γ -substituted allenates with various aryl imines (Table 2). The reactions of γ -isopropyl allenate provided the corresponding 3-pyrrolines in excellent yields (Table 2, entries 1–3). The reactions of γ -*tert*-butyl allenate with various aryl imines led to quantitative yields of all products (Table 2, entries 9–16).

Scheme 1 presents a proposed mechanism for the formation of 2,5-disubstituted-3-pyrrolines from γ -substituted allenates and imines. Nucleophilic addition of the phosphine catalyst to the allenate **1** results in the intermediate zwitterion **4**, which then adds to the imine **5** to produce the phosphonium amide zwitterion **6**. A 5-endo-trig cyclization of the amide anion to the vinyl phosphonium cation results in the formation of the ylide **7**. After proton transfer to generate the β -phosphonium enolate intermediate **8**, liberation of the catalyst, through β -elimination of the phosphine, yields the final 2,5-disubstituted-pyrroline-3-carboxylate (**2**).

In summary, various 2,5-disubstituted 3-pyrrolines are readily available from the reactions of γ -substituted allenates with aryl imines in the presence of catalytic amounts of a trialkylphosphine. These phosphine-catalyzed reactions are highly efficient, diastereoselective, operationally simple, and executed under mild conditions.

Acknowledgments

This research was supported by the U.S. National Institutes of Health (O.K.: R01GM071779, P41GM081282).

References

2. Lu K, Kwon O. *Org Synth.* 2009; 86:212. [PubMed: 20161208]
3. For reviews on phosphine-catalyzed reactions, see: Lu X, Zhang C, Xu Z. *Acc Chem Res.* 2001; 34:535. [PubMed: 11456471] Valentine DH, Hillhouse JH. *Synthesis.* 2003; 3:317. Methot JL, Roush WR. *Adv Synth Catal.* 2004; 346:1035. Lu X, Du Y, Lu C. *Pure Appl Chem.* 2005; 77:1985. Nair V, Menon RS, Sreekanth AR, Abhilash N, Biju AT. *Acc Chem Res.* 2006; 39:520. [PubMed: 16906748] Denmark SE, Beutner GL. *Angew Chem, Int Ed.* 2008; 47:1560. Ye L, Zhou J, Tang Y. *Chem Soc Rev.* 2008; 37:1140. [PubMed: 18497927] Kwong CKW, Fu MY, Lam CSL, Toy PH. *Synthesis.* 2008; 2307. Aroyan CE, Dermenci A, Miller SJ. *Tetrahedron.* 2009; 65:4069. Cowen BJ, Miller SJ. *Chem Soc Rev.* 2009; 38:3102. [PubMed: 19847345] Marinetti A, Voituriez A. *Synlett.* 2010:174. Beata K. *Cent Eur J Chem.* 2010:1147. Xu S, He Z. *Sci Sinica Chim.* 2010; 40:856.
4. (a) Zhu XF, Henry CE, Wang J, Dudding T, Kwon O. *Org Lett.* 2005; 7:1387. [PubMed: 15787513] (b) Zhu XF, Schaffner AP, Li RC, Kwon O. *Org Lett.* 2005; 7:2977. [PubMed: 15987184] (c) Dudding T, Kwon O, Mercier E. *Org Lett.* 2006; 8:3643. [PubMed: 16898781] (d) Zhu XF, Henry CE, Kwon O. *J Am Chem Soc.* 2007; 129:6722. [PubMed: 17488018] (e) Henry CE, Kwon O. *Org Lett.* 2007; 9:3069. [PubMed: 17629288] (f) Creech GS, Kwon O. *Org Lett.* 2008; 10:429. [PubMed: 18173275] (g) Creech GS, Zhu X, Fonovic B, Dudding T, Kwon O. *Tetrahedron.* 2008; 64:6935. [PubMed: 19606204] (h) Sriramurthy V, Barcan GA, Kwon O. *J Am Chem Soc.* 2007; 129:12928. [PubMed: 17924625] (i) Guo H, Xu Q, Kwon O. *J Am Chem Soc.* 2009; 131:6318. [PubMed: 19374356]
5. (a) Zhang C, Lu X. *J Org Chem.* 1995; 60:2906. (b) Xu Z, Lu X. *Tetrahedron Lett.* 1999; 40:549. (c) Xia Y, Liang Y, Chen Y, Wang M, Jiao L, Huang F, Liu S, Li Y, Yu Z. *J Am Chem Soc.* 2007;

- 129:3470. [PubMed: 17319666] (d) Mercier E, Fonovic B, Henry C, Kwon O, Dudding T. *Tetrahedron Lett.* 2007; 48:3617.
6. (a) Zhu G, Chen Z, Jiang Q, Xiao D, Cao P, Zhang X. *J Am Chem Soc.* 1997; 119:3836. (b) Wilson JE, Fu GC. *Angew Chem, Int Ed.* 2006; 45:1426. (c) Cowen BJ, Miller SJ. *J Am Chem Soc.* 2007; 129:10988. [PubMed: 17711278] (d) Fang Y, Jacobsen EN. *J Am Chem Soc.* 2008; 130:5660. [PubMed: 18393504] (e) Voituriez A, Panossian A, Fleury-Bregeot N, Retailleau P, Marinetti A. *J Am Chem Soc.* 2008; 130:14030. [PubMed: 18834118]
7. (a) Du Y, Lu X. *J Org Chem.* 2003; 68:6463. [PubMed: 12895091] (b) Wang J, Krische MJ. *Angew Chem, Int Ed.* 2003; 42:5855. (c) Pham TQ, Pyne SG, Skelton BW, White AH. *J Org Chem.* 2005; 70:6369. [PubMed: 16050699]
8. Zhu X, Lan J, Kwon O. *J Am Chem Soc.* 2003; 125:4716. [PubMed: 12696883] . (b) Reference 2.
9. Tran YS, Kwon O. *Org Lett.* 2005; 7:4289. [PubMed: 16146409]
10. Wurz RP, Fu GC. *J Am Chem Soc.* 2005; 127:12234. [PubMed: 16131196]
11. Tran YS, Kwon O. *J Am Chem Soc.* 2007; 129:12632. [PubMed: 17914823]
12. Wang T, Ye S. *Org Lett.* 2010; 12:4168. [PubMed: 20712333]
13. Zhu X, Henry CE, Kwon O. *Tetrahedron.* 2005; 61:6276.
14. For examples of 3-pyrrolines used in natural product syntheses and in pharmaceutical analogues, see: Burley I, Hewson AT. *Tetrahedron Lett.* 1994; 35:7099. Huwe CM, Blechert S. *Tetrahedron Lett.* 1995; 36:1621. Green MP, Prodger JC, Hayes CJ. *Tetrahedron Lett.* 2002; 43:6609.
15. (a) Castellano S, Fiji HDG, Kinderman SS, Watanabe M, de Leon P, Tamanoi F, Kwon O. *J Am Chem Soc.* 2007; 129:5843. [PubMed: 17439124] (b) Watanabe M, Fiji HDG, Guo L, Chan L, Kinderman SS, Slamin DJ, Kwon O, Tamanoi F. *J Biol Chem.* 2008; 283:9571. [PubMed: 18230616] (c) Lu J, Chan L, Fiji HDG, Dahl R, Kwon O, Tamanoi F. *Mol Cancer Ther.* 2009; 8:1218. [PubMed: 19417142]

Biographies

Ohyun Kwon was born in South Korea in 1968. She received her B.S. and M.S. in chemistry (with Eun Lee) from Seoul National University in 1991 and 1993, respectively. She came to the US in 1993 and obtained her PhD (with Samuel J. Danishefsky) from Columbia University in 1998. After a postdoctoral stint in the laboratory of Stuart L. Schreiber at Harvard University, she began her independent career in 2001 as an Assistant Professor at the University of California, Los Angeles (UCLA). In 2008, she was promoted to Associate Professor with tenure at UCLA. Her research involves the development of phosphine-catalyzed reactions and their application to natural product synthesis and chemical biology.

Ian P. Andrews was born in Monterey, California in 1983. He received his B.S. degree from the University of California, San Diego in 2005. He was a summer intern at Pfizer inc. working in medicinal chemistry on the discovery of new carbonic anhydrase inhibitors. He worked at Trinlink Biotechnologies in San Diego before pursuing a Ph.D. at the University of California, Los Angeles. He is currently conducting research in the Laboratory of Dr. Ohyun Kwon at UCLA.

John T. Colyer earned his B.S. in chemistry from Indiana University-Purdue University Indianapolis in 2000 working under the guidance of Professor William H. Moser. He completed his M.S. in 2004 at the University of Arizona under the guidance of Professor Michael P. Doyle, where he studied dirhodium (II) carboxamidate catalysis. In 2004 he joined the Chemical Process Research & Development department at Amgen in Thousand Oaks, CA.

Oliver R. Thiel studied chemistry at the Technical University Munich, Germany and completed a Diploma thesis on rhodium-catalyzed hydroaminations under supervision of Professor Matthias Beller. He then pursued a Ph.D. (1998–2001) at the Max-Planck-Institut für Kohlenforschung Mülheim, Germany under guidance of Professor Alois Fürstner, exploring RCM-reactions in natural product synthesis. After a postdoctoral appointment (2001–2003) at Stanford University with Professor Barry M. Trost, Oliver joined the Chemical Process Research & Development department at Amgen in Thousand Oaks, CA.

where he has been involved in the development of synthetic processes of numerous clinical candidates.

Appendix. Chemical Abstracts Nomenclature; (Registry Number)

Ethyl 5,5-dimethylhexa-2,3-dienoate: 2,3-Hexadienoic acid, 5,5-dimethyl-, ethyl ester; (35802-59-6)

Ethyl (triphenylphosphoranylidene)acetate: Acetic acid, 2-(triphenylphosphoranylidene)-, ethyl ester; (1099-45-2)

3,3-Dimethylbutyryl chloride: Butanoyl chloride, 3,3-dimethyl-; (7065-046-05)

Triethylamine: Ethanamine, *N,N*-diethyl-; (121-44-8)

Ethyl 5-(*tert*-butyl)-2-phenyl-1-tosyl-3-pyrroline-3-carboxylate: 1*H*-Pyrrole-3-carboxylic acid, 5-(1,1-dimethylethyl)-2,5-dihydro-1-[(4-methylphenyl)sulfonyl]-2-phenyl-, ethyl ester, (2*R*,5*R*)-rel-; (861668-04-4)

(*E*)-*N*-Benzylidene-4-methylbenzenesulfonamide: Benzenesulfonamide, 4-methyl-*N*-(phenylmethylene)-, [*N*(*E*)]-; (51608-60-7)

Tributylphosphine; (998-40-3)

Figure 1.
Geranylgeranyltransferase type-I inhibitor P61A6.

Scheme 1.

Proposed Mechanism for the Phosphine-Catalyzed [3 + 2] Annulations of γ -Substituted Allenates and Imines

Table 1

Synthesis of Pyrrolines **2** From γ -Substituted Allenates^a

Entry	R	Time (hr)	Product	Yield (%) ^b	2/3 ^c
1	Methyl (1a)	2	2a/3a	89	91:9
2	Ethyl (1b)	5	2b/3b	99	95:5
3	Propyl (1c)	5	2c/3c	98	96:4
4	Isopropyl (1d)	5	2d	99	100:0
5	<i>tert</i> -Butyl (1)	8	2	99	100:0
6	Phenyl (1e)	1.5	2e	99	100:0

^aReactions were performed on 1 mmol scale.

^bIsolated yield.

^cDetermined using ¹H NMR spectroscopy.

Table 2

Reactions of γ -Substituted Allenates With Aryl Imines^a

Entry	R	Ar	Product	Yield (%) ^b
1	Isopropyl (1d)	2-FC ₆ H ₄	2f	97
2	Isopropyl (1d)	3-BrC ₆ H ₄	2g	95
3	Isopropyl (1d)	4-CF ₃ C ₆ H ₄	2h	96
4	Isopropyl (1d)	4-(<i>i</i> -Pr)C ₆ H ₄	2i	96
5	Phenyl (1e)	2-ClC ₆ H ₄	2j	99
6	Phenyl (1e)	3-ClC ₆ H ₄	2k	99
7	Phenyl (1e)	4-FC ₆ H ₄	2l	99
8	Phenyl (1e)	4-MeOC ₆ H ₄	2m	99
9	<i>tert</i> -Butyl (1)	2-ClC ₆ H ₄	2n	>99
10	<i>tert</i> -Butyl (1)	3-ClC ₆ H ₄	2o	>99
11	<i>tert</i> -Butyl (1)	3,4-Cl ₂ C ₆ H ₃	2p	>99
12	<i>tert</i> -Butyl (1)	4-CNC ₆ H ₄	2q	>99
13	<i>tert</i> -Butyl (1)	4-FC ₆ H ₄	2r	>99
14	<i>tert</i> -Butyl (1)	4-MeC ₆ H ₄	2s	>99
15	<i>tert</i> -Butyl (1e)	4-MeOC ₆ H ₄	2t	>99
16	<i>tert</i> -Butyl (1e)	1-Naphthyl	2u	>99

^a Reactions performed on 1 mmol scale.^b Isolated yield.