Does Well-Child Care Have a Future in Pediatrics?

AUTHORS: Tumaini R. Coker, MD, MBA,^a Tainayah Thomas, MPH,^{b,c} and Paul J. Chung, MD, MS^c

^aUCLA/RAND Prevention Research Center, Los Angeles, California;

^bDepartment of Health Policy and Management, UCLA School of Public Health, Los Angeles, California; and ^cDepartment of Pediatrics, Mattel Children's Hospital, UCLA David Geffen School of Medicine, Los Angeles, California

KEY WORDS

chronic disease, health economics, primary care, well-child care

ABBREVIATIONS

ACO—Accountable Care Organization

SES—Socioeconomic Status

WCC—well-child care

www.pediatrics.org/cgi/doi/10.1542/peds.2013-0252f

doi:10.1542/peds.2013-0252f

Accepted for publication Jan 23, 2013

Address correspondence to Tumaini R. Coker, MD, MBA, UCLA/RAND Prevention Research Center, 10960 Wilshire Blvd, Suite 1550, Los Angeles, CA 90024. E-mail: tcoker@mednet.ucla.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2013 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: No external funding.

abstract

The most common adult chronic diseases affect 1 in 3 adults and account for more than three-quarters of US health care spending. The major childhood drivers of adult disease are distinctly nonmedical: poverty, poor educational outcomes, unhealthy social and physical environments, and unhealthy lifestyle choices. Ideally, well-child care (WCC) would address these drivers and help create healthier adults with more productive lives and lower health care costs. For children without serious acute and chronic medical problems, however, traditional pediatric preventive services may be largely ineffective in addressing the outcomes that really matter; that is, improving lifelong health and reducing the burden of adult chronic disease. In this article, we examine what role WCC has in addressing the major childhood drivers of adult disease and consider various models for the future of WCC within pediatrics. *Pediatrics* 2013;131:S149–S159

Well-child care (WCC) is the foundation of pediatric primary care for most children in the United States. Pediatricians provide the vast majority of this care to children from infancy through adolescence across more than 20 WCC visits. Through these visits, pediatricians have a unique opportunity to identify and address important social, developmental, behavioral, and health issues that could have significant and long-lasting effects on children's lives as adults. In this article, we discuss how this opportunity is routinely squandered, not because pediatricians are inadequate, but because WCC is not designed to address the most critical prevention goals that can affect child health and well-being over a lifetime. First, we review the adult chronic diseases that dominate the overall health care landscape. Next, we discuss the modifiable childhood drivers of these adult chronic diseases that any meaningfully effective WCC system must aim to address. Finally, we discuss the shortcomings of today's WCC in addressing these childhood drivers and present alternative models that may represent WCC's best hope for the 21st century. Although adult health care expenditures already dwarf health care expenditures for children, the potential for child health and well-being to alter adult trajectories presents a tantalizing return on investment that has yet to be fully realized.

In 2011, health care expenditures totaled nearly 18% of the nation's gross domestic product, up from 12% in 1990 and 7% in 1970.^{1,2} It is unclear whether the United States can sustain indefinite growth in health care expenditures as a percentage of gross domestic product; such a trend potentially endangers nonhealth government expenditures; reduces nonhealth private sector growth; and undermines our ability to control state and federal budgets. The ability of the Affordable Care Act to

restrain health care expenditures long-term is equally unclear, and increases in expenditures may accelerate at least temporarily as uninsured Americans gain access.³

Two long-term changes have been cited as key culprits of health care spending growth: the increases in adult obesity and chronic disease.⁴ The proportion of obese adults has more than doubled since 1980, and with that sharp rise, the proportions of adults with related chronic conditions including heart disease, hypertension, and diabetes have increased.^{5–7} Other chronic diseases affecting adult morbidity and mortality, including cancer and mental disorders, have also risen.⁴ Chronic disease management and treatment currently account for 78% of US health care expenditures.⁸

Pediatricians generally assume that they are uniquely positioned to affect a child's lifelong health trajectory. Under their care, children could enter adulthood with a greatly reduced risk of developing many chronic diseases, thus not only reducing the expenditures needed to manage those conditions but also creating a healthier, more productive workforce.

But is this assumption correct? Are pediatricians actually equipped to make a substantial difference in the lives of most children? For the relatively few children who have serious acute or chronic illnesses, pediatricians obviously serve an important role. For the majority of children, however, most of the major drivers of adult disease are distinctly nonmedical (poverty, poor educational outcomes, unhealthy social and physical environments, and unhealthy lifestyle choices).⁹ Physicians and policy makers in the health care system often consider such children to be "healthy," but given the high prevalence of these conditions and their potentially devastating lifelong health impact, it could be argued that these

healthy but at-risk children constitute a population of equal or greater importance for aggressive systematic intervention.

Of course, that is why we have WCC: to improve the health and well-being of children and their families, to give families tools to help children reach their full potential, and to increase the likelihood that children will grow into healthy, happy, and productive adults. Although improving health and well-being means more than reducing the risk of adult chronic disease, it is difficult to conceive of improving health and well-being without reducing this risk.

In this article, we argue that, as currently structured and delivered, WCC, for most at-risk children, does not effectively address these key determinants of child health and well-being that eventually determine adult health. The question, therefore, may not be whether pediatricians can tweak an existing WCC system to maximize benefits. The question may be whether the system, as currently structured, should exist at all.

MODIFIABLE CHILDHOOD DRIVERS OF ADULT CHRONIC DISEASE

Most morbidity and mortality in US adults are directly related to just a handful of chronic diseases. The top 5 causes of death for US adults have been fairly stable over the past 30 years; heart disease, cancer, chronic lower respiratory disease, stroke, and accidents have been the leading causes of death since 1979.¹⁰ The 7 most common chronic diseases are cancer, diabetes, hypertension, stroke, heart disease, pulmonary conditions, and mental illness. More than 1 in 3 US adults report having at least 1 of these conditions, and the combined lost productivity and treatment costs of these diseases have been estimated at \$1.3 trillion annually. These costs are projected to rise

rapidly and steeply as our population continues to age.¹¹

Many of these chronic diseases have their roots in childhood. The main modifiable childhood drivers of these diseases, however, seem not to be childhood illnesses but nonmedical factors such as socioeconomic status (SES), especially poverty and educational attainment; environmental exposures; and health-related behaviors.^{9,12}

Socioeconomic Status: Poverty and Educational Attainment

Poverty and educational attainment are closely related to health; many studies have demonstrated a socioeconomic gradient for health outcomes in childhood and adulthood.^{12–14}

There is compelling evidence linking childhood and adolescent SES with adult health.^{15–18} Childhood poverty and low parental educational attainment are associated with higher risk of death and disease in adulthood,^{15,16} including increased cardiovascular morbidity and mortality, respiratory disease, diabetes, cancers, and poor health-related behaviors such as smoking.¹⁹ College graduates have fewer risky health-related behaviors and other risk factors for the most common adult chronic diseases.^{20,21} Adults with greater educational attainment are less likely to be obese, less likely to be current smokers, and more likely to report vigorous physical activity.²¹ Studies also link educational attainment to lower risk of coronary heart disease and all-cause mortality in adults.^{22–25}

Although the association between childhood SES and adult health is well established, the mechanisms behind this association are complex and poorly understood. In the United States, children in lower SES families often have poorer access to quality health care, resulting in more unmet needs and foregone services that might place

these children at greater risk for health problems as adults.^{26–28} However, SES health disparities persist stubbornly in the face of improvements to health care quality and access.^{17,29} The link between childhood SES and adult health remains robust even in countries with universal access to health care.³⁰

This link points to a more insidious explanation for the childhood SES–adult health association: children in lower SES families are exposed to different neighborhood influences, family environments, housing conditions, educational opportunities, and food choices, which, taken together, may have an impact on future adult health through unhealthy exposures, cumulative stress, and adverse developmental trajectories. A recent study demonstrated a link between the proportion of childhood spent in poverty through age 9 years and elevated allostatic load (a marker of physiologic stress that includes blood pressure, BMI, and epinephrine, norepinephrine, and cortisol levels) at age 17 years.³¹ Lower childhood SES also puts children at risk for exposures in both the physical environment (eg, air pollution) and the social environment (eg, child abuse and neglect)⁹ that can lead to poorer adult health.

Environmental Exposures

Convincing evidence links adult morbidity and premature mortality from chronic disease to adverse physical and social environmental exposures early in life.³² With respect to the social environment, repeated exposure to childhood traumatic events has been associated with chronic disease and cardiovascular risk factors in adulthood. These adverse childhood exposures include child abuse and neglect; domestic violence; and parental substance abuse, mental health disorder, and imprisonment.^{33,34}

There is also growing evidence that exposure to adverse physical environments, both prenatally and postnatally, can contribute to chronic disease risk in adulthood. Fetal growth retardation, low birth weight, and poor weight gain postnatally have been linked to cardiovascular disease and related chronic diseases (ie, stroke, hypertension, type 2 diabetes) in adults.^{35–37} Childhood lead exposure has been associated with central nervous system deficits that persist into adulthood, and childhood environmental tobacco exposure can lead to cardiovascular changes that mirror those found in smokers.^{38–42}

The “built environment” (the constructed indoor and outdoor spaces in the child’s community⁴³) may be another important childhood determinant of adult health. Built environments can confer a number of advantages during childhood, including availability of parks, fresh fruits and vegetables, quality schools, good air quality, safe areas to play, and social support. Low neighborhood-level poverty and other indicators of neighborhood advantage have been linked to child and adult health.^{44–46} A 38-year longitudinal study of neighborhood quality during childhood indicated a positive association with self-reported health during adulthood.⁴⁷ Other elements of the built environment, including design (eg, close proximity of homes, schools, business, and parks to each other; the presence of sidewalks), can increase physical activity among residents.^{43,48} Although there has been no thorough examination of the relationship between the childhood built environment and adult health, the relationship of the built environment with physical activity, social capital, positive mental health outcomes, and lower levels of obesity suggests its potential for long-term impact on health.^{48–50}

Health-Related Behaviors

Four health-related behaviors are major culprits of population-level morbidity

and mortality among US adults: (1) tobacco use; (2) excessive alcohol consumption; (3) poor nutrition; and (4) physical inactivity.⁵¹ These health risk behaviors can start in childhood and are often firmly established by the end of adolescence. Approximately 90% of adult smokers started smoking before the age of 18 years. Reducing the number of adolescents that enter adulthood as smokers has led to a dramatic decrease in the proportion of US adults who smoke.⁵² Still, 10.7% of US teenagers reported tobacco use in 2010; additionally, 13.6% reported alcohol use.⁵³ Both of these risk behaviors often establish themselves before adulthood. Early initiation of alcohol drinking and problem drinking not only poses immediate health risks to adolescents and to society (eg, binge drinking, drunk driving) but also dramatically increases the risk of alcohol dependence during adulthood.⁵⁴

The statistics of the obesity epidemic in this country are well-known; the proportion of obese adults in the United States has doubled in the past 30 years and has tripled among children. Analyses of birth cohorts indicate that the age at which any cohort can expect to have a 20% obesity rate is continually getting younger. The 1976–1985 birth cohort reached an obesity prevalence of 20% by age 20 to 29 years, whereas that proportion was not reached until ages 50 to 59 years for the 1926–1935 birth cohort.⁵⁵ Data also suggest that overweight and obese children are much more likely to become obese adults than normal-weight children.⁵⁶ There is evidence that healthy eating habits and high levels of physical activity established during childhood and early adolescence predict healthy eating and physical activity during adulthood.⁵⁷

HOW DOES TODAY'S WCC MEASURE UP?

Ideally, WCC would directly address these major childhood drivers of adult

disease. From birth to age 21 years, American Academy of Pediatrics guidelines for preventive care recommend that children receive 29 well-child visits. Eleven of these visits are scheduled in the first 3 years of life.⁵⁸ These visits present an opportunity to address these childhood drivers of adult disease and to bring healthier, more productive individuals into adulthood. However, in actual practice, WCC seldom addresses these childhood drivers of adult disease, and the evidence suggests that typical WCC visits do not substantially improve outcomes in these areas.

Poverty

Twenty-two percent of US children live in poverty.⁵⁹ WCC may be able to identify families that have additional needs because of poverty, but there are few WCC services that aim to reduce the negative consequences of poverty on child health and well-being. Inquiries into poverty-related family risk factors, such as homelessness and food insecurity, demonstrate that WCC can identify some families living in poverty. In addition, psychosocial assessment, recommended at every WCC visit from 2 months to 21 years of age, may allow clinicians to recognize stressors associated with poverty and the resultant health-related needs. Some pediatricians may direct families to programs that reduce the impact of poverty on child and maternal nutrition and health care, and connect parents with social services and community resources to address family needs related to poverty, such as housing, employment, and affordable child care.⁵⁸

Parent-reported screening tools for assessment and referral of poverty-related concerns have been developed and tested for use in WCC,^{60,61} but none have been widely adopted, and studies suggest that these social

concerns are usually not elicited during WCC visits. A study of parents in 2 urban pediatric clinics found that, although 82% of families reported having at least 1 psychosocial family concern (eg, housing, food, income insecurity), nearly one-half also reported having at least 1 unmet referral need.⁶¹

Educational Attainment

Developmental surveillance and screening of children throughout infancy, childhood, and adolescence is a well-accepted WCC guideline. Well-child visits present opportunities to identify younger children who may be at risk for developmental delay and older children who may be at risk for poor educational outcomes due to learning disabilities, attention-deficit/hyperactivity disorder, or other developmental, behavioral, and social problems. Despite this, many children at risk for poor educational outcomes are not identified at well-visits. Just one-half of parents nationwide report that their physicians ask about their developmental concerns, and less than one-fifth of children receive standardized screening for developmental delay.^{62,63} There are significant delays in the diagnosis of developmental delay and in the receipt of early-intervention services for many young children.^{64–67} Furthermore, data suggest that children at risk for developmental, behavioral, and social delays may actually receive a lower quality of preventive care services than children who are not at risk.⁶⁸

Comprehensive WCC services may also include screening and intervention for school failure, as well as a general surveillance for achievement at every visit. More than two-thirds of eighth-graders test below proficiency level for math or reading, and 40% of US high school seniors have less than a basic level of achievement on national science examinations.^{69,70} Successful interventions for school failure require more than simply asking how a child is

doing in school or providing brief counseling; successful interventions generally require multiple visits and close communication with school personnel.⁷¹ WCC may have the potential to improve educational outcomes in children, but as currently provided, there is often little systematic or intensive collaboration or integration with schools or school services.

Environmental Exposures

Physical environmental exposures that WCC addresses include lead and tobacco exposure. Bright Futures guidelines recommend lead screening at 12 and 24 months of age, or in accordance with state law.⁵⁸ Tobacco exposure screening is recommended at every visit. A meta-analysis of parent smoking cessation interventions reported a modest 4% increase in the parental quit rate among intervention group parents; moreover, physician-delivered and clinic-based interventions were generally even less effective.⁷²

WCC likewise generally fails to undertake broader assessments of the physical environment, such as addressing the built environment. Pediatricians may be able to help parents understand how the built environment can affect their child's health and can advocate for improvements to the built environment in their local community.⁴³ However, the level of community knowledge, engagement, and participation necessary for a pediatrician to intervene successfully on behalf of their patients may be out of reach for some, and perhaps many, pediatric practices.

The level of pediatrician engagement with community organizations needed to address social environmental exposures can be just as challenging. Many families leave well-child visits without needed strategies or referrals for critical psychosocial family problems ranging from parental intimate partner violence to maternal depression.

Problem-specific tools to screen for these exposures have been developed and tested; research indicates that these problem-specific tools may be more effective than global psychosocial screeners for sensitive family psychosocial issues.⁷³ The resources needed to maintain consistent surveillance, standardized screenings, and targeted intervention for a range of family psychosocial problems, however, extend well beyond the capacities of most pediatric practices.

Health-Related Behaviors

One of the key challenges to addressing concerns related to poverty, educational attainment, and environmental exposures through WCC is that, even with time and incentives to screen children for these important concerns, most pediatricians do not have the training, skill, or capacity to provide intensive services on an ongoing basis and are mainly reliant on referrals. In contrast, health-related behavior change has been more traditionally viewed as within the domain of the pediatrician. Standard recommendations for WCC include screening and counseling on a range of health behaviors, including those linked to chronic diseases in adulthood. As described earlier, the health-related behaviors during childhood that could have the greatest impact on the leading causes of adult morbidity and mortality are: (1) tobacco use; (2) excessive alcohol consumption; (3) poor nutrition; and (4) physical inactivity. The evidence, however, that office-based counseling or guidance during ordinary WCC reduces or prevents these health-related behaviors is weak.

Adolescent Smoking Prevention and Cessation

There are few data to suggest that including simple, brief counseling in WCC visits is beneficial in smoking

prevention or cessation among adolescents. In a randomized controlled trial of a pediatric practice-based adolescent smoking prevention and cessation intervention, the intervention was modestly successful in preventing smoking initiation but was not successful in increasing smoking cessation after a 12-month period.⁷⁴ Moreover, the intervention required a significant amount of clinic time, including counseling by the pediatric provider, followed by a 15- to 30-minute counseling session by a peer counselor, and 4 follow-up telephone counseling sessions. In a systematic review of smoking prevention interventions delivered by health care providers, only 1 of 4 articles meeting inclusion criteria showed a significant difference between the intervention and control groups.⁷⁵ A more recent review examined 24 trials that included >5000 adolescents.⁷⁶ Interventions deemed effective were generally complex and not easily performed in a typical primary care visit with a physician.

Alcohol Use

Successful interventions for the prevention of alcohol use in adolescents generally also require a much more comprehensive approach than would be practical in a typical well-child visit.^{77–79} Universal screening and counseling/guidance for alcohol use is often recommended during adolescent preventive visits, but data suggest that most pediatricians do not screen or counsel adequately, and there are few trials evaluating the effectiveness of alcohol-use screening and intervention during preventive visits.^{80,81} One recent study tested an alcohol screening and brief primary care office intervention (2–3 minutes of physician counseling) that used a nonrandomized, asynchronous study design.⁸² At 12-month follow-up, there was no difference in cessation between the intervention and control groups. The nondrinkers in the intervention group, however, did have

a modestly lower risk of initiation at 12-month follow-up, with a 6% difference between the 2 groups in percentage of adolescents reporting initiating alcohol use.

Obesity, Unhealthy Eating, and Physical Inactivity

Similarly, successful obesity prevention and treatment strategies are generally not simple, office-based interventions that can be easily included in a well-child visit. Systematic reviews identify successful interventions as programs that have substantial intensity, incorporate healthy eating at school, provide sessions for physical activity, include parent support and family-focused activities, and encourage community practices regarding healthy eating and physical activity.⁸³ Although pediatricians may have an important role in obesity prevention and treatment, traditional WCC may not be a particularly useful venue except as an opportunity to screen and refer to an intensive program. Success requires a multidisciplinary approach that includes multiple community settings (eg, home, school, clinic) and in general will reach well beyond WCC.^{84,85}

WHAT IS THE FUTURE FOR WCC?

It is admittedly unfair to argue that WCC does not adequately address fundamental societal problems when, in many respects, other components of society have not done any better. But understanding the value of health care services today is critical, and gauging the value of WCC has always been undermined by uncertainty regarding which outcomes are important. If we believe that preventing or reducing the impact of poverty, poor educational outcomes, unhealthy social and physical environments, and unhealthy lifestyle choices is critically important for lifelong health, then we should evaluate

WCC, in part, by its ability to contribute to that end.

WCC, as currently provided for most children, is a time-compressed, resource-poor, often superficial tour through haphazardly selected topics, led by pediatricians with highly variable levels of expertise in addressing the major childhood drivers of adult disease. The question that remains unanswered is whether pediatricians are willing or able to redesign their practice of health care enough to change this reality. High-quality WCC may be able to play a role in preventing the negative impact of childhood poverty on children's health and well-being, enhancing educational outcomes for children, improving social and physical environments, and encouraging healthy behaviors, but can WCC ever be as central a hub for health promotion as our field has wanted it to be?

There may be several options for the future of WCC, but they likely fit into 1 of 2 basic choices: (1) radically change the model of WCC within the field of pediatrics so that it can actually address the critical nonmedical childhood drivers of adult health; or (2) rethink whether pediatrician-provided WCC should exist at all. Different models of care may be more efficient or effective in addressing the major childhood drivers of adult disease than current common practices; a few are summarized in the following discussion.^{86–89}

One-Stop Shopping Approach

In this model of care, the pediatrician's office or clinic remains the major location for WCC, but a number of other professionals participate to provide services that are not in the area of the clinician's expertise. For example, at a well-child visit, families would have ready access to nonphysicians who could more directly address concerns related to each of the childhood drivers

of adult disease, perhaps including a social worker (childhood poverty and environmental exposures), early childhood development and school specialists (educational attainment), and a health educator (health-related behaviors). This 1-stop shopping approach could bring various professionals under 1 roof, within the framework of a unified team. There might also be other members of the team to supplement these services, such as a nutritionist, family counselor, and a legal aid specialist. Such a model would require fundamental changes in financing to accommodate multiple specialists and services, which poses a serious barrier. Moreover, the model brings into sharp relief the question of what exactly would be left for pediatricians to do.

Community Connections Approach

This approach changes the pediatrician's role in WCC from the central provider to the key coordinator in an integrated community care system for WCC. Children and families would have access to social services, developmental screening and intervention, educational support services, health education, and other services through a network that includes community-based organizations, schools, clinics, social service agencies, recreation centers, parks, and others. This suite of resources may not be under 1 roof, but its components would be linked to each other through the pediatrician in a patient-centered medical home previously characterized in the literature but substantially more expansive than in most current practices.^{90,91} Although financing might not be as directly problematic in this model (assuming that each organization in the network could continue to tap its existing funding streams), its success would depend on the richness of the community network. Moreover, financing concerns might be replaced

by equally problematic communication and data-sharing issues.

Either of these systems (1-stop shopping or community connections) might create a WCC system that could address the major childhood drivers of adult disease. In these models, however, the pediatrician remains the central provider, gatekeeper, coordinator, or organizer of WCC services. Successful examples in limited domains do exist. In Healthy Steps for Young Children, the pediatrician and a child developmental specialist (a nurse, social worker, or early childhood educator) provide WCC in partnership, leading to more positive parenting practices and improved parent experiences with care.^{88,92,93} Medical–legal partnerships link legal services with primary care to identify and intervene when families require legal services for poverty or social environment concerns, such as housing or food insecurity.^{94–96} The Reach Out and Read program has become a basic staple of WCC; this modest intervention improves the home learning environment for children by promoting early literacy habits, particularly for children who might otherwise not have an optimal home learning environment.^{97–99} Proven comprehensive models, however, remain elusive. Promise Neighborhoods, based on the success of the Harlem Children's Zone, integrates and coordinates social, community, educational, and health support for children, using effective schools as the central hub. These types of shared, integrated models may be necessary to fully address the childhood drivers of adult health.^{100,101}

What breadth and depth of training are required to equip a generation of pediatricians not only to provide comprehensive, high-quality, and coordinated care addressing the sometimes complex medical needs of acutely or chronically ill children but also to coordinate effective preventive care

addressing the nonmedical risks related to poverty, low educational achievement, adverse social and physical environmental exposures, and unhealthy lifestyle choices? That these 2 equally important and equally difficult areas of need often coexist within individual patients does not alter the fact that the required skill sets are highly disparate. The recognition that the current image of a high-quality pediatrician requires these divergent skill sets is not new; it has been discussed in the pediatric literature for years and has been an explicit goal of pediatric residency education.^{102,103} As a profession, however, we have not yet addressed the distinct possibility that, for many pediatricians, such an aspiration may be unrealistic.

Pediatricians as Medical Specialists

In several other developed nations, pediatricians are not the primary providers of routine prevention and health promotion services for children. Most children and their families receive health education, anticipatory guidance, developmental screening and surveillance, immunizations, and psychosocial screening and services from public health nurses and general practitioners, working alone or in collaboration with pediatricians. Pediatricians generally are used for consultation or referral for children with acute or chronic medical and developmental/behavioral needs^{87,104,105}; developed nations that use this or a similar model of child health rank higher than the United States on several measures of child health and well-being.^{106,107} In this model, WCC would no longer remain in the purview of primary care pediatrics; currently, however, it is not clear that any other single professional field (eg, public health nursing) has or will have the capacity to manage comprehensive WCC in the

United States substantially better. Therefore, regardless of who serves as the primary contact, a multidisciplinary, largely nonmedical approach would seem necessary.

In this model of care, WCC “centers” would replace the preventive care function of the pediatrician's office or clinic. These centers would be staffed by the same type of multidisciplinary team described earlier in the pediatrics-based WCC models but without the pediatrician. Parents would receive preventive health services from a team of child professionals that might include a public health nurse, child health education specialist, and an education/school specialist (eg, early childhood educator). Similar models have been used in other developed nations,⁸⁷ but such a model's success in the United States would be dependent on restructured reimbursement systems for WCC, redesigned professional education and training around child health for nonphysicians, and a reimagined role for pediatricians as a substantially smaller field of medical care providers. Finally, it would require a strategically designed, multiyear workforce and cultural transition for both families and pediatricians from pediatric-based WCC to pediatric-supervised WCC centers and then finally to team-based, nonphysician-coordinated WCC at WCC centers. Whether this is an effective or viable model for the United States remains unclear. Equally unclear is whether pulling away from direct involvement in major childhood determinants of lifelong health is a defeat that pediatricians should even accept.

In any of these possible models, a new system for reimbursement would be needed to encourage linkages with and utilization of various providers and community-based organizations to address these nonmedical drivers of health. One potentially viable option would be a pediatric-specific accountable care

organization (ACO). An ACO is an integrated system that uses a network of providers, including primary care physicians, subspecialists, and hospitals, to provide care to its patients. This provider network is responsible for the quality, costs, and overall care of its members, and it shares in the savings accrued from improvements in the quality and efficiency of care.¹⁰⁸ Although much of the recent action around ACOs has been focused on Medicare beneficiaries, some pediatric Medicaid-focused ACOs have emerged.¹⁰⁹ To support the key elements of WCC discussed earlier, and to address these 4 main childhood determinants of adult health, a pediatric ACO would have to explicitly include incentives, or even requirements, for provider networks to provide families with access to providers and programs related to education, social, and other related community-level services that may not typically be considered in the usual medical model of pediatric WCC. For example, a pediatric ACO could require that a social worker, legal aid partner, and early childhood education specialist be part of the network for any ACO that serves some minimum number of Medicaid-insured children, and that these additional providers would bill the ACO for services provided. A pediatric ACO could be structured for use in any of the 3 models for pediatric WCC described earlier. Given the additional resources

and lengthy time horizons entailed in preventing adult disease, larger initial investments would likely be necessary, and shared savings might need to be based on achievement of intermediate outcomes and expected, rather than realized, savings due to downstream improvements in adult health. How these savings would be estimated and how payers would be incentivized to provide such payments remain unclear, but some type of integration with adult care payment systems might be necessary.

CONCLUSIONS

Effective prevention and health promotion for children will need to address the critical “nonmedical” childhood drivers of adult health. Few approaches would keep the pediatrician as a central player in these promotion efforts, and all of them would require major restructuring at all levels. Pediatric residency training programs, for instance, are not presently equipped to train young pediatricians to effectively address major nonmedical issues, with the possible exception of health-related behaviors. Much of pediatric primary care training is appropriately dedicated to ensuring that pediatricians are ready to tackle the complex acute and chronic concerns of children with special health care needs. Changing the training paradigm

to cover both medical and nonmedical issues would most likely require a substantially longer training period, new teachers, and new investments in resident education. In addition, WCC reimbursement would have to be completely redesigned to incentivize payers and providers to address these nonmedical concerns in a way that is effective and patient centered, a change that would undoubtedly cost more than usual WCC and require considerable investment in community organizations, many of which may already be under-resourced and overburdened. A real opportunity to reverse the rising costs of adult chronic disease, however, presents an undeniably attractive long-term return on investment that pediatrics currently does not offer.

Primary care pediatrics must radically change if it hopes to serve a central role in prevention and health promotion for children. If pediatrics fails to take up this challenge to redesign our current child preventive health care system, our specialty may find itself losing relevance and influence in the lives of the children and families we serve. That is a possibility that pediatricians will have to assess (or will ultimately have assessed for them) before anyone can determine what the future of pediatrics really is.

REFERENCES

1. Kaiser Family Foundation. Health care spending in the United States and selected OECD countries. Snapshots: health care costs. Available at: www.kff.org/insurance/snapshot/oeod042111.cfm. Accessed October 11, 2012
2. Altarum Institute. Spending Brief. Health sector economic indicators. Available at: www.altarum.org/research-initiatives-health-systems-health-care/altarum-center-for-studying-health-spending/health-indicator-reports. Accessed October 11, 2012
3. Keehan SP, Cuckler GA, Sisko AM, et al. National health expenditure projections: modest annual growth until coverage expands and economic growth accelerates. *Health Aff (Millwood)*. 2012;31(7):1600–1612
4. Thorpe KE, Philyaw M. The medicalization of chronic disease and costs. *Annu Rev Public Health*. 2012;33:409–423
5. Flegal KM, Carroll MD, Kit BK, Ogden CL. Prevalence of obesity and trends in the distribution of body mass index among US adults, 1999–2010. *JAMA*. 2012;307(5):491–497
6. Flegal KM, Carroll MD, Ogden CL, Johnson CL. Prevalence and trends in obesity among US adults, 1999–2000. *JAMA*. 2002; 288(14):1723–1727
7. Bodenheimer T, Chen E, Bennett HD. Confronting the growing burden of chronic disease: can the US health care workforce do the job? *Health Aff (Millwood)*. 2009;28(1):64–74
8. Anderson G. *Chronic Care: Making the Case for Ongoing Care*. Princeton, NJ: Robert Wood Johnson Foundation; 2010

9. Forrest CB, Riley AW. Childhood origins of adult health: a basis for life-course health policy. *Health Aff (Millwood)*. 2004;23(5):155–164
10. Hoyert DL. 75 Years of Mortality in the United States 1935–2010. Washington, DC: US Department of Health and Human Services; 2012
11. DelVol R, Bedroussian A. *An Unhealthy America: The Economic Burden of Chronic Disease*. Santa Monica, CA: Milken Institute; 2007
12. Marmot M. *Closing the Gap in a Generation: Health Equity Through Action on the Social Determinants of Health*. Geneva, Switzerland: World Health Organization; 2008
13. Adler NE, Ostrove JM. Socioeconomic status and health: what we know and what we don't. *Ann N Y Acad Sci*. 1999;896:3–15
14. Loucks EB, Lynch JW, Pilote L, et al. Life-course socioeconomic position and incidence of coronary heart disease: the Framingham Offspring Study. *Am J Epidemiol*. 2009;169(7):829–836
15. Galobardes B, Lynch JW, Davey Smith G. Childhood socioeconomic circumstances and cause-specific mortality in adulthood: systematic review and interpretation. *Epidemiol Rev*. 2004;26:7–21
16. Galobardes B, Lynch JW, Smith GD. Is the association between childhood socioeconomic circumstances and cause-specific mortality established? Update of a systematic review. *J Epidemiol Community Health*. 2008;62(5):387–390
17. Cohen S, Janicki-Deverts D, Chen E, Matthews KA. Childhood socioeconomic status and adult health. *Ann N Y Acad Sci*. 2010;1186:37–55
18. Pollitt RA, Rose KM, Kaufman JS. Evaluating the evidence for models of life course socioeconomic factors and cardiovascular outcomes: a systematic review. *BMC Public Health*. 2005;5:7
19. Gilman SE, Abrams DB, Buka SL. Socioeconomic status over the life course and stages of cigarette use: initiation, regular use, and cessation. *J Epidemiol Community Health*. 2003;57(10):802–808
20. Mirowsky J, Ross C. *Education, Social Status, and Health*. New York, NY: Walter de Gruyter, Inc; 2003
21. Baum S, Ma J, Payea K. *Education Pays: The Benefits of Higher Education for Individuals and Society*. New York, NY: College Board Advocacy and Policy Center; 2010
22. González MA, Rodríguez Artalejo F, Calero JR. Relationship between socioeconomic status and ischaemic heart disease in cohort and case-control studies: 1960–1993. *Int J Epidemiol*. 1998;27(3):350–358
23. Kaplan GA, Keil JE. Socioeconomic factors and cardiovascular disease: a review of the literature. *Circulation*. 1993;88(4 pt 1):1973–1998
24. Harper S, Lynch J, Smith GD. Social determinants and the decline of cardiovascular diseases: understanding the links. *Annu Rev Public Health*. 2011;32:39–69
25. Albert MA, Glynn RJ, Buring J, Ridker PM. Impact of traditional and novel risk factors on the relationship between socioeconomic status and incident cardiovascular events. *Circulation*. 2006;114(24):2619–2626
26. Mayer ML, Skinner AC, Slifkin RT; National Survey of Children With Special Health Care Needs. Unmet need for routine and specialty care: data from the National Survey of Children With Special Health Care Needs. *Pediatrics*. 2004;113(2). Available at: www.pediatrics.org/cgi/content/full/113/2/e109
27. Honberg L, McPherson M, Strickland B, Gage JC, Newacheck PW. Assuring adequate health insurance: results of the National Survey of Children with Special Health Care Needs. *Pediatrics*. 2005;115(5):1233–1239
28. Strickland BB, Jones JR, Ghandour RM, Kogan MD, Newacheck PW. The medical home: health care access and impact for children and youth in the United States. *Pediatrics*. 2011;127(4):604–611
29. Adler NE, Newman K. Socioeconomic disparities in health: pathways and policies. *Health Aff (Millwood)*. 2002;21(2):60–76
30. Mackenbach JP, Stirbu I, Roskam AJ, et al; European Union Working Group on Socioeconomic Inequalities in Health. Socioeconomic inequalities in health in 22 European countries. *N Engl J Med*. 2008;358(23):2468–2481
31. Evans GW, Kim P. Childhood poverty and young adults' allostatic load: the mediating role of childhood cumulative risk exposure. *Psychol Sci*. 2012;23(9):979–983
32. Shonkoff JP, Boyce WT, McEwen BS. Neuroscience, molecular biology, and the childhood roots of health disparities: building a new framework for health promotion and disease prevention. *JAMA*. 2009;301(21):2252–2259
33. Anda RF, Felitti VJ, Bremner JD, et al. The enduring effects of abuse and related adverse experiences in childhood. A convergence of evidence from neurobiology and epidemiology. *Eur Arch Psychiatry Clin Neurosci*. 2006;256(3):174–186
34. Felitti VJ, Anda RF, Nordenberg D, et al. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. The Adverse Childhood Experiences (ACE) Study. *Am J Prev Med*. 1998;14(4):245–258
35. Barker DJ. A new model for the origins of chronic disease. *Med Health Care Philos*. 2001;4(1):31–35
36. Barker DJP, Osmond C, Forsén TJ, Kajantie E, Eriksson JG. Trajectories of growth among children who have coronary events as adults. *N Engl J Med*. 2005;353(17):1802–1809
37. Barker D. Commentary: birthweight and coronary heart disease in a historical cohort. *Int J Epidemiol*. 2006;35:886–887
38. Mazumdar M, Bellinger DC, Gregas M, Abanilla K, Bacic J, Needleman HL. Low-level environmental lead exposure in childhood and adult intellectual function: a follow-up study. *Environ Health*. 2011;10:24
39. Needleman HL, Schell A, Bellinger D, Leviton A, Allred EN. The long-term effects of exposure to low doses of lead in childhood. An 11-year follow-up report. *N Engl J Med*. 1990;322(2):83–88
40. Suzuki T, Tomiyama H, Higashi Y. Vascular dysfunction even after 20 years in children exposed to passive smoking: alarming results and need for awareness. *Arterioscler Thromb Vasc Biol*. 2012;32(4):841–842
41. Kallio K, Jokinen E, Raitakari OT, et al. Tobacco smoke exposure is associated with attenuated endothelial function in 11-year-old healthy children. *Circulation*. 2007;115(25):3205–3212
42. Celermajer DS, Adams MR, Clarkson P, et al. Passive smoking and impaired endothelium-dependent arterial dilatation in healthy young adults. *N Engl J Med*. 1996;334(3):150–154
43. Tester JM; Committee on Environmental Health. The built environment: designing communities to promote physical activity in children. *Pediatrics*. 2009;123(6):1591–1598
44. Leventhal T, Brooks-Gunn J. Moving to opportunity: an experimental study of neighborhood effects on mental health. *Am J Public Health*. 2003;93(9):1576–1582
45. Pickett KE, Pearl M. Multilevel analyses of neighbourhood socioeconomic context and health outcomes: a critical review. *J Epidemiol Community Health*. 2001;55(2):111–122
46. Gordon-Larsen P, Nelson MC, Page P, Popkin BM. Inequality in the built environment underlies key health disparities

- in physical activity and obesity. *Pediatrics*. 2006;117(2):417–424
47. Vartanian TP, Houser L. The effects of childhood neighborhood conditions on self-reports of adult health. *J Health Soc Behav*. 2010;51(3):291–306
 48. Cummins SK, Jackson RJ. The built environment and children's health. *Pediatr Clin North Am*. 2001;48(5):1241–1252, x
 49. Papas MA, Alberg AJ, Ewing R, Helzlsouer KJ, Gary TL, Klassen AC. The built environment and obesity. *Epidemiol Rev*. 2007; 29:129–143
 50. Renalds A, Smith TH, Hale PJ. A systematic review of built environment and health. *Fam Community Health*. 2010;33(1):68–78
 51. Mokdad AH, Marks JS, Stroup DF, Gerberding JL. Actual causes of death in the United States, 2000. *JAMA*. 2004;291 (10):1238–1245
 52. US Department of Health and Human Services. *Preventing Tobacco Use Among Youth and Young Adults: A Report of the Surgeon General*. Rockville, MD: US Department of Health and Human Services; Centers for Disease Control and Prevention; National Center for Chronic Disease Prevention and Health Promotion; Office on Smoking and Health; 2012
 53. Substance Abuse and Mental Health Services Administration. *Results From the 2010 National Survey on Drug Use and Health: Summary of National Findings*. Rockville, MD: HHS Publication No. (SMA) 11-4658; 2011
 54. Dawson DA, Goldstein RB, Chou SP, Ruan WJ, Grant BF. Age at first drink and the first incidence of adult-onset DSM-IV alcohol use disorders. *Alcohol Clin Exp Res*. 2008;32:2149–2160
 55. Lee JM, Pilli S, Gebremariam A, et al. Getting heavier, younger: trajectories of obesity over the life course. *Int J Obes (Lond)*. 2010;34(4):614–623
 56. Freedman DS, Khan LK, Dietz WH, Srinivasan SR, Berenson GS. Relationship of childhood obesity to coronary heart disease risk factors in adulthood: the Bogalusa Heart Study. *Pediatrics*. 2001;108(3):712–718
 57. Daniels SR, Pratt CA, Hayman LL. Reduction of risk for cardiovascular disease in children and adolescents. *Circulation*. 2011;124(15):1673–1686
 58. Hagan J, Shaw J, Duncan P, eds. *Bright Futures Guidelines for Health Supervision of Infants, Children, and Adolescents*. 3rd ed. Elk Grove, IL: American Academy of Pediatrics; 2008
 59. Child Trends. Children in poverty. Available at: www.childtrendsdatabank.org/?q=node/221. Accessed May 8, 2012
 60. Garg A, Butz AM, Dworkin PH, Lewis RA, Thompson RE, Serwint JR. Improving the management of family psychosocial problems at low-income children's well-child care visits: the WE CARE Project. *Pediatrics*. 2007;120(3):547–558
 61. Fleegler EW, Lieu TA, Wise PH, Muret-Wagstaff S. Families' health-related social problems and missed referral opportunities. *Pediatrics*. 2007;119(6). Available at: www.pediatrics.org/cgi/content/full/119/6/e1332
 62. Guerrero AD, Rodríguez MA, Flores G. Disparities in provider elicitation of parents' developmental concerns for US children. *Pediatrics*. 2011;128(5):901–909
 63. Bethell C, Reuland C, Schor E, Abrahms M, Halfon N. Rates of parent-centered developmental screening: disparities and links to services access. *Pediatrics*. 2011; 128(1):146–155
 64. Mann JR, Crawford S, Wilson L, McDermott S. Does race influence age of diagnosis for children with developmental delay? *Disabil Health J*. 2008;1(3):157–162
 65. Rosenberg SA, Zhang D, Robinson CC. Prevalence of developmental delays and participation in early intervention services for young children. *Pediatrics*. 2008; 121(6). Available at: www.pediatrics.org/cgi/content/full/121/6/e1503
 66. Mandell DS, Listerud J, Levy SE, Pinto-Martin JA. Race differences in the age at diagnosis among Medicaid-eligible children with autism. *J Am Acad Child Adolesc Psychiatry*. 2002;41(12):1447–1453
 67. Mandell DS, Novak MM, Zubritsky CD. Factors associated with age of diagnosis among children with autism spectrum disorders. *Pediatrics*. 2005;116(6):1480–1486
 68. Coker TR, Shaikh Y, Chung PJ. Parent-reported quality of preventive care for children at-risk for developmental delay. *Acad Pediatr*. 2012;12(5):384–390
 69. US Department of Education National Center for Education Statistics. National Assessment of Educational Progress. Available at: <http://nces.ed.gov/nationsreportcard>. Accessed July 24, 2012
 70. Child Trends. Science proficiency. Available at: www.childtrendsdatabank.org/?q=node/257. Accessed May 8, 2012
 71. Byrd RS. School failure: assessment, intervention, and prevention in primary pediatric care. *Pediatr Rev*. 2005;26(7): 233–243
 72. Tanski SE, Klein JD, Winickoff JP, Auinger P, Weitzman M. Tobacco counseling at well-child and tobacco-influenced illness visits: opportunities for improvement. *Pediatrics*. 2003;111(2). Available at: www.pediatrics.org/cgi/content/full/111/2/E162
 73. Garg A, Dworkin PH. Applying surveillance and screening to family psychosocial issues: implications for the medical home. *J Dev Behav Pediatr*. 2011;32(5): 418–426
 74. Pbert L, Flint AJ, Fletcher KE, Young MH, Druker S, DiFranza JR. Effect of a pediatric practice-based smoking prevention and cessation intervention for adolescents: a randomized, controlled trial. *Pediatrics*. 2008;121(4). Available at: www.pediatrics.org/cgi/content/full/121/4/e738
 75. Christakis DA, Garrison MM, Ebel BE, Wiehe SE, Rivara FP. Pediatric smoking prevention interventions delivered by care providers: a systematic review. *Am J Prev Med*. 2003;25(4):358–362
 76. Grimshaw GM, Stanton A. Tobacco cessation interventions for young people. *Cochrane Database Syst Rev*. 2010;(4): CD003289
 77. Lemstra M, Bennett NR, Neudorf C, Kunst A, Nannapaneni U, Warren LM, Kershaw T, Scott CR. A meta-analysis of marijuana and alcohol use by socio-economic status in adolescents aged 10-15 years. *Can J Public Health*. 2008 May-Jun;99(3):172–177
 78. Stevens MM, Olson AL, Gaffney CA, Tosteson TD, Mott LA, Starr PA. A pediatric, practice-based, randomized trial of drinking and smoking prevention and bicycle helmet, gun, and seatbelt safety promotion. *Pediatrics*. 2002;109(3):490–497
 79. Foxcroft DR, Ireland DJ, Lister-Sharp DJ, Lowe G, Breen R. Longer-term primary prevention for alcohol misuse in young people: a systematic review. *Addiction*. 2003;98(4):397–411
 80. Millstein SG, Marcell AV. Screening and counseling for adolescent alcohol use among primary care physicians in the United States. *Pediatrics*. 2003;111(1):114–122
 81. Kulig JW; American Academy of Pediatrics Committee on Substance Abuse. Tobacco, alcohol, and other drugs: the role of the pediatrician in prevention, identification, and management of substance abuse. *Pediatrics*. 2005;115(3):816–821
 82. Harris SK, Csémy L, Sherritt L, et al. Computer-facilitated substance use screening and brief advice for teens in primary care: an international trial. *Pediatrics*. 2012; 129(6):1072–1082
 83. Waters E, de Silva-Sanigorski A, Hall BJ, et al. Interventions for preventing obesity in children. *Cochrane Database Syst Rev*. 2011;(12):CD001871
 84. Pratt CA, Stevens J, Daniels S. Childhood obesity prevention and treatment:

- recommendations for future research. *Am J Prev Med*. 2008;35(3):249–252
85. Hopkins KF, Decristofaro C, Elliott L. How can primary care providers manage pediatric obesity in the real world? *J Am Acad Nurse Pract*. 2011;23(6):278–288
 86. Zuckerman B, Parker S. Preventive pediatrics—new models of providing needed health services. *Pediatrics*. 1995;95(5):758–762
 87. Kuo AA, Inkelas M, Lotstein DS, Samson KM, Schor EL, Halfon N. Rethinking well-child care in the United States: an international comparison. *Pediatrics*. 2006;118(4):1692–1702
 88. Minkovitz CS, Strobino D, Mistry KB, et al. Healthy Steps for Young Children: sustained results at 5.5 years. *Pediatrics*. 2007;120(3). Available at: www.pediatrics.org/cgi/content/full/120/3/e658
 89. Halfon N, DuPlessis H, Inkelas M. Transforming the US child health system. *Health Aff (Millwood)*. 2007;26(2):315–330
 90. Medical Home Initiatives for Children With Special Needs Project Advisory Committee. American Academy of Pediatrics. The medical home. *Pediatrics*. 2002;110(1 pt 1):184–186
 91. Patient-Centered Primary Care Collaborative. Joint principles of the patient-centered medical home. Available at: www.pcpcc.net/node/14. Accessed June 28, 2010
 92. Zuckerman B, Parker S, Kaplan-Sanoff M, Augustyn M, Barth MC. Healthy Steps: a case study of innovation in pediatric practice. *Pediatrics*. 2004;114(3):820–826
 93. Piotrowski CC, Talavera GA, Mayer JA. Healthy Steps: a systematic review of a preventive practice-based model of pediatric care. *J Dev Behav Pediatr*. 2009;30(1):91–103
 94. Sandel M, Hansen M, Kahn R, et al. Medical-legal partnerships: transforming primary care by addressing the legal needs of vulnerable populations. *Health Aff (Millwood)*. 2010;29(9):1697–1705
 95. Beck AF, Klein MD, Schaffzin JK, Tallent V, Gillam M, Kahn RS. Identifying and treating a substandard housing cluster using a medical-legal partnership. *Pediatrics*. 2012;130(5):831–838
 96. National Center for Medical Legal Partnership. The National Center for Medical Legal Partnership: raising the bar for health. Available at: www.medical-legalpartnership.org/. Accessed November 1, 2012
 97. Needlman R, Toker KH, Dreyer BP, Klass P, Mendelsohn AL. Effectiveness of a primary care intervention to support reading aloud: a multicenter evaluation. *Ambul Pediatr*. 2005;5(4):209–215
 98. Zuckerman B. Promoting early literacy in pediatric practice: twenty years of reach out and read. *Pediatrics*. 2009;124(6):1660–1665
 99. Weitzman CC, Roy L, Walls T, Tomlin R. More evidence for reach out and read: a home-based study. *Pediatrics*. 2004;113(5):1248–1253
 100. Komro KA, Flay BR, Biglan A; Promise Neighborhoods Research Consortium. Creating nurturing environments: a science-based framework for promoting child health and development within high-poverty neighborhoods. *Clin Child Fam Psychol Rev*. 2011 Jun;14(2):111–134
 101. US Department of Education. The Promise Neighborhoods program. Available at: www2.ed.gov/programs/promiseneighborhoods/index.html. Accessed November 1, 2012
 102. Leslie L, Rappo P, Abelson H, et al. Final report of the FOPE II Pediatric Generalists of the Future Workgroup. *Pediatrics*. 2000;106(5):1199–1223
 103. Hicks PJ, Englander R, Schumacher DJ, Burke A, Benson BJ, Guralnick S, Ludwig S, Carraccio C. Pediatrics milestone project: next steps toward meaningful outcomes assessment. *J Grad Med Educ*. 2010;2:577–584
 104. Williams BC, Miller CA. American Academy of Pediatrics. Preventive health care for young children: findings from a 10-country study and directions for United States policy. *Pediatrics*. 1992;89(5 pt 2):981–998
 105. Chaulk CP. Preventive health care in six countries: models for reform? *Health Care Financ Rev*. 1994;15(4):7–19
 106. UNICEF. *Child Poverty in Perspective: An Overview of Child Well-Being in Rich Countries, Innocenti Report Card 7*. Florence, Italy: UNICEF Innocenti Research Centre; 2007. Available at: www.unicef.org/media/files/ChildPovertyReport.pdf. Accessed February 18, 2013
 107. OECD. OECD family database. Available at: www.oecd.org/social/family/database. Accessed May 8, 2012
 108. Goldsmith J. Accountable care organizations: the case for flexible partnerships between health plans and providers. *Health Aff (Millwood)*. 2011;30(1):32–40
 109. Nationwide Children's. America's largest pediatric accountable care organization. Available at: www.nationwidechildrens.org/largest-pediatric-accountable-care-organization. Accessed November 8, 2012