Published in final edited form as:

Met Ions Life Sci. 2014 ; 14: 71–97. doi:10.1007/978-94-017-9269-1_4.

Investigations of the Efficient Electrocatalytic Interconversions of CO₂ and CO by Nickel-containing Carbon monoxide Dehydrogenases

Vincent Wang, Stephen W Ragsdale, and Fraser A. Armstrong

[†]Department of Chemistry, Inorganic Chemistry Laboratory, University of Oxford, South Park Road, Oxford, OX1 3QR, U.K

[‡]Department of Biological Chemistry, University of Michigan, Ann Arbor, Michigan 48109-0606, USA

Fraser A. Armstrong: fraser.armstrong@chem.ox.ac.uk

Abstract

Carbon monoxide dehydrogenases (CODH) play an important role in utilizing carbon monoxide (CO) or carbon dioxide (CO₂) in the metabolism of some microorganisms. Two distinctly different types of CODH are distinguished by the elements constituting the active site. A Mo-Cu containing CODH is found in some aerobic organisms, whereas a Ni-Fe containing CODH (henceforth simply Ni-CODH) is found in some anaerobes. Two members of the simplest class (IV) of Ni-CODH behave as efficient, reversible electrocatalysts of CO₂/CO interconversion when adsorbed on a graphite electrode. Their intense electroactivity sets an important benchmark for the standard of performance at which synthetic molecular and material electrocatalysts comprised of suitably attired abundant first-row transition elements must be able to operate. Investigations of CODHs by protein film electrochemistry (PFE) reveal how the enzymes respond to the variable electrode potential that can drive CO₂/CO interconversion in each direction, and identify the potential thresholds at which different small molecules, both substrates and inhibitors, enter or leave the catalytic cycle. Experiments carried out on a much larger (Class III) enzyme CODH/ACS, in which CODH is complexed tightly with acetylCoA synthase, show that some of these characteristics are retained, albeit with much slower rates of interfacial electron transfer, attributable to the difficulty in making good electronic contact at the electrode. The PFE results complement and clarify investigations made using spectroscopic investigations.

1. Direct CO₂/CO Interconversions in Biology

Finding new ways of using renewable energy to reduce carbon dioxide (CO₂) to fuels and thus supplement biological photosynthetic CO₂ fixation, is a major scientific challenge with huge implications for future civilizations. Green plants and many microorganisms generally fix CO₂ by photosynthesis – a process that is familiar to everyone: however, some microorganisms use a totally different system for fixing CO₂ and use pathways in which carbon monoxide (CO) is an important intermediate. Moreover, some anaerobic desulfuricants, such as *Desulfotomaculum carboxydivorans*, hydrogenogens such as *Carboxydotherrmus hydrogenoformans* (*Ch*), some phototrophs, such as *Rhodospirillum rubrum* (*Rr*), CO₂-utilizing microorganisms and aerobic carboxidotrophs, such as

Oligotropha carboxidovorans, are able to use CO as their sole carbon and energy source. These alternative pathways have been documented in various review articles [1–6]. The metalloenzymes involved are called carbon monoxide dehydrogenases (CODH) and they provide particularly useful insight into the chemical principles underlying catalysis of carbon dioxide activation. The primary reaction being catalyzed is the half-cell reaction that is written as Equation 1.

Two chemically distinct types of CODH are distinguished by the structure of their active site. Some aerobes use enzymes containing a sulfido-linked Cu-S-Mo(pyranopterin) center [5, 7]; on the other hand, some anaerobic microorganisms, such as *Moorella thermoacetica* (*Mt*) and *Carboxydotherrmus hydrogenoformans* (*Ch*) use enzymes in which the active site is a distorted [Ni₃Fe-4S] cubane-like cluster with a pendant (dangling) Fe [8]. The entire [Ni₄Fe-4S] cluster is referred to as the ‘C-cluster’ and the corresponding enzymes are known as Ni-CODHs. The Ni-CODH family is extended to include the class of enzymes known as CODH/ACS, in which a Ni-CODH is strongly associated with an acetyl CoA synthase. In acetogenic bacteria, CODH-ACS catalyzes C-C bond formation using CO (formed from CO₂) and a methyl group (donated by B₁₂) to form acetyl-CoA (Equation 2): in methanogenic archaea, Ni-CODH coupled to acetyl-CoA decarbonylase/synthase (ACDS) cleaves acetyl-CoA to CO₂ and methane [9].

This article focuses on the anaerobic Ni-CODHs, which have been divided into four classes in terms of the metabolic role and subunit components of the enzymes, as depicted, cartoonwise, in Figure 1 [10]. The classification can be confusing, as Roman numerals are used to denote both class and isozymes within a class, but hopefully this will not present a problem. Class I enzymes consist of five subunits (αβγδε) and are found in autotrophic methanogens, where acetyl-CoA is synthesized from CO₂ and H₂. Class II enzymes are found in acetoclastic methanogens and catalyze the decarbonylation of acetyl-CoA to CO₂, coenzyme A (CoA) and a methyl group which is transferred to tetrahydrosarcinapterin. Class III enzymes are found in homoacetogens and are sub-classified as two independent enzymes, a α₂β₂ tetramer (CODH/ACS) and a γδ heterodimer (CoFeSP): the catalytic center for synthesizing acetyl-CoA in tetrameric CODH/ACS is the same as found in Class I enzymes. Finally, Class IV enzymes consist only of the CODH entity, a monofunctional α₂ dimer that catalyzes the conversion of CO to CO₂ in the anaerobic CO-dependent energy metabolism of bacteria and archaea. The structure of a Class IV CODH is shown in Figure 2, along with the structure of a Class III CODH/ACS.

Important examples of all five Ni-CODHs are provided by the thermophilic bacterium *Carboxydotherrmus hydrogenoformans* (*Ch*) which can grow on CO as the sole carbon and energy source [11]. The first of these, referred to as CODH I_{Ch} (the Roman letter does not refer to the classification mentioned above) is involved in energy conversion and delivers electrons derived from CO oxidation to a hydrogenase which evolves H₂ [12]. In

Rhodospirillum rubrum, an enzyme very closely related to CODH I_{Ch} forms part of a well-characterized CODH-hydrogenase complex [13]. In contrast, the role of CODH II_{Ch} remains unclear despite this enzyme being the most structurally well characterized by x-ray crystallography [8, 14, 15]. The sequence identity and similarity between CODH I_{Ch} and CODH II_{Ch} are 58.3% and 73.9%, respectively. The third member, CODH III_{Ch}, is complexed tightly with ACS as CODH/ACS, as mentioned above, and catalyzes acetyl-CoA synthesis [16]. Of the remaining examples, CODH IV_{Ch} is suggested to be associated with a multisubunit enzyme complex for oxidative stress response based on genomic analysis, while the biological role of CODH V_{Ch} remains unclear.

The ease by which these enzymes interconvert CO₂ and CO has attracted intense interest from both chemists and biochemists, who have applied a variety of spectroscopic and structural methods in efforts to establish a firm mechanistic understanding. The aim of this chapter is to describe how the application of protein film electrochemistry (PFE) has added to this understanding[17–19]. But first we will summarize some of the structural and spectroscopic information that has been available now for several years.

2. Structures of Ni-containing carbon monoxide dehydrogenases

Several crystal structures of NiFe-containing CODH (Class IV) or CODH/ACS (Class III) from different organisms have been solved [4, 8, 15, 20–22]. All Class IV enzymes have a dimeric structure (shown in Figure 2) in which each monomer contains a unique active site (called the C-cluster) which is a [Ni₄Fe-4S] (or [NiFe-5S]) cubane cluster linked to an extra-cuboidal pendant or ‘dangling’ Fe. A [4Fe-4S] cluster (called the B-cluster) is located about 10 Å from each C-cluster, although these are coordinated by the other subunit. Finally, a single [4Fe-4S] cluster (called the D-cluster) lying about 10 Å from each B-cluster and close to the protein surface, is coordinated by both subunits. The distances suggest immediately that the B-cluster and D-cluster convey electrons between the C-cluster and an external physiological redox partner, which is probably a ferredoxin.

Several redox states of the C-cluster involved in the mechanism of CO/CO₂ interconversion by CODH have been identified [23]. These are known as C_{ox}, C_{red1}, C_{int}, and C_{red2} in order of decreasing oxidation level. Various structures of the active site of CODH II_{Ch} are shown in Figure 3. Those shown in Figure 3a and 3b were obtained at two different reduction potentials (–320 mV, and –600 mV with CO₂ present) [8]. The –320 mV structure shows that an O-donor (it is assumed this is hydroxide) binds to the pendant Fe atom and the Ni atom is coordinated by three sulfido ligands from the [3Fe-4S] core with a distorted T-shaped coordination geometry. The structure (3b) of CO₂-bound CODH II_{Ch} obtained by incubating CODH II_{Ch} with NaHCO₃ at –600mV reveals further that the C-atom from CO₂ binds to the Ni-atom at a distance of 1.96 Å, completing a distorted square-planar geometry, and one of the O-atoms of the bound CO₂ binds to the pendant Fe at a distance of 2.05 Å. These results suggest a mechanism in which CO oxidation involves nucleophilic attack on CO-bound Ni by a OH[–] ligand that is coordinated to the pendant Fe, thereby yielding the Ni-(CO(OH)) intermediate that is detected. In the reverse reaction, the pendant Fe atom abstracts an O-atom from the C-coordinated CO₂ via a proton-coupled two-electron process, and leaves CO bound to Ni with OH[–] on the pendant Fe. This picture of the mechanism is

included in Scheme 1 [4]. Spectroscopic data add further structural definition to this description, while PFE data discussed below how the reactions with substrates and inhibitors depend on potential.

From potentiometric redox titrations, it is known that the inactive and EPR-silent state, C_{ox} , is reduced by one electron to give the active state C_{red1} , which exhibits a characteristic EPR signal, $g_{av} \sim 1.82$ ($g = 2.01, 1.81, 1.65$) [23]. Based on the structural evidence described above, CO binds to the Ni atom in the C_{red1} state to start the catalytic cycle for CO oxidation, the eventual result of which is that CO_2 is released leaving the two-electron reduced state C_{red2} . The C-cluster is re-reduced back to C_{red1} by two one-electron transfers, via an EPR-silent state known as C_{int} . It is very significant that C_{red2} displays an EPR signal with $g_{av} \sim 1.86$ ($g = 1.97, 1.87, 1.75$) [23]. Based on earlier Mössbauer spectroscopy data, Lindahl suggested that C_{ox} should be assigned as $[Ni^{2+}Fe_p^{3+}]:[3Fe-4S]^1 - (Fe_p = \text{pendant Fe site})$ and C_{red1} should be assigned as $[Ni^{2+}Fe_p^{2+}]:[3Fe-4S]^1 -$ [24]. The fact that C_{red1} and C_{red2} differ by two electrons but both show an EPR spectrum with $g_{av} < 2$ suggests that the underlying electronic structure of the $[3Fe-4S]$ core fragment is unchanged between the two states. At least one electron must therefore be transferred at the Ni subsite, since it is unclear how a two-electron change could be accommodated at the pendant Fe that is also coordinated by sulphide, histidine-N and cysteine-S. The question then arises: what formal redox changes actually occur at Ni? If the pendant Fe remains as Fe^{2+} , both electrons must be transferred at the Ni subsite. Two alternative descriptions of the Ni atom in the C_{red2} state have been suggested: Ni(0) or the isoelectronic protonated site formulated as a nickel hydrido species Ni(II)-H [8, 25]. Further light on this question stems from Ni K- or L-edge X-ray absorption spectroscopy (XAS) studies on CODH $_{Rr}$ and CODH II $_{Ch}$ in differing redox states [26–28]. Results from as-isolated CODH II $_{Ch}$, CO-treated CODH II $_{Ch}$, dithionite-reduced CODH II $_{Ch}$, indigo-carmin-oxidized CODH $_{Rr}$, dithionite-reduced CODH II $_{Rr}$ and CO-treated CODH $_{Rr}$, all suggest that the Ni subsite in the C-cluster in these samples is present as Ni(II) regardless of redox state.

The reduction potential for the two-electron interconversion between C_{red1} and C_{red2} is approximately -520mV according to EPR potentiometric titrations [29]. The reduced B-cluster exhibits the typical EPR signal of a $[4Fe-4S]^{1+}$ cluster ($g_{av} \sim 1.94$) and the reduction potential for the $[4Fe-4S]^{2+/1+}$ couple is -440 mV in the enzyme from *Moorella thermoacetica* and -418 mV in the enzyme from *Rhodospirillum rubrum* [29, 30]. Significantly, the D-cluster remained undetected until the first crystal structure of CODH was solved. One possibility is that this cluster had escaped detection because the reduced $[4Fe-4S]^{1+}$ state has $S > 1/2$, but studies of CODH $_{Rr}$ by MCD and resonance Raman suggest that the D-cluster remains in the oxidized $[4Fe-4S]^{2+}$ state even at -520 mV [31]. This information will be important when we consider the results from PFE experiments later.

Cyanide, isoelectronic with CO, is a well-studied inhibitor of CODH [32–34]. Several CODH crystal structures from different species with cyanide or CO bound to the Ni atom in the C-cluster have been determined: that of CN-CODH I $_{Ch}$ is included in Figure 3 [14]. A consensus is not achieved: the CN-bound CODH/ACS $_{Mt}$ and CO(formyl)-bound ACDS/CODH $_{Mb}$ reveal, respectively, a *bent* NC-Ni structure with bond angle $\sim 114^\circ$ and a *bent* OC-Ni structure with bond angle $\sim 107^\circ$ [20, 35]. The OH^- ligand remains on the pendant Fe

in both structures. Compared with the structure of CO₂-bound CODH II_{Ch}, the N-atom in CN-bound CODH/ACS_{Mt} and the O-atom in CO-bound ACDS/CODH_{Mb} each overlay with the corresponding O-atom from CO₂ in CO₂-bound CODH II_{Ch} whereas the C-atom in both CN-bound ACS/CODH_{Mt} and CO-bound ACDS/CODH_{Mb} is displaced from its position in CO₂-bound CODH II_{Ch} [8]. The shift in C-atom position could facilitate the nucleophilic attack by OH[−]. Interestingly, the crystal structure of CODH II_{Ch} incubated with *n*-butyl isocyanide at −320mV reveals that the C-atom from the plausible product “*n*-butyl-isocyanate” binds to the Ni with a distorted tetrahedral coordination geometry [36]. A second *n*-butyl-isocyanide is found in the putative gas channel in CODH II_{Ch}.

Introduction of cyanide to a sample of CODH/ACS_{Mt} poised in the C_{red1} state results in a new EPR signal with $g_{av} = 1.72$ ($g = 1.87, 1.78, 1.55$) [23, 33, 34, 37]. In contrast, no change in the spectrum is observed when cyanide is introduced to the C_{red2} state [37]. Further studies of ¹³CN-bound CODH/ACS_{Mt} by ENDOR revealed a doublet peak in the C_{red1} state - early evidence that cyanide (¹³C has $I = 1/2$) binds directly to the C-cluster [38, 39]. Experiments to evaluate if NCO[−] is an inhibitor indicated that it binds also to C_{red1} even (since it is an analogue of CO₂ not CO) it should display behaviour opposite to that of CN[−]: this result was puzzling, but as we explain below, it was clarified by PFE experiments.

Some crystal structures of CODH II_{Ch} revealed a sulfide bridge between the Ni atom and the pendant Fe atom, as shown in Figure 3d, and this observation led to suggestions that an additional (5th) sulfide is necessary for catalysis [15, 40]. In place of inorganic sulfide, the S-atom from cys531 was found to bridge the Ni atom and the pendant Fe atom in the crystal structure of CODH_{Rr} [21]. However, crystal structures from recombinant CODH II expressed in *E. coli* and other CODH crystal structures from *Moorella thermoacetica* and *Methanosarcina barkeri* revealed no presence of a 5th sulfido ligand [20, 35]. The CO oxidation activities of CODH measured by solution assays in the presence of sodium sulfide showed varying results between different species and redox conditions [33, 41]. The role of sulfide is considered later when we discuss the PFE results.

Scheme 1 highlights a challenge for investigating enzymes such as CODHs that catalyse redox reactions via rapid passage through a series of intermediates, the presence (lifetime) of each of which depends on how fast or favourable are the electron transfers that determine their existence. Direct coordination of a reactant or inhibitor, or release of a product, should be selective for a particular oxidation state: the question is how to control and fine-tune the relative amounts of each state during catalysis and simultaneously observe the effect on rate when reactant or inhibitor are introduced. Not all inhibitors bind to active states: some exert their influence by facilitating redox processes that lead to inactive states. Here, an immediate problem is posed by the fact that C_{ox} is EPR-silent, thus denying us a substantial amount of spectroscopic information; the same is true for C_{int}, which is particularly elusive.

3. Protein Film Electrochemistry

Protein film electrochemistry (PFE) refers to a suite of dynamic electrochemical techniques that address enzyme molecules directly attached onto an electrode surface, allowing catalytic activity to be recorded (as current) as a continuous function of electrode potential

and/or time [42–45]. The importance of steady-state activity relating directly to steady-state current at any given potential is a unique aspect of PFE. The concept is depicted (idealistically) in Figure 4. The electrode provides a wide and continuous range of potential that is very difficult to achieve using conventional methods; for example catalytic CO₂ reduction by CODH requires potentials more negative than –0.5 V at pH=7.0, and it is difficult to obtain such a negative potential by chemical mediators or titrants, let alone provide a continuous scale. A pyrolytic graphite ‘edge’ (PGE) electrode is particularly suitable: it is easily polished by abrasion (an aqueous alumina slurry) to expose a pristine surface and it has a very wide potential region, extending from approximately +1 V to –1 V vs the standard hydrogen electrode (SHE) at neutral pH.

Many enzymes are now established to be excellent electrocatalysts – defined here as catalysts for half-cell reactions that occur at an electrode [46, 47]. Enzymes are typically bound at the electrode by simple adsorption (exploiting a rough electrode surface that contacts the enzyme at different points and maximizes hydrophobic and electrostatic interactions) or by chemical attachment that may involve engineering the enzyme to place specific residues in suitable positions for linkage formation. It is essential that the enzyme contacts the electrode surface in such a way that electron relay centers within the enzyme, such as the FeS clusters in CODHs, lie within a sufficiently short distance to allow fast electron tunnelling. For class IV CODHs, the obvious entry and exit site is the exposed D-cluster, as emphasised in Figure 2.

Dynamic electrochemical techniques include cyclic voltammetry (the basic search tool, in which current is monitored as the potential is scanned) and chronoamperometry (initiating a reaction at a fixed electrode potential, either by stepping to that potential or injecting a reactant, and monitoring the current as a function of time). Because a change in current corresponds directly to a change in activity, chronoamperometry is an excellent technique for measuring how fast inhibitors are bound or released, all under strictly controlled potential conditions.

In terms of operational advantage, PFE normally needs a relatively tiny amount of enzyme sample, about pmole/cm² – orders of magnitude smaller than spectroscopy. Because the enzyme is immobilized on the electrode (area typically 0.03 cm²) the same sample can be successively exposed to solutions having different pH values and inhibitor/substrate concentrations. For studying air-sensitive enzymes such as Ni-CODH, the sealed electrochemical cell is housed in a glove box. The electrode is normally rotated at variable high speed, which in addition to assisting the transport of reactants to the enzyme, ensures that dispersion of product can be controlled – the latter factor allowing product inhibition to be investigated easily. If the substrate is a gas, this is introduced into the headspace of the electrochemical cell and it is easy to replace substrates in the solution by purging with inert gas.

Cyclic voltammetry provides a direct, continuous ‘spectrum’ of the potential dependence. Importantly, when the enzyme catalysis approaches electrochemical reversibility, the trace cuts across the potential axis at the equilibrium value and just a tiny potential bias either side of this value switches the reaction between reduction and oxidation. Such reversibility is

found with many enzymes. The ratio of oxidation and reduction currents measured at appropriate potential values yields the so-called ‘catalytic bias’ – the tendency of the enzyme to operate preferentially in a particular direction [47]. Overlay of scans in each direction shows that electrocatalysis is at steady state, whereas hysteresis is a sign that the enzyme alters its activity on a timescale that is slow relative to the rate that the potential is scanned.

Fact that the catalytic activity is observed as current and measured across a range of potentials means that each enzyme produces a spectrum-like fingerprint of its characteristics, but PFE rarely allows determination of absolute steady-state catalytic rates because the electroactive coverage is generally unknown. In order to determine the electroactive coverage of an enzyme it is necessary to detect so-called ‘non-turnover’ signals due to stoichiometric electron exchanges with relay centers such as Fe-S clusters. These signals are peak-like in each scan direction and the electroactive coverage is estimated from the areas under the peaks. Turnover frequency k_{cat} is then calculated from the current (i) using the relationship

$$i = k_{\text{cat}} n F A \Gamma \quad (3)$$

where n is the number of electrons transferred, F is the Faraday constant, A is the electrode area and Γ is the electroactive coverage. As a guide, it is considered very difficult to observe a non-turnover signal if Γ is below 10^{-12} moles cm^{-2} . Consequently, with $A = 0.03 \text{ cm}^2$, a current of $10 \text{ }\mu\text{A}$ corresponds to a lower limit for k_{cat} in the region of 2000 s^{-1} .

The difficulty in determining absolute steady-state activities contrasts with the ease by which transient reaction rates are measured at a constant potential. The rate of a change in activity is measured from the time course of the current following a perturbation such as potential step or injection of reagent. The amplitude of the change is proportional to catalytic activity and electroactive coverage, but the rate should be independent of these variables. The chronoamperometry experiment examines, directly, the rate of change of rate – a measurement that is equivalent to recording the second derivative of a normal kinetic plot of concentration vs time.

4. Carbon monoxide dehydrogenases as electrocatalysts

Based upon the structure of CODH II_{Ch} , it is clear from Figure 2 that the D-cluster should be a feasible entry/exit point for electrons, and the enzyme should be electroactive when attached to a rough electrode surface at which productive orientations are statistically likely.

4.1 The electrocatalytic voltammograms of Class IV Enzymes

As shown in Figure 5A, a cyclic voltammogram of CODH I_{Ch} adsorbed on a PGE electrode displays reversible electrocatalysis: under a gas mixture of composition 50% $\text{CO}/50\% \text{CO}_2$, the trace cuts through the potential axis at the expected equilibrium potential and a detectable net current in each direction is easily achieved with a minimal overpotential [17, 18]. Under the same conditions, CODH II_{Ch} shows very little CO_2 reduction, but a much larger reduction current is observed if the atmosphere is changed to 100% CO_2 , showing that

the marked bias toward CO oxidation displayed by CODH II_{Ch} is due, at least in part, to CO being a much stronger product inhibitor of CO₂ reduction for that isozyme [19]. Features of the complex wavelshape are explained in terms of a model that takes into consideration the potential at which electrons enter and leave the enzyme (via the D-cluster) and potential-dependent interconversions between different catalytic states [47]. The potential of the center that serves as the electron entry/exit site is an important determinant of catalytic bias, and maximum reversibility is obtained when this value matches that of the reaction being catalysed. For CODH I_{Ch} and CODH II_{Ch} where the obvious electron entry/exit site is the D cluster, it is noteworthy that the reduction potential of the D-cluster has not been determined. However, a rationale for the observation of reversible electrocatalysis in terms of the model shows that this value must lie close to -0.5 V.

At high potentials, both CODH I_{Ch} and CODH II_{Ch} become inactivated as the active site transforms into the C_{ox} state: this is seen particularly clearly for CODH II_{Ch} which shows a re-activation 'peak' at approximately -0.2 V [19]. Generally, inactivation is slow and its rate is independent of potential (consistent with the rate being determined by a chemical process rather than electron transfer) whereas re-activation is fast and controlled by electron transfer, hence the sharp transition.

Before proceeding further, it is worthwhile drawing comparisons with conventional methods of measuring activity. The standard procedure for determining the activity of CO oxidation by Ni-Fe CODH is to measure the color change that occurs when colorless oxidized methylviologen (MV²⁺) is reduced to the blue reduced MV⁺ radical by electron transfer from CO-reduced CODH [48–51]. Measuring CO₂ reduction is much more problematic because it is necessary to use a fast electron donor that is more reducing than CO: it is obviously difficult to generate and stabilize such a donor in aqueous solution. Both $V_{\max}$ and K_M are sensitive to the driving force (electrode potential) and this fact is easy to see from cyclic voltammograms measured for a range of substrate concentrations [52].

4.2 The electrocatalytic voltammograms of Class III enzymes (CODH/ACS)

Figure 5B shows the same kind of experiment carried out with a film of CODH/ACS_{Mt} on a PGE electrode. This enzyme is much larger than CODH I or II, the two CODH subunits being flanked by two ACS subunits, as shown in Figure 2. In contrast to the smaller enzymes, the cyclic voltammogram of CODH/ACS_{Mt} shows no CO₂ reduction, even when no CO is present (100 % CO₂); instead the scan reveals a CO oxidation current that increases slowly as the potential is raised, indicative of sluggish electron transfer. The CO oxidation current at 0 V is about two orders of magnitude smaller than that typically observed for CODH I_{Ch} and CODH II_{Ch} under similar conditions. The sluggish electron transfer may reflect the much larger size of CODH/ACS_{Mt} as the flanking ACS subunits could prevent the D-cluster from achieving such a close approach to the electrode surface. What is not clear though is why CO₂ reduction is not observed, when it is fully expected in view of the physiological function of the enzyme. One possibility is that electrons cannot use the D-cluster which may become obstructed or held too far from the electrode, but instead enter the catalytic cycle at a higher potential than with CODH I_{Ch} or CODH II_{Ch}.

The reducing power is thus dissipated and electron transfer to the C-cluster becomes an unfavourable step.

To identify whether ACS influences the performance of CODH, an experiment was performed in which several chemical reagents, such as sodium dodecyl sulfate (which separates CODH and ACS partially), 1, 10-phenanthroline, (which inhibits the active site in ACS) and acetyl-CoA (the product of the reaction carried out by CODH/ACS_{Mt}) were added [53–55]. However, an electrocatalytic current due to CO₂ reduction was still not observed, so this observation remains a puzzle. The turnover frequency for CO₂ reduction by CODH/ACS_{Mt} is normally determined by measuring the CO product binding to haemoglobin and a rate constant of approximately 1.3 s⁻¹ was reported [56]. However, several studies have led to the picture that during synthesis of acetyl-CoA, CO arising from reduction of CO₂ diffuses through a channel in the enzyme complex to reach the A-cluster without escaping [57, 58].

5. Potential-dependent reactions with inhibitors

The catalytic current provides an important observable (commonly called a ‘handle’) for investigating the kinetics and potential dependence of the reactions with inhibitors, at a level of kinetic detail that is difficult to achieve by conventional methods. Some of the opportunities for inhibitors to bind are shown in Scheme 1 and elaborated upon (in an electrochemical sense) in Scheme 2. Protein film electrochemistry is able to reveal and clarify the different ways that these small molecules interrupt catalysis. We will discuss next how cyanide, isoelectronic with CO, mainly inhibits CO oxidation, whereas cyanate, isoelectronic with CO₂, targets CO₂ reduction. Sulfide is also an inhibitor but is unusual because it acts only under oxidizing conditions. Thiocyanate and azide are also considered.

5.1 How Class IV CODHs respond to cyanide

Qualitative details of the enzyme’s inhibition by cyanide are revealed in the cyclic voltammograms shown in Figure 6, which were recorded for CODH I_{Ch} and CODH I_{Ch} [18, 19]. A complication here is that cyanide evaporates quite rapidly from the electrochemical cell as HCN, but key characteristics are easily seen and extracted. The important observations are summarized as follows. When CN⁻ is injected at -0.3 V, during CO oxidation, the current drops: the reaction is more rapid for CODH I. Injecting CN⁻ instead at the limit of very negative potential, at which CO₂ reduction is observed, then scanning in a positive direction, CN⁻ binds once the electrode potential becomes more positive than -750 mV, resulting in complete inhibition of CO₂ reduction (within a narrow potential window). Continuing the scan to more positive potential, it is clear that CO oxidation is completely blocked. The activity remains at zero during the remaining cycle, until the electrode potential passes down through -0.6 V, at which point the CO₂ reduction activity starts to increase strongly. The rapid release of cyanide when the electrode potential is taken below -0.6 V is an observation that is unique to PFE, because conventional kinetic assays rarely explore such highly reducing conditions and lack the ability to record the activity simultaneously as the potential is scanned.

The data are interpreted in terms of the differing dominance of the two major redox states of the C-cluster, C_{red1} and C_{red2} as the potential is varied. At potentials above -0.6 V, C_{red1} dominates and CN^- is tightly bound, blocking catalysis. At electrode potentials below -0.6 V, C_{red2} becomes the dominant catalytic species, and recovery of activity that occurs during the scan in this very negative potential region reveals that CN^- is released. During the experiment, which is carried out at a very slow scan rate, the cyanide is lost as HCN which is removed in the gas flow; hence the effect diminishes with time. The preferential binding to C_{red1} perfectly complements the EPR data which showed that only the spectrum of this state is affected by CN^- [37]. The kinetics of binding and release of CN^- are measured quantitatively by chronoamperometric experiments, in which the time dependence of the increase or decrease in current is recorded at a constant potential. Some exemplary experiments are shown in Figure 7. The top panels show the rates at which CN^- reacts to inhibit CO oxidation and CO₂ reduction by CODH II_{Ch}. The lower panels show a more complicated series of injection and potential step, to compare the rates of reductive reactivation of CODH I_{Ch} and CODH II_{Ch} upon stepping from $+0.14$ V to -0.76 V. Reductive release of CN^- from CODH II_{Ch} is much faster and occurs within the step time.

5.2 How class IV CODHs respond to cyanate

Inhibition by cyanate (NCO^-), an analogue of CO₂, shows the opposite trend to that observed for CN^- : it binds tightly at potentials below -0.5 V but is released at higher potentials (Figure 8) [18, 19]. The effect is such that CODH functions as a unidirectional catalyst in the presence of cyanate, and CO oxidation is virtually unaffected apart from the requirement for a small overpotential before the activity climbs. The sharp potential dependence shows that NCO^- reacts selectively with C_{red2} , the small overpotential reflecting the additional energy required to dislodge the inhibitor by oxidation.

The binding and release of CN^- and NCO^- are orders of magnitude slower than the catalytic turnover frequencies of their substrate counterparts CO and CO₂, thus showing that these analogues, while having very similar shapes and size, fall far short of being really good analogues [19]. The differences must ultimately relate to acid-base (proton-transfer) properties, overall electrostatic effects and subtle effects of frontier orbitals. Comparisons between CODH I_{Ch} and CODH II_{Ch} show that rates of both inhibition by CN^- and reactivation are higher for CODH II_{Ch} even though it binds CN^- more tightly in a thermodynamic sense, mirroring the stronger binding of CO that makes it such a potent product inhibitor of this isozyme [19].

5.3 How class IV CODHs respond to sulfide and thiocyanate

Sulfide inhibits CODH_{Ch} rapidly when the potential is more positive than -50 mV but no reaction is observed at more negative potentials, at which CO oxidation occurs unaffected by its presence [18]. This observation is shown in Figure 9, which shows that sulfide is not a competitive inhibitor like CN^- , but instead reacts to stabilise an inactive state at a higher oxidation level. This state must be analogous to but not identical with C_{ox} because the reactivation potential of the sulfido product is much more negative. It is therefore unlikely that sulfide can be an activator of CODH catalysis, as was suggested by earlier work. An interesting observation is made when thiocyanate, NCS^- , is introduced. In marked contrast

to NCO^- , NCS^- does not inhibit CO_2 reduction; instead there is partial inhibition of CO oxidation and an oxidized inactive state is formed that does not activate at the same potential as C_{ox} , but at a more negative potential, similar to that observed when sulfide is present. One possibility is that NCS^- also reacts to stabilise an inactive oxidised state by leaving a bridging S, either as sulfide or an intact S-bound NCS^- .

The main differences between CODH I_{Ch} and CODH II_{Ch} are the stronger inhibition of CODH II_{Ch} by CO product and cyanide [19]. These discoveries may shed light on the possible role of CODH II_{Ch} in biological systems. Interestingly, sulfide is a similar inhibitor of [NiFe]-hydrogenases, binding to and stabilizing the inactive Ni(III) form rather than acting competitively in the active potential region. One aspect here is that sulfide is a good bridging ligand, which CN^- is not, so it can form the Ni-S-Fe linkage that is observed in some crystal structures.

Figure 10 summarises the ‘redox spectrum’ for activities and inhibition of CODH I_{Ch} .

6. Demonstrations of technological significance

The high electrocatalytic activities of Class IV CODHs match those observed with many hydrogenases – a fact that inspired an aesthetically interesting experiment to mimic the physiological process in which CO oxidation is coupled to H_2 evolution [59]. This process is the biological counterpart of the industrially important ‘water gas shift’ (WGS) reaction that requires high temperatures. Notably, CO as a component of syngas, can be used to synthesis liquid hydrocarbons by reaction with H_2 in the Fisher-Tropsch process [60]. In the enzyme coupled version, a hydrogenase (Hyd-2) from *Escherichia coli* and CODH I_{Ch} are both adsorbed onto conducting graphite platelets, formed by grinding a piece of pyrolytic graphite. In microorganisms, CO-reduced CODH I_{Ch} produces electrons that are transferred one at a time via a ferredoxin to a membrane-bound hydrogenase for H_2 evolution [48]. In the system of two enzymes on the graphite platelets, CO-reduced CODH releases electrons to Hyd-2 for H_2 evolution via conduction across the graphite platelets. The turnover frequency is 2.5 sec^{-1} at 30°C , which is comparable to catalysts operated at high temperatures in industry. Results are shown in Figure 11.

The development of efficient photo-catalysts for CO_2 reduction plays an important role in converting CO_2 to carbon-based fuel in a clean and sustainable way. Experiments have been carried out in which CODH I_{Ch} is attached to semiconductor nanoparticles, such as TiO_2 and CdS, to study the visible light-driven reduction of CO_2 to CO as illustrated in Figure 12 [61, 62]. For TiO_2 (anatase) the bandgap is 3.1 eV which lies in the UV region, so a Ru-photosensitizer was co-attached to the nanoparticle along with CODH I_{Ch} . Upon illumination, the turnover frequency for CO production is 0.14 s^{-1} at pH 6 and 30°C . In the CdS system, a photosensitizer is not necessary because the band gap lies in the visible region. For the hybrid CODH-CdS nanorod system, the turnover frequency for CO production is 1.23 s^{-1} [62].

We have already discussed how CODH I_{Ch} and CODH II_{Ch} attached to graphite electrodes show high activities of CO oxidation and CO_2 reduction with a minimal overpotential [17]. The catalytic bias is largely determined by the fact that the reduction potential of the D-

cluster lies in the region of the potential for the CO₂/CO redox couple [47]. When TiO₂ or CdS are used as electrode materials, the electrocatalytic activity is biased instead strongly in favour of CO₂ reduction, an observation that is explained by the potential dependence of electron availability in the material. Semiconductors have a low carrier density and, in simplistic physics terms, the availability of electrons for transfer into a catalyst is governed by the position of the flatband potential E_{FB} . Both TiO₂ and CdS are *n*-type semiconductors and a potential more negative than E_{FB} is required for electrons to become concentrated close to the surface and available for transfer. Above E_{FB} no current flows. Results comparing CODH I_{Ch} attached to PGE, CdS thin film and TiO₂ thin film electrodes are shown in Figure 13. In contrast to the result observed for PGE, only CO₂ reduction is observed at CdS and TiO₂. The experiments demonstrate how to rectify reversible electrocatalysts for the fuel-forming direction since the most efficient catalysts for solar fuel production should operate close to reversible potentials [63].

7. Conclusions

The linear CO₂ molecule is difficult to reduce, but enzymes provide a superb example of how this chemistry has been perfected by evolution. The active site of CODH reveals simple tricks that could be mimicked by chemists, although it is going to be challenging to synthesise the precise, outer-shell environment that nature so easily rolls out during transcription. Protein film electrochemistry has proved to be a valuable tool for separating the different reactions as a smooth function of potential, as it has been for hydrogenases. The PFE results have yielded important information – in the ‘potential domain’ – to complement and clarify the picture built up from structure and spectroscopy.

References

1. Ragsdale SW, Pierce E. Acetogenesis and the Wood-Ljungdahl pathway of CO₂ fixation. *Biochimica Et Biophysica Acta-Proteins and Proteomics*. 2008; 1784(12):1873–1898.
2. Ferry JG. How to Make a Living by Exhaling Methane. *Annual Review of Microbiology*, Vol 64, 2010. 2010; 64:453–473.
3. Oelgeschlaeger E, Rother M. Carbon monoxide-dependent energy metabolism in anaerobic bacteria and archaea. *Archives of Microbiology*. 2008; 190(3):257–269. [PubMed: 18575848]
4. Can M, Armstrong FA, Ragsdale SW. Structure, Function, and Mechanism of the Nickel Metalloenzymes, CO Dehydrogenase, and Acetyl-CoA Synthase. *Chemical Reviews*. 2014
5. Meyer O, Schlegel HG. BIOLOGY OF AEROBIC CARBON-MONOXIDE OXIDIZING BACTERIA. *Annual Review of Microbiology*. 1983; 37:277–310.
6. Bender G, et al. Metal centers in the anaerobic microbial metabolism of CO and CO₂. *Metallomics*. 2011; 3(8):797–815. [PubMed: 21647480]
7. Dobbek H, et al. Catalysis at a dinuclear CuSMo(=O)OH cluster in a CO dehydrogenase resolved at 1.1-angstrom resolution. *Proceedings of the National Academy of Sciences of the United States of America*. 2002; 99(25):15971–15976. [PubMed: 12475995]
8. Jeoung JH, Dobbek H. Carbon dioxide activation at the Ni,Fe-cluster of anaerobic carbon monoxide dehydrogenase. *Science*. 2007; 318:1461–1464. [PubMed: 18048691]
9. Grahame DA. CATALYSIS OF ACETYL-COA CLEAVAGE AND TETRAHYDROSARCINAPTERIN METHYLATION BY A CARBONMONOXIDE DEHYDROGENASE-CORRINOID ENZYME COMPLEX. *Journal of Biological Chemistry*. 1991; 266(33):22227–22233. [PubMed: 1939246]

10. Lindahl PA, Chang B. The evolution of acetyl-CoA synthase. *Origins of Life and Evolution of the Biosphere*. 2001; 31(4–5):403–434. [PubMed: 11599178]
11. Ragsdale SW. Life with carbon monoxide. *Critical Reviews in Biochemistry and Molecular Biology*. 2004; 39(3):165–195. [PubMed: 15596550]
12. Soboh B, Linder D, Hedderich R. Purification and catalytic properties of a CO-oxidizing : H₂-evolving enzyme complex from *Carboxydothermus hydrogenoformans*. *European Journal of Biochemistry*. 2002; 269(22):5712–5721. [PubMed: 12423371]
13. Fox JD, et al. Characterization of the region encoding the CO-induced hydrogenase of *Rhodospirillum rubrum*. *Journal of Bacteriology*. 1996; 178(21):6200–6208. [PubMed: 8892819]
14. Jeoung JH, Dobbek H. Structural Basis of Cyanide Inhibition of Ni, Fe-Containing Carbon Monoxide Dehydrogenase. *Journal of the American Chemical Society*. 2009; 131(29):9922–9923. [PubMed: 19583208]
15. Dobbek H, et al. Carbon monoxide induced decomposition of the active site Ni-4Fe-5S cluster of CO dehydrogenase. *Journal of the American Chemical Society*. 2004; 126(17):5382–5387. [PubMed: 15113209]
16. Svetlitchnyi V, et al. A functional Ni-Ni- 4Fe-4S cluster in the monomeric acetyl-CoA synthase from *Carboxydothermus hydrogenoformans*. *Proceedings of the National Academy of Sciences of the United States of America*. 2004; 101(2):446–451. [PubMed: 14699043]
17. Parkin A, et al. Rapid and efficient electrocatalytic CO₂/CO interconversions by *Carboxydothermus hydrogenoformans* CO dehydrogenase I on an electrode. *Journal of the American Chemical Society*. 2007; 129(34):10328–10329. [PubMed: 17672466]
18. Wang VCC, et al. A Unified Electrocatalytic Description of the Action of Inhibitors of Nickel Carbon Monoxide Dehydrogenase. *Journal of the American Chemical Society*. 2013; 135(6): 2198–2206. [PubMed: 23368960]
19. Wang VCC, Ragsdale SW, Armstrong FA. Investigations of Two Bidirectional Carbon Monoxide Dehydrogenases from *Carboxydothermus hydrogenoformans* by Protein Film Electrochemistry. *ChemBioChem*. 2013; 14(14):1845–1851. [PubMed: 24002936]
20. Kung Y, et al. Crystallographic Snapshots of Cyanide- and Water-Bound C-Clusters from Bifunctional Carbon Monoxide Dehydrogenase/Acetyl-CoA Synthase. *Biochemistry*. 2009; 48(31):7432–7440. [PubMed: 19583207]
21. Drennan CL, et al. Life on carbon monoxide: X-ray structure of *Rhodospirillum rubrum* Ni-Fe-S carbon monoxide dehydrogenase. *Proceedings of the National Academy of Sciences of the United States of America*. 2001; 98(21):11973–11978. [PubMed: 11593006]
22. Darnault C, et al. Ni-Zn- Fe-4-S-4 and Ni-Ni- Fe-4-S-4 clusters in closed and open subunits of acetyl-CoA synthase/carbon monoxide dehydrogenase. *Nature Structural Biology*. 2003; 10(4): 271–279.
23. Lindahl PA. Implications of a carboxylate-bound C-cluster structure of carbon monoxide dehydrogenase. *Angewandte Chemie-International Edition*. 2008; 47(22):4054–4056.
24. Lindahl PA. The Ni-containing carbon monoxide dehydrogenase family: Light at the end of the tunnel? *Biochemistry*. 2002; 41(7):2097–2105. [PubMed: 11841199]
25. Amara P, et al. Carbon Monoxide Dehydrogenase Reaction Mechanism: A Likely Case of Abnormal CO₂ Insertion to a Ni-H- Bond. *Inorganic Chemistry*. 2011; 50(5):1868–1878. [PubMed: 21247090]
26. Gu WW, et al. CO-Induced structural rearrangement of the c cluster in *Carboxydothermus hydrogenoformans* CO dehydrogenase - Evidence from NiK-edge X-ray absorption spectroscopy. *Biochemistry*. 2004; 43(28):9029–9035. [PubMed: 15248760]
27. Tan GO, et al. On the Structure of the Nickel Iron Sulfur Center of the Carbon-Monoxide Dehydrogenase from *Rhodospirillum-Rubrum* - an X-Ray Absorption-Spectroscopy Study. *Proceedings of the National Academy of Sciences of the United States of America*. 1992; 89(10): 4427–4431. [PubMed: 1584775]
28. Ralston CY, et al. Characterization of heterogeneous nickel sites in CO dehydrogenases from *Clostridium thermoaceticum* and *Rhodospirillum rubrum* by nickel L-edge X-ray spectroscopy. *Journal of the American Chemical Society*. 2000; 122(43):10553–10560.

29. Lindahl PA, Munck E, Ragsdale SW. CO DEHYDROGENASE FROM CLOSTRIDIUM-THERMOACETICUM - EPR AND ELECTROCHEMICAL STUDIES IN CO₂ AND ARGON ATMOSPHERES. *Journal of Biological Chemistry*. 1990; 265(7):3873–3879. [PubMed: 2154491]
30. Spangler NJ, et al. Spectroelectrochemical characterization of the metal centers in carbon monoxide dehydrogenase (CODH) and nickel-deficient CODH from *Rhodospirillum rubrum*. *Journal of Biological Chemistry*. 1996; 271(14):7973–7977. [PubMed: 8626477]
31. Craft JL, Ludden PW, Brunold TC. Spectroscopic studies of nickel-deficient carbon monoxide dehydrogenase from *Rhodospirillum rubrum*: Nature of the iron-sulfur clusters. *Biochemistry*. 2002; 41(5):1681–1688. [PubMed: 11814363]
32. Seravalli J, et al. MECHANISM OF CO OXIDATION BY CARBONMONOXIDE DEHYDROGENASE FROM CLOSTRIDIUM-THERMOACETICUM AND ITS INHIBITION BY ANIONS. *Biochemistry*. 1995; 34(24):7879–7888. [PubMed: 7794899]
33. Ha SW, et al. Interaction of potassium cyanide with the Ni-4Fe-5S active site cluster of CO dehydrogenase from *Carboxydotherrmus hydrogenoformans*. *Journal of Biological Chemistry*. 2007; 282(14):10639–10646. [PubMed: 17277357]
34. Ensign SA, Hyman MR, Ludden PW. NICKEL-SPECIFIC, SLOW-BINDING INHIBITION OF CARBON-MONOXIDE DEHYDROGENASE FROM RHODOSPIRILLUM-RUBRUM BY CYANIDE. *Biochemistry*. 1989; 28(12):4973–4979. [PubMed: 2504285]
35. Gong W, et al. Structure of the alpha(2)epsilon(20) Ni-dependent CO dehydrogenase component of the *Methanosarcina barkeri* acetyl-CoA decarbonylase/synthase complex. *Proceedings of the National Academy of Sciences of the United States of America*. 2008; 105(28):9558–9563. [PubMed: 18621675]
36. Jeoung J-H, Dobbek H. n-Butyl isocyanide oxidation at the NiFe₄S₄OH_x cluster of CO dehydrogenase. *Journal of Biological Inorganic Chemistry*. 2012; 17(2):167–173. [PubMed: 21904889]
37. Anderson ME, Lindahl PA. ORGANIZATION OF CLUSTERS AND INTERNAL ELECTRON PATHWAYS IN CO DEHYDROGENASE FROM CLOSTRIDIUM-THERMOACETICUM - RELEVANCE TO THE MECHANISM OF CATALYSIS AND CYANIDE INHIBITION. *Biochemistry*. 1994; 33(29):8702–8711. [PubMed: 8038160]
38. Anderson ME, et al. IDENTIFICATION OF A CYANIDE BINDING-SITE IN CO DEHYDROGENASE FROM CLOSTRIDIUM-THERMOACETICUM USING EPR AND ENDOR SPECTROSCOPIES. *Journal of the American Chemical Society*. 1993; 115(25):12204–12205.
39. DeRose VJ, et al. A multinuclear ENDOR study of the C-cluster in CO dehydrogenase from *Clostridium thermoaceticum*: Evidence for H_xO and histidine coordination to the Fe₄S₄ center. *Journal of the American Chemical Society*. 1998; 120(34):8767–8776.
40. Dobbek H, et al. Crystal structure of a carbon monoxide dehydrogenase reveals a Ni-4Fe-5S cluster. *Science*. 2001; 293(5533):1281–1285. [PubMed: 11509720]
41. Feng J, Lindahl PA. Effect of sodium sulfide on Ni-containing carbon monoxide dehydrogenases. *Journal of the American Chemical Society*. 2004; 126(29):9094–9100. [PubMed: 15264843]
42. Vincent KA, Parkin A, Armstrong FA. Investigating and exploiting the electrocatalytic properties of hydrogenases. *Chemical Reviews*. 2007; 107:4366–4413. [PubMed: 17845060]
43. Leger C, Bertrand P. Direct electrochemistry of redox enzymes as a tool for mechanistic studies. *Chemical Reviews*. 2008; 108(7):2379–2438. [PubMed: 18620368]
44. Vincent KA, Armstrong FA. Investigating metalloenzyme reactions using electrochemical sweeps and steps: Fine control and measurements with reactants ranging from ions to gases. *Inorganic Chemistry*. 2005; 44(4):798–809. [PubMed: 15859247]
45. Leger C, et al. Enzyme electrokinetics: Using protein film voltammetry to investigate redox enzymes and their mechanisms. *Biochemistry*. 2003; 42(29):8653–8662. [PubMed: 12873124]
46. Armstrong FA, Hirst J. Reversibility and efficiency in electrocatalytic energy conversion and lessons from enzymes. *Proceedings of the National Academy of Sciences of the United States of America*. 2011; 108(34):14049–14054. [PubMed: 21844379]

47. Hexter SV, et al. Electrocatalytic mechanism of reversible hydrogen cycling by enzymes and distinctions between the major classes of hydrogenases. *Proceedings of the National Academy of Sciences of the United States of America*. 2012; 109(29):11516–11521. [PubMed: 22802675]
48. Svetlitchnyi V, et al. Two membrane-associated NiFeS-carbon monoxide dehydrogenases from the anaerobic carbon-monoxide-utilizing eubacterium *Carboxydotherrmus hydrogenoformans*. *Journal of Bacteriology*. 2001; 183(17):5134–5144. [PubMed: 11489867]
49. Ragsdale SW, et al. Properties of Purified Carbon Monoxide Dehydrogenase from *Clostridium thermoaceticum*, a Nickel, Iron-Sulfur Protein. *Journal of Biological Chemistry*. 1983; 258(4): 2364–2369. [PubMed: 6687389]
50. Shin WS, Lindahl PA. LOW-SPIN QUANTITATION OF NIFEC EPR SIGNAL FROM CARBON-MONOXIDE DEHYDROGENASE IS NOT DUE TO DAMAGE INCURRED DURING PROTEIN-PURIFICATION. *Biochimica Et Biophysica Acta*. 1993; 1161(2–3):317–322. [PubMed: 8381672]
51. Heo JY, et al. Evidence for a ligand CO that is required for catalytic activity of CO dehydrogenase from *Rhodospirillum rubrum*. *Biochemistry*. 2000; 39(27):7956–7963. [PubMed: 10891076]
52. Lukey MJ, et al. How *Escherichia coli* is equipped to oxidize hydrogen under different redox conditions (vol 285, pg 3928, 2010). *Journal of Biological Chemistry*. 2010; 285(26):20421–20421.
53. Xia JQ, et al. Carbon monoxide dehydrogenase from *Clostridium thermoaceticum*: Quaternary structure, stoichiometry of its SDS-induced dissociation, and characterization of the faster-migrating form. *Biochemistry*. 1996; 35(6):1965–1971. [PubMed: 8639680]
54. Shin W, Lindahl PA. DISCOVERY OF A LABILE NICKEL ION REQUIRED FOR CO ACETYL-COA EXCHANGE ACTIVITY IN THE NIFE COMPLEX OF CARBON-MONOXIDE DEHYDROGENASE FROM *CLOSTRIDIUM-THERMOACETICUM*. *Journal of the American Chemical Society*. 1992; 114(24):9718–9719.
55. Ragsdale SW, Wood HG. ACETATE BIOSYNTHESIS BY ACETOGENIC BACTERIA - EVIDENCE THAT CARBON-MONOXIDE DEHYDROGENASE IS THE CONDENSING ENZYME THAT CATALYZES THE FINAL STEPS OF THE SYNTHESIS. *Journal of Biological Chemistry*. 1985; 260(7):3970–3977. [PubMed: 2984190]
56. Kumar M, Lu WP, Ragsdale SW. BINDING OF CARBON-DISULFIDE TO THE SITE OF ACETYL-COA SYNTHESIS BY THE NICKEL-IRON-SULFUR PROTEIN, CARBON-MONOXIDE DEHYDROGENASE, FROM *CLOSTRIDIUM-THERMOACETICUM*. *Biochemistry*. 1994; 33(32):9769–9777. [PubMed: 8068656]
57. Maynard EL, Lindahl PA. Evidence of a molecular tunnel connecting the active sites for CO₂ reduction and acetyl-CoA synthesis in acetyl-CoA synthase from *Clostridium thermoaceticum*. *Journal of the American Chemical Society*. 1999; 121(39):9221–9222.
58. Seravalli J, Ragsdale SW. Channeling of carbon monoxide during anaerobic carbon dioxide fixation. *Biochemistry*. 2000; 39(6):1274–1277. [PubMed: 10684606]
59. Lazarus O, et al. Water-Gas Shift Reaction Catalyzed by Redox Enzymes on Conducting Graphite Platelets. *Journal of the American Chemical Society*. 2009; 131(40):14154–14155. [PubMed: 19807170]
60. James OO, et al. Increasing carbon utilization in Fischer-Tropsch synthesis using H₂-deficient or CO₂-rich syngas feeds. *Fuel Processing Technology*. 2010; 91(2):136–144.
61. Woolerton TW, et al. CO₂ photoreduction at enzyme-modified metal oxide nanoparticles. *Energy & Environmental Science*. 2011; 4(7):2393–2399.
62. Chaudhary YS, et al. Visible light-driven CO₂ reduction by enzyme coupled CdS nanocrystals. *Chemical Communications*. 2012; 48(1):58–60. [PubMed: 22083268]
63. Bachmeier A, et al. How Light-Harvesting Semiconductors Can Alter the Bias of Reversible Electrocatalysts in Favor of H₂ Production and CO₂ Reduction. *Journal of the American Chemical Society*. 2013; 135(40):15026–15032. [PubMed: 24070184]
64. Wang VCC, Ragsdale SW, Armstrong FA. Investigations of Two Bidirectional Carbon Monoxide Dehydrogenases from *Carboxydotherrmus hydrogenoformans* by Protein Film Electrochemistry. *Chembiochem : a European journal of chemical biology*. 2013; 14(14):1845–1851. [PubMed: 24002936]

65. Wang VCC, et al. A Unified Electrocatalytic Description of the Action of Inhibitors of Nickel Carbon Monoxide Dehydrogenase. *Journal of the American Chemical Society*. 2013; 135(6): 2198–2206. [PubMed: 23368960]

Class I (methanogens) and Class II (acetoclastic methanogens)

Class III (homoacetogen)

Class IV

Figure 1.

Four different classes of anaerobic NiFe-containing CODH are classified by the subunit composition and biological roles. The Upper case letters denote the type of metal cluster. The A-cluster is the active site for acetyl-CoA synthesis. The C-cluster is the active site in which CO/CO₂ conversion occurs. Several [Fe₄S₄] clusters are required for electron transfers in enzymes (B-cluster) and Class I and Class II enzymes have two additional [4Fe-4S] clusters (E and F clusters) in the enzyme complex. Co refers to the cobalt-corrinoid serving as a methyl carrier

Figure 2. CODES TO BE GIVEN

3D-structures of carbon monoxide dehydrogenases that have been studied by PFE. *Upper left*: CODH II_{Ch} showing how a relay of FeS clusters leads from the exposed D-cluster to two C-clusters, each one housed in one of the subunits. *Lower left*: same structure viewed facing up from the D-cluster. *Right*: CODH/ACS shown with the CODH viewed in same way as for lower-left view of CODH II_{Ch}.

Figure 3. The active site of CODH II_{Ch} seen through various crystal structures. (a) –600mV with CO₂, (b) –320mV, (c) –320mV with cyanide and (d) CO-reduced CODH. Two positions are found for the dangling iron atom in the crystal structure, which is labelled Fe_{1a} and Fe_{1b} respectively. The pdb codes are shown in each case.

Figure 4.
Concept of PCE, applied to CODH. Squares represent the FeS clusters D, B and C.

Figure 5.

Electrocatalysis by carbon monoxide dehydrogenases adsorbed on a PGE rotating disk electrode in the presence of CO₂ and CO. A. CODH I_{Ch} and CODH II_{Ch} under 50% CO and 50% CO₂. Experimental Conditions: 25 °C, 0.2 M MES buffer (pH 7.0), electrode rotation rate 3500 rpm, and scan rate 2 mV s⁻¹. Figure is reproduced from ref [19]. B.

Figure 6.

A potential-domain dynamic picture of the effect of cyanide on catalytic properties of CODH. *Left panels.* Inhibition of CO oxidation activity of CODH I_{Ch} and CODH II_{Ch} by cyanide under 100 % CO. An aliquot of KCN stock solution (giving a final concentration of 1mM in the electrochemical cell) was injected during the second cycle. Conditions: 25 °C, 0.2 M MES buffer (pH=7.0), electrode rotation rate, 3500 rpm, scan rate, 1mV sec⁻¹. *Right panels.* Inhibition of CO₂ reduction activity of CODH I_{Ch} and CODH II_{Ch} by CN⁻ under 100% CO₂, showing that CN⁻ does not bind under strongly reducing conditions. An aliquot of KCN stock solution (giving a final concentration of 1mM in the electrochemical cell) was injected during the second cycle. Note that a more negative potential (by approximately 70 mV) is required to reductively reactivate CODH II_{Ch}. Conditions: 25 °C, 0.2 M MES buffer (pH=7.0), electrode rotation rate, 3500 rpm, scan rate, 1mV sec⁻¹.

Figure 5.

Chronoamperometric measurements of the inactivation (Figure 5a and b) and re-activation (Figure 5c and d) rate of cyanide-inhibited CODH I_{Ch} and CODH II_{Ch} . The inactivation rate of CODH II_{Ch} by cyanide was measured at -460 mV (CO oxidation, Figure 5a) and -560 mV (CO_2 reduction, Figure 5b). A final concentration of 0.5 mM cyanide in the electrochemical cell was used to measure the half-life time for inactivation. Cyanide release from CODH II_{Ch} (Figure 5d) at -760 mV is much faster than the instrumental response. Conditions: 25°C , 0.2 M MES buffer ($\text{pH}=7.0$), and rotation rate 3500 rpm .

Figure 8.

Inhibition of CODH I_{Ch} by cyanate. Potassium cyanate was injected (final concentration in the solution is 6.67mM) into the electrochemical cell under gas atmosphere at 100% CO_2 (Figure 3a) and 50% CO, 50% CO_2 (Figure 3b). The experimental condition: 25°C, 0.2M MES buffer (pH=7.0), rotation rate 3500 rpm and scan rate $1mV s^{-1}$.

Figure 9.

Reactions of carbon monoxide dehydrogenase with sulfide and thiocyanate. *Upper.* Cyclic voltammograms showing the reaction of CODH I_{Ch} and CODH II_{Ch} with sulfide: An aliquot of sodium sulfide stock solution (giving 1 mM final concentration) was injected into the electrochemical cell. Experimental conditions: 25 °C, 0.2 M MES buffer (pH 7.0), rotation rate 3500 rpm, scan rate: 1 mV s⁻¹. Figure is reproduced from ref [64] *Lower.* Inhibition of CODH by thiocyanate. Thiocyanate was injected into the electrochemical cell to give a final

concentration of 6.6 mM. Experimental conditions: 25 °C, 0.2 M MES buffer (pH 7.0), rotation rate 3500 rpm, scan rate: 1 mV s⁻¹.

Figure 10.
Potential dependence of binding of inhibitors to CODHI_{Ch} .

Figure 11.

Water gas shift reaction catalyzed by hydrogenase and CODH co-adsorbed on conducting graphite particles. Plot shows production of H₂ and depletion of CO over the course of 55 h, with fresh CO injections of 600 μL at the points indicated.

Figure 12.

Upper. Cartoon showing a hybrid system for light-driven CO_2 reduction using CODH on CdS nanoparticles. Numbers refer to potentials in V vs SHE. *Lower.* Comparison of the performance of CODH adsorbed on various CdS nanoparticles: NR = nanorods, QD = quantum dots, CdS_{calc} = calcined sample. Adapted from Reference $\times$ with permission.

Figure 13.

Cyclic voltammograms (scan rate 10 mV s^{-1}) of electrocatalysis by CODH I_{Ch} adsorbed at PGE (upper), CdS thin film (middle) and TiO_2 thin film (lower) electrodes, scanned in 0.2 M MES, pH 6.0. Voltammograms recorded under 100% CO_2 are depicted in blue, and those recorded under 50% CO , 50% CO_2 are shown in red.

Scheme 1.

Two possible mechanisms of Ni-Fe CODH proposed by (a) Dobbek and Lindahl, in which the Ni subsite in C_{red2} is formally Ni(0) or (b) as proposed by Fontecilla-Camps et al[25], in which the Ni subsite in C_{red2} is bonded to a H atom, making what is formally a Ni(II)-H species. The species 'B' refers to the amino acid that accepts/donates the proton, possibly His or Lys. The red colors represent substrate-mimic inhibitors in which cyanide (CN^-) inhibits the C_{red1} state and cyanate (NCO^-) inhibits the C_{red2} state.

Scheme 2.

Summary of the interceptions of the catalytic cycle of CODH by small molecule inhibitors. The potential -520 mV is the standard potential for the CO_2/CO half cell reaction at pH 7.0. The potentials -50 mV and -250 mV are the values observed for re-activation of enzyme with and without sulfide. Reprinted from ref [65]