Published in final edited form as:

Environ Res. 2014 November ; 135: 221–226. doi:10.1016/j.envres.2014.09.014.

Exposure to Airborne Polycyclic Aromatic Hydrocarbons During Pregnancy and Risk of Preterm Birth

Amy M. Padula, Ph.D., M.Sc.¹, Elizabeth M. Noth, Ph.D., M.P.H.², S. Katharine Hammond, Ph.D., M.S.², Fred W. Lurmann, M.S.³, Wei Yang, M.D., M.S., M.A.¹, Ira B. Tager, M.D., M.P.H.², and Gary M. Shaw, Dr.P.H.¹

¹ Stanford University, School of Medicine, Department of Pediatrics, Stanford, CA

² University of California, Berkeley, School of Public Health, Division of Environmental Health Sciences, Berkeley, CA

³ Sonoma Technology, Inc., Petaluma, CA

Abstract

Background—Preterm birth is an important marker of health and has a prevalence of 12-13% in the U.S. Polycyclic aromatic hydrocarbons (PAHs) are a group of organic contaminants that form during the incomplete combustion of hydrocarbons, such as coal, diesel and gasoline. Studies suggest that exposure to PAHs during pregnancy is related to adverse birth outcomes. The aim of this study is to evaluate the association between exposure to PAHs during the pregnancy and preterm birth.

Methods—The study population included births from years 2001-2006 of women whose maternal residence was within 20 km of the primary monitoring site in Fresno, California. Data in the Fresno area were used to form a spatio-temporal model to assign daily exposure to PAHs with 4, 5, or 6 rings at the maternal residence throughout pregnancy of all of the births in the study area. Gestational age at birth and relevant covariates were extracted from the birth certificate.

Results—We found an association between PAHs during the last 6 weeks of pregnancy and birth at 20-27 weeks (OR=2.74; 95% CI: 2.24-3.34) comparing the highest quartile to the lower three. The association was consistent when each quartile was compared to the lowest (OR_{2nd}=1.49, 95% CI: 1.08-2.06; OR_{3rd}=2.63, 95% CI: 1.93-3.59; OR_{4th}=3.94, 95% CI: 3.03-5.12). Inverse associations were also observed for exposure to PAHs during the entire pregnancy and the first trimester and birth at 28-31 weeks and 20-27 weeks.

Conclusion—An association between PAH exposure during the 6 weeks before delivery and early preterm birth was observed. However, the inverse association with early preterm birth offers an unclear, and potentially complex, inference of these associations.

© 2014 Elsevier Inc. All rights reserved.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

outdoor air pollution; prematurity; polycyclic aromatic hydrocarbons; pregnancy

INTRODUCTION

Preterm birth is an important marker of health in the neonatal period, through childhood and possibly into adulthood. In the U.S., 12-13% of births are preterm (1) – a public health challenge that costs society at least \$26 billion a year (2). Preterm birth is generally defined as birth at <37 weeks gestation; however, it can be and should be classified into finer gestational periods. Several studies have examined the relation between air pollution and preterm birth, many of which have found associations with criteria pollutants (monitored by the U.S. Environmental Protection Agency) including particulate matter (PM) with aerodynamic diameter <2.5 and <10 microns (PM_{2.5} and PM₁₀) (3-5).

Important components of PM_{2.5} and PM₁₀ include polycyclic aromatic hydrocarbons (PAHs), a group of organic contaminants that form during combustion of hydrocarbons, such as coal, diesel, gasoline, wood, cigarettes and food items and food preparation (6). PAHs are ubiquitous environmental contaminants that can be due to occupational, dietary or airborne sources. The current study focuses on ambient PAHs found in the environment primarily as a result of automobile exhaust and industrial emissions.

Long-term exposure to ambient PAHs has been associated with increased risks of morbidity (7-13) and mortality (14, 15)(16, 17). . Perera *et al.* proposed that placental transfer of PAHs to the fetus have an impact on fetal development (18). Evidence from experimental studies demonstrate that benzo[a]pyrene (BaP) crosses the placenta (19) and has been classified by the U.S. Environmental Protection Agency (EPA) as a probable human carcinogen. Furthermore, animal studies have indicated PAHs may influence gestational length and intrauterine growth (20) and human studies have associated urinary and blood biomarkers of exposures to PAHs with birth weight (21) and gestational length (22).

PAH exposure is ubiquitous and detected in pregnant women (23, 24). Studies suggest that exposure to ambient PAH during pregnancy is related to adverse birth outcomes including low birth weight (25, 26), preterm birth (27, 28), intrauterine growth retardation (28, 29), and early pregnancy fetal death (30). However, a lack of adequate air monitoring data has limited our understanding of the effects of airborne PAHs on adverse birth outcomes. A recent study of traffic-related air toxics and preterm birth recommended future studies to focus on 1) PAHs and 2) accurate modeling of both local and regional spatial and temporal distributions, and incorporation of source information (27).

The current study extends previous work that found associations between PM_{2.5} and PM₁₀ and preterm birth, with stronger associations with earlier gestational ages (*i.e.*, <27 weeks gestation) in the San Joaquin Valley (5). Here we evaluate associations between exposure to ambient PAHs during specific periods of pregnancy and preterm birth in Fresno, California during years 2001-2006. Fresno is a highly polluted city in California with a high prevalence of adverse health outcomes including premature mortality and asthma (31).

METHODS

Study population

The study population is a subset of the Study of Air pollution, Genetics and Early life events (SAGE). The entire study population has been described previously (5, 32). The population subset was selected because of its proximity to intensive monitoring of PAHs in the Fresno area during these years.

Outcome ascertainment

Analyses were limited to singleton births between 20 and 42 weeks gestation and birth weight between 500 and 5000 grams. Preterm birth was defined by gestational age at birth as determined from the last menstrual period on the birth certificate. Preterm birth is a complex phenomenon and a simple dichotomous definition has been argued to be too simplistic for etiologic studies owing to the heterogeneity that has been observed with this outcome (33). Four categories of preterm birth were created based on gestational ages: 20-27 weeks, 28-31 weeks, 32-33 weeks and 34-36 weeks. Term births (*i.e.*, 37-42 weeks) were considered the reference in all analyses.

Exposure assessment

Maternal residence at birth locations, as determined from birth certificates, were geocoded (point addresses) with Arc GIS software (Environmental Systems Research Institute, Redlands, CA, version 9). Addresses were corrected with ZP4 software (Semaphore Corporation, Aptos, CA) in ArcView and SAS (SAS Institute Inc., Cary, NC, version 9.2).

PAH exposure assignments were made if the geocoded residences were within 20 km (12.4 miles) of the First Street U.S. EPA air monitoring site in Fresno, California and were available for at least 75% of the exposure period during pregnancy. The study area included urban, suburban and rural areas. A combination of data including measurements made at the EPA air monitoring site of 4, 5 and 6 ring PAHs (PAH456) between 2000-2006, intensive air pollution sampling conducted at 83 locations in Fresno between February 2002 and February 2003, meteorological data, source data, and other spatial variables were used to form a spatio-temporal model to assign daily exposure to PAHs at the maternal residence throughout pregnancy of all births in the study area (34).

The model estimates the sum of PAHs including fluoranthene, benz[a]anthracene, chrysene, benzo[a]pyrene, benzo[b]fluoranthene, benzo[k]fluoranthene, benzo[ghi]perylene, indeno[1,2,3-cd]pyrene, and dibenz[a,h]anthracene. These PAHs represent semi-volatile PAHs of the original 16 priority PAHs that were identified by the U.S. EPA. This model explained 81% of the between-residence variability and 18% of the within-residence variability as measured by mixed-effects model variance components. Main sources of PAHs in Fresno are the freeways, major arterials, agricultural burns, residential fireplace and woodburning stove use, and heating (35). More detail on field collection, measurement results, and modeling can be found in Noth et al. (34).

Covariates

Variables from birth certificates included in analyses were: infant birth weight, maternal age (<20, 20-35, >35 years), maternal race/ethnicity (White, Hispanic, African-American, Asian, other), maternal education (no high school, some high school, some college, bachelors or other degree), parity (0, >1), prenatal care (initiated in first trimester or not), Medi-Cal (Medicaid) or other government program payment of birth expenses, infant sex, and birth year (2001-2006).

Socioeconomic status (SES), such as poverty and unemployment, has been associated with adverse birth outcomes (36). Furthermore, SES has been identified as an effect modifier in the relationship between air pollution and adverse birth outcomes (5, 37-40). Based on measures implemented in a previous study (40), we created an indicator variable for low SES that was defined as unemployment >10 percent, income from public assistance >15 percent, and families below poverty level >20 percent in the 2000 U.S. Census at the block group level (40, 41). Such an indicator may not pertain directly to any individual, but provides contextual information about study population neighborhoods. This research was approved by the University of California, Berkeley Office for Protection of Human Subjects, Stanford University Institutional Review Board, and the California State Committee for the Protection of Human Subjects.

Statistical Analyses

First, second, and third pregnancy trimesters were defined as gestational weeks 1-13, 14-26, and 27 to birth, respectively. The exposure period for pregnancy began 14 days after the first day of the last menstrual period. Additionally, we calculated metrics for the last 6 weeks of pregnancy (birth minus 42 days). We used logistic regression to examine the association between the highest quartile of PAH456 compared to the lower three and each of the four gestational definitions of preterm birth (20-27 weeks, 28-31 weeks, 32-33 weeks and 34-26 weeks) versus term (37-42 weeks). Exposure periods of the term births were truncated to match the same period as the comparison period-length of the preterm births.

The distributions of several covariates were also examined in relation to the PAHs and preterm birth distributions: maternal age, maternal race/ethnicity, maternal education, parity, prenatal care, Medi-Cal birth expenses payment, infant sex, neighborhood SES, season of conception, and birth year.

Exposures were categorized as “high” if they were above the 75%-ile of entire pregnancy averages. First, second, and third pregnancy trimesters were defined as gestational weeks 1-13, 14-26, and 27 to birth, respectively. We used logistic regression to examine associations between PAH456 during each trimester and the last 6 weeks of pregnancy and the four gestational definitions of prematurity (20-27 weeks, 28-31 weeks, 32-33 weeks and 34-36 weeks) versus term (37-42 weeks). Previous analyses of this cohort indicated the last 6 weeks of pregnancy may be a critical period for the association between particulate matter and early preterm birth (5).

We adjusted models for the following covariates: maternal age, race/ethnicity, education, prenatal care in the first trimester, and Medi-Cal payment of birth costs. We stratified by

neighborhood SES and season of conception to determine whether either factor modified an effect of PAH exposures on preterm birth. Analyses were restricted to births without reported maternal diabetes (types 1 and 2, and gestational) or hypertension (chronic, gestational and preeclampsia) owing to the known association between these morbidities and preterm birth.

All analyses were conducted using SAS 9.3 (SAS Institute Inc., Cary, NC, 2012).

RESULTS

The characteristics of the study population of 42,904 births are presented in Table 1 by gestational age category. Most study mothers were Hispanic, had Medi-Cal payment of birth costs, and had at least a high school education.

PAH exposures during pregnancy had a median of 3.6 ng/m³ with an interquartile range of 1.6 ng/m³ and a range of 12.4 ng/m³. PAH exposures over the entire pregnancy were correlated with outdoor PM_{2.5} ($\rho=0.53$) and PM₁₀ ($\rho=0.38$) as well as traffic density within 300m of the home ($\rho=0.30$) (ρ is defined as the Pearson correlation coefficient). The distribution of PAH exposures by gestational age category is presented in Table 2.

Table 3 displays the relation between PAH exposures during different periods of pregnancy and the four gestational week definitions of preterm birth. Associations were observed between PAH exposures during the last 6 weeks of pregnancy and birth at 20-27 weeks (OR=2.74; 95% CI: 2.24-3.34) comparing the highest quartile to the lower three quartiles. When examined for an exposure-response, the association increased across each quartile of PAH exposures. Inverse associations were also observed for exposure to PAHs during the entire pregnancy and the first trimester and birth at 28-31 weeks and 20-27 weeks.

Stratification by neighborhood SES showed little differences between groups. For births conceived during summer (June-August), associations were statistically significant for each category of preterm birth and exposure to PAHs during the last 6 weeks with odds ratios ranging from 2.5 (95% CI: 2.1, 2.9) for birth at 34-36 weeks to 10.2 (95% CI: 7.8, 13.4) for birth 20-27 weeks (Table 4). Inverse associations were observed for those conceived in the spring (March-May).

DISCUSSION

Our study shows an association between elevated levels of PAH₄₅₆ during the last 6 weeks of pregnancy and early preterm birth (i.e., <27 weeks gestation) in Fresno, California with risk increasing with increasing quartiles of exposure. However, inverse associations with early preterm birth were also observed for the first trimester and entire pregnancy metrics of PAH exposures. The biologic explanation for the latter is unknown. It could be hypothesized that highly exposed pregnancies were spontaneously aborted during the first trimester creating a bias among the survived births; however, we do not have the data in this study to address such a question.

Ambient PAH levels vary by season in Fresno, with higher levels in the winter. Preterm birth also varies by season, especially in Fresno during these years, but it is unknown whether such preterm birth variation is because of PAHs or an unknown and highly correlated factor. It is unclear if or when a critical period may exist for risk of preterm birth; though this study and its larger cohort have found the last 6 weeks (which also corresponds to the second trimester for early preterm birth) to have the strongest associations.

A few previous studies of PAH exposures and preterm birth have been conducted. A recent study found an association between PAH exposure (as predicted by an EPA Chemical Mass Balance model) and preterm birth in another geographic area of California, i.e., Los Angeles (27). However, no clear pattern indicating specific gestational periods of greater susceptibility were observed in this population.

A study in New York City found associations between PAH exposure during pregnancy and preterm birth among African Americans; however, the associations were not seen among Dominicans. A 1-In-unit increase in prenatal PAH exposure was associated with a 5-fold greater risk of preterm birth for African Americans (28).

Our study is subject to certain limitations. Estimation of personal exposure to air pollutants using ambient monitoring data suffers from potential exposure misclassification due to intra-urban variability in outdoor concentrations of pollutants (42). This misclassification would likely bias our estimates toward the null since the quartiles held a generally consistent exposure response relationship. Although we have precise information of the maternal residence at birth, we do not have data on residential mobility or workplace location. We lack data on other sources of PAHs including cigarette smoke (active and passive), dietary ingestion and indoor or occupational sources such as cooking. There are additional potential confounders for which we do not have data, including maternal pre-pregnancy BMI and nutrition.

The strengths of our study include comprehensive spatio-temporal modeling of PAHs over 6 years in a highly exposed region of the country. The model was able to show both temporal and spatial variation by using a mixed-effects regression model based on measurement data combined with source, land use, and neighborhood characteristics. The model generated the daily spatial distribution of outdoor residential PAH₄₅₆ concentrations in Fresno, as well as individual exposure estimates for each participant. Daily estimates were combined to provide pregnancy-specific exposure estimates. Previous work describing the spatial distribution of PAHs has been limited primarily to source-to-receptor studies (43, 44), small neighborhood-sized areas (45) or to large-scale regional or national annual distributions (46-50). While each of these approaches is useful for certain situations, such as for policy recommendations or long-term exposure studies, they cannot address the impact of short-term exposure on acute health outcomes as our daily model can.

In conclusion, ambient PAH exposure in the last 6 weeks of pregnancy may be associated with increased risk of early preterm birth. However, given the mixed results, more studies are needed to determine the relationship between PAHs and early preterm birth.

ACKNOWLEDGEMENTS

This work was supported by the National Institute for Environmental Health Science (R21ES014891; P20ES018173; P01ES022849; K99ES021470) and the March of Dimes Prematurity Center at Stanford University. This publication was made possible by US EPA STAR Grant RD83459601 and RD83543501. Its contents are solely the responsibility of the grantee and do not necessarily represent the official views of the US EPA. Further, the US EPA does not endorse the purchase of any commercial products or services mentioned in the publication.

Abbreviations

CI	confidence interval
PAHs	polycyclic aromatic hydrocarbons
PM	particulate matter
SAGE	study of air pollution, genetics and the environment
SES	socioeconomic status

References

1. Goldenberg RL, Culhane JF, Iams JD, Romero R. Epidemiology and causes of preterm birth. *Lancet*. Jan 5; 2008 371(9606):75–84. PubMed PMID: 18177778. [PubMed: 18177778]
2. Behrman, RE.; Butler, AS., editors. Preterm Birth: Causes, Consequences and Prevention. Institute of Medicine, Committee on Understanding Premature Birth and Assuring Healthy Outcomes; Washington, D.C.: 2007.
3. Stieb DM, Chen L, Eshoul M, Judek S. Ambient air pollution, birth weight and preterm birth: A systematic review and meta-analysis. *Environ Res*. Aug; 2012 117:100–11. PubMed PMID: 22726801. Epub 2012/06/26. eng. [PubMed: 22726801]
4. Shah PS, Balkhair T. Air pollution and birth outcomes: A systematic review. *Environ Int*. Feb; 2010 37(2):498–516. PubMed PMID: 21112090. Epub 2010/11/30. Eng. [PubMed: 21112090]
5. Padula AM MK, Tager IB, Hammond SK, Lurmann F, Yang W, Stevenson D, Shaw GM. Traffic-Related Air Pollution and Risk of Preterm Birth in the San Joaquin Valley of California. (Under Review).
6. Bostrom CE, Gerde P, Hanberg A, Jernstrom B, Johansson C, Kyrklund T, et al. Cancer risk assessment, indicators, and guidelines for polycyclic aromatic hydrocarbons in the ambient air. *Environmental health perspectives*. Jun; 2002 110(Suppl 3):451–88. PubMed PMID: 12060843. Pubmed Central PMCID: 1241197. [PubMed: 12060843]
7. Gale SL, Noth EM, Mann J, Balmes J, Hammond SK, Tager IB. Polycyclic aromatic hydrocarbon exposure and wheeze in a cohort of children with asthma in Fresno, CA. *J Expo Sci Environ Epidemiol*. Jul; 2012 22(4):386–92. PubMed PMID: 22549720. Epub 2012/05/03. eng. [PubMed: 22549720]
8. Nadeau K, McDonald-Hyman C, Noth EM, Pratt B, Hammond SK, Balmes J, et al. Ambient air pollution impairs regulatory T-cell function in asthma. *J Allergy Clin Immunol*. Oct; 2010 126(4): 845–52. e10. PubMed PMID: 20920773. Epub 2010/10/06. eng. [PubMed: 20920773]
9. Gauderman WJ, McConnell R, Gilliland F, London S, Thomas D, Avol E, et al. Association between air pollution and lung function growth in southern California children. *Am J Respir Crit Care Med*. Oct; 2000 162(4 Pt 1):1383–90. PubMed PMID: 11029349. Epub 2000/10/13. eng. [PubMed: 11029349]
10. Jedrychowski W, Flak E, Mroz E. The adverse effect of low levels of ambient air pollutants on lung function growth in preadolescent children. *Environmental health perspectives*. Aug; 1999 107(8):669–74. PubMed PMID: 10417366. Pubmed Central PMCID: 1566490. [PubMed: 10417366]
11. Miller RL, Garfinkel R, Horton M, Camann D, Perera FP, Whyatt RM, et al. Polycyclic aromatic hydrocarbons, environmental tobacco smoke, and respiratory symptoms in an inner-city birth

- cohort. *Chest*. Oct; 2004 126(4):1071–8. PubMed PMID: 15486366. Pubmed Central PMCID: 2223076. [PubMed: 15486366]
12. Perera FP, Rauh V, Whyatt RM, Tsai WY, Tang D, Diaz D, et al. Effect of prenatal exposure to airborne polycyclic aromatic hydrocarbons on neurodevelopment in the first 3 years of life among inner-city children. *Environmental health perspectives*. Aug; 2006 114(8):1287–92. PubMed PMID: 16882541. Pubmed Central PMCID: 1551985. [PubMed: 16882541]
 13. Rundle A, Hoepner L, Hassoun A, Oberfield S, Freyer G, Holmes D, et al. Association of childhood obesity with maternal exposure to ambient air polycyclic aromatic hydrocarbons during pregnancy. *American journal of epidemiology*. Jun 1; 2012 175(11):1163–72. PubMed PMID: 22505764. Pubmed Central PMCID: 3491973. [PubMed: 22505764]
 14. Martin S, Dawidowski L, Mandalunis P, Cereceda-Balic F, Tasat DR. Characterization and biological effect of Buenos Aires urban air particles on mice lungs. *Environ Res*. Nov; 2007 105(3):340–9. PubMed PMID: 17628521. [PubMed: 17628521]
 15. Pei XH, Nakanishi Y, Inoue H, Takayama K, Bai F, Hara N. Polycyclic aromatic hydrocarbons induce IL-8 expression through nuclear factor kappaB activation in A549 cell line. *Cytokine*. Sep 7; 2002 19(5):236–41. PubMed PMID: 12393170. [PubMed: 12393170]
 16. Hoshuyama T, Pan G, Tanaka C, Feng Y, Yu L, Liu T, et al. Mortality of iron-steel workers in Anshan, China: a retrospective cohort study. *International journal of occupational and environmental health*. Jul-Sep; 2006 12(3):193–202. PubMed PMID: 16967824. [PubMed: 16967824]
 17. Matsumoto Y, Ide F, Kishi R, Akutagawa T, Sakai S, Nakamura M, et al. Aryl hydrocarbon receptor plays a significant role in mediating airborne particulate-induced carcinogenesis in mice. *Environmental science & technology*. May 15; 2007 41(10):3775–80. PubMed PMID: 17547212. [PubMed: 17547212]
 18. Perera FP, Mooney LA, Stampfer M, Phillips DH, Bell DA, Rundle A, et al. Associations between carcinogen-DNA damage, glutathione S-transferase genotypes, and risk of lung cancer in the prospective Physicians' Health Cohort Study. *Carcinogenesis*. Oct; 2002 23(10):1641–6. PubMed PMID: 12376472. [PubMed: 12376472]
 19. Mathiesen L, Rytting E, Mose T, Knudsen LE. Transport of benzo[alpha]pyrene in the dually perfused human placenta perfusion model: effect of albumin in the perfusion medium. *Basic & clinical pharmacology & toxicology*. Sep; 2009 105(3):181–7. PubMed PMID: 19508462. [PubMed: 19508462]
 20. Detmar J, Rennie MY, Whiteley KJ, Qu D, Taniuchi Y, Shang X, et al. Fetal growth restriction triggered by polycyclic aromatic hydrocarbons is associated with altered placental vasculature and AhR-dependent changes in cell death. *American journal of physiology Endocrinology and metabolism*. Aug; 2008 295(2):E519–30. PubMed PMID: 18559983. [PubMed: 18559983]
 21. Pedersen M, Schoket B, Godschalk RW, Wright J, von Stedingk H, Tornqvist M, et al. Bulky dna adducts in cord blood, maternal fruit-and-vegetable consumption, and birth weight in a European mother-child study (NewGeneris). *Environmental health perspectives*. Oct; 2013 121(10):1200–6. PubMed PMID: 23906905. Pubmed Central PMCID: 3801201. [PubMed: 23906905]
 22. Singh VK, Patel DK, Jyoti, Ram S, Mathur N, Siddiqui MK. Blood levels of polycyclic aromatic hydrocarbons in children and their association with oxidative stress indices: an Indian perspective. *Clinical biochemistry*. Feb; 2008 41(3):152–61. PubMed PMID: 18154730. [PubMed: 18154730]
 23. Woodruff TJ, Zota AR, Schwartz JM. Environmental chemicals in pregnant women in the United States: NHANES 2003-2004. *Environmental health perspectives*. Jun; 2011 119(6):878–85. PubMed PMID: 21233055. Pubmed Central PMCID: 3114826. [PubMed: 21233055]
 24. Nethery E, Wheeler AJ, Fisher M, Sjodin A, Li Z, Romanoff LC, et al. Urinary polycyclic aromatic hydrocarbons as a biomarker of exposure to PAHs in air: a pilot study among pregnant women. *J Expo Sci Environ Epidemiol*. Jan-Feb; 2012 22(1):70–81. PubMed PMID: 21915154. [PubMed: 21915154]
 25. Choi H, Jedrychowski W, Spengler J, Camann DE, Whyatt RM, Rauh V, et al. International studies of prenatal exposure to polycyclic aromatic hydrocarbons and fetal growth. *Environmental health perspectives*. Nov; 2006 114(11):1744–50. PubMed PMID: 17107862. Pubmed Central PMCID: 1665416. [PubMed: 17107862]

26. Perera FP, Rauh V, Tsai WY, Kinney P, Camann D, Barr D, et al. Effects of transplacental exposure to environmental pollutants on birth outcomes in a multiethnic population. *Environmental health perspectives*. Feb; 2003 111(2):201–5. PubMed PMID: 12573906. Pubmed Central PMCID: 1241351. [PubMed: 12573906]
27. Wilhelm M, Ghosh JK, Su J, Cockburn M, Jerrett M, Ritz B. Traffic-related air toxics and preterm birth: a population-based case-control study in Los Angeles County, California. *Environmental health : a global access science source*. 2011; 10:89. PubMed PMID: 21981989. Pubmed Central PMCID: 3204282. [PubMed: 21981989]
28. Choi H, Rauh V, Garfinkel R, Tu Y, Perera FP. Prenatal exposure to airborne polycyclic aromatic hydrocarbons and risk of intrauterine growth restriction. *Environmental health perspectives*. May; 2008 116(5):658–65. PubMed PMID: 18470316. Pubmed Central PMCID: 2367680. [PubMed: 18470316]
29. Dejmeek J, Solansky I, Benes I, Lenicek J, Sram RJ. The impact of polycyclic aromatic hydrocarbons and fine particles on pregnancy outcome. *Environmental health perspectives*. Dec; 2000 108(12):1159–64. PubMed PMID: 11133396. Pubmed Central PMCID: 1240197. [PubMed: 11133396]
30. Wu J, Hou H, Ritz B, Chen Y. Exposure to polycyclic aromatic hydrocarbons and missed abortion in early pregnancy in a Chinese population. *The Science of the total environment*. May 1; 2010 408(11):2312–8. PubMed PMID: 20219237. [PubMed: 20219237]
31. Place Matters for Health in the San Joaquin Valley: Ensuring Opportunities for Good Health for All. Joint Center for Political and Economic Studies; Washington, D.C.: Mar. 2012 Report No
32. Padula AM, Mortimer K, Hubbard A, Lurmann F, Jerrett M, Tager IB. Exposure to traffic-related air pollution during pregnancy and term low birth weight: estimation of causal associations in a semiparametric model. *American journal of epidemiology*. Nov 1; 2012 176(9):815–24. PubMed PMID: 23045474. Pubmed Central PMCID: 3571254. [PubMed: 23045474]
33. Zhang, J.; Savitz, DA. Duration of gestation and timing of birth.. In: Buck Louis, GM.; Platt, RW., editors. *Reproductive and Perinatal Epidemiology*. Oxford University Press; New York: 2011. p. 152–67.
34. Noth EM, Hammond K, Biging GS, Tager IB. A spatial-temporal regression model to predict daily outdoor residential PAH concentrations in an epidemiologic study in Fresno, CA. *Atmospheric Environment*. 2011; 45:2394–403.
35. Schauer JJ, Cass GR. Source apportionment of wintertime gas-phase and particle phase air pollutants using organic compounds as tracers. *Environmental science & technology*. 2000; 34:1821–32.
36. Hubbard A, van der Laan MJ. Population intervention methods. *Biometrika*. 2008; 95:35–77. [PubMed: 18629347]
37. Yi O, Kim H, Ha E. Does area level socioeconomic status modify the effects of PM(10) on preterm delivery? *Environ Res*. Jan; 110(1):55–61. PubMed PMID: 19878932. Epub 2009/11/03. eng. [PubMed: 19878932]
38. Zeka A, Melly SJ, Schwartz J. The effects of socioeconomic status and indices of physical environment on reduced birth weight and preterm births in Eastern Massachusetts. *Environ Health*. 2008; 7:60. PubMed PMID: 19032747. eng. [PubMed: 19032747]
39. Genereux M, Auger N, Goneau M, Daniel M. Neighbourhood socioeconomic status, maternal education and adverse birth outcomes among mothers living near highways. *J Epidemiol Community Health*. Aug; 2008 62(8):695–700. PubMed PMID: 18621954. Epub 2008/07/16. eng. [PubMed: 18621954]
40. Ponce N, Hoggatt K, Wilhelm M, Ritz B. Preterm birth: the interaction of traffic-related air pollution with economic hardship in Los Angeles neighborhoods. *Am J Epidemiol*. Jul 15; 2005 162(2):140–8. Epub 2005 Jun 22. [PubMed: 15972941]
41. Census USBot. Census of population and housing. Summary tape file 3: technical documentation. Vol. 2000. Bureau of the Census; Washington D.C.: 2000. p. 2000
42. Briggs DJ, de Hoogh C, Gulliver J, Wills J, Elliott P, Kingham S, et al. A regression-based method for mapping traffic-related air pollution: application and testing in four contrasting urban

- environments. *The Science of the total environment*. May 15; 2000 253(1-3):151–67. PubMed PMID: 10843339. [PubMed: 10843339]
43. Vardoulakis S, Gonzalez-Flesca N, Fisher BEA, Pericleous K. Spatial variability of air pollution in the vicinity of a permanent monitoring station in central Paris. *Atmospheric Environment*. 2005; 39:2725–36.
44. Li C-T, Lin Y-C, Lee W-J, Tsai P-J. Emission of polycyclic aromatic hydrocarbons and their carcinogenic potencies from cooking sources to the urban atmosphere. *Environmental Health Perspectives*. 2003; 111(4):483–7. [PubMed: 12676603]
45. Levy JI, Houseman EA, Spengler JD, Loh P, Ryan L. Fine particulate matter and polycyclic aromatic hydrocarbon concentration patterns in Roxbury, Massachusetts: a community-based GIS analysis. *Environmental Health Perspectives*. 2001; 109(4):341–7. [PubMed: 11335181]
46. Guo H, Lee SC, Ho KF, Wang XM, Zou SC. Particle-associated polycyclic aromatic hydrocarbons in urban air of Hong Kong. *Atmospheric Environment*. Dec; 2003 37(38):5307–17. PubMed PMID: ISI:000186602000002.
47. Lang C, Tao S, Zhang G, Fu J, Simonich S. Outflow of polycyclic aromatic hydrocarbons from Guangdong, Southern China. *Environ Sci Technol*. 2007; 41:8370–5. [PubMed: 18200865]
48. Prevedouros K, Jones KC, Sweetman AJ. Modelling the atmospheric fate and seasonality of polycyclic aromatic hydrocarbons in the UK. *Chemosphere*. 2004; 56:195–208. [PubMed: 15172593]
49. Manoli E, Kouras A, Samara C. Profile analysis of ambient and source emitted particle-bound polycyclic aromatic hydrocarbons from three sites in northern Greece. *Chemosphere*. 2004; 56:867–78. [PubMed: 15261533]
50. Thornhill DA, de Foy B, Herndon SC, Onasch TB, Wood EC, Zavala M, et al. Spatial and temporal variability of particulate polycyclic aromatic hydrocarbons in Mexico City. *Atmospheric Chemistry and Physics*. 2008; 8(12):3093–105.

Highlights

- Spatio-temporal models estimated exposure to ambient PAH during pregnancy.
- PAH exposure in the end of pregnancy was associated with early preterm birth.
- Seasonal differences in PAH and preterm birth are apparent in Fresno, California.

Table 1

Characteristics of the study population, total and by preterm birth (%)

Characteristic	Gestational age in weeks					Total n=42,904
	37-42 n=37,158	34-36 n=3940	32-33 n=757	28-31 n=605	20-27 n=444	
Maternal age (years)						
<20	14.0	15.7	14.7	18.4	17.6	14.3
20-24	30.3	30.4	32.4	32.7	34.9	30.4
25-29	27.0	25.7	23.8	22.8	23.0	26.7
30-34	18.5	16.9	16.8	14.1	15.1	18.2
35	10.2	11.3	12.4	12.1	9.5	10.4
Maternal race/ethnicity						
White	26.8	22.6	18.9	15.2	14.6	26.0
Hispanic	53.8	54.9	56.7	55.7	59.5	54.0
Black	7.0	9.2	11.6	13.4	8.3	7.4
Asian	10.7	11.6	10.8	13.6	14.2	10.8
Other	1.8	1.7	2.0	2.2	3.4	1.8
Maternal education						
<High school	10.2	9.8	12.4	12.2	13.1	10.3
High school	51.2	57.4	59.1	65.0	63.3	52.2
Some college	21.9	20.2	18.8	16.5	16.2	21.6
College or other degree	15.4	11.0	8.2	4.5	5.4	14.6
Missing	1.3	1.6	1.6	1.8	2.0	1.3
Initiation of prenatal care						
Before 1 st trimester	88.2	83.9	80.3	77.9	75.9	87.4
After 1 st trimester	11.2	14.8	17.8	18.8	18.9	11.9
Missing	0.6	1.3	1.9	3.3	5.2	0.7
Parity						
0	35.5	31.0	30.5	26.9	29.5	34.9
1	64.5	69.0	69.5	73.1	70.5	65.2
Infant sex						
Male	51.1	54.6	52.1	54.7	58.3	51.6
Female	48.9	45.4	48.0	45.3	41.7	48.4
Low neighborhood SES	30.4	34.8	38.2	41.8	42.1	31.2
Medi-Cal payment	58.6	65.4	68.3	74.4	77.7	59.8
Cesarean section	26.9	28.8	34.2	33.2	36.7	27.4
Season of conception						
Winter (December-February)	25.5	23.0	19.7	15.9	11.5	24.9
Spring (March-May)	24.8	26.0	26.6	19.2	14.4	24.8
Summer (June-August)	23.9	24.4	30.4	45.5	60.8	24.8

Characteristic	Gestational age in weeks					Total n=42,904
	37-42 n=37,158	34-36 n=3940	32-33 n=757	28-31 n=605	20-27 n=444	
Fall (September-November)	25.7	26.7	23.4	19.5	13.3	25.6
Year of birth						
2001	11.4	9.7	7.0	8.4	5.4	11.1
2002	14.7	14.4	12.4	9.8	5.9	14.5
2003	15.8	15.5	13.1	12.4	9.5	15.6
2004	17.4	17.1	15.3	9.4	10.1	17.2
2005	19.5	20.9	21.5	28.3	38.5	20.0
2006	21.1	22.4	30.7	31.7	30.6	21.7

Table 2

Median (interquartile range) of PAHs (ng/m³) for each exposure period by gestational age categories.

Exposure period	Gestational age in weeks					Total
	37-42	34-36	32-33	28-31	20-27	
Entire pregnancy	3.57 (1.59)	3.49 (1.66)	3.43 (1.66)	3.41 (1.48)	3.22 (1.30)	3.55 (1.59)
1 st trimester	3.18 (1.70)	3.14 (1.63)	3.07 (1.73)	2.97 (1.41)	2.74 (1.15)	3.17 (1.69)
2 nd trimester	3.11 (1.58)	3.12 (1.60)	3.22 (1.60)	3.44 (1.52)	3.50 (1.44)	3.12 (1.58)
3 rd trimester	3.09 (1.50)	3.06 (1.60)	3.06 (1.56)	3.27 (1.53)	3.06 (1.74)	3.09 (1.51)
Last 6 weeks	3.01 (1.69)	3.02 (1.66)	3.06 (1.66)	3.28 (1.46)	4.11 (2.12)	3.03 (1.69)

Table 3

Adjusted^a odds of birth at each gestational age for exposure to the highest quartile versus lower three quartiles of PAHs during each exposure period and 95% confidence intervals compared to term (gestational age at 37-42 weeks).

PAH exposure period	Gestational age in weeks									
	34-36		32-33		28-31		20-27			
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
Entire Pregnancy	0.99	0.92	1.07	0.93	0.79	1.11	0.76	0.62	0.51	0.39
1 st trimester	0.92	0.84	1.00	0.87	0.72	1.05	0.56	0.44	0.42	0.31
2 nd trimester	0.98	0.89	1.07	1.03	0.85	1.24	1.02	0.83	1.10	0.81
3 rd trimester	1.00	0.92	1.09	0.99	0.81	1.19	1.00	0.81	NC	--
Last 6 weeks	0.96	0.88	1.05	0.97	0.80	1.16	0.86	0.70	2.74	2.24

Gestation at 20-27 weeks PAHs during last 6 weeks			95% CI	
2 nd quartile	1.49		1.08	2.06
3 rd quartile	2.63		1.93	3.59
4 th quartile	3.94		3.03	5.12

Abbreviations: PAHs, polycyclic aromatic hydrocarbons; OR, odds ratio; CI, confidence interval

^aModels adjusted for maternal age, education, race/ethnicity, prenatal care in the first trimester and payment of birth expenses

Table 4

Season of conception-stratified, adjusted^a odds of birth at each gestational age for exposure to the highest quartile versus lower three quartiles of PAHs during each exposure period and 95% confidence intervals compared to term (gestational age at 37-42 weeks).

Season of conception	PAH exposure period	Gestational age											
		34-36 weeks		32-33 weeks		28-31 weeks		20-27 weeks					
		OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI				
Winter	Entire Pregnancy	0.97	0.80	1.19	1.07	1.70	2.43	1.52	3.87	1.05	0.47	2.37	
	1 st trimester	0.80	0.68	0.95	0.94	0.64	1.37	1.07	0.65	1.76	0.63	0.30	1.29
	2 nd trimester	1.00	0.69	1.43	0.36	0.09	1.47	1.81	0.78	4.19	0.95	0.13	7.09
	3 rd trimester	0.82	0.57	1.18	1.80	0.96	3.38	1.31	0.53	3.26	NC	--	--
	Last 6 weeks	0.57	0.39	0.83	1.53	0.84	2.80	1.21	0.52	2.79	0.78	0.19	3.26
Spring	Entire Pregnancy	0.66	0.54	0.80	0.41	0.25	0.68	0.20	0.08	0.50	0.15	0.04	0.63
	1 st trimester	1.20	0.87	1.65	1.22	0.64	2.34	0.43	0.11	1.75	0.85	0.20	3.52
	2 nd trimester	0.87	0.62	1.22	1.41	0.79	2.51	0.18	0.03	1.32	0.49	0.07	3.56
	3 rd trimester	0.76	0.66	0.87	0.47	0.34	0.64	0.74	0.50	1.10	NC	--	--
	Last 6 weeks	0.50	0.44	0.58	0.27	0.20	0.37	0.27	0.18	0.41	0.04	0.01	0.11
Summer	Entire Pregnancy	1.15	0.99	1.33	0.91	0.68	1.21	0.42	0.31	0.59	0.18	0.11	0.28
	1 st trimester	1.06	0.76	1.50	0.91	0.48	1.74	0.22	0.07	0.70	0.22	0.07	0.69
	2 nd trimester	0.93	0.80	1.08	0.72	0.54	0.96	0.40	0.30	0.53	0.47	0.32	0.67
	3 rd trimester	1.25	1.07	1.45	1.15	0.85	1.57	0.78	0.58	1.03	NC	--	--
	Last 6 weeks	2.45	2.09	2.88	2.95	2.17	4.02	2.17	1.62	2.9	10.2	7.8	13.4
Fall	Entire Pregnancy	1.08	0.94	1.23	1.22	0.90	1.67	1.14	0.78	1.69	2.73	1.57	4.74
	1 st trimester	0.97	0.83	1.12	1.23	0.87	1.73	0.88	0.58	1.33	1.75	0.95	3.23
	2 nd trimester	0.93	0.79	1.08	0.86	0.59	1.25	1.02	0.67	1.55	0.65	0.29	1.44
	3 rd trimester	0.86	0.60	1.24	0.77	0.31	1.89	0.24	0.03	1.73	NC	--	--
	Last 6 weeks	0.77	0.58	1.04	0.70	0.34	1.42	0.13	0.02	0.95	3.80	1.93	7.5

Abbreviations: PAHs, polycyclic aromatic hydrocarbons; OR, odds ratio; CI, confidence interval

^aModels adjusted for maternal age, education, race/ethnicity, prenatal care in the first trimester and payment of birth expenses