Published in final edited form as:

. 2014 May 1; 45(3): 364–374. doi:10.1007/s13592-013-0244-3.

The Architecture of the Pollen Hoarding Syndrome in Honey Bees: Implications for Understanding Social Evolution, Behavioral Syndromes, and Selective Breeding

Olav Rueppell^{1,*}

¹Department of Biology, The University of North Carolina at Greensboro, 312 Eberhart Building, 321 McIver Street, Greensboro, North Carolina, 27403, USA

Abstract

Social evolution has influenced every aspect of contemporary honey bee biology, but the details are difficult to reconstruct. The reproductive ground plan hypothesis of social evolution proposes that central regulators of the gonotrophic cycle of solitary insects have been coopted to coordinate social complexity in honey bees, such as the division of labor among workers. The predicted trait associations between reproductive physiology and social behavior have been identified in the context of the pollen hoarding syndrome, a larger suite of interrelated traits. The genetic architecture of this syndrome is characterized by a partially overlapping genetic architecture with several consistent, pleiotropic QTL. Despite these central QTL and an integrated hormonal regulation, separate aspects of the pollen hoarding syndrome may evolve independently due to peripheral QTL and additionally segregating genetic variance. The characterization of the pollen hoarding syndrome has also demonstrated that this syndrome involves many non-behavioral traits, which may be the case for numerous “behavioral” syndromes. Furthermore, the genetic architecture of the pollen hoarding syndrome has implications for breeding programs for improving honey health and other desirable traits: If these traits are comparable to the pollen hoarding syndrome, consistent pleiotropic QTL will enable marker assisted selection, while sufficient additional genetic variation may permit the dissociation of trade-offs for efficient multiple trait selection.

Keywords

Apis mellifera; Pleiotropy; Behavioral Syndrome; Honey Bee Health; Social Behavior; Correlated Evolution; Artificial Selection; Ovary

Behavioral Syndromes and Pleiotropy in Social Insects

Behavioral traits have generally been perceived as highly plastic and independent of other traits because they are the outcome of a complex central nervous system that integrates numerous external and internal influences. However, our increasing understanding of the proximate mechanisms of behavior has led to the realization that different behavioral traits are commonly correlated with each other. Correlation of different behavioral traits or the

*Corresponding author: olav_rueppell@uncg.edu, FAX: (1) 336-334 5839, Phone: (1) 336-256 2591.

same trait across different contexts has been defined as a behavioral syndrome to parallel the concept of life history syndromes (Sih, Bell et al., 2004). If the behavioral syndrome in question has a genetic basis, it parallels pleiotropy, the genetic description for one or multiple genes concomitantly influencing several phenotypes.

Studies of behavioral syndromes and studies of pleiotropy have focused on the resulting trade-offs between multiple correlated traits and the possibility of evolutionary constraint due to genetic covariance (Agrawal and Stinchcombe, 2009). The simple logic behind the trade-off is that an advantageous change in trait A may result in a non-advantageous change in the correlated trait B, potentially constraining the optimization of both traits (Price and Langen, 1992). While the possibility of such trade-offs through pleiotropy has long been realized in the fields of quantitative genetics (Lande, 1982) and life history theory (Williams, 1957; Stearns, 1989), the ideas has only recently found traction in the study of behavior (Sih, Bell et al., 2004; Jandt, Bengtson et al., 2013). The correlation between traits depends on the proximate mechanisms and genetic architecture that are responsible for their inter-dependence. Strong genetic covariance, high penetrance, and low intra-individual plasticity increase the possibility of trade-offs. Thus, the underlying genetic architecture of any syndrome of correlated traits is central to understanding the evolution of these traits (Roff, 2011).

Social insects are particularly interesting because of their strong phenotypic differentiation among closely related individuals and the additional colony level that results from the integration of many individuals. The integration of individuals into a colony leads to a new layer of behavioral syndrome and potential constraint (Jandt, Bengtson et al., 2013), particularly when individuals are closely related and therefore genetically similar. Similar to a genetic correlation between the two sexes (Delph, Arntz et al., 2010), genetic correlations between different castes and colony members may constrain the evolutionary optimization of specific phenotypes. The phenotypic plasticity that leads to differences among closely related individuals has presumably evolved to reduce constraints imposed by pleiotropy and facilitate social evolution (Gadagkar, 1997).

Honey bees are a highly derived clade and numerous changes have occurred during their social evolution from a solitary ancestor (Arias and Sheppard, 2005). It is difficult to determine how social evolution and adaptations to the new social organization have progressed in honey bees because fossil evidence does not resolve many behaviors (Engel, 1998). Furthermore, causes and consequences of social evolution are hard to establish because honey bees present only one singular transition to sociality (Arias and Sheppard, 2005). Therefore, the phenotypic and genotypic architectures of the present species need to inform us about plausible processes that have led to honey bee sociality. One hypothesis that is testable by studying current species is the reproductive ground plan hypothesis of social evolution in honey bees (Amdam, Norberg et al., 2004). The reproductive ground plan hypothesis proposes that the gonotrophic cycling of a hypothetical solitary ancestor has been coopted by social evolution to facilitate intra-group specialization: The temporal phenotypic plasticity between reproductive and non-reproductive states has been transformed into stable differences among individuals (Page and Amdam, 2007; Amdam and Page, 2010). The reproductive ground plan hypothesis predicts that phenotypic and genetic associations that

characterize the insect gonotrophic cycle will still be apparent as sets of correlated traits or syndromes in modern honey bees (Wang, Amdam et al., 2009; Graham, Munday et al., 2011). Below, I describe how the phenotypic and genetic architecture of the pollen hoarding syndrome (PHS) supports the reproductive ground plan hypothesis and what the results suggest more generally about behavioral syndromes and the prospects for selective breeding in honey bees.

Phenotypic Associations of the Pollen Hoarding Syndrome

Social insects, including the Western Honey Bee, *Apis mellifera* (L), protect and regulate their colony environment to rear brood and store food. The two principal food sources, pollen and nectar, are brought into the colony by foragers that may specialize on one of these resources or collect both (Fewell and Page, 1993). In contrast to stored nectar or honey (Fewell and Winston, 1996), the amount of pollen in the honey bee hive is tightly regulated (Fewell and Winston, 1992; Dreller, Page et al., 1999). Artificial selection has repeatedly demonstrated a substantial genetic component for the amount of pollen stored, or pollen hoarding (Hellmich, Kulinčević et al., 1985; Page and Fondrk, 1995). Pollen hoarding is a complex social trait because the amount of pollen stored in the hive is a function of available space, pollen collection, and consumption, and the active regulation occurs by adjusting the level of pollen collection by the foragers (Dreller and Tarpy, 2000). Therefore, it is not surprising to find artificial selection for pollen hoarding associated with corresponding changes in the probability of individual forager to collect pollen and their pollen load sizes (Fewell and Page 2000; Pankiw and Page 2001). Consistently, workers from the strain selected for high pollen hoarding performed more recruitment dances for pollen sources and also follow such dances more (Waddington, Nelson et al., 1998; Page, Rueppell et al., 2012).

However, detailed studies of the high and low pollen hoarding strains have revealed a number of correlated changes in other traits with less obvious connections to pollen hoarding (Page, Rueppell et al., 2012). These include behavioral traits, such as an earlier transition from in-hive to foraging activities (Pankiw and Page, 2001), higher locomotor activity after emergence (Humphries, Fondrk et al., 2005), increased sensitivity to several stimuli (Tsuruda and Page, 2009), and increased learning performance (Scheiner, Page et al., 2001) in the strain selected for high pollen hoarding. Workers from this strain are also smaller (Linksvayer, Fondrk et al., 2009), die earlier (Amdam, Nilsen et al., 2007; Rueppell, Bachelier et al., 2007), complete pupation at a later age (Amdam, Page et al., 2010), and have larger ovaries (Amdam, Csondes et al., 2006). At the physiological level, the high pollen hoarding workers differ in brain biochemistry (Humphries, Muller et al., 2003) and the endocrine dynamics of vitellogenin, juvenile hormone, and ecdysteroids (Amdam, Nilsen et al., 2007; Page and Amdam, 2007; Amdam, Page et al., 2010).

A substantial part of the phenotypic trait associations of the PHS has also been found in unselected honey bees in the commercial North American bee population (Amdam, Csondes et al., 2006; Page, Rueppell et al., 2012), Africanized honey bees (Pankiw, 2003; Page, Rueppell et al., 2012), and in the closely related *Apis cerana* (Rueppell, Hunggims et al., 2008). Thus, the connections between the studied traits are robust, suggesting co-regulation

at a fundamental level (Amdam, Norberg et al., 2004). Presumably, the basic relation between nutrition and reproduction preceded the evolution of social honey bees and has been sophisticated to regulate certain aspects of their social organization (Toth and Robinson, 2007). This concept of cooption of reproductive regulatory modules by social evolution has been proposed as the reproductive ground plan hypothesis of social evolution in honey bees (Amdam, Norberg et al., 2004; Amdam and Page, 2010).

Genetic Architecture of the Pollen Hoarding Syndrome

The phenotypic studies of the PHS and associated phenotypes have been accompanied from their beginning by genetic analyses, yielding an understanding of the genetic architecture of the behavioral syndrome from quantitative trait loci mapping, transcriptome analyses, and candidate gene studies.

Quantitative Trait Loci

The first detailed genetic study of the pollen hoarding trait revealed two quantitative trait loci (QTL: *pln1* and *pln2*) that together explained 59 % of the pollen hoarding variation in a backcross between the selected strains (Hunt, Page Jr. et al., 1995). In a second mapping study of a cross between these selected strains after several more generations confirmed one of these QTL and identified another (*pln3*) explaining 10% of the observed variation (Page, Fondrk et al., 2000). Finally, a third study confirmed *pln1*, *pln2*, and *pln3* by demonstrating an effect on individual foraging behavior, reported a fourth QTL (*pln4*), and demonstrated complex interactions between these loci (Rüppell, Pankiw et al., 2004).

As predicted based on the phenotypic associations, the *pln* QTL also were shown to have pleiotropic effects on other aspects of the PHS. Either alone or interactively, *pln1–pln3* affected the age of first foraging (Rueppell, Pankiw et al., 2004) and all four *pln* QTL had an effect on the sucrose responsiveness of honey bees (Rueppell, Chandra et al., 2006). Most importantly for the reproductive ground plan hypothesis, ovary size of worker honey bees was affected by *pln2*, *pln3* (Wang, Amdam et al., 2009), and *pln4* (Ihle, Rueppell et al., in preparation), and hormonal dynamics were affected by *pln3* (Ihle, Rueppell et al., in preparation). In all instances, mapping studies of these other traits have revealed additional QTL that usually exceeded the pleiotropic effects of the *pln* QTL in the specific crosses of the high and low pollen hoarding strains: Three additional QTL for the age of first foraging were identified (Rueppell, Pankiw et al., 2004; Rueppell, 2009), one for sucrose responsiveness (Rueppell, Chandra et al., 2006), and three for worker ovary size (Rueppell, Metheny et al., 2011). One QTL for the age of first foraging (*aff2*) also showed an effect on worker ovary size and two worker ovary size QTL showed effects on hormonal dynamics (Ihle, Rueppell et al., in preparation). Single effects of the *pln3* and *aff3* QTL on the life expectancy of honey bee workers have also been demonstrated (Dixon, McQuage et al., 2012). However, most conceivable tests for pleiotropy at the QTL level remain to be explored.

Outside the selected pollen hoarding strains, *pln1* was confirmed to affect pollen hoarding in a cross between European and Africanized honey bees, (Page, Fondrk et al., 2000). Analyses of two parallel backcrosses between these two populations in a later experiment have further

demonstrated overlap between the previously identified behavioral QTL and QTL for worker ovary size: Markers for two of the three age of first foraging QTL (*aff2* and *aff4*) and two of the four pollen hoarding QTL (*pln1* and *pln2*) showed significant effects on ovary size (Graham, Munday et al., 2011). The high genetic overlap between reproductive and social traits supports the reproductive ground plan hypothesis of social evolution in honey bees (Wang, Amdam et al., 2009; Graham, Munday et al., 2011) and suggests a partially overlapping genetic architecture of the different aspects of the PHS with a set of common core regulators (Page, Rueppell et al., 2012).

The identified QTL do not represent specific genes but designate genome intervals that contain positional candidate genes (Hunt, Amdam et al., 2007). The exceptionally high recombination rate of the honey bee genome typically results in small QTL regions (Hunt, Amdam et al., 2007), facilitating the identification of the genetic element that is responsible for each QTL effect. The PHS is one of very few examples where the connection from QTL to candidate genes has been made (see below). However, this is not true for all QTL and complementary data, such as gene expression patterns, are a powerful complement to QTL maps.

Transcriptome Studies

Complementary to QTL mapping, the study of transcriptome patterns generates functional candidate genes for complex traits that could be cause or consequence of the phenotype studied. Among the traits of the PHS, the gene expression changes that are related to the transition from in-hive tasks to foraging have been analyzed in most detail. Multiple gene expression signatures for the behavioral transition from in-hive tasks to foraging exist (Whitfield, Ben-Shahar et al., 2006) and some of the central functional candidate genes (e.g. the MAP-kinase *ERK7*) have also been suggested by the *aff* QTL (Rueppell, 2009). The PHS links into the central endocrine core of the regulation of this life history transition with juvenile hormone and vitellogenin co-regulation, affecting insulin-like signaling (Amdam, Nilsen et al., 2007; Nelson, Ihle et al., 2007; Nilsen, Ihle et al., 2011). However, the transcriptome studies provide little corroborating evidence for an involvement of these endocrine regulators.

The second trait of the PHS that has been studied by transcriptomic approaches is the ovary. Representational difference analysis of the developing ovary in worker and queen destined larvae revealed a small number of differentially expressed genes (Humann and Hartfelder, 2011). Two of these transcripts were long non-coding RNAs that are located near the *aff2* QTL that has repeatedly exhibited an influence on worker ovary size (Graham, Munday et al., 2011; Ihle, Rueppell et al., in preparation). However, sequence analysis of a part of both genes failed to identify allelic variants in the original mapping population that may be responsible for the QTL effects (B. Vannasane and O. Rueppell, unpublished data). Transcriptome comparisons of the adult worker ovary between the high and low pollen hoarding strains revealed more than 2000 differentially expressed genes with significant overlap to a similar number differentially expressed genes in the honey bee brain in response to hormonal treatment and the transition from in-hive to forager status (Wang, Kocher et al., 2012). Although these numbers of differentially expressed transcripts are frustratingly large

in the context of searching for singular candidate genes, overlap between multiple studies (Whitfield, Ben-Shahar et al., 2006) and the combination of positional and functional candidates improve the prospects of finding the underlying molecular causes of the observable social phenotypes.

Candidate Genes

Informed by the genomic studies (Hunt, Amdam et al., 2007) and theoretical considerations (Amdam, Norberg et al., 2004), several specific genes have been studied in the context of the PHS. The vitellogenin gene encodes a principal egg yolk protein that has been co-opted by social evolution to fulfill additional functions (Amdam, Norberg et al., 2003). Although the vitellogenin gene is not located in any of the QTL intervals, it has been studied in great detail and RNAi-mediated gene knockdown experiments show extensive effects on central traits of the PHS, such as gustatory responsiveness, foraging specialization, and the age of transitioning to foraging activity (Amdam, Norberg et al., 2006; Amdam, Nilsen et al., 2007; Ihle, Page et al., 2010). Vitellogenin effects on the PHS are not localizing to the described QTL, suggesting that vitellogenin levels may be controlled by segregating genetic variation in other genome locations.

Multiple genes in the insulin-like signaling pathway have been identified as candidate genes for the *pln* QTL (Hunt, Amdam et al., 2007), and three genes have been experimentally associated with the PHS: Expression differences of nuclear hormone receptor homolog *HR46* and phosphoinositide-dependent kinase-1 (*PDK1*) were associated with the selected strains and directly with worker ovary size (Wang, Amdam et al., 2009). Additionally, down-regulation of the insulin receptor substrate (*IRS*) causes honey bee foragers to collect more pollen relative to nectar without affecting sucrose responsiveness (Wang, Mutti et al., 2010). Vitellogenin levels affect insulin-like signaling in honey bees (Nilsen, Ihle et al., 2011) and the inverse seems also likely because insulin-like signaling affects juvenile hormone (Mutti, Dolezal et al., 2011), which may decrease vitellogenin levels (Page, Rueppell et al., 2012).

Conclusions

The PHS relates to many behavioral and life history traits in honey bees although its initial study began with a specific behavioral trait, pollen hoarding. The artificial selection program generating the high and low pollen hoarding lines has proven to be a powerful investigative tool (Conner, 2003). Particularly, the search for mechanisms explaining this specific phenotype led to the discovery that numerous other traits are involved, resulting in the best characterization of a behavioral syndrome in social insects so far. The mechanistic emphasis of the described studies has resulted in an integrative understanding of the proximate causation of the PHS (Page and Amdam, 2007; Page, Rueppell et al., 2012). However, the fitness consequences of the coordinated specialization of honey bee workers within colonies remain to be explored. The PHS involves numerous developmental, physiological, behavioral, morphological, and life history traits (Page, Rueppell et al., 2012). Specifically, the connections between reproductive physiology and behavioral specialization have been predicted by the reproductive ground plan hypothesis and the direct phenotypic and genetic

correlations support the reproductive ground plan hypothesis of social evolution in honey bees. The reproductive ground plan hypothesis may exemplify a more general pattern of social evolution by modification of fundamental, preexisting life history regulators. The consistency and number of relations among traits and genetic effects is remarkable, given that honey bees cannot be inbred and all complex traits of the PHS depend on the interaction of the current genetic variation at multiple loci with a changing environment.

Implications for Studies of Behavioral Syndromes

The studies of the PHS at the phenotypic and genetic level have demonstrated a central role for reproductive physiology in social evolution (Amdam, Csondes et al., 2006; Page and Amdam, 2007). Even though the PHS initially was described as a set of correlated behavioral traits (Page and Fondrk, 1995; Page and Erber, 2002), the underlying hormonal differences affect also development, physiology, and life history (Page, Rueppell et al., 2012). Studies of the PHS exemplify the connection between different levels of biological organization from molecular differences in single genetic regulators to colony-level resource allocation changes. Thus, the PHS may also be conceptualized as a life history syndrome instead of a behavioral syndrome. This conclusion prompts the general question how much the concept of behavioral syndromes differs from the concept of established syndromes in ecology and evolution (Pianka, 1970). Life history syndromes inevitably involve correlated behavioral traits. It should be realized that behavior integrates with most other aspects of biology, such as reproductive and nutritional physiology (Page and Amdam, 2007; Toth and Robinson, 2007). The study of behavioral syndromes that fail to make these connections will only represent an incremental advance over the analysis of single behavioral traits in behavioral ecology.

The growing understanding of the genetic architectures and underlying mechanisms of animal behavior needs to be integrated with phenotypic studies measuring correlations among different behavioral contexts. The relatively weak but significant behavioral correlations that are characterized in many studies of behavioral syndromes (Zsolt Garamszegi, Marko et al., 2012) suggest partially overlapping genetic architectures that are similar to the PHS (Rueppell, Pankiw et al., 2004; Rueppell, Chandra et al., 2006; Graham, Munday et al., 2011) and can be explored to identify central control elements (Hunt, Amdam et al., 2007; Wang, Amdam et al., 2009; Page, Rueppell et al., 2012). Due to its complexity and plasticity, natural behavior is more challenging to understand than most other kinds of traits, but the example of the PHS provides a cause for optimism beyond the study of social insects.

Implications for Selective Breeding for Honey Bee Health

Pleiotropy and genetic covariance among traits may prevent optimization of single traits by natural or artificial selection. The large number of traits integrated in the PHS and complex correlations across castes (Rueppell, Fondrk et al., 2006; Page, Fondrk et al., 2012) suggest that fundamental processes underlie the PHS that prevents independent evolution. Alternatively, the coordinated multi-trait evolution of the different members of the honey bee colonies may have been beneficial, leading to trait linkage and pleiotropy (Lande, 1984).

Most traits of the PHS are of little direct interest to the applied problems of declining honey bee health (vanEngelsdorp and Meixner, 2010) and selective breeding to increase the quality of honey bees for pollination and general apiculture (Rinderer, Harris et al., 2010). Disease syndromes, such as colony collapse disorder (vanEngelsdorp, Evans et al., 2009) or brood disease syndrome (vanEngelsdorp, Tarpy et al., 2013), also combine multiple disease symptoms but are conceptually unrelated to life history and behavioral syndromes discussed above. Nevertheless, the study of the PHS and the artificially selected high and low pollen hoarding strains for over twenty years (Page, Rueppell et al., 2012) provides important lessons for the current efforts to identify genes for disease resistance and apply this knowledge in selective breeding programs. One direct lesson is the potential for individual drone selection via artificial insemination based on male-female trait correlations (Rueppell, Fondrk et al., 2006; Page, Fondrk et al., 2012).

Honey bees have been bred for resistance against the ectoparasitic *Varroa* mite and commercially available stocks in the U.S. include the Russian honey bees (Bourgeois and Rinderer, 2009), the Minnesota Hygienic honey bees (Spivak, Reuter et al., 2009), and the *Varroa* Sensitive Hygiene stock (Harris, 2007). These selection programs have all been successful, relying on different selection and breeding strategies. Although the principal mechanism of resistance of at least two of these stocks appears to be hygienic behavior (Harris, 2007), no “hygienic syndrome” has emerged and multiple mechanisms of resistance are possible (Rinderer, Harris et al., 2010). The existence of a “hygienic syndrome” cannot be excluded and a detailed characterization of these mite-resistant stocks with respect to a wider array of life history and behavioral traits remains to be performed. A direct test failed to link hygienic behavior to the PHS (Goode, Huber et al., 2006), even though Africanized honey bees combine increased mite resistance (Camazine, 1986) with multiple aspects of the PHS (Pankiw, 2003; Graham, Munday et al., 2011).

In order to enable marker assisted selection (Rinderer, Harris et al., 2010), QTL studies of the hygienic behavior of the Minnesota Hygienic (Lapidge, Oldroyd et al., 2002; Oxley, Spivak et al., 2010) and the *Varroa* Sensitive Hygiene stocks (Tsuruda, Harris et al., 2012) have been conducted. Additional QTL studies have targeted mite grooming behavior (Arechavaleta-Velasco, Alcala-Escamilla et al., 2012) and mite reproductive success (Behrens, Huang et al., 2011). A QTL search was also performed for resistance to chalkbrood (*Ascosphaera apis*) (Holloway, Sylvester et al., 2012). However, over 70 different honey bee diseases are known (Schmid-Hempel, 1998) and studies of the genetic architecture of honey bee resistance to most of them are still lacking, even though a genetic basis for resistance against most diseases probably exists (Kulincevic and Rothenbuhler, 1975; Bailey and Ball, 1991; Palmer and Oldroyd, 2003; Huang, Kryger et al., 2012).

The studies of the PHS have demonstrated that QTL effects can be relatively robust across time and different mapping populations (Page, Fondrk et al., 2000; Rueppell, Pankiw et al., 2004; Ihle, Rueppell et al., in preparation), which increases the prospects for marker associated selection of honey bees. However, the QTL studies of *Varroa* resistant stocks have not yielded consistent QTL and show no overlap with transcriptome analyses (Le Conte, Alaux et al., 2011; Tsuruda, Harris et al., 2012). A tighter control of breeding and the identification of causative SNPs may increase the prospects for consistent findings in honey

bee health research and enable marker assisted selection. Marker assisted selection is a highly desirable breeding strategy in honey bees, given the practical limits to breeding and numerous desirable traits that could be selected for in addition to disease resistance (Laidlaw and Page, 1997). Genetic studies of the PHS have revealed also pleiotropy and epistasis (Rüppell, Pankiw et al., 2004), non-additivity and genetic heterogeneity (Linksvayer, Rueppell et al., 2009; Graham, Munday et al., 2011), and complex trait relations (Page, Fondrk et al., 2012), which may limit the success of phenotypic selection. In contrast, marker assisted selection may increase long-term breeding success by enabling breeders to maintain and combine specific genetic variants and to cross “adaptive valleys” from one local optimum to another on the selective landscape (O'Hagan, Knowles et al., 2012).

The prospects for multi-trait selection also relate to another central question for the field of honey bee health research raised by studies of the PHS: How much overlap in the resistance to different honey bee diseases exists and is there a “healthy” syndrome in honey bees? Hygienic behavior counteracts a series of brood diseases (Spivak and Gilliam, 1998) but how other desirable traits relate to this behavior and to each other remains to be studied. The phenotypic and genomic studies of trait relations are important to determine the prospects for combined selection on multiple traits. Comparisons between European and Africanized honey bees suggest potential trade-offs between disease resistance and less desirable behavioral traits, such as defensiveness. However, it is unclear whether these traits are genetically linked or could be dissociated by selective breeding. Studies of the PHS have shown that correlated sets of complex traits most likely have a partially overlapping genetic architecture consisting of central and peripheral regulators. Depending on the trait correlations and the genetic architecture, either the central or the peripheral regulators will present the basis for progress during social evolution and for selective breeding.

Acknowledgments

I thank David Tarpy and Stan Schneider for the invitation to write this review. I am greatly indebted to Robert Page for introducing me to many fascinating problems that relate to pollen hoarding behavior in honey bees. I would also like to acknowledge the financial support of the National Institutes of Health grant (R15GM102753), the United States Department of Agriculture (2010-65104-20533), and the North American Pollinator Protection Campaign.

References

- Agrawal AF, Stinchcombe JR. How much do genetic covariances alter the rate of adaptation? *Proc R Soc B-Biol Sci.* 2009; 276(1659):1183–1191.
- Amdam GV, Csondes A, et al. Complex social behaviour derived from maternal reproductive traits. *Nature.* 2006; 439:76–78. [PubMed: 16397498]
- Amdam GV, Nilsen KA, et al. Variation in endocrine signaling underlies variation in social life history. *Am Nat.* 2007; 170(1):37–46. [PubMed: 17853990]
- Amdam GV, Norberg K, et al. Reproductive ground plan may mediate colony-level selection effects on individual foraging behavior in honey bees. *Proc Natl Acad Sci U S A.* 2004; 101(31):11350–11355. [PubMed: 15277665]
- Amdam GV, Norberg K, et al. Social exploitation of vitellogenin. *Proc Natl Acad Sci U S A.* 2003; 100(4):1799–1802. [PubMed: 12566563]
- Amdam GV, Norberg K, et al. Downregulation of vitellogenin gene activity increases the gustatory responsiveness of honey bee workers (*Apis mellifera*). *Behavioral Brain Research.* 2006; 169(2): 201–205.

- Amdam GV, Page RE. The developmental genetics and physiology of honeybee societies. *Anim Behav.* 2010; 79(5):973–980. [PubMed: 20514137]
- Amdam GV, Page RE, et al. Hormone response to bidirectional selection on social behavior. *Evol Dev.* 2010; 12(5):428–436. [PubMed: 20883212]
- Arechavala-Velasco ME, Alcala-Escamilla K, et al. Fine-scale linkage mapping reveals a small set of candidate genes influencing honey bee grooming behavior in response to *Varroa* mites. *PLoS One.* 2012; 7(11):e47269. [PubMed: 23133594]
- Arias MC, Sheppard WS. Phylogenetic relationships of honey bees (Hymenoptera:Apinae:Apini) inferred from nuclear and mitochondrial DNA sequence data. *Mol Phylogenet Evol.* 2005; 37(1): 25–35. [PubMed: 16182149]
- Bailey, L.; Ball, BV. *Honey Bee Pathology*. Academic Press; London, UK: 1991.
- Behrens D, Huang Q, et al. Three QTL in the honey bee *Apis mellifera* L. suppress reproduction of the parasitic mite *Varroa destructor*. *Ecol Evol.* 2011; 1(4):451–458. [PubMed: 22393513]
- Bourgeois AL, Rinderer TE. Genetic characterization of Russian Honey Bee stock selected for improved resistance to *Varroa destructor*. *J Econ Entomol.* 2009; 102(3):1233–1238. [PubMed: 19610443]
- Camazine S. Differential reproduction of the mite, *Varroa jacobsoni* (Mesostigmata, Varroidae), on Africanized and European honey bees (Hymenoptera, Apidae). *Ann Entomol Soc Am.* 1986; 79(5):801–803.
- Conner JK. Artificial selection: A powerful tool for ecologists. *Ecology.* 2003; 84(7):1650–1660.
- Delph LF, Arntz AM, et al. The genomic architecture of sexual dimorphism in the dioecious plant *Silene latifolia*. *Evolution.* 2010; 64(10):2873–2886. [PubMed: 20550575]
- Dixon LR, McQuage MR, et al. Pleiotropy of segregating genetic variants that affect honey bee worker life expectancy. *Exp Gerontol.* 2012; 47(8):631–637. [PubMed: 22664574]
- Dreller C, Page RE, et al. Regulation of pollen foraging in honeybee colonies: effects of young brood, stored pollen, and empty space. *Behav Ecol Sociobiol.* 1999; 45:227–233.
- Dreller C, Tarpy DR. Perception of the pollen need by foragers in a honeybee colony. *Anim Behav.* 2000; 59(1):91–96. [PubMed: 10640370]
- Engel MS. Fossil honey bees and evolution in the genus *Apis* (Hymenoptera : Apidae). *Apidologie.* 1998; 29(3):265–281.
- Fewell JH, Page RE. Genotypic variation in foraging responses to environmental stimuli by honey bees, *Apis mellifera*. *Experientia.* 1993; 49(12):1106–1112.
- Fewell JH, Winston ML. Colony state and regulation of pollen foraging in the honey-bee, *Apis mellifera*. *L Behav Ecol Sociobiol.* 1992; 30:387–393.
- Fewell JH, Winston ML. Regulation of nectar collection in relation to honey storage levels by honey bees, *Apis mellifera*. *Behav Ecol.* 1996; 7(3):286–291.
- Gadagkar R. The evolution of caste polymorphism in social insects: genetic release followed by diversifying evolution. *Journal of Genetics.* 1997; 76(3):167–179.
- Goode K, Huber Z, et al. Hygienic behavior of the honey bee (*Apis mellifera*) is independent of sucrose responsiveness and foraging ontogeny. *Horm Behav.* 2006; 49(3):391–397. [PubMed: 16225878]
- Graham AM, Munday MD, et al. Support for the reproductive ground plan hypothesis of social evolution and major QTL for ovary traits of Africanized worker honey bees (*Apis mellifera* L). *BMC Evol Biol.* 2011; 11:95. [PubMed: 21489230]
- Harris JW. Bees with *Varroa* Sensitive Hygiene preferentially remove mite infested pupae aged <= five days post capping. *Journal of Apicultural Research.* 2007; 46(3):134–139.
- Hellmich RL, Kulincevic JM, et al. Selection for high and low pollen-hoarding honey bees. *J Hered.* 1985; 76(3):155–158.
- Holloway B, Sylvester HA, et al. Association of single nucleotide polymorphisms to resistance to chalkbrood in *Apis mellifera*. *Journal of Apicultural Research.* 2012; 51(2):154–163.
- Huang Q, Kryger P, et al. Survival and immune response of drones of a Nosemosis tolerant honey bee strain towards *N. ceranae* infections. *J Invertebr Pathol.* 2012; 109(3):297–302. [PubMed: 22285444]

- Humann FC, Hartfelder K. Representational Difference Analysis (RDA) reveals differential expression of conserved as well as novel genes during caste-specific development of the honey bee (*Apis mellifera* L.) ovary. *Insect Biochem Molec.* 2011; 41(8):602–612.
- Humphries MA, Fondrk MK, et al. Locomotion and the pollen hoarding behavioral syndrome of the honeybee (*Apis mellifera* L.). *J Comp Physiol A.* 2005; 191(7):669–674.
- Humphries MA, Muller U, et al. PKA and PKC content in the honey bee central brain differs in genotypic strains with distinct foraging behavior. *J Comp Physiol A.* 2003; 189(7):555–562.
- Hunt GJ, Amdam GV, et al. Behavioral genomics of honeybee foraging and nest defense. *Naturwissenschaften.* 2007; 94(4):247–267. [PubMed: 17171388]
- Hunt GJ, Page RE Jr, et al. Major quantitative trait loci affecting honey bee foraging behavior. *Genetics.* 1995; 141:1537–1545. [PubMed: 8601492]
- Ihle KE, Page RE, et al. Genotype effect on regulation of behaviour by vitellogenin supports reproductive origin of honeybee foraging bias. *Anim Behav.* 2010; 79(5):1001–1006. [PubMed: 20454635]
- Ihle KE, Rueppell O, et al. QTL for ovary size and juvenile hormone response to Vg-RNAi knockdown. in preparation.
- Jandt JM, Bengston S, et al. Behavioural syndromes and social insects: personality at multiple levels. *Biol Rev Online.* 2013
- Kulincevic JM, Rothenbuhler WC. Selection for resistance and susceptibility to hairless-black syndrome in the honeybee. *J Invertebr Pathol.* 1975; 25(3):289–295. [PubMed: 1141705]
- Laidlaw, HH.; Page, RE. *Queen Rearing and Bee Breeding.* Wicwas Press; Cheshire, CT: 1997.
- Lande R. A quantitative genetic theory of life history evolution. *Ecology.* 1982; 63(3):607–615.
- Lande R. The genetic correlation between characters maintained by selection, linkage and inbreeding. *Genet Res.* 1984; 44(3):309–320. [PubMed: 6530140]
- Lapidge KL, Oldroyd BP, et al. Seven suggestive quantitative trait loci influence hygienic behavior of honey bees. *Naturwissenschaften.* 2002; 89(12):565–568. [PubMed: 12536279]
- Le Conte Y, Alaux C, et al. Social immunity in honeybees (*Apis mellifera*): transcriptome analysis of varroa-hygienic behaviour. *Insect Molecular Biology.* 2011; 20(3):399–408. [PubMed: 21435061]
- Linksvayer TA, Fondrk MK, et al. Honeybee social regulatory networks are shaped by colony-level selection. *Am Nat.* 2009; 173(3):E99–E107. [PubMed: 19140771]
- Linksvayer TA, Rueppell O, et al. The genetic basis of transgressive ovary size in honey bee workers. *Genetics.* 2009; 183:693–707. [PubMed: 19620393]
- Mutti NS, Dolezal AG, et al. IRS and TOR nutrient-signaling pathways act via juvenile hormone to influence honey bee caste fate. *J Exp Biol.* 2011; 214(23):3977–3984. [PubMed: 22071189]
- Nelson CM, Ihle KE, et al. The gene vitellogenin has multiple coordinating effects on social organization. *Plos Biology.* 2007; 5(3):e62. [PubMed: 17341131]
- Nilsen KA, Ihle KE, et al. Insulin-like peptide genes in honey bee fat body respond differently to manipulation of social behavioral physiology. *J Exp Biol.* 2011; 214(9):1488–1497. [PubMed: 21490257]
- O'Hagan S, Knowles J, et al. Exploiting genomic knowledge in optimising molecular breeding programmes: algorithms from evolutionary computing. *PLoS One.* 2012; 7(11)
- Oxley PR, Spivak M, et al. Six quantitative trait loci influence task thresholds for hygienic behaviour in honeybees (*Apis mellifera*). *Mol Ecol.* 2010; 19(7):1452–1461. [PubMed: 20298472]
- Page RE, Amdam GV. The making of a social insect: developmental architectures of social design. *Bioessays.* 2007; 29:334–343. [PubMed: 17373656]
- Page RE, Erber J. Levels of behavioral organization and the evolution of division of labor. *Naturwissenschaften.* 2002; 89:91–106. [PubMed: 12046634]
- Page RE, Fondrk MK. The effects of colony level selection on the social organization of honey bee (*Apis mellifera* L.) colonies - colony level components of pollen hoarding. *Behav Ecol Sociobiol.* 1995; 36(2):135–144.
- Page RE, Fondrk MK, et al. Genetic dissection of honeybee (*Apis mellifera* L.) foraging behavior. *J Hered.* 2000; 91(6):474–479. [PubMed: 11218085]

- Page RE, Fondrk MK, et al. Complex pleiotropy characterizes the pollen hoarding syndrome in honey bees (*Apis mellifera* L). *Behav Ecol Sociobiol.* 2012; 66(11):1459–1466. [PubMed: 23226916]
- Page RE, Rueppell O, et al. Genetics of reproduction and regulation of honeybee (*Apis mellifera* L.) social behavior. *Annual Review of Genetics.* 2012; 46:97–119.
- Palmer KA, Oldroyd BP. Evidence for intra-colonial genetic variance in resistance to American foulbrood of honey bees (*Apis mellifera*): further support for the parasite/pathogen hypothesis for the evolution of polyandry. *Naturwissenschaften.* 2003; 90(6):265–268. [PubMed: 12835837]
- Pankiw T. Directional change in a suite of foraging behaviors in tropical and temperate evolved honey bees (*Apis mellifera* L). *Behav Ecol Sociobiol.* 2003; 54(5):458–464.
- Pankiw T, Page RE. Genotype and colony environment affect honeybee (*Apis mellifera* L.) development and foraging behavior. *Behav Ecol Sociobiol.* 2001; 51:87–94.
- Pianka ER. On r- and K-selection. *Am Nat.* 1970; 104(940):592–597.
- Price T, Langen T. Evolution of correlated characters. *Trends Ecol Evol.* 1992; 7(9):307–310. [PubMed: 21236041]
- Rinderer TE, Harris JW, et al. Breeding for resistance to *Varroa destructor* in North America. *Apidologie.* 2010; 41(3):409–424.
- Roff, DA. Genomic insights into life history evolution. In: Flatt, T.; Heyland, A., editors. *Mechanisms of Life History Evolution.* Oxford University Press; Oxford, UK: 2011. p. 11-25.
- Rueppell O. Characterization of quantitative trait loci for the age of first foraging in honey bee workers. *Behavior Genetics.* 2009; 39:541–553. [PubMed: 19449161]
- Rueppell O, Bachelier C, et al. Regulation of life history determines lifespan of worker honey bees (*Apis mellifera* L). *Exp Gerontol.* 2007; 42:1020–1032. [PubMed: 17689041]
- Rueppell O, Chandra SBC, et al. The genetic architecture of sucrose responsiveness in the honey bee (*Apis mellifera* L). *Genetics.* 2006; 172:243–251. [PubMed: 16172502]
- Rueppell O, Fondrk MK, et al. Male maturation response to selection of the pollen-hoarding syndrome in honey bees (*Apis mellifera* L). *Animal Behavior.* 2006; 71:227–234.
- Rueppell O, Hunggins E, et al. Association between larger ovaries and pollen foraging in queenless *Apis cerana* workers supports the reproductive ground - plan hypothesis of social evolution. *J Insect Behav.* 2008; 21:317–321.
- Rueppell O, Metheny JD, et al. Genetic architecture of ovary size and asymmetry in European honeybee workers. *Heredity.* 2011; 106:894–903. [PubMed: 21048673]
- Rueppell O, Pankiw T, et al. The genetic architecture of the behavioral ontogeny of foraging in honey bee workers. *Genetics.* 2004; 167:1767–1779. [PubMed: 15342515]
- Rüppell O, Pankiw T, et al. Pleiotropy, epistasis and new QTL: the genetic architecture of honey bee foraging behavior. *J Hered.* 2004; 95:481–491. [PubMed: 15475393]
- Scheiner R, Page RE, et al. The effects of genotype, foraging role, and sucrose responsiveness on the tactile learning performance of honey bees (*Apis mellifera* L). *Neurobiology of Learning and Memory.* 2001; 76(2):138–150. [PubMed: 11502146]
- Schmid-Hempel, P. *Parasites in Social Insects.* Princeton University Press; Princeton, NJ: 1998.
- Sih A, Bell A, et al. Behavioral syndromes: an ecological and evolutionary overview. *Trends Ecol Evol.* 2004; 19:372–378. [PubMed: 16701288]
- Spivak M, Gilliam M. Hygienic behaviour of honey bees and its application for control of brood diseases and *Varroa* Part I. Hygienic behaviour and resistance to American foulbrood. *Bee World.* 1998; 79(3):124–134. 169–186.
- Spivak M, Reuter GS, et al. The future of the MN Hygienic Stock of bees is in good hands! *American Bee Journal.* 2009; 149(10):965–967.
- Stearns SC. Trade-offs in life-history evolution. *Functional Ecology.* 1989; 3(3):259–268.
- Toth AL, Robinson GE. Evo-devo and the evolution of social behavior. *Trends in Genetics.* 2007; 23(7):334–341. [PubMed: 17509723]
- Tsuruda JM, Harris JW, et al. High-resolution linkage analyses to identify genes that influence *Varroa* Sensitive Hygiene behavior in honey bees. *PLoS One.* 2012; 7(11)

- Tsuruda JM, Page RE. The effects of foraging role and genotype on light and sucrose responsiveness in honey bees (*Apis mellifera* L). Behavioural Brain Research. 2009; 205(1):132–137. [PubMed: 19643145]
- vanEngelsdorp D, Evans JD, et al. Colony Collapse Disorder: A descriptive study. PLoS ONE. 2009; 4(8):e6481. [PubMed: 19649264]
- vanEngelsdorp D, Meixner MD. A historical review of managed honey bee populations in Europe and the United States and the factors that may affect them. Journal of Invertebrate Pathology. 2010; 103:S80–S95. [PubMed: 19909973]
- vanEngelsdorp D, Tarpy DR, et al. Idiopathic brood disease syndrome and queen events as precursors of colony mortality in migratory beekeeping operations in the eastern United States. Prev Vet Med. 2013; 108(2-3):225–233. [PubMed: 22939774]
- Waddington KD, Nelson M, et al. Effects of pollen quality and genotype on the dance of foraging honey bees. Animal Behavior. 1998; 56:35–39.
- Wang Y, Amdam GV, et al. PDK1 and HR46 gene homologs tie social behavior to ovary signals. PLoS ONE. 2009; 4(4):e4899. [PubMed: 19340296]
- Wang Y, Kocher SD, et al. Regulation of behaviorally associated gene networks in worker honey bee ovaries. J Exp Biol. 2012; 215(1):124–134. [PubMed: 22162860]
- Wang Y, Mutti NS, et al. Down-regulation of honey bee IRS gene biases behavior toward food rich in protein. PLoS Genet. 2010; 6(4):e1000896. [PubMed: 20369023]
- Whitfield CW, Ben-Shahar Y, et al. Genomic dissection of behavioral maturation in the honey bee. Proc Natl Acad Sci U S A. 2006; 103(44):16068–16075. [PubMed: 17065327]
- Williams GC. Pleiotropy, natural selection, and the evolution of senescence. Evolution. 1957; 11:398–411.
- Zsolt Garamszegi L, Marko G, et al. A meta-analysis of correlated behaviours with implications for behavioural syndromes: mean effect size, publication bias, phylogenetic effects and the role of mediator variables. Evol Ecol. 2012; 26(5):1213–1235.