Diagnostic Capability of Biological Markers in Assessment of Obstructive Sleep Apnea: A Systematic Review and Meta-Analysis

Graziela De Luca Canto, DDS, MSc, PhD^{1,2}; Camila Pachêco-Pereira, DDS²; Secil Aydinöz, MD^{3,4}; Paul W. Major, DDS, MSc, FRCD(C)²; Carlos Flores-Mir, DDS, DSc, FRCD(C)²; David Gozal, MD⁴

¹Department of Dentistry, Federal University of Santa Catarina, Florianópolis, SC, Brazil; ²School of Dentistry, Faculty of Medicine and Dentistry, University of Alberta, Canada; ³GATA Haydarpasa Teaching Hospital, Istanbul, Turkey; ⁴Department of Pediatrics, University of Chicago, Chicago, IL

Objective: The purpose of this systematic review is to evaluate the diagnostic value of biological markers (exhaled breath condensate, blood, salivary and urinary) in the diagnosis of OSA in comparison to the gold standard of nocturnal PSG.

Methods: Studies that differentiated OSA from controls based on PSG results, without age restriction, were eligible for inclusion. The sample of selected studies could include studies in obese patients and with known cardiac disease. A detailed individual search strategy for each of the following bibliographic databases was developed: Cochrane, EMBASE, MEDLINE, PubMed, and LILACS. The references cited in these articles were also crosschecked and a partial grey literature search was undertaken using Google Scholar. The methodology of selected studies was evaluated using the 14-item Quality Assessment Tool for Diagnostic Accuracy Studies.

Results: After a two-step selection process, nine articles

were identified and subjected to qualitative and quantitative analyses. Among them, only one study conducted in children and one in adults found biomarkers that exhibit sufficiently satisfactory diagnostic accuracy that enables application as a diagnostic method for OSA.

Conclusion: Kallikrein-1, uromodulin, urocortin-3, and orosomucoid-1 when combined have enough accuracy to be an OSA diagnostic test in children. IL-6 and IL-10 plasma levels have potential to be good biomarkers in identifying or excluding the presence of OSA in adults.

Keywords: biological markers, diagnosis, sleep apnea syndromes, review

Citation: De Luca Canto G, Pachêco-Pereira C, Aydinöz S, Major PW, Flores-Mir C, Gozal D. Diagnostic capability of biological markers in assessment of obstructive sleep apnea: a systematic review and meta-analysis. *J Clin Sleep Med* 2015;11(1):27–36.

Obstructive sleep apnea (OSA) has become widely recognized as a potential cause of significant morbidity in both children and adults.^{1,2} OSA symptoms include habitual snoring and reporting of disturbed unrefreshing sleep, frequently accompanied by excessive daytime sleepiness, and daytime neurobehavioral problems.³ The increasing understanding, awareness and familiarity with OSA has resulted in an ever expanding spectrum of OSA-associated morbidities that encompasses not only the central nervous system (cognitive, mood disturbances, and behavioral deficits), but affects also many other organ systems, ultimately imposing substantial increases in healthcare costs, as well as adverse outcomes.^{4–7}

Among the prototypic risk factors associated with OSA, adenotonsillar hypertrophy, obesity, craniofacial and anatomical anomalies, and neuromuscular disorders, seemingly interact to a greater or lesser extent among patients, leading to the putative assumption that multiple clinical phenotypes exist and potentially merit divergent therapeutic approaches better tailored at the constellation of pathophysiological mechanisms leading to OSA in these clinical clusters.³ The prevalence of OSA is markedly variable both during childhood (1% to 5%) and during adulthood (4% to 15%), with major contributions of age, gender, and ethnicity.^{1,8–11} However, it is clear that independently of

BRIEF SUMMARY

Current Knowledge/Study Rationale: The purpose of this systematic review was to evaluate the diagnostic properties of markers in biological samples, such as in exhaled breath condensate, blood, saliva, and urine, and compare their predictive characteristics to the gold standard in the diagnosis of OSA—nocturnal PSG.

Study Impact: A substantial number of studies have been published in the literature in the quest for diagnostic biomarkers of OSA in both children and adults; however, most of the explored approaches do not identify definitive biomarkers, and only a small number of candidates appears promising and merits further research.

whether we consider the lowest or the highest estimated prevalence reported for any population, OSA is a frequent condition that imposes a high degree of disease burden, thereby requiring timely diagnosis and effective treatment.

An overnight in-laboratory polysomnographic evaluation (PSG) remains the gold standard diagnostic method for OSA at any age.^{3,12} Unfortunately, overnight PSGs are onerous, labor-intensive, may impose substantial inconvenience to the child and caretakers, and are variably accessible around the world. Waiting time between referral for evaluation to diagnosis may commonly take 3–6 months across the United States and even

longer elsewhere.¹³ Although the PSG is employed as the gold standard for diagnosing the vast majority of sleep disorders, the relative complexity of PSG application and the inherent costs associated with PSG has spurred the quest for alternative diagnostic methods.¹³ Among these, simple approaches such as questionnaires with or without medical history and physical examination, audiotaping, videotaping, pulse oximetry, abbreviated polysomnography (aPSG), home-based polygraphy, or multichannel recordings have all been assessed, albeit with variable success.^{12,14–18} However, among the alternative diagnostic tools, special interest has recently centered on the identification of biomarkers.

A biomarker is a “biological molecule found in blood, other body fluids, or tissues that is a sign of a normal or abnormal processes, or of a condition or disease.”¹³ Gene expression arrays have revealed significant and reproducible changes in a restricted number of genes that could enable discriminatory ability in the recognition of OSA. Similarly, a number of serum and urinary proteins have been identified that display favorable significant receiver-operator properties towards the diagnosis of OSA.¹³ Provided that acceptable sensitivity and specificity are achieved, a unique set of disease biomarkers would enable greatly simplified, user-friendly, and context-relevant approaches to the diagnosis of OSA in the future.¹⁹ Over the last 14 years, a substantial number of studies have tackled the identification of an ideal biomarker for OSA, and although, there is still no simple and useful disease marker panel for OSA available, considerable progress has been accomplished and merits critical review and scrutiny.¹⁹ Therefore the purpose of this systematic review was to critically evaluate the diagnostic properties of markers in biological samples, such as in exhaled breath condensate (EBC), blood, saliva, and urine, and compare their predictive characteristics to the gold standard—nocturnal PSG. We further aimed to formulate potential future exploratory research directions aiming at advancing this promising area of clinical translation in sleep medicine.

METHODS

This systematic review was done adhering to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses PRISMA Checklist.²⁰

Diagnostic Terminology

All terms that mean obstructive sleep apnea (OSA), including sleep disordered breathing (SDB), sleep-related breathing disorder (SRBD) and obstructive sleep apnea syndrome (OSAS) were standardized as OSA.

Protocol and Registration

The systematic review protocol was registered at the international prospective register of systematic reviews (PROSPERO). The number of register is CRD42014007427.

Study Design

A systematic review of human studies that evaluated the diagnostic value of biological markers (blood, EBC, salivary, and urinary) in the diagnosis of OSA was undertaken.

Eligibility Criteria

Inclusion Criteria

Studies that differentiated the OSA group from controls based on full PSG results, without age restriction, were eligible for inclusion. The sample of selected studies could include studies in obese patients and in those with known cardiac disease.

Retained articles included only those studies whose primary objective was to identify biomarkers in subjects with OSA confirmed by overnight PSG. Only studies in English, Portuguese and Spanish languages were considered.

Exclusion Criteria

Reviews, letters, conference abstracts, and personal opinions were not considered.

Studies using daytime PSG, home-based PSG or multichannel polygraphic recordings were also excluded. Studies using biomarkers to detect the presence of OSA-associated morbidities (cognitive or behavioral deficits, excessive daytime sleepiness, cardiovascular or metabolic end-organ dysfunction) were excluded. In addition, studies in which the clinical cohort included craniofacial, genetic syndromes, neuromuscular diseases, or patients with a primary disease for which OSA prevalence is being investigated, such as patients with kidney disease or rheumatologic conditions were also discarded. In phase 2, we excluded studies that did not report sensitivity and specificity or in which the data presented did not enable these assessments to be extrapolated.

Information Sources

Detailed individual search strategies for each of the following bibliographic databases were developed: Cochrane, EMBASE, MEDLINE, PubMed, and LILACS. A partial grey literature search was taken using Google Scholar. The end search date was January 3, 2014, and an updated search was completed on March 20, 2014, across all databases. The references cited in the selected articles were also checked for any incremental references that could have been inadvertently omitted during the electronic database searches.

Search

Appropriate truncation and word combinations were selected and adapted for each database search (see **Appendix 1**). All references were managed by reference manager software (RefWorks-COS, ProQuest, Bethesda, MD), and duplicate hits were removed.

Study Selection

The selection was completed in 2 phases.

In phase 1, two reviewers independently reviewed the titles and abstracts of all identified electronic database citations (GDL and CPP). The following criteria were applied to select studies in phase 1: studies with an objective of identifying biomarkers in subjects with OSA confirmed by full overnight PSG. A third author (SA) was involved when disagreements emerged among the 2 initial evaluators. Any studies that did not fulfill the inclusion criteria were discarded.

In phase 2, the following selection criteria were applied to the full articles to confirm their eligibility: only studies that reported sensitivity and specificity or in which the data presented

enabled these diagnostic assessments to be extrapolated were selected. The same 2 reviewers (GDL and CPP) independently participated in phase 2. The reference list of all included articles was critically assessed by one examiner (GDL). The articles that were selected were then read by both examiners (GDL and CPP). Any disagreement in either phase was resolved by discussion and mutual agreement among the 3 reviewers (GDL, CPP, SA). A fourth author with extensive experience in sleep medicine and biomarker discovery (DG) was involved when controversy arose in the process of reaching a final decision. Final selection was always based on the full-text of the publication.

Data Collection Process

One author (GDL) collected the required information from the selected articles. A second author (CPP) crosschecked all the collected information and confirmed its accuracy. Again, any disagreement in either phase was resolved by discussion and mutual agreement among the 3 reviewers (GDL, CPP, SA). The fourth author was involved as required, to enable formulation of the final decision (DG).

Data Items

For all of the included studies the following information was recorded: author(s), year of publication, country, sample size, age, type of biomarkers, apnea hypopnea index used to define OSA from the PSG, name of biomarkers, and results (including sensitivity and specificity). If the required data were not complete, attempts were made to contact the authors to retrieve the missing information.

Risk of Bias in Individual Studies

The methodology of selected studies was evaluated using the 14-item Quality Assessment Tool for Diagnostic Accuracy Studies (QUADAS).²¹ Two reviewers (GDL and CPP) scored each item as “yes,” “no,” or “unclear” and assessed independently the quality of each included study. Disagreement between the 2 reviewers was resolved by a third reviewer (CFM).

Summary Measures

Sensitivity and specificity of biomarkers as diagnostic tests against PSG were considered as the main outcomes.

Synthesis of Results

The diagnostic capability of the identified biomarkers against PSG was combined through a meta-analysis following the appropriate Cochrane guidelines.²² Review Manager 5.2 (RevMan 5.2, The Nordic Cochrane Centre, Copenhagen, Denmark) was used to construct receiver operating characteristic (ROC) graphs and forest plots as part of the meta-analysis. Some of the required data were calculated by the authors.

Risk of Bias across Studies

To decrease the heterogeneity, the studies were separated in 3 groups according to age (children or adults) and biomarker characteristics (single or combined biomarkers).

Additional Analyses

Additional analysis was performed using positive predictive value (PPV), negative predictive value (NPV), positive

likelihood ratio (LR+), negative likelihood Ratio (LR-), diagnostic odds ratio (DOR), and Youden's Index. The cutoff values used to interpret these data are presented in **Appendix 2**.

RESULTS

Study Selection

A flowchart describing the process of identification, inclusion, and exclusion of studies is shown in **Figure 1**. A total of 141 articles were retrieved during phase 1 selection. Thereafter, 132 studies were excluded due to different reasons (see **Appendix 3**). Only 9 articles were finally included in the qualitative and quantitative synthesis. Eight of those^{23–31} were initially identified from the main electronic search; only one³¹ was directly received from expert sources.

Study Characteristics

From the 9 selected studies, 4 were conducted in children^{23,24,30,31} and 5 in adults.^{25–29} The studies in children were conducted in 2 different countries: Hungary²³ and United States.^{24,30,31} The following PSG-based criteria were used for OSA: AHI $\geq$ 1/h TST,²³ AHI $\geq$ 2/h TST,³¹ AHI $>$ 2/h TST,²⁴ AHI $>$ 5/h TST.³⁰ Two of these studies tested urinary biomarkers against PSG,^{24,31} one tested blood-based biomarkers,³⁰ and one evaluated EBC.²³ The sample size ranged from 28 to 120 subjects.^{23,24} A summary of the study descriptive characteristics can be found in **Table 1**.

The studies conducted in adults were conducted in Brazil,²⁵ China,²⁸ Germany,²⁹ Thailand,²⁶ and Turkey.²⁷ The following definitions for OSA were used: AHI $>$ 5/h TST,²⁹ AHI $\geq$ 5/h TST,^{26–28} and AHI $\geq$ 15/h TST.²⁵ All of the studies^{25,27–29} evaluated blood biomarkers. One of the studies appraised both blood and EBC.²⁶ The sample size ranged from 63²⁸ to 1,021²⁵ participants. A summary of the study descriptive characteristics can be found in **Table 2**.

Risk of Bias within Studies

The studies were very homogeneous—all had high methodological quality, even though none of the studies fulfilled all methodological quality criteria. In 8 studies, the QUADAS²¹ criteria were fulfilled in 78.6%. In another study,²⁵ QUADAS criteria were met in 85.7% (**Appendix 4**).

Results of Individual Studies

Although the studies used different types of biomarkers and reported different sensitivity and specificity all 9 articles concluded that biomarkers had the capacity to correctly classify OSA and non-OSA subjects.

Synthesis of Results

To improve our interpretation of results, the studies were clustered in 3 groups, according to the sample and the index test: using only one biomarker in children or adults, and using combined biomarkers in children. Diagnostic tables were constructed using the data extracted from each article (**Tables 3, 4, 5**). In these tables, all accuracy measurements (sensitivity, specificity, PPV, NPV, LR+, LR-, DOR, and Youden's Index) are presented. Some studies^{25–27} provided more than one accuracy measurement. Therefore those findings are reported twice in the same table. From the 4 studies

Figure 1—Flow diagram of literature search and selection criteria.

Adapted from PRISMA.

conducted in children, a total of 258 subjects were assessed. From the 5 studies conducted in adults, 1,458 subjects were evaluated. The total sample for this meta-analysis was 1,716 subjects.

The diagnostic accuracy (sensitivity, specificity, and 95% confidence interval) of the studies included in a meta-analysis is shown in **Figures 2** and **3**. The sensitivity and specificity for different selected studies varied substantially from 43% to 100%, and from 45% to 100%, respectively. Only 5 studies reported excellent sensitivity: Li et al. (100%),²⁶ Gozal et al. (95%),²⁴ Shah et al. (93%),³⁰ Guo et al. (91%),²⁸ and Kheirandish-Gozal et al. (82%).³¹ From these 5 studies, only Gozal et al.²⁴ and Li et al.²⁶ also reported excellent specificity (both 97%).

Risk of Bias across Studies

The main methodological limitations of the studies were related to poor reporting for items 1 (Was the spectrum of patients representative of the patients who will receive the test in clinical practice?), and for items 10 and 11 (blind interpretation of the reference and index test). The complete item list analyzed is presented in **Appendix 3**.

Additional Analysis

Only one pediatric²⁴ and one adult study²⁶ reported LR values considered excellent DTA. Thioredoxin (TRX),²⁸ kalikrein-1,²⁴ uromodulin,²⁴ urocortin-3,²⁴ orosomucoid-1,²⁴

Table 1—Summary of study descriptive characteristics of included studies (children).

Year	Author	Country	Cases	Controls	Age range, years	Type of Biomarker	OSA* diagnostic criteria at PSG	Biomarker	Main Conclusion
2006	Shah et al. ³⁰	United States	OSA (n = 20)	Non-OSA with HS (n = 20)	3–12	Blood	AHI > 5	Proteomic patterns	Proteomic profiling of serum samples in OSA children revealed differential expression of circulating proteins that may provide useful future diagnostic approaches.
2009	Gozal et al. ²⁴	United States	Children with HS and suspected OSA: (n = 60/32 male)	Children from community (n = 30/16 male) + Children with HS and suspected OSA: PS (n = 30/16 male)	2–9	Urine	OAI > and/or OAI > 2	Kallikrein-1 Uromodulin Urocortin-3 Orosomucoid-1	Kallikrein-1, uromodulin, urocortin-3, and orosomucoid-1 may be potential biomarkers to identify OSA and non-OSA children.
2013	Benedek et al. ²³	Hungary	OSA (n = 18)	Non-OSA with HS (n = 10)	8.5	EBC	AHI ≥ 1	VOCs mixtures	VOCs mixtures can be distinguished in OSA children from controls.
2013	Kheirandish-Gozal et al. ³¹	United States	OSA (n = 50) Matched with controls for age, sex, BMI and ethnicity	Non-OSA (n = 20)	3–12	Urine	AHI ≥ 2	Urinary Neurotransmitters	Combinatorial approaches of selected neurotransmitters in urine may predict OSA in children.

*All terms that mean obstructive sleep apnea (SDB, SRDB, OSAS) were standardized as OSA. AHI, apnea-hypopnea index; BMI, body mass index; EBC, exhaled breath condensate; GABA, γ -aminobutyric acid; HS, habitual snoring; OAI, obstructive apnea-hypopnea index; OAI, obstructive apnea index; OSA, obstructive sleep apnea; PS, primary snoring; PSG, polysomnography; ROC, receiver operating characteristic; VOCs, complex volatile organic compounds.

proteomic patterns,³⁰ and urinary neurotransmitters³¹ had accuracy enough to be an acceptable diagnostic test.

Regarding PPV values, taurine,³¹ the combined biomarkers tested by Gozal et al.²⁴ had the highest PPV values among pediatric studies (100% and 97%). In studies conducted in adults interleukin-6 (IL-6),²⁶ creatine phosphokinase (CK),²⁹ interleukin-10 (IL-10),²⁶ and TRX²⁸ had the highest PPV values.

The combined biomarkers tested by Gozal et al.²⁴, γ -aminobutyric acid,³² and phenylethylamine (PEA)³¹ had the highest NPV (100%) in pediatric studies, while IL-6 and IL-10²⁶ had the highest NPV in adults ones (100%).

The combined biomarkers tested by Gozal et al.²⁴ and the 2 biomarkers tested by Li et al.²⁶ reported excellent Youden's Index (0.97, 1.00, 0.97).

Five studies^{24,26,28,30,31} reported the highest diagnostic odds ratio (DOR). The results reported when the biomarkers were combined in Gozal et al.²⁴ and Kheirandish-Gozal et al.³¹ showed better accuracy results than when they were tested

individually (DTA measurements are presented in **Tables 3** and **4**).

In summary, only the biomarkers tested by Gozal et al.²⁴ satisfied the criteria required for an excellent diagnostic test in children. Kheirandish-Gozal et al.³¹ and Shah et al.³⁰ satisfied the criteria for acceptable diagnostic test in children.

In adults, the study conducted by Li et al.²⁶ that tested IL-6 and IL-10 could be considered an excellent diagnostic test.

DISCUSSION

Summary of Evidence

This systematic review investigated the available evidence on the diagnostic capability of biomarkers for the diagnosis of OSA. The actual gold standard for OSA diagnosis, i.e., the overnight PSG, has several important limitations: (a) it is potentially stressful, (b) requires sleep outside the home environment, (c)

Table 2—Summary of study descriptive characteristics of included studies (adults).

Year	Author	Country	Cases	Controls	Mean age, years	Type of Biomarker	OSA* diagnostic criteria at PSG	Biomarker	Main Conclusion
2006	Lentini et al. ²⁹	Germany	OSA suspected: Mild to Moderate OSA (n = 93/17 male) Severe OSA (n = 89/71 male)	OSA suspected: (n = 19/8 male)	54.9	Blood	AHI > 5	CK levels	Mild-to-moderate elevation in CK level was predictive of OSA. The application of CPAP therapy in OSA patients resulted in a significant decrease in CK level.
2007	Ursavas et al. ²⁷	Turkey	Moderate-to-severe OSA (n = 39/30 male) Matched for age, gender, BMI, smoking history, and CVD with controls.	Non-OSA (n = 34/23 male)	50.5 [#]	Blood	AHI ≥ 5	ICAM-1 VCAM-1	ICAM-1 and VCAM-1 may be potential biomarkers to identify OSA and non-OSA adults.
2009	Li et al. ²⁶	Thailand	Patients with EDS and loud snoring: Mild OSA (n = 22) Moderate OSA (n = 22) Severe OSA (n = 24)	Patients with EDS and loud snoring: Smoker control group (n = 10) Non-OSA (n = 22)	44 [#]	Blood EBC	AHI ≥ 5	8-isoprostane IL-6 TNF-α IL-10	EBC IL-6 and IL-10 have potential to predict severity of OSA in non-smoker OSA suspects.
2013	Guo et al. ²⁸	China	OSA suspected: Mild OSA (n = 14) Moderate OSA (n = 11) Severe OSA (n = 29)	OSA suspected (n = 9)	47.8 [#]	Blood	AHI ≥ 5	TRX	Plasma TRX level is associated with OSA severity and may be used as a severity indicator of OSA.
2013	Hirotsu et al. ²⁵	Brazil	Volunteers: OSA adults (n = 339/18.8% male)	Volunteers: (n = 682/25.9% male)	44.6 [#]	Blood	AHI ≥ 15	Uric acid	A strong association was found between uric acid levels and OSA, although acid uric levels do not qualify for a biomarker alone.

*All terms that mean obstructive sleep apnea (SDB, SRDB, OSAS) were standardized as OSA. [#] Mean calculated by author. AHI, apnea-hypopnea index; BMI, body mass index; CK, creatine phosphokinase; CPAP, continuous positive airway pressure; CVD, cardiovascular disease; EBC, exhaled breath condensate; EDS, excessive daytime sleepiness; ICAM-1, intercellular adhesion molecule 1; IL-6, interleukin-6; IL-10, interleukin-10; OAH, obstructive apnea-hypopnea index; OSA, obstructive sleep apnea; PS, primary snoring; PSG, polysomnography; ROC, receiver operating characteristic; TNF-α, tumor necrosis factor α; TRX, thioredoxin; VCAM-1, vascular cell adhesion molecule-1.

may not be widely available; and (d) is expensive.³³ Therefore, development of simple, cheap, and reliable diagnostic tools that would at least permit large scale screening of at-risk populations, and enable accurate identification of the subjects with definitive disease or with definitive absence of disease would potentially revolutionize the field.¹³ This urgent need to find an ideal biomarker for OSA could explain the large number of studies about this topic published since 2000. We found a large number of studies in phase 1 screening process. Unfortunately, 106 studies were excluded because they did not report sensitivity and specificity. Without these values it is impossible to properly assess the real diagnostic capability of any alternative test. Brockman et al.³³ emphasizes that the lack of important information in DTA publications is sobering, as clear guidelines for reporting validity measures of alternative exploratory diagnostic methods were

published in 2003, and encourage future authors of DTA studies to follow these recommendations.

Before we analyze our results, it is important emphasize that there was wide variation in the OSA diagnostic criteria employed by the pediatric studies. The AHI was the most frequently used diagnostic PSG measure of OSA severity. However, the use of AHI was associated with two major limitations. Firstly, the clinically accepted consensus for the cutoff AHI value for either the presence or absence of OSA remains unresolved. Secondly, no widely accepted agreement has been reached regarding whether children with PSG-based AHI values between the “normal cutoff” and 5/h TST should undergo surgical adenotonsillectomy.¹⁹ Based on these considerations, it becomes apparent that the definitive diagnosis of OSA solely based on the low-end spectrum of the PSG-based measures (e.g., AHI, RDI, OAH) is difficult if not

Table 3—Diagnostic test accuracy (children).

Author	Age (mean or range in years)	Sample Size (N)	OSA/ NON-OSA	Biomarker	Sensitivity/ Specificity (%)	PPV/NPV (%) [#]	LR+/LR- [#]	DOR [#]	Youden's Index Value [#]
Benedek et al. ²³	8.5	28	18/10	VOCs mixtures	78/70	82/64	2.59/0.31	8.35	0.48
Gozal et al. ²⁴	2–9	120	60/60	Kallikrein-1	97/77	81/96	4.22/0.04	105.50	0.74
Gozal et al. ²⁴	2–9	120	60/60	Uromodulin	88/80	81/87	4.40/0.15	29.33	0.63
Gozal et al. ²⁴	2–9	120	60/60	Urocortin-3	94/79	82/93	4.48/0.08	56.00	0.73
Gozal et al. ²⁴	2–9	120	60/60	Orosomucoid-1	90/83	84/89	5.29/0.12	44.08	0.73
Kheirandish-Gozal et al. ³¹	6.3	70	50/20	GABA	100/45	82/100	1.82/zero	∞	0.45
Kheirandish-Gozal et al. ³¹	6.3	70	50/20	PEA	100/65	88/100	2.86/zero	∞	0.65
Kheirandish-Gozal et al. ³¹	6.3	70	50/20	Taurine	88/100	100/77	∞/0.12	∞	0.88
Shah et al. ³⁰	3–12	40	20/20	Proteomic Patterns	93/90	90/93	9.30/0.08	116.25	0.83

[#] Data not available in the original article. The authors calculated data from information available in the article. OSA, obstructive sleep apnea; PPV, positive predictive value; NPV, negative predictive value; LR+, positive likelihood ratio; LR-, negative likelihood ratio; DOR, diagnostic odds ratio; VOCs, complex volatile organic compounds; GABA, γ-aminobutyric acid; PEA, β-phenylethylamine.

Table 4—Measurements for combined biomarkers (children).

Author	Age (range in years)	Sample Size (N)	OSA/ NON-OSA	Biomarker	Sensitivity/ Specificity (%)	PPV/NPV (%) [#]	LR+/LR- [#]	DOR [#]	Youden's Index Value [#]
Gozal et al. ²⁴	2–9	120	60/60	Kallikrein-1 Uromodulin Urocortin-3 Orosomucoid-1	100/97	97/100	28.60/zero	∞	0.97
Kheirandish-Gozal et al. ³¹	6.3	70	50/20	Urinary Neurotransmitters	82/90	95/67	8.20/0.20	41.00	0.72

[#] Data not available in the original article. The authors calculated data from information available in the article. OSA, obstructive sleep apnea; PPV, positive predictive value; NPV, negative predictive value; LR+, positive likelihood ratio; LR-, negative likelihood ratio; DOR, diagnostic odds ratio.

Table 5—Diagnostic test accuracy (adults).

Author	Mean age, years	Sample Size (N)	OSA/ NON-OSA	Biomarker	Sensitivity/ Specificity (%)	PPV/NPV (%) [#]	LR+/LR- [#]	DOR [#]	Youden's Index Value [#]
Guo et al. ²⁸	47.8 [#]	63	54/9	TRX	91/78	96/58	4.08	34.00	0.69
Hirotsu et al. ²⁵	44.6 [#]	456 (male)	192/264	Uric acid	62/57	51/67	1.44	0.02	0.26
Hirotsu et al. ²⁵	44.6 [#]	565 (female)	147/418	Uric acid	63/64	38/64	1.75	3.01	0.27
Lentini et al. ²⁹	54.9	201	182/19	CK	43/95	99/14	8.14	13.57	0.38
Li et al. ²⁶	44.0 [#]	100	68/32	IL-6	100/100	100/100	∞	∞	1.00
Li et al. ²⁶	44.0 [#]	100	68/32	IL-10	100/97	99/100	32.00	∞	0.97
Ursavas et al. ²⁷	50.5 [#]	73	39/34	ICAM-1	69/82	81/70	3.92	10.59	0.51
Ursavas et al. ²⁷	50.5 [#]	73	39/34	VCAM-1	74/65	71/69	2.11	5.41	0.39

[#] Data not available in the original article. The authors calculated data from information available in the article. OSA, obstructive sleep apnea; PPV, positive predictive value; NPV, negative predictive value; LR+, positive likelihood ratio; LR-, negative likelihood ratio; DOR, diagnostic odds ratio; CK, creatine phosphokinase; TRX, thioredoxin; IL-6, interleukin-6; IL-10, interleukin-10; ICAM-1, intercellular adhesion molecule 1; VCAM-1, vascular cell adhesion molecule-1.

Figure 2—Forest plot with diagnostic test accuracy (sensitivity, specificity, and 95% confidence interval) of each study.

(A) Studies in children that analyzed each biomarker individually. (B) Studies in children that combined three or four biomarkers in one analysis. (C) Studies in adults. TP, true positive; FP, false positive; FN, false negative; TN, true negative.

Figure 3—Receiver operating characteristic (ROC) curves for each group.

(A) Studies in children that analyzed each biomarker individually. (B) Studies in children that combined three or four biomarkers in one analysis. (C) Studies in adults.

impossible. Similar, albeit less vague overlap exists among adult patients, even if the PSG criteria for the presence of OSA have been more firmly established and accepted around the world.³⁴

Although we found only nine eligible studies, this meta-analysis is informative, because by combining the available data it increases the sample size to 258 children and 1,458 adults. The

results of meta-analysis showed that five studies^{24,26,28–31} provide acceptable metrics enabling identification of those who really suffered from OSA (true positive), while two of them^{24,26} performed well for identification of those who did not have OSA (true-negative). The LR values confirm that both had excellent DTA. Similarly, PPV and NPV values showed that five studies^{24,26,28,29,31}

performed acceptably in identifying OSA subjects. Two studies^{24,26} were also good in identifying OSA and non-OSA subjects, thereby concurring with sensitivity and specificity numbers.

Also, the combined biomarker approaches tested by Gozal et al.²⁴ and the two biomarkers tested by Li et al.²⁶ reported excellent Youden's Index (0.97, 1.00, 0.97), the latter indicative of high accuracy.²²

The DOR for three pediatric^{24,30,31} and two adult studies^{26,28} indicate that the biomarkers tested in children (combined kallikrein-1, uromodulin, urocortin-3, orosomucoid-1, proteomic patterns, and urinary neurotransmitters) and in adults (IL-6, IL-10, TRX) had better discriminatory test performance. Is it important to emphasize that the results reported when the biomarkers were combined in Gozal et al.²⁴ and Kheirandish-Gozal et al.³¹ showed better accuracy measurements than when the biomarkers tested in these studies were analyzed individually (Table 3).

In summary, only the putative biomarkers tested in Gozal et al.²⁴ satisfied the required criteria for an excellent diagnostic test in children. Gozal et al.²⁴ investigated urinary biomarkers in 60 OSA patients, 30 primary snorers, and 30 healthy controls in order to identify urinary protein clusters that were highly sensitive and specific for OSA. They found that unique sets of proteins were either increased or decreased in the urine of OSA children, and that their combined ROC curve analysis using four candidate proteins simultaneously provided a near-perfect DTA (close to 100%). Another useful set of different biomarkers was subsequently identified by Kheirandish-Gozal et al.,³¹ who examined urinary neurotransmitters in 50 OSA and 20 controls. They reported an overnight increase in epinephrine and norepinephrine levels in children with OSA, while taurine levels were decreased. Using combinatorial approaches and cutoff values for overnight changes of these four neurotransmitters enabled a good prediction of OSA. Also, Shah et al.³⁰ evaluated the proteomic patterns of 20 children with OSA and of 20 children with habitual primary snoring but no evidence of OSA using surface-enhanced laser desorption/ionization time of flight mass spectrometry. The proteomic patterns were capable of diagnosing OSA with 93% sensitivity and 90% specificity. However, their methodological approaches did not allow for identification of the actual candidates, such that this work remains a proof of principle rather than provide yet other defined biomarker candidates.

In adults, the study conducted by Li et al.,²⁶ which tested IL-6 and IL-10, could be considered an excellent diagnostic test. The study aimed to identify the best biomarker, either single or in combination, with best cost-effectiveness ratio. The authors analyzed 8-isoprostane, IL-6, TNF- α , and IL-10 in the EBC and serum of OSA, non-OSA, and healthy smoking subjects. These investigators reported that levels, in both EBC and serum, differed significantly across the four biomarkers tested.

Overall Assessment

A previous review focused on different pediatric OSA diagnostic tests³³ have identified several approaches that putatively provide either acceptable or excellent DTA in the prediction of OSA. These tests have included sleep lab-based polygraphy, anterior rhinomanometry, and urinary biomarkers. However, the authors³³ stated that there was still insufficient evidence to

recommend any of these alternative tests to PSG for diagnosis of pediatric OSA.

The current systematic meta-analysis indicates that although all selected articles concluded that biomarkers could be useful to reliably diagnose OSA, not all approaches can actually be used as viable or definitive biomarkers. Only the combination of kallikrein-1, uromodulin, urocortin-3, and orosomucoid-1 displayed sufficient accuracy to be considered an OSA diagnostic test in children. In contrast, the combination of urinary neurotransmitters³¹ and of the serum proteomic patterns³⁰ displayed acceptable accuracy to serve as a screening test in children. In adults, IL-6 and IL-10 show favorable potential to become a good biomarker to identify OSA and non-OSA subjects.

Limitations

Except for one study,²⁵ the other studies used a sample from sleep center or subjects with OSA symptoms. This can affect the prevalence, which can bias the sensitivity and specificity of the biomarker-based test. Thus, we do not know if the tests would respond similarly when applied to the general population. Other identified limitations in the published studies were: lack of a masked interpretation of the reference and index test and no clear information regarding how many investigators analyzed the test data or if their techniques were calibrated. Finally, 106 potential biomarkers studies had to be excluded due to lack of DTA values, suggesting that if such DTA assessments were available, the present conclusions could be markedly affected, further reinforcing the need for standardized reporting of predictive DTA values. Notwithstanding such considerations, the current findings are encouraging toward the implementation of biomarkers in the diagnosis of OSA, and prompted us to perform a relatively simplistic financial cost analysis of potential savings embedded in such a clinically based approach. For example, assuming that a combinatorial multiple biomarker-based assay would be required, and estimating that the global cost of such assay would amount to one-fourth of the cost of a PSG, then application of the biomarker-based approach would be economically advantageous if < 25% of the biomarker test results would be equivocal, thereby necessitating a PSG. Similar models can be implemented using various cost estimates, with obviously, more favorable "equivocal result" rates still being financially viable if the assay costs are lower.

CONCLUSIONS

Kallikrein-1, uromodulin, urocortin-3, and orosomucoid-1 have enough accuracy to be used as an OSA diagnostic test in children when used in combination.

Plasma IL-6 and IL-10 levels are potentially promising to become a good biomarker aiming to identify adult individuals with and without OSA.

REFERENCES

1. Lumeng JC, Chervin RD. Epidemiology of pediatric obstructive sleep apnea. *Proc Am Thorac Soc* 2008;5:242-52.
2. Jean-Louis G, Zizi F, Brown D, Ogedegbe G, Borer J, McFarlane S. Obstructive sleep apnea and cardiovascular disease: evidence and underlying mechanisms. *Minerva Pneumol* 2009;48:277-93.

3. Marcus CL, Brooks LJ, Draper KA, et al. Diagnosis and management of childhood obstructive sleep apnea syndrome. *Pediatrics* 2012;130:576–84.
4. Ali SS, Oni ET, Warraich HJ, et al. Systematic review on noninvasive assessment of subclinical cardiovascular disease in obstructive sleep apnea: new kid on the block! *Sleep Med Rev* 2014;18:379–91.
5. Lal C, Strange C, Bachman D. Neurocognitive impairment in obstructive sleep apnea. *Chest* 2012;141:1601–10.
6. Tan HL, Gozal D, Kheirandish-Gozal L. Obstructive sleep apnea in children: a critical update. *Nat Sci Sleep* 2013;5:109–23.
7. Drager LF, Togeiro SM, Polotsky VY, Lorenzi-Filho G. Obstructive sleep apnea: a cardiometabolic risk in obesity and the metabolic syndrome. *J Am Coll Cardiol* 2013;62:569–76.
8. Bixler EO, Vgontzas AN, Lin HM, et al. Sleep disordered breathing in children in a general population sample: prevalence and risk factors. *Sleep* 2009;32:731–6.
9. Yamagishi K, Ohira T, Nakano H, et al. Cross-cultural comparison of the sleep-disordered breathing prevalence among Americans and Japanese. *Eur Respir J* 2010;36:379–84.
10. Ralls FM, Grigg-Damberger M. Roles of gender, age, race/ethnicity, and residential socioeconomic in obstructive sleep apnea syndromes. *Curr Opin Pulm Med* 2012;18:568–73.
11. Bixler EO, Vgontzas AN, Lin HM, et al. Sleep disordered breathing in children in a general population sample: prevalence and risk factors. *Sleep* 2009;32:731–6.
12. Spruyt K, Gozal D. Screening of pediatric sleep-disordered breathing: a proposed unbiased discriminative set of questions using clinical severity scales. *Chest* 2012;142:1508–15.
13. Gozal D. Serum, urine, and breath-related biomarkers in the diagnosis of obstructive sleep apnea in children: is it for real? *Curr Opin Pulm Med* 2012;18:561–7.
14. Lamm C, Mandeli J, Kattan M. Evaluation of home audiotapes as an abbreviated test for obstructive sleep apnea syndrome (OSAS) in children. *Pediatr Pulmonol* 1999;27:267–72.
15. Sivan Y, Kornecki A, Schonfeld T. Screening obstructive sleep apnoea syndrome by home videotape recording in children. *Eur Respir J* 1996;9:2127–31.
16. Brouillette RT, Morielli A, Leimanis A, Waters KA, Luciano R, Ducharme FM. Nocturnal pulse oximetry as an abbreviated testing modality for pediatric obstructive sleep apnea. *Pediatrics* 2000;105:405–12.
17. Finn D, McNally P. Estimated population prevalence of obstructive sleep apnoea in a community of German third graders. *Eur Respir J* 2011;37:975; author reply 75–6.
18. Andreu AL, Chiner E, Sancho-Chust, JN et al. Effect of an ambulatory diagnostic and treatment programme in patients with sleep apnoea. *Eur Respir J* 2012;39:305–12.
19. Wong TK. The search on an ideal disease marker for childhood obstructive sleep apnea syndrome. *Sleep* 2011;34:133–4.
20. Moher D, Liberati A, Tetzlaff J, Altman DG; Prisma Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Int J Surg* 2010;8:336–41.
21. Whiting P, Rutjes AW, Reitsma JB, Bossuyt PM, Kleijnen J. The development of QUADAS: a tool for the quality assessment of studies of diagnostic accuracy included in systematic reviews. *BMC Med Res Methodol* 2003;3:25.
22. Deeks JJ, Bossuyt P, Gatsonis C, eds. *Cochrane Handbook for Systematic Reviews of Diagnostic Test Accuracy* Version 1.0. The Cochrane Collaboration, 2010. Available from: <http://srdta.cochrane.org/>.
23. Benedek P, Lazar Z, Bikov A, Kunos L, Katona G, Horvath I. Exhaled biomarker pattern is altered in children with obstructive sleep apnoea syndrome. *Int J Pediatr Otorhinolaryngol* 2013;77:1244–7.
24. Gozal D, Jortani S, Snow AB, et al. Two-dimensional differential in-gel electrophoresis proteomic approaches reveal urine candidate biomarkers in pediatric obstructive sleep apnea. *Am J Respir Crit Care Med* 2009;180:1253–61.
25. Hirotsu C, Tufik S, Guindalini C, Mazzotti DR, Bittencourt LR, Andersen ML. Association between uric acid levels and obstructive sleep apnea syndrome in a large epidemiological sample. *PLoS One* 2013;8:e66891.
26. Li Y, Chongsuvivatwong V, Geater A, Liu A. Exhaled breath condensate cytokine level as a diagnostic tool for obstructive sleep apnea syndrome. *Sleep Med* 2009;10:95–103.
27. Ursavas A, Karadag M, Rodoplu E, Yilmaztepe A, Oral HB, Gozu RO. Circulating ICAM-1 and VCAM-1 levels in patients with obstructive sleep apnea syndrome. *Respiration* 2007;74:525–32.
28. Guo Q, Wang Y, Li QY, Li M, Wan HY. Levels of thioredoxin are related to the severity of obstructive sleep apnea: based on oxidative stress concept. *Sleep Breath* 2013;17:311–6.
29. Lentini S, Manka R, Scholtyssek S, Stoffel-Wagner B, Luderitz B, Tasci S. Creatine phosphokinase elevation in obstructive sleep apnea syndrome: an unknown association? *Chest* 2006;129:88–94.
30. Shah ZA, Jortani SA, Tauman R, Valdes R Jr, Gozal D. Serum proteomic patterns associated with sleep-disordered breathing in children. *Pediatr Res* 2006;59:466–70.
31. Kheirandish-Gozal L, McManus CJ, Kellermann GH, Samiei A, Gozal D. Urinary neurotransmitters are selectively altered in children with obstructive sleep apnea and predict cognitive morbidity. *Chest* 2013;143:1576–83.
32. Takagi T, Morser J, Gabazza EC, et al. The coagulation and protein C pathways in patients with sleep apnea. *Lung* 2009;187:209–13.
33. Brockmann PE, Schaefer C, Poets A, Poets CF, Urschitz MS. Diagnosis of obstructive sleep apnea in children: a systematic review. *Sleep Med Rev* 2013;17:331–40.
34. Nerfeldt P, Aoki F, Friberg D. Polygraphy vs. polysomnography: missing osas in symptomatic snorers—a reminder for clinicians. *Sleep Breath* 2014;18:297–303.

SUBMISSION & CORRESPONDENCE INFORMATION

Submitted for publication July, 2014

Submitted in final revised form August, 2014

Accepted for publication September, 2014

Address correspondence to: David Gozal, MD, Department of Pediatrics, The University of Chicago; 5721 S. Maryland Avenue, MC 8000, Suite K-160, Chicago, IL 60637; Tel: (773) 702-3360; Fax: (773) 926-0756; Email: dgozal@uchicago.edu

DISCLOSURE STATEMENT

This was not an industry supported study. Dr. Gozal is supported in part by NIH grants RO1 HL-65270 and P50 HL-107160. The authors have indicated no financial conflicts of interest. This research was performed at the Department of Dentistry, University of Alberta, Canada.

Appendix 1—Search.

Database	Search
Cochrane (March 20, 2014) EMBASE (1974-March 20, 2014) MEDLINE (1948-March 20, 2014) PubMed (March 20, 2014)	(sleep apnea or sleep apnoea or sleep apnea, obstructive or obstructive sleep apnea or obstructive sleep apnoea) and (exhaled condensate biomarker* or salivary biomarker* or urinary biomarker* or blood biomarker* or serum biomarker* or biomarker*)
LILACS (March 20, 2014)	"apnea" or "apneia" and "marcadores biológicos"
Google Scholar (March 20, 2014)	"biomarkers and sleep apnea" (without patents or citations)

Appendix 2

Test indicators	Data analysis	References
DOR	The value of a DOR ranges from 0 to infinity, with higher values indicating better discriminatory test performance. A value of 1 means that a test does not discriminate between patients with the disorder and those without it. Values lower than 1 point to improper test interpretation (more negative tests among the diseased).	Glas et al. ¹
LR	LR+ >3 and an LR- <0.3 – acceptable diagnostic test accuracy (DTA) LR+ >10 and LR- <0.1 – excellent DTA.	Brockmann et al. ²
Sensitivity	>80% excellent, 70-80% good, 60-69% fair, <60% poor	No consensus in this regard exists in the literature.
Specificity	>90% excellent, 80-90% good, 70-79% fair, <70% poor	No consensus in this regard exists in the literature.
Youden's Index	Youden's Index values close to 1 indicate high accuracy; a value of zero is equivalent to uninformed guessing and indicates that a test has no diagnostic value.	Deeks et al. ³

REFERENCES

1. Glas AS, Lijmer JG, Prins MH, Bonsel GJ, Bossuyt PM. The diagnostic odds ratio: a single indicator of test performance. *J Clin Epidemiol* 2003;56:1129-35.
2. Brockmann PE, Schaefer C, Poets A, Poets CF, Urschitz MS. Diagnosis of obstructive sleep apnea in children: a systematic review. *Sleep Med Rev* 2013;17:331-40.
3. Deeks JJ, Bossuyt P, Gatsonis C, eds. *Cochrane Handbook for Systematic Reviews of Diagnostic Test Accuracy* Version 1.0. The Cochrane Collaboration, 2010. Available from: <http://srdta.cochrane.org/>.

Appendix 3—Excluded articles and reasons for exclusion.

Author/Year	Reason for exclusion *	Author/Year	Reason for exclusion *	Author/Year	Reason for exclusion *
Aihara et al. 2013 ¹	9	Khalyfa et al. 2012 ⁴⁵	9	Patel et al. 2009 ⁸⁹	4
Akinnusi et al. 2011 ²	9	Kheirandish-Gozal 2014 ⁴⁶	9	Peled et al. 2007 ⁹⁰	9
Alzogaibi et al. 2005 ³	9	Kheirandish-Gozal et al. 2006 ⁴⁷	9	Petrosyan et al. 2008 ⁹¹	9
Antonopoulou et al. 2008 ⁴	9	Kim et al. 2009 ⁴⁸	9	Phillips et al. 2007 ⁹²	9
Arias et al. 2008 ⁵	9	Kim et al. 2010 ⁴⁹	9	Pinto et al. 2013 ⁹³	9
Bhushan et al. 2011 ⁶	9	Kim et al. 2012 ⁵⁰	2	Przybylowski et al. 2006 ⁹⁴	5
Braga et al. 2006 ⁷	9	Kim et al. 2013 ⁵¹	9	Punjabi et al. 2007 ⁹⁵	9
Bratel et al. 1999 ⁸	6	Kishida et al. 2014 ⁵²	8	Roche et al. 2009 ⁹⁶	2
Burioka et al. 2008 ⁹	9	Kohler et al. 2011 ⁵³	9	Rubinsztajn et al. 2006 ⁹⁷	5
Calvin et al. 2010 ¹⁰	1	Krishna et al. 2006 ⁵⁴	9	Ryan et al. 2007 ⁹⁸	9
Carpagnano et al. 2002 ¹¹	9	Kuramoto et al. 2009 ⁵⁵	9	Salord 2014 ⁹⁹	2
Carpagnano et al. 2003 ¹²	9	Kurt et al. 2013 ⁵⁶	9	Schulz et al. 2002 ¹⁰⁰	9
Chin et al. 2000 ¹³	9	Ladesich et al. 2011 ⁵⁷	9	Shamsuzzaman et al. 2002 ¹⁰¹	9
Cholidou et al. 2013 ¹⁴	2	Lam et al. 2009 ⁵⁸	9	Shi et al. 2013 ¹⁰²	9
Christou et al. 2002 ¹⁵	9	Larkin et al. 2005 ⁵⁹	9	Simiakakis et al. 2012 ¹⁰³	9
Chung et al. 2013 ¹⁶	9	Lavie et al. 2002 ⁶⁰	9	Sokucu et al. 2012 ¹⁰⁴	9
Cintra et al. 2011 ¹⁷	9	Lederer et al. 2009 ⁶¹	9	Stats et al. ¹⁰⁵	4
Cofta et al. 2013 ¹⁸	9	Lee et al. 2010 ⁶²	9	Stefanini et al. 2012 ¹⁰⁶	9
Constantinidis et al. 2008 ¹⁹	9	Lee et al. 2012 ⁶³	9	Steiropoulos et al. 2010 ¹⁰⁷	9
Culla et al. 2010 ²⁰	2	Li et al. 2006 ⁶⁴	9	Sukegawa et al. 2005 ¹⁰⁸	9
DeBoer 2012 ²¹	9	Li et al. 2008 ⁶⁵	9	Svensson et al. 2012 ¹⁰⁹	9
Duru et al. 2012 ²²	9	Li et al. 2008 ⁶⁶	9	Takahashi et al. 2008 ¹¹⁰	9
El-Solh 2002 ²³	2	Li et al. 2010 ⁶⁷	9	Tauman et al. 2004 ¹¹¹	9
Feng et al. 2012 ²⁴	9	Lin et al. 2013 ⁶⁸	5	Tauman et al. 2007 ¹¹²	9
Ferrarini et al. 2013 ²⁵	9	Loubaki et al. 2008 ⁶⁹	7	Ting et al. 2009 ¹¹³	9
Goldbart et al. 2006 ²⁶	9	Lui et al. 2009 ⁷⁰	9	Tual-Chalot et al. 2014 ¹¹⁴	9
Gozal et al. 2002 ²⁷	9	Makino et al. ⁷¹	7	Ucar et al. 2009 ¹¹⁵	9
Gozal et al. 2007 ²⁸	2	Malakasioti et al. 2012 ⁷²	9	Uysal et al. 2014 ¹¹⁶	2
Gozal et al. 2008 ²⁹	9	Mancuso et al. 2012 ⁷³	9	Van Hoorenback ¹¹⁷	6
Gozal et al. 2010 ³⁰	2	Mehra et al. 2006 ⁷⁴	9	Vavougios et al. 2014 ¹¹⁸	9
Gozal et al. 2013 ³¹	9	Montgomery-Downs et al. 2006 ⁷⁵	9	Villa et al. 2014 ¹¹⁹	9
Guilleminault et al. 2004 ³²	9	Murase et al. 2013 ⁷⁶	9	Wang et al. 2005 ¹²⁰	5
Guven et al. 2012 ³³	9	Norman et al. 2008 ⁷⁷	9	Wang et al. 2010 ¹²¹	2
Hira et al. 2012 ³⁴	9	Ntalapascha et al. 2013 ⁷⁸	9	Wang et al. 2012 ¹²²	4
Htoo et al. 2006 ³⁵	9	O'Brien et al. 2006 ⁷⁹	9	Wang et al. 2013 ¹²³	9
Imagawa et al. 2004 ³⁶	9	Ohga et al. 2003 ⁸⁰	9	Wang et al. 2013 ¹²⁴	9
Jurado-Gamez et al. 2011 ³⁷	9	Osorio 2014 ⁸¹	2	Wang et al. 2014 ¹²⁵	9
Jurado-Gamez et al. 2012 ³⁸	9	Oyama et al. 2012 ⁸²	2	Yamauchi et al. 2005 ¹²⁶	9
Kaditis et al. 2005 ³⁹	9	Ozben et al. 2013 ⁸³	9	Ye et al. 2007 ¹²⁷	9
Kaditis et al. 2007 ⁴⁰	9	Pallayova et al. 2010 ⁸⁴	9	Ye et al. 2010 ¹²⁸	9
Kaditis et al. 2009 ⁴¹	9	Pallayova et al. 2011 ⁸⁵	9	Yokoe et al. 2003 ¹²⁹	9
Kaditis et al. 2009 ⁴²	9	Papaioannou et al. 2012 ⁸⁶	9	Zamarron et al. 2008 ¹³⁰	9
Kanbay et al. 2008 ⁴³	9	Park et al. 2013 ⁸⁷	9	Zamarron et al. 2011 ¹³¹	9
Khalyfa et al. 2011 ⁴⁴	3	Patacchioli et al. 2014 ⁸⁸	9	Zhang et al. 2013 ¹³²	9

Appendix 3 (continued)—Excluded articles and reasons for exclusion.

* **Reasons for exclusion:** (1) Central apnea (n = 1), (2) OSA with morbidities (n = 12), (3) Not OSA patients (n = 1), (4) Different target condition (n = 3), (5) Not English, Spanish or Portuguese (n = 4), (6) Not full overnight PSG (n = 2), (7) Not biomarkers study (n = 2), (8) Review (n = 1), (9) Not reported sensitivity and specificity.

REFERENCES

- Aihara K, Oga T, Chihara Y, et al. Analysis of systemic and airway inflammation in obstructive sleep apnea. *Sleep Breath* 2013;17:597-604.
- Akinnusi ME, Laporta R, El-Sohl AA. Lectin-like oxidized low-density lipoprotein receptor-1 modulates endothelial apoptosis in obstructive sleep apnea. *Chest* 2011;140:1503-10.
- Alzoghbi MA, Bahammam AS. Lipid peroxides, superoxide dismutase and circulating IL-8 and GCP-2 in patients with severe obstructive sleep apnea: a pilot study. *Sleep Breath* 2005;9:119-26.
- Antonopoulou S, Loukides S, Papatheodorou G, Roussos C, Alchanatis M. Airway inflammation in obstructive sleep apnea: is leptin the missing link? *Respir Med* 2008;102:1399-405.
- Arias MA, Garcia-Rio F, Alonso-Fernandez A, et al. CPAP decreases plasma levels of soluble tumour necrosis factor-alpha receptor 1 in obstructive sleep apnoea. *Eur Respir J* 2008;32:1009-15.
- Bhushan B, Khalyfa A, Spruyt K, et al. Fatty-acid binding protein 4 gene polymorphisms and plasma levels in children with obstructive sleep apnea. *Sleep Med* 2011;12:666-71.
- Braga CW, Martinez D, Wofchuk S, Portela LV, Souza DO. S100B and NSE serum levels in obstructive sleep apnea syndrome. *Sleep Med* 2006;7:431-5.
- Bratel T, Wennlund A, Carlstrom K. Pituitary reactivity, androgens and catecholamines in obstructive sleep apnoea. Effects of continuous positive airway pressure treatment (CPAP). *Respir Med* 1999;93:1-7.
- Burioka N, Miyata M, Fukuoka Y, Endo M, Shimizu E. Day-night variations of serum interleukin-6 in patients with severe obstructive sleep apnea syndrome before and after continuous positive airway pressure (CPAP). *Chronobiol Int* 2008;25:827-34.
- Calvin AD, Somers VK, Steensma DP, et al. Advanced heart failure and nocturnal hypoxaemia due to central sleep apnoea are associated with increased serum erythropoietin. *Eur J Heart Fail* 2010;12:354-9.
- Carpagnano GE, Kharitonov SA, Resta O, Foschino-Barbaro MP, Gramiccioni E, Barnes PJ. Increased 8-isoprostane and interleukin-6 in breath condensate of obstructive sleep apnea patients. *Chest* 2002;122:1162-7.
- Carpagnano GE, Kharitonov SA, Resta O, Foschino-Barbaro MP, Gramiccioni E, Barnes PJ. 8-Isoprostane, a marker of oxidative stress, is increased in exhaled breath condensate of patients with obstructive sleep apnea after night and is reduced by continuous positive airway pressure therapy. *Chest* 2003;124:1386-92.
- Chin K, Nakamura T, Shimizu K, et al. Effects of nasal continuous positive airway pressure on soluble cell adhesion molecules in patients with obstructive sleep apnea syndrome. *Am J Med* 2000;109:562-7.
- Cholidou KG, Kostakis ID, Manali ED, et al. Calprotectin: a protein related to cardiovascular risk in adult patients with obstructive sleep apnea. *Cytokine* 2013;61:917-23.
- Christou K, Moulas AN, Pastaka C, Gourgoulis KI. Antioxidant capacity in obstructive sleep apnea patients. *Sleep Med* 2003;4:225-8.
- Chung F, Chau E, Yang Y, Liao P, Hall R, Mokhlesi B. Serum bicarbonate level improves specificity of STOP-Bang screening for obstructive sleep apnea. *Chest* 2013;143:1284-93.
- Cintra F, Tufik S, D'Almeida V, et al. Cysteine: a potential biomarker for obstructive sleep apnea. *Chest* 2011;139:246-52.
- Cofta S, Wysocka E, Dziegielewska-Gesiak S, et al. Plasma selectins in patients with obstructive sleep apnea. *Adv Exp Med Biol* 2013;756:113-9.
- Constantinidis J, Ereliadis S, Angouridakis N, Konstantinidis I, Vital V, Angouridakis C. Cytokine changes after surgical treatment of obstructive sleep apnoea syndrome. *Eur Arch Otorhinolaryngol* 2008;265:1275-9.
- Culla B, Guida G, Brussino L, et al. Increased oral nitric oxide in obstructive sleep apnoea. *Respir Med* 2010;104:316-20.
- Deboer MD, Mendoza JP, Liu L, Ford G, Yu PL, Gaston BM. Increased systemic inflammation overnight correlates with insulin resistance among children evaluated for obstructive sleep apnea. *Sleep Breath* 2012;16:349-54.
- Duru S, Hikmet Firat I, Colak N, Ginis Z, Delibasi T, Ardic S. Serum S100B protein: a useful marker in obstructive sleep apnea syndrome. *Neurol Neurochir Pol* 2012;46:450-5.
- El-Sohl AA, Mador MJ, Sikka P, Dhillon RS, Amsterdam D, Grant BJ. Adhesion molecules in patients with coronary artery disease and moderate-to-severe obstructive sleep apnea. *Chest* 2002;121:1541-7.
- Feng X, Li P, Zhou C, Jia X, Kang J. Elevated levels of serum chemerin in patients with obstructive sleep apnea syndrome. *Biomarkers* 2012;17:248-53.
- Ferrarini A, Ruperez FJ, Erazo M, et al. Fingerprinting-based metabolomic approach with LC-MS to sleep apnea and hypopnea syndrome: a pilot study. *Electrophoresis* 2013;34:2873-81.
- Goldbart AD, Krishna J, Li RC, Serpero LD, Gozal D. Inflammatory mediators in exhaled breath condensate of children with obstructive sleep apnea syndrome. *Chest* 2006;130:143-8.
- Gozal D, Lipton AJ, Jones KL. Circulating vascular endothelial growth factor levels in patients with obstructive sleep apnea. *Sleep* 2002;25:59-65.
- Gozal D, Crabtree VM, Sans Capdevila O, Witcher LA, Kheirandish-Gozal L. C-reactive protein, obstructive sleep apnea, and cognitive dysfunction in school-aged children. *Am J Respir Crit Care Med* 2007;176:188-93.
- Gozal D, Serpero LD, Sans Capdevila O, Kheirandish-Gozal L. Systemic inflammation in non-obese children with obstructive sleep apnea. *Sleep Med* 2008;9:254-9.
- Gozal D, Serpero LD, Kheirandish-Gozal L, Capdevila OS, Khalyfa A, Tauman R. Sleep measures and morning plasma TNF-alpha levels in children with sleep-disordered breathing. *Sleep* 2010;33:319-25.
- Gozal D, Kheirandish-Gozal L, Bhattacharjee R, Molero-Ramirez H, Tan HL, Bandla HP. Circulating adiponectin concentrations in pediatric obstructive sleep apnea: potential relevance to endothelial function. *J Pediatr* 2013;163:1122-6.
- Guilleminault C, Kirisoglu C, Ohayon MM. C-reactive protein and sleep-disordered breathing. *Sleep* 2004;27:1507-11.
- Guven SF, Turkmani MH, Ciftci B, Ciftci TU, Erdogan Y. The relationship between high-sensitivity C-reactive protein levels and the severity of obstructive sleep apnea. *Sleep Breath* 2012;16:217-21.
- Hira HS, Shukla A, Kaur A, Kapoor S. Serum uric acid and lactate levels among patients with obstructive sleep apnea syndrome: which is a better marker of hypoxemia? *Ann Saudi Med* 2012;32:37-42.
- Htoo AK, Greenberg H, Tongia S, et al. Activation of nuclear factor kappaB in obstructive sleep apnea: a pathway leading to systemic inflammation. *Sleep Breath* 2006;10:43-50.
- Imagawa S, Yamaguchi Y, Ogawa K, et al. Interleukin-6 and tumor necrosis factor-alpha in patients with obstructive sleep apnea-hypopnea syndrome. *Respiration* 2004;71:24-9.
- Jurado-Gamez B, Fernandez-Marin MC, Gomez-Chaparro JL, et al. Relationship of oxidative stress and endothelial dysfunction in sleep apnoea. *Eur Respir J* 2011;37:873-9.
- Jurado-Gamez B, Cabrera CB, Ballesteros LC, et al. Association of cellular adhesion molecules and oxidative stress with endothelial function in obstructive sleep apnea. *Intern Med* 2012;51:363-8.
- Kaditis AG, Alexopoulos EI, Damani E, et al. Obstructive sleep-disordered breathing and fasting insulin levels in nonobese children. *Pediatr Pulmonol* 2005;40:515-23.
- Kaditis AG, Alexopoulos EI, Kalampouka E, et al. Nocturnal change of circulating intercellular adhesion molecule 1 levels in children with snoring. *Sleep Breath* 2007;11:267-74.
- Kaditis AG, Alexopoulos E, Chaidas K, et al. Urine concentrations of cysteinyl leukotrienes in children with obstructive sleep-disordered breathing. *Chest* 2009;135:1496-501.
- Kaditis AG, Alexopoulos EI, Damani E, et al. Urine levels of catecholamines in Greek children with obstructive sleep-disordered breathing. *Pediatr Pulmonol* 2009;44:38-45.
- Kanbay A, Kokturk O, Ciftci TU, Tavil Y, Bukan N. Comparison of serum adiponectin and tumor necrosis factor-alpha levels between patients with and without obstructive sleep apnea syndrome. *Respiration* 2008;76:324-30.
- Khalyfa A, Gharib SA, Kim J, et al. Peripheral blood leukocyte gene expression patterns and metabolic parameters in habitually snoring and non-snoring children with normal polysomnographic findings. *Sleep* 2011;34:153-60.
- Khalyfa A, Kheirandish-Gozal L, Capdevila OS, Bhattacharjee R, Gozal D. Macrophage migration inhibitory factor gene polymorphisms and plasma levels in children with obstructive sleep apnea. *Pediatr Pulmonol* 2012;47:1001-11.
- Kheirandish-Gozal L, Peris E, Wang Y, et al. Lipopolysaccharide-binding protein plasma levels in children: effects of obstructive sleep apnea and obesity. *J Clin Endocrinol Metab* 2014;99:656-63.

Appendix 3 (continued)—Excluded articles and reasons for exclusion.

47. Kheirandish-Gozal L, Capdevila OS, Tauman R, Gozal D. Plasma C-reactive protein in nonobese children with obstructive sleep apnea before and after adenotonsillectomy. *J Clin Sleep Med* 2006;2:301-4.
48. Kim J, Bhattacharjee R, Dayyat E, et al. Increased cellular proliferation and inflammatory cytokines in tonsils derived from children with obstructive sleep apnea. *Pediatr Res* 2009;66:423-8.
49. Kim J, Bhattacharjee R, Snow AB, Capdevila OS, Kheirandish-Gozal L, Gozal D. Myeloid-related protein 8/14 levels in children with obstructive sleep apnoea. *Eur Respir J* 2010;35:843-50.
50. Kim J, Bhattacharjee R, Khalyfa A, et al. DNA methylation in inflammatory genes among children with obstructive sleep apnea. *Am J Respir Crit Care Med* 2012;185:330-8.
51. Kim J, Gozal D, Bhattacharjee R, Kheirandish-Gozal L. TREM-1 and pentraxin-3 plasma levels and their association with obstructive sleep apnea, obesity, and endothelial function in children. *Sleep* 2013;36:923-31.
52. Kishida K, Funahashi T, Shimomura I. Adiponectin as a routine clinical biomarker. Best practice & research. *Clin Endocrinol Metab* 2014;28:119-30.
53. Kohler M, Stoewhas AC, Ayers L et al. Effects of continuous positive airway pressure therapy withdrawal in patients with obstructive sleep apnea: a randomized controlled trial. *Am J Respir Crit Care Med* 2011;184:1192-9.
54. Krishna J, Shah ZA, Merchant M, Klein JB, Gozal D. Urinary protein expression patterns in children with sleep-disordered breathing: preliminary findings. *Sleep Med* 2006;7:221-7.
55. Kuramoto E, Kinami S, Ishida Y, Shiotani H, Nishimura Y. Continuous positive nasal airway pressure decreases levels of serum amyloid A and improves autonomic function in obstructive sleep apnea syndrome. *Int J Cardiol* 2009;135:338-45.
56. Kurt OK, Yildiz N. The importance of laboratory parameters in patients with obstructive sleep apnea syndrome. *Blood Coagul Fibrinolysis* 2013;24:371-4.
57. Ladesich JB, Pottala JV, Romaker A, Harris WS. Membrane level of omega-3 docosahexaenoic acid is associated with severity of obstructive sleep apnea. *J Clin Sleep Med* 2011;7:391-6.
58. Lam JC, Yan CS, Lai AY, et al. Determinants of daytime blood pressure in relation to obstructive sleep apnea in men. *Lung* 2009;187:291-8.
59. Larkin EK, Rosen CL, Kirchner HL, et al. Variation of C-reactive protein levels in adolescents: association with sleep-disordered breathing and sleep duration. *Circulation* 2005;111:1978-84.
60. Lavie L, Kraiczi H, Hefetz A, et al. Plasma vascular endothelial growth factor in sleep apnea syndrome: effects of nasal continuous positive air pressure treatment. *Am J Crit Care Med* 2002;165:1624-8.
61. Lederer DJ, Jelic S, Basner RC, Ishizaka A, Bhattacharya J. Circulating KL-6, a biomarker of lung injury, in obstructive sleep apnoea. *Eur Respir J* 2009;33:793-6.
62. Lee LA, Chen NH, Huang CG, Lin SW, Fang TJ, Li HY. Patients with severe obstructive sleep apnea syndrome and elevated high-sensitivity C-reactive protein need priority treatment. *Otolaryngol Head Neck Surg* 2010;143:72-7.
63. Lee SD, Ju G, Choi JA, Kim JW, Yoon IY. The association of oxidative stress with central obesity in obstructive sleep apnea. *Sleep Breath* 2012;16:511-7.
64. Li AM, Chan MH, Chan DF, et al. Insulin and obstructive sleep apnea in obese Chinese children. *Pediatr Pulmonol* 2006;41:1175-81.
65. Li Y, Chongsuvivatwong V, Geater A, Liu A. Are biomarker levels a good follow-up tool for evaluating obstructive sleep apnea syndrome treatments? *Respiration* 2008;76:317-23.
66. Li AM, Chan MH, Yin J, et al. C-reactive protein in children with obstructive sleep apnea and the effects of treatment. *Pediatr Pulmonol* 2008;43:34-40.
67. Li AM, Ng C, Ng SK, et al. Adipokines in children with obstructive sleep apnea and the effects of treatment. *Chest* 2010;137:529-35.
68. Lin QC, Xie HS, Liu XJ, Zhou JL, Zhao JM. [Relationship between obstructive sleep apnea-hypopnea syndrome and high sensitivity C-reactive protein in non-obese subjects]. *Zhonghua Yi Xue Za Zhi* 2013;93:2355-8.
69. Loubaki L, Jacques E, Semlali A, Biardel S, Chakir J, Series F. Tumor necrosis factor-alpha expression in uvular tissues differs between snorers and apneic patients. *Chest* 2008;134:911-8.
70. Lui MM, Lam JC, Mak HK, et al. C-reactive protein is associated with obstructive sleep apnea independent of visceral obesity. *Chest* 2009;135:950-6.
71. Makino S, Handa H, Suzukawa K, et al. Obstructive sleep apnoea syndrome, plasma adiponectin levels, and insulin resistance. *Clin Endocrinol* 2006;64:12-9.
72. Malakasioti G, Alexopoulos E, Befani C, et al. Oxidative stress and inflammatory markers in the exhaled breath condensate of children with OSA. *Sleep Breath* 2012;16:703-8.
73. Mancuso M, Bonanni E, LoGerfo A, et al. Oxidative stress biomarkers in patients with untreated obstructive sleep apnea syndrome. *Sleep Med* 2012;13:632-6.
74. Mehra R, Storer-Isser A, Kirchner HL, et al. Soluble interleukin 6 receptor: a novel marker of moderate to severe sleep-related breathing disorder. *Arch Intern Med* 2006;166:1725-31.
75. Montgomery-Downs HE, Krishna J, Roberts LJ 2nd, Gozal D. Urinary F2-isoprostane metabolite levels in children with sleep-disordered breathing. *Sleep Breath* 2006;10:211-5.
76. Murase K, Mori K, Yoshimura C, et al. Association between plasma neutrophil gelatinase associated lipocalin level and obstructive sleep apnea or nocturnal intermittent hypoxia. *PLoS One* 2013;8:e54184.
77. Norman D, Bardwell WA, Arosemena F, et al. Serum aminotransferase levels are associated with markers of hypoxia in patients with obstructive sleep apnea. *Sleep* 2008;31:121-6.
78. Ntalapascha M, Makris D, Kyparos A, et al. Oxidative stress in patients with obstructive sleep apnea syndrome. *Sleep Breath* 2013;17:549-55.
79. O'Brien LM, Serpero LD, Tauman R, Gozal D. Plasma adhesion molecules in children with sleep-disordered breathing. *Chest* 2006;129:947-53.
80. Ohga E, Tomita T, Wada H, Yamamoto H, Nagase T, Ouchi Y. Effects of obstructive sleep apnea on circulating ICAM-1, IL-8, and MCP-1. *J Appl Physiol* 2003;94:179-84.
81. Osorio RS, Ayappa I, Mantua J, et al. The interaction between sleep-disordered breathing and apolipoprotein E genotype on cerebrospinal fluid biomarkers for Alzheimer's disease in cognitively normal elderly individuals. *Neurobiol Aging* 2014;35:1318-24.
82. Oyama J, Yamamoto H, Maeda T, Ito A, Node K, Makino N. Continuous positive airway pressure therapy improves vascular dysfunction and decreases oxidative stress in patients with the metabolic syndrome and obstructive sleep apnea syndrome. *Clin Cardiol* 2012;35:231-6.
83. Ozben S, Guvenc TS, Huseyinoglu N, et al. Low serum copeptin levels in patients with obstructive sleep apnea. *Sleep Breath* 2013;17:1187-92.
84. Pallayova M, Steele KE, Magnuson TH, et al. Sleep apnea predicts distinct alterations in glucose homeostasis and biomarkers in obese adults with normal and impaired glucose metabolism. *Cardiovasc Diabetol* 2010;9:83.
85. Pallayova M, Steele KE, Magnuson TH, et al. Sleep apnea determines soluble TNF-alpha receptor 2 response to massive weight loss. *Obes Surg* 2011;21:1413-23.
86. Papaioannou I, Twigg GL, Kemp M, et al. Melatonin concentration as a marker of the circadian phase in patients with obstructive sleep apnoea. *Sleep Med* 2012;13:167-71.
87. Park CS, Guilleminault C, Hwang SH, Jeong JH, Park DS, Maeng JH. Correlation of salivary cortisol level with obstructive sleep apnea syndrome in pediatric subjects. *Sleep Med* 2013;14:978-84.
88. Patacchioli FR, Tabarrini A, Ghiciuc CM, et al. Salivary biomarkers of obstructive sleep apnea syndrome in children. *Pediatr Pulmonol* 2014;49:1145-52.
89. Patel SR, Zhu X, Storer-Isser A et al. Sleep duration and biomarkers of inflammation. *Sleep* 2009;32:200-4.
90. Peled N, Shitrit D, Bendayan D, Peled E, Kramer MR. Association of elevated levels of vascular endothelial growth factor in obstructive sleep apnea syndrome with patient age rather than with obstructive sleep apnea syndrome severity. *Respiration* 2007;74:50-5.
91. Petrosyan M, Perraki E, Simoes D, et al. Exhaled breath markers in patients with obstructive sleep apnoea. *Sleep Breath* 2008;12:207-15.
92. Phillips CL, Yang Q, Williams A, et al. The effect of short-term withdrawal from continuous positive airway pressure therapy on sympathetic activity and markers of vascular inflammation in subjects with obstructive sleep apnoea. *J Sleep Res* 2007;16:217-25.
93. Pinto P, Barbara C, Montserrat JM, et al. Effects of CPAP on nitrate and norepinephrine levels in severe and mild-moderate sleep apnea. *BMC Pulm Med* 2013;13:13.
94. Przybylowski T, Bielicki P, Kumor M et al. [Exhaled nitric oxide in patients with obstructive sleep apnea syndrome]. *Pneumonol Alergol Pol* 2006;74:21-5.
95. Punjabi NM, Beamer BA. C-reactive protein is associated with sleep disordered breathing independent of adiposity. *Sleep* 2007;30:29-34.
96. Roche F, Gaspoz JM, Pichot V, et al. Association between C-reactive protein and unrecognised sleep-disordered breathing in the elderly. *Eur Respir J* 2009;33:797-803.
97. Rubinsztajn R, Kumor M, Byskiniewicz K, Chazan R [The influence of 3 weeks therapy with continuous positive airway pressure on serum leptin and homocysteine concentration in patients with obstructive sleep apnea syndrome]. *Pneumonol Alergol Pol* 2006;74:63-7.

Appendix 3 (continued)—Excluded articles and reasons for exclusion.

98. Ryan S, Nolan GM, Hannigan E, Cunningham S, Taylor C, McNicholas WT. Cardiovascular risk markers in obstructive sleep apnoea syndrome and correlation with obesity. *Thorax* 2007;62:509-14.
99. Salord N, Gasa M, Mayos M, et al. Impact of OSA on biological markers in morbid obesity and metabolic syndrome. *J Clin Sleep Med* 2014;10:263-70.
100. Schulz R, Hummel C, Heinemann S, Seeger W, Grimminger F. Serum levels of vascular endothelial growth factor are elevated in patients with obstructive sleep apnea and severe nighttime hypoxia. *Am J Respir Crit Care Med* 2002;165:67-70.
101. Shamsuzzaman AS. Elevated C-Reactive protein in patients with obstructive sleep apnea. *Circulation* 2002;105:2462-64.
102. Shi YK, Chen JX, Huang Y, Li AY. Serum S100A12 levels are associated with the presence and severity of obstructive sleep apnea syndrome in male patients. *Sleep Breath* 2014;18:269-74.
103. Simiakakis M, Kapsimalis F, Chaligiannis E, Loukides S, Sitaras N, Alchanatis M. Lack of effect of sleep apnea on oxidative stress in obstructive sleep apnea syndrome (OSAS) patients. *PloS One* 2012;7:e39172.
104. Sokucu SN, Karasulu L, Dalar L, Seyhan EC, Altin S. Can red blood cell distribution width predict severity of obstructive sleep apnea syndrome? *J Clin Sleep Med* 2012;8:521-5.
105. Staats R, Stoll P, Zingler D, Virchow JC, Lommatsch M. Regulation of brain-derived neurotrophic factor (BDNF) during sleep apnoea treatment. *Thorax* 2005;60:688-92.
106. Stefanini Dde O, Barros EL, Stefanini R, Pradella-Hallinan ML, Pignatari SS, Fujita RR. Comparing the clinical profile of non obese children with sleep apnea and snoring. *Braz J Otorhinolaryngol* 2012;78:22-6.
107. Steiropoulos P, Papanas N, Nena E, et al. Inflammatory markers in middle-aged obese subjects: does obstructive sleep apnea syndrome play a role? *Mediators Inflamm* 2010;2010:675320.
108. Sukegawa M, Noda A, Sugiura T, et al. Assessment of continuous positive airway pressure treatment in obstructive sleep apnea syndrome using 24-hour urinary catecholamines. *Clin Cardiol* 2005;28:519-22.
109. Svensson M, Venge P, Janson C, Lindberg E. Relationship between sleep-disordered breathing and markers of systemic inflammation in women from the general population. *J Sleep Res* 2012;21:147-54.
110. Takahashi K, Chin K, Nakamura H, et al. Plasma thioredoxin, a novel oxidative stress marker, in patients with obstructive sleep apnea before and after nasal continuous positive airway pressure. *Antioxid Redox Signal* 2008;10:715-26.
111. Tauman R, Ivanenko A, O'Brien LM, Gozal D. Plasma C-reactive protein levels among children with sleep-disordered breathing. *Pediatrics* 2004;113:e564-9.
112. Tauman R, Serpero LD, Capdevila OS, et al. Adipokines in children with sleep disordered breathing. *Sleep* 2007;30:443-9.
113. Ting H, Lo HS, Chang SY, et al. Post- to pre-overnight sleep systolic blood pressures are associated with sleep respiratory disturbance, pro-inflammatory state and metabolic situation in patients with sleep-disordered breathing. *Sleep Med* 2009;10:720-5.
114. Tual-Chalot S, Gagnadoux F, Trzepizur W, Priou P, Andriantsitohaina R, Martinez MC. Circulating microparticles from obstructive sleep apnea syndrome patients induce endothelin-mediated angiogenesis. *Biochim Biophys Acta* 2014;1842:202-7.
115. Ucar ZZ, Taymaz Z, Erbaycu AE, Kirakli C, Tuksavul F, Guclu SZ. Nocturnal hypoxia and arterial lactate levels in sleep-related breathing disorders. *South Med J* 2009;102:693-700.
116. Uysal A, Liendo C, McCarty DE et al. Nocturnal hypoxemia biomarker predicts sleepiness in patients with severe obstructive sleep apnea. *Sleep Breath* 2014;18:77-84.
117. Van Hoorenbeeck K, Franckx H, Debode P, et al. Weight loss and sleep-disordered breathing in childhood obesity: effects on inflammation and uric acid. *Obesity* 2012;20:172-7.
118. Vavougiou G, Pastaka C, Tsilioni I, et al. The DJ-1 protein as a candidate biomarker in obstructive sleep apnea syndrome. *Sleep Breath*. 2014 Feb 15. [Epub ahead of print].
119. Villa MP, Supino MC, Fedeli S, et al. Urinary concentration of 8-isoprostane as marker of severity of pediatric OSAS. *Sleep Breath*. 2014 Jan 17. [Epub ahead of print].
120. Wang YN, Yang Y, Luo YQ, Chen LL. [Effects of nasal continuous positive airway pressure short-term treatment on C-reactive protein and intercellular adhesion molecule-1 in patients with obstructive sleep apnea-hypopnea syndrome]. *Zhong Nan Da Xue Xue Bao Yi Xue Ban* 2005;30:666-9.
121. Wang Y, Wang JJ, Zhao MQ, Liu SM, Li YZ. Changes of serum brain-derived neurotrophic factor in children with obstructive sleep apnoea-hypopnoea syndrome following adenotonsillectomy. *J Int Med Res* 2010;38:1942-51.
122. Wang W, Redline S, Khoo MC. Autonomic markers of impaired glucose metabolism: effects of sleep-disordered breathing. *J Diabetes Sci Technol* 2012;6:1159-71.
123. Wang Q, Feng X, Zhou C, Li P, Kang J. Decreased levels of serum omentin-1 in patients with obstructive sleep apnoea syndrome. *Ann Clin Biochem* 2013;50:230-5.
124. Wang Y, Li Y, Chen P, Luo Y, Yang Y, Yang Y. Elevated fractalkine in patients with obstructive sleep apnea hypopnea syndrome. *Sleep Breath* 2013;17:203-8.
125. Wang X, Xing GH. Serum YKL-40 concentrations are elevated and correlated with disease severity in patients with obstructive sleep apnea syndrome. *Scand J Clin Lab Invest* 2014;74:74-8.
126. Yamauchi M, Nakano H, Maekawa J et al. Oxidative stress in obstructive sleep apnea. *Chest* 2005;127:1674-9.
127. Ye J, Liu H, Li Y, Liu X, Zhu JM. Increased serum levels of C-reactive protein and matrix metalloproteinase-9 in obstructive sleep apnea syndrome. *Chinese Med J* 2007;120:1482-6.
128. Ye L, Ma GH, Chen L, et al. Quantification of circulating cell-free DNA in the serum of patients with obstructive sleep apnea-hypopnea syndrome. *Lung* 2010;188:469-74.
129. Yokoe T. Elevated levels of C-Reactive Protein and Interleukin-6 in patients with obstructive sleep apnea syndrome are decreased by nasal continuous positive airway pressure. *Circulation* 2003;107:1129-34.
130. Zamarron C, Ricoy J, Riveiro A, Gude F. Plasminogen activator inhibitor-1 in obstructive sleep apnea patients with and without hypertension. *Lung* 2008;186:151-6.
131. Zamarron C, Riveiro A, Gude F. Circulating levels of vascular endothelial markers in obstructive sleep apnoea syndrome. Effects of nasal continuous positive airway pressure. *Arch Med Sci* 2011;7:1023-8.
132. Zhang XB, Lin QC, Deng CS, Chen GP, Cai ZM, Chen H. Elevated serum cystatin C in severe OSA younger men without complications. *Sleep Breath* 2013;17:235-41.

Appendix 4—QUADAS criteria fulfilled.

Item	Benedek et al. [23] *	Gozal et al. [24]	Guo et al. [28]	Hirotsu et al. [25]	Kheirandish-Gozal et al. [31]	Lentini et al. [29]	Li et al. [26]	Shah et al. [30]	Ursavas et al. [27]
1. Was the spectrum of patients representative of the patients who will receive the test in the practice?	N	N	N	N	N	N	N	N	N
2. Were selection criteria clearly described?	Y	Y	Y	Y	Y	Y	Y	Y	Y
3. Is the reference standard likely to correctly classify the target condition?	Y	Y	Y	Y	Y	Y	Y	Y	Y
4. Is the time period between reference standard and index test short enough to be reasonably sure that the target condition did not change between the two tests?	Y	Y	Y	Y	Y	Y	Y	Y	Y
5. Did the whole sample or a random selection of the sample, receive verification using a reference standard of diagnosis?	Y	Y	Y	Y	Y	Y	Y	Y	Y
6. Did patients receive the same reference standard regardless of the index test results?	Y	Y	Y	Y	Y	Y	Y	Y	Y
7. Was the reference standard independent of the index test (i.e. the index test did not form part of the reference standard)?	Y	Y	Y	Y	Y	Y	Y	Y	Y
8. Was the execution of the index test described in sufficient detail to permit replication of the test?	Y	Y	Y	Y	Y	Y	Y	Y	Y
9. Was the execution of the reference standard describes in sufficient detail to permit its replication?	Y	Y	Y	Y	Y	Y	Y	Y	Y
10. Were the index test results interpreted without knowledge of the results of the reference standard?	U	U	U	U	U	U	U	U	U
11. Were the reference standard results interpreted without knowledge of the results of the index test?	U	U	U	U	U	U	U	U	U
12. Were the same clinical data available when test results were interpreted as would be available when the test used in practice?	Y	Y	Y	Y	Y	Y	Y	Y	Y
13. Were uninterpretable/ intermediate test results reported?	Y	Y	Y	Y	Y	Y	Y	Y	Y
14. Were withdrawals from the study explained?	Y	Y	Y	Y	Y	Y	Y	Y	Y
QUADAS CRITERIA FULFILLED (%)	78.6	78.6	78.6	85.7	78.6	78.6	78.6	78.6	78.6

Y, yes; N, no; U, unclear. *Numbers in brackets refer to reference number in original manuscript.