Chemotherapy-Induced Nausea and Vomiting #285

Nishant Tageja, MD and Hunter Groninger, MD, FAAHPM

Background

CLINICAL TRIALS BASED ON PRECLINICAL STUDIES have better defined the role of pharmacotherapies in controlling both acute and delayed onset chemotherapy-induced nausea and vomiting (CINV).^{1–6} This *Fast Fact* offers a critical appraisal of CINV treatment guidelines, specifically considering new antiemetic drugs.

Pathophysiology and Targeted Pharmacotherapy

As a response to chemotherapy, substance P, cholecystokinin, and 5-hydroxytryptamine (5-HT₃) are segregated from enterochromaffin cells in the gastrointestinal mucosa.⁷ 5-HT₃ and NK-1 receptors in vagal afferent nerves, along with central 5-HT₂, NK-1, dopamine, serotonin, histamine, GABA, and cannabinoid receptors in the brain stem chemoreceptor trigger zone and vomiting center, process the emetogenic stimuli leading to nausea and/or vomiting.^{8–10}

Classification

Historically, CINV has been classified according to time of onset.

- **Acute CINV** occurs within minutes to several hours after chemotherapy administration, reaching maximal intensity after 5–6 hours, and usually resolving within 24 hours.^{11,12}
- **Delayed CINV** occurs more than 24 hours (peaking at 48–72 hours) after chemotherapy.^{11,12}
- **Anticipatory emesis** precedes drug administration, and generally occurs after a previous negative vomiting experience with chemotherapy.¹³

Drug and guideline development have focused on the degree of emetogenicity of a chemotherapy regimen: highly emetogenic chemotherapy (HEC) drugs (CINV in at least

90% of patients after chemotherapy); moderately emetogenic (MEC) drugs (30%–90%); low emetogenic (LEC) drugs (10%–30%); and minimal emetogenicity (<10%). Unfortunately, most of the research has focused on preventing episodes of vomiting. Less is known about reducing nausea independent of vomiting.

Guidelines: Highly Emetogenic Chemotherapy

Expert guidelines recommend similar prophylaxis regimens:^{4–6}

- Glucocorticoid: dexamethasone 12 mg PO/IV day 1, then 8 mg PO days 2–4.
- 5-HT₃-receptor antagonist PO/IV on day 1 (e.g., palonosetron, granisetron). First-generation 5-HT₃ receptor antagonists are no longer recommended for routine use post-chemotherapy for delayed CINV.¹⁴
- NK-1 antagonist: aprepitant 125 mg PO day 1, then 80 mg PO days 2–4, or fosaprepitant 150 mg IV day 1 only.

Key points:

- A meta-analysis in 2000 concluded there is strong evidence for glucocorticoids' efficacy for prophylaxis of acute and delayed CINV with both HEC and MEC.²
- First-generation 5-HT₃ receptor antagonists (e.g., ondansetron, granisetron, dolasetron, tropisetron) are effective in controlling acute CINV when combined with corticosteroids. None is clearly superior over the other.¹
- Palonosetron (a second-generation 5-HT₃ receptor antagonist) has a longer half-life (about 40 hr), stronger binding affinity for the receptor, and does not cause QT prolongation. Some guidelines prefer palonosetron to first-generation 5-HT₃ antagonists;^{5,6} however, key comparative studies did not include NK-1 antagonists, which are now widely used.

Fast Facts and Concepts are edited by Drew A. Rosielle, MD (University of Minnesota Medical School and Fairview Health Services) and Sean Marks, MD (Medical College of Wisconsin), with the generous support of a volunteer peer review editorial board, and are made available online by the Center to Advance Palliative Care (www.capc.org). *Fast Facts and Concepts* are editorially independent of the Center to Advance Palliative Care, and the authors of each individual *Fast Fact* are solely responsible for that *Fast Fact's* content. The full set of *Fast Facts* are available at www.capc.org, along with contact information, and how to reference *Fast Facts*.

Copyright: All *Fast Facts and Concepts* are published under a Creative Commons Attribution-Noncommercial 4.0 International Copyright (creativecommons.org/licenses/by-nc/4.0/). *Fast Facts* can only be copied and distributed for noncommercial, educational purposes. If you adapt or distribute a *Fast Fact*, let us know!

Disclaimer: *Fast Facts and Concepts* provide educational information for health care professionals. This information is not medical advice. *Fast Facts* are not always continually updated, and new safety information may emerge after a *Fast Fact* is published. Health care providers should always exercise their own independent clinical judgment. Some *Fast Facts* cite the use of a product in a dosage, for an indication, or in a manner other than that recommended in the product labeling. Accordingly, the official prescribing information should be consulted before any such product is used.

- NK-1 receptor antagonists aprepitant and fosaprepitant have been approved for prevention of acute and delayed CINV from HEC and MEC. Fosaprepitant, a water soluble pro-drug that is converted to aprepitant, has a longer half-life and more convenient dosing but is not otherwise known to be superior to aprepitant.^{15,16}
- For patients with refractory CINV following HEC, consider substituting olanzapine for the NK-1 antagonist.¹⁷
- Evidence for integrative medicine strategies was not strong enough to be included in current CINV guidelines.

Guidelines: Moderately Emetogenic Chemotherapy

For MEC drug regimens, expert guidelines recommend a combination of corticosteroid and 5-HT₃ receptor antagonist to manage acute CINV. As noted above, certain guidelines recommend palonosetron over first-generation 5-HT₃ antagonists, although the evidence behind this is modest (level IIB evidence). For refractory CINV following MEC, consider adding a NK-1 antagonist.¹⁸

Guidelines: Low and Minimally Emetogenic Chemotherapy

For patients on LEC, a single antiemetic agent such as 5-HT₃-receptor antagonist, dexamethasone, or a phenothiazine should be used prior to chemotherapy; and an antiemetic following chemotherapy should be administered only if indicated. No antiemetic should be routinely administered prior to minimally emetogenic chemotherapy.

Anticipatory Nausea

Patients should be counseled regarding the risks of CINV prior to the first course of chemotherapy to prevent anticipatory CINV. Prophylactic antiemetic therapy should be begun and the use of anti-anxiety medications should be considered prior to the first cycle of chemotherapy to ensure optimal outcomes and prevent anticipatory CINV. Although clinical data is limited, mind-body therapies (e.g., hypnosis, biofeedback, relaxation training, guided imagery) can help prevent or lessen anticipatory CINV.¹⁹

Breakthrough Nausea

Breakthrough nausea/vomiting, defined as an event that occurs despite best preventative therapy, probably will not respond to the same class of drugs as used during unsuccessful prophylaxis. When patients develop nausea or vomiting post-chemotherapy on days 1 to 5 despite adequate prophylaxis, consider a three-day regimen of oral olanzapine (10 mg PO daily) or oral metoclopramide (10 mg PO tid).²⁰ A blinded randomized control trial (RCT) found olanzapine to be more effective than metoclopramide in this situation.²⁰ Although evidence is limited, acupuncture/acupressure may be a useful adjunct strategy to manage acute chemotherapy-induced vomiting.²¹

References

1. Jordan K, Hinke A, Grothey A, et al.: A meta-analysis comparing the efficacy of four 5-HT₃-receptor antagonists for acute chemotherapy-induced emesis. *Support Care Cancer* 2007;15:1023–1033.
2. Ioannidis JP, Hesketh PJ, Lau J: Contribution of dexamethasone to control of chemotherapy-induced nausea and vomiting: A meta-analysis of randomized evidence. *J Clin Oncol* 2000;18:3409–3422.
3. Grunberg S, Chua D, Maru A, et al.: Single-dose fosaprepitant for the prevention of chemotherapy-induced nausea and vomiting associated with cisplatin therapy: Randomized, double-blind study protocol—EASE. *J Clin Oncol* 2011;29:1495–1501.
4. Roila F, Herrstedt J, Aapro M, et al.: Guideline update for MASCC and ESMO in the prevention of chemotherapy- and radiotherapy-induced nausea and vomiting: Results of the Perugia consensus conference. *Ann Oncol* 2010;21:232–243.
5. Basch E, Prestrud AA, Hesketh PJ, et al.: Antiemetic American Society Clinical Oncology clinical practice guideline update. *J Clin Oncol* 2011;29:4189–4198.
6. National Comprehensive Cancer Network: NCCN Clinical Practice Guidelines in Oncology version 1. Antiemesis. National Comprehensive Cancer Network, 2012. www.nccn.org/professionals/physician_gls/f_guidelines.asp#antiemesis. (Last accessed June 20, 2013.)
7. Stern RM, Koch KL, Andrews PLR (eds): *Nausea: Mechanisms and Management*. New York: Oxford University Press, 2011.
8. Frame DG: Best practice management of CINV in oncology patients: Physiology and treatment of CINV: Multiple neurotransmitters and receptors and the need for combination therapeutic approaches. *J Support Oncol* 2010;8(2 suppl 1):5–9.
9. Hesketh PJ, Van Belle S, Aapro M, et al.: Differential involvement of neurotransmitters through the time course of cisplatin-induced emesis as revealed by therapy with specific receptor antagonists. *Eur J Cancer* 2003;39:1074–1080.
10. Darmani NA, Chebolu S, Amos B, Alkam T: Synergistic antiemetic interactions between serotonergic 5-HT₃ and tachykinergic NK1-receptor antagonists in the least shrew (*Cryptotis parva*). *Pharmacol Biochem Behav* 2011;99:573–579.
11. Kris MG, Gralla RJ, Clark RA, et al.: Incidence, course, and severity of delayed nausea and vomiting following the administration of high-dose cisplatin. *J Clin Oncol* 1985;3:1379–1384.
12. Roila F, Boschetti E, Tonato M, et al.: Predictive factors of delayed emesis in cisplatin-treated patients and antiemetic activity and tolerability of metoclopramide or dexamethasone: A randomized single-blind study. *Am J Clin Oncol* 1991;14:238–242.
13. Moher D, Arthur AZ, Pater JL: Anticipatory nausea and/or vomiting. *Cancer Treat Rev* 1984;11:257–264.
14. Geling O, Eichler HG: Should 5-hydroxytryptamine-3 receptor antagonists be administered beyond 24h after chemotherapy to prevent delayed emesis? Systematic re-evaluation of clinical evidence and drug cost implications. *J Clin Oncol* 2005;23:1289–1294.
15. Lasseter KC, Gambale J, Jin B, et al.: Tolerability of fosaprepitant and bioequivalency to aprepitant in healthy subjects. *J Clin Pharmacol* 2007;47:834–840.
16. Van Laere K, De Hoon J, Bormans G, et al.: Equivalent dynamic human brain NK1-receptor occupancy following single-dose i.v. fosaprepitant vs. oral aprepitant as assessed by PET imaging. *Clin Pharmacol Ther* 2012;92:243–250.

17. Navari RM, Gray SE, Kerr AC: Olanzapine versus aprepitant for the prevention of chemotherapy-induced nausea and vomiting: A randomized phase III trial. *J Support Oncol* 2011;9:188–195.
18. Rapoport BL, Jordan K, Boice JA, et al.: Aprepitant for the prevention of chemotherapy-induced nausea and vomiting associated with a broad range of moderately emetogenic chemotherapies and tumor types: A randomized, double-blind study. *Support Care Cancer* 2010;18:423–431.
19. Mansky PJ, Wallerstedt DB: Complementary medicine in palliative care and cancer symptom management. *Cancer J* 2006;12:425–431.
20. Navari RM, Gray SE, Kerr AC: The use of olanzapine versus metoclopramide for the treatment of breakthrough chemotherapy-induced nausea and vomiting in patients receiving highly emetogenic chemotherapy. *Support Care Cancer* 2013;21:1655–1663.
21. Bao T: Use of acupuncture in the control of chemotherapy-induced nausea and vomiting. *J Natl Compr Canc Netw* 2009;7:606–612.

Address correspondence to:

Hunter Groninger, MD, FAAHPM

Pain and Palliative Care Service

Clinical Center, National Institutes of Health

Building 10, Room 2-1733

Bethesda, MD 20892

E-mail: hunter.groninger@nih.gov