Published in final edited form as:

J Mater Chem B Mater Biol Med. 2014 October 14; 2(38): 6634–6638. doi:10.1039/C4TB01109K.

A Benzothiazole-Based Sensor for Pyrophosphate (PPi) and ATP: Mechanistic Insight for Anion-Induced ESIPT Turn-On

Junfeng Wang^a, Xiumin Liu^a, and Yi Pang^{a,b}

Yi Pang: yp5@uakron.edu

^aDepartment of Chemistry, The University of Akron, Akron, Ohio 44325 U.S.A

^bMaurice Morton Institute of Polymer Science, The University of Akron, Akron, Ohio 44325 U.S.A

Abstract

A benzothiazole derivative **2** bearing two 2,2'-dipicolylamine (DPA) groups was examined for its zinc-binding and subsequent anion sensing properties. The study revealed the anion sensing mechanism of polyphosphate anions via sequential binding to two zinc centers, on the basis of both ¹H NMR and mass spectral evidences. The mechanistic insight would provide valuable information for the future design of new excited state intramolecular proton transfer (ESIPT) sensors. In addition, the zinc complex exhibited the solvent-switchable selectivity, responding to pyrophosphate (PPi) in EtOH but adenosine 5'-triphosphate (ATP) in water.

Design of fluorescent chemosensors for biologically relevant anions remains to be a challenging topic.^{1–3} This is due to the lack of general design principle that can reliably translate an anion-binding event to a large fluorescent signal response.⁴ One of the most successful strategies for anion recognition has been the use of coordination complexes that bear one or two vacant coordination sites for binding anionic guests.^{5–8} This strategy can be realized *via* a proper arrangement of two zinc centers, which can selective bind anions such as pyrophosphate (PPi)^{9–16} and adenosine 5'-triphosphate (ATP).^{4,17,18} However, this binding pattern usually induced little spectral shift in the fluorescence signal, which lowers the sensitivity and hampers their practical applications. An ideal fluorescent sensor should exhibit a great fluorescence turn-on with a large spectral shift, in order to minimize the interference from the free dye.¹⁹

The excited state intramolecular proton transfer (ESIPT) has shown to be an attractive mechanism, as the ESIPT ON-OFF could induce a large spectral shift.²⁰ Few attempts have been made to incorporate the ESIPT mechanism in the PPi sensor scheme on the basis of either mononuclear (**1a**-Zn),²¹ which responds to diphosphate ions (H₂P₂O₇²⁻ and ADP), or binuclear zinc complex (**1b**-Zn, where R = “-CH₂DPA-Zn”).²² Herein, we report the ESIPT properties of sensor **2**, a thiazole analogue of **1b**, where the sulfur heteroatom perturbs the ESIPT and related chemical processes. Interestingly, the sensor exhibited the solvent-switchable recognition for pyrophosphate (in alcohol) and ATP (in water). The anion recognition is accompanied with a large spectral shift (~95 nm shift from blue to green),

which is useful for naked eye detection. In addition, the heteroatom substitution makes the two zinc binding sites in **2** distinguishable from each other, thereby allowing close examination of the sensing mechanism step-by-step. The result led to a better understanding about the ESIPT turn-on.

Formation of ligand-zinc complex

Ligand **2** was synthesized as a light yellow syrup-like oil by using a similar strategy with a modified procedure (See ESI, Scheme S1).^{21,22} Our initial attention was paid to the process of zinc binding-induced ESIPT turning-off, by addition of different equiv. of $\text{Zn}(\text{NO}_3)_2$ to **2**. When the first equiv. of Zn^{2+} was added, the absorption peak ($\lambda_{\text{max}} = 343 \text{ nm}$) was red-shifted to $\lambda_{\text{max}} = 367 \text{ nm}$, as a consequence of deprotonation $\text{Ph-OH} \rightarrow \text{Ph-O}^-$ (Figure 1a).^{21,22} The observation clearly indicated that the first Zn^{2+} predominantly bound to “the site I” to disable the ESIPT, which caused the blue fluorescent shift with great fluorescence turn-on (Figure 1b). The second equiv. of Zn^{2+} induced nearly no spectral shift in the absorption, as the Zn^{2+} cation was bound to the second DPA group, resulting in the formation of complex **4**. The binding of the second Zn^{2+} further increased the fluorescence, which is consistent with the previous data (by removing the weak PET effect from the DPA).²¹ Fluorescence quantum yields of **2** ($\phi_{\text{fl}} = 0.01$) and **4** ($\phi_{\text{fl}} = 0.17$) were calculated by using quinine sulfate as reference.

^1H NMR titration further confirmed the assumption. The proton signals at $\sim 8.5 \text{ ppm}$ were attributed to two overlapping doublet from the characteristic pyridine protons H_a and H_a' , which are associated with different zinc binding sites in **2**. Addition of the first equiv. Zn^{2+} only caused downfield shift of H_a at the site **I** (from 8.53 to 8.70), and the second equiv. Zn^{2+} gradually induced downfield shift of H_a' at the site **II** (from 8.56 to 8.60). Similarly, the same trend was observed from the mass experiments (Fig. S1, ESI). Upon addition of the first equiv. of Zn^{2+} , the signal of ligand **2** (TOF-MS- ES^+ at peaks $[\text{2}+\text{H}^+]^+ = 650.3146$ and $[\text{2}+\text{Na}^+]^+ = 672.2971$) almost disappeared (ESI Fig S1) with the formation of a new peak corresponding to $[\text{2}+\text{Zn}^{2+}-\text{H}^+]^+ = 712.2447$. Two-dimensional ESI-traveling wave ion mobility mass spectrometry (TWIM-MS) further revealed that the mass peak at $m/z = 712$ included two signals with drift time at 2.89 ms and 4.96 ms, which were assumed to be **3a** and **3b**, respectively (Fig. 3). It appeared that the mono charged complex **3a** with the drift time 4.96 ms was predominant. Upon addition of the second equiv. Zn^{2+} , the signals of complexes **3** gradually disappeared, and the di-zinc complex **4** was detected as the predominant species with three positive charges: $[\text{2}+2\text{Zn}^{2+}-\text{H}^+]^{3+} = 258.7352$ (Fig. S2, ESI). All the data pointed to that the zinc complex with four coordination (at the site I) is easier to form than that with three coordination (at the site II).

Binding to phosphate anions

The anion response of the resulting zinc complex **4** was examined by addition of 5.0 equiv. of different anions in EtOH. Much to our delight, **4** showed very good selectivity toward H_2PPI , giving the ESIPT turn-on with bright green emission (Figure 4a and 4c). Interestingly, **4** showed selectivity toward H_2ATP over the other anions in aqueous (Fig. 4b–c). The mass spectra from the solution of “**4** + H_2PPI ” in EtOH detected the major species

with $m/z = 952.1031$ and 974.1014 , which matched the isotope patterns for $[\mathbf{5} + \text{H}^+]^+ = 952.0404$ ($\text{C}_{39}\text{H}_{36}\text{N}_7\text{O}_8\text{P}_2\text{SZn}_2$) and $[\mathbf{5} + \text{Na}^+]^+ = 974.0224$ ($\text{C}_{39}\text{H}_{36}\text{N}_7\text{NaO}_8\text{P}_2\text{SZn}_2$). The observation pointed to the formation of **5**, which is essential to turn on the ESIPT.

Clear differentiation of ^1H NMR signals at two different zinc binding sites allowed us to further examine the details of the pyrophosphate binding. Upon addition of H_2PPi to the zinc complex **4** (i.e. 2-Zn_2), both H_a and H_a' (at 8.6 and 8.7 ppm) disappeared simultaneously to form a new broad peak at ~ 8.8 ppm. The spectral shift suggests that H_2PPi was binding to both site **I** (from H_a shift) and site **II** (from H_a' shift) (Fig. 5a). The observation supports the formation of **5** (Scheme 3), which was the key for ESIPT turn-on mechanism. The PPi adduct **6** might also be possible, since PPi anion can bind to a mono-nuclear zinc center.²¹ The control experiment was carried out by reaction of zinc complex **3a** with PPi, which would give **7**. The ^1H NMR of **7** showed that the H_a at site **I** would have different pattern if PPi was bonded to only one zinc cation (Figure 5b). The result thus consistently pointed to the formation of **5**, and ruled out the presence of **6**.

As seen in Figure 4b–c, the zinc complex showed interesting response to H_2ATP in aqueous solution ($\text{pH} \approx 4.0$ for 2mM H_2ATP). The mass spectrum of the aqueous solution detected the formation of a major species at $m/z = 1281.2192$ and 1303.2233 , which matched the isotope pattern of intermediate $[\mathbf{8} + \text{H}^+]^+$ and $[\mathbf{8} + \text{Na}^+]^+$ respectively (ESI, Fig. S3), also showing the binding of H_2ATP to both binding site **I** and **II**.

The stability of complex **4** was further examined in buffered aqueous at different pH. The study revealed that complex **4** was quite stable in buffered aqueous at different pH (ranges from $\text{pH} = 4.7$ to 7.7) (Fig. 7). When $\text{pH} = 4.7$, the ESIPT turn-on could be accomplished by H_2ATP (Fig. 8). The selectivity **4** to phosphate anions was examined. The result showed that H_2ATP exhibited a stronger tendency to turn-on ESIPT over other phosphate anions such as H_2PPi (Fig. 9), revealing the same trend as observed in DI water (Fig. 4b). The reversal of selectivity in different solvent, i.e. H_2PPi in EtOH and H_2ATP in water (or acidic buffer), showed that the anion binding-induced ESIPT turn-on was quite sensitive to the solvent environment, which should be further examined. Clearly, the current sensor worked better under the acidic conditions, as the ESIPT turn-on requires an H^+ to generate the phenol: $\text{Ph-O}^- + \text{H}^+ \rightarrow \text{Ph-OH}$.

In conclusion, benzothiazole derivative **2** bearing two DPA groups was examined for its zinc-binding and subsequent anion sensing properties. The binuclear zinc complex was formed *via* sequential zinc chelating at two different DPA groups. Spectral data analysis, including UV-vis, fluorescence, ^1H NMR and ESI-Mass spectra, suggested that the polyphosphate anions was likely binding to two zinc centers during the sensing process. Unexpectedly, the zinc complex **4** showed very good selectivity toward H_2PPi in EtOH and H_2ATP in H_2O , respectively, over the other anions. The intriguing selectivity is accompanied with ESIPT turn-on (from blue color to green color) to generate a large spectral shift, making it possible for naked-eye detection. The solvent-switchable selectivity suggests that the ESIPT probe is quite sensitive to the conditions used, and has the potential to differentiate structurally similar anions (PPi and ATP). The study also revealed the anion

sensing mechanism, on the basis of both ^1H NMR and mass spectral evidences. The mechanistic insight would provide useful guide for the future design of new ESIPT sensors.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work was supported by National Institute of Health (Grant No: 1R15EB014546-01A1). We also thank the Ohio Department of Development and Coleman endowment from the University of Akron for partial support.

Notes and references

1. O'Neil EJ, Smith BD. Coord. Chem. Rev. 2006; 250:3068.
2. Martínez-Máñez R, Sancenón F. Chem. Rev. 2003; 103:4419. [PubMed: 14611267]
3. Beer PD, Gale PA. Angew. Chem. Int. Ed. 2001; 40:486.
4. Ojida A, Takashima I, Kohira T, Nonaka H, Hamachi I. J. Am. Chem. Soc. 2008; 130:12095. [PubMed: 18700758]
5. Beer PD, Bayly SR. Top. Curr. Chem. 2005; 255:125.
6. Steed JW. Chem. Soc. Rev. 2009; 38:506. [PubMed: 19169464]
7. O'Neil EJ, Smith BD. Coord. Chem. Rev. 2006; 250:3068.
8. Fabbrizzi L, Licchelli M, Taglietti A. Dalton Trans. 2003:3471.
9. Ojida A, Mito-oka Y, Inoue Ma, Hamachi I. J. Am. Chem. Soc. 2002; 124:6256. [PubMed: 12033851]
10. Lee DH, Im JH, Son SU, Chung YK, Hong JI. J. Am. Chem. Soc. 2003; 125:7752. [PubMed: 12822964]
11. Ojida A, Mito-oka Y, Sada K, Hamachi I. J. Am. Chem. Soc. 2004; 126:2454. [PubMed: 14982454]
12. Ojida A, Nonaka H, Miyahara Y, Tamaru S, Sada K, Hamachi I. Angew. Chem, Int. Ed. 2006; 45:5518.
13. McDonough MJ, Reynolds AJ, Lee WYG, Jolliffe KA. Chem. Commun. 2006; 2006:2971.
14. Lee HN, Xu Z, Kim SK, Swamy KMK, Kim Y, Kim SJ, Yoon J. J. Am. Chem. Soc. 2007; 129:3828. [PubMed: 17348656]
15. Rhee HW, Lee CR, Cho SH, Song MR, Cashel M, Choy HE, Seok YJ, Hong JI. J. Am. Chem. Soc. 2008; 130:784. [PubMed: 18166051]
16. Shin IS, Bae SW, Kim H, Hong JI. Anal. Chem. 2010; 82:8259. [PubMed: 20843022]
17. Ngo HT, Liu X, Jolliffe KA. Chem. Soc. Rev. 2012; 41:4928. [PubMed: 22688834]
18. Kurishita Y, Kohira T, Ojida A, Hamachi I. J. Am. Chem. Soc. 2010; 132:13290. [PubMed: 20812684]
19. (a) Peng XJ, Song FL, Lu EH, Wang YN, Zhou W, Fan JL, Gao YL. J. Am. Chem. Soc. 2005; 127:4170. [PubMed: 15783189] (b) Tolosa, L.; Nowaczyk, K.; Lakowicz, J. An Introduction to Laser Spectroscopy. 2nd ed.. New York: Kluwer; 2002. (c) Zhang Z, Achilefu S. Org. Lett. 2004; 6:2067. [PubMed: 15176820] (d) Wang JF, Pang Y. RSC Advances. 2013; 3:10208.
20. (a) Taki M, Wolford JL, O'Halloran TV. J. Am. Chem. Soc. 2004; 126:712. [PubMed: 14733534] (b) Hu R, Feng J, Hu DH, Wang SQ, Li SY, Li Y, Yang GQ. Angew. Chem., Int. Ed. 2010; 49:4915. (c) Wu JS, Liu WM, Ge JC, Zhang HY, Wang PF. Chem. Soc. Rev. 2011; 40:3483. [PubMed: 21445455] (d) Xu Z, Xu L, Zhou J, Xu YF, Zhu WP, Qian XH. Chem. Commun. 2012; 48:10871. (e) Kim T, Kang HJ, Han G, Chung SJ, Kim Y. Chem. Commun. 2009:5985. (f) Wang JF, Li YB, Duah E, Paruchuri S, Zhou DM, Pang Y. J. Mater. Chem. B. 2014; 2:2008.
21. Wang JF, Chen WH, Liu XM, Wesdemiotis C, Pang Y. J. Mater. Chem. B. 2014; 2:3349.

22. Chen W, Xing Y, Pang Y. Org. Lett. 2011; 13:1262. (b) The sensor **1b-Zn** was found to respond also to HADP in ethanol solution.

Fig. 1. UV-vis (left) and fluorescence (right) titration of **2** (10 μM) upon addition of different equiv. of Zn^{2+} in EtOH.

Fig. 2.

 ^1H NMR titration of 2 upon addition of different equiv. of Zn^{2+} in CD_3OD .

Fig. 3.

Two-dimensional ESI TWIM MS plot for m/z at 712. Ion mobility separation was effected using a traveling wave height of 9V and a traveling wave velocity of 350 m/s. The precursor ion from m/z at 712 gives rise to signals at 2.89 and 4.96 ms; on the basis of the isotope patterns observed, the signals at 2.89 and 4.96 ms agree well with $[(2+\text{Zn}^{2+})_2-2\text{H}^+]^{2+}$ (**3b**) and $[2+\text{Zn}^{2+}-\text{H}^+]^+$ (**3a**).

Fig. 4.

Fluorescence response of Zn complex **4** (10 μM) with different anions (50 μM) in ethanol (a) and different anions (2 mM) in water (b), while the dye was excited at the isobestic point ~ 357 nm. The fluorescence image (c) shows the response of **4** to various phosphate anions in ethanol (top) and DI water (bottom).

Fig. 5.
 ^1H NMR spectra of **4** (a) and **3** (b) upon addition of H_2PPI in CD_3OD .

Fig. 6.
Possible formation of H₂ATP adducts from zinc complexes in water.

Fig. 7. Fluorescence intensity of **4** in PBS buffer at different pH. The inset plot the stability of the complex at different pH (4.7–7.7).

Fig. 8.

Fluorescence intensity of **4** (10 μM) in PBS buffer at different pH points, upon addition of 2 mM of H₂ATP.

Fig. 9. Fluorescence intensity of **4** (10 μ M) in PBS buffer (pH = 4.7), upon addition of 2 mM of phosphate anions.

Scheme 1.

Structure of **1**, the zinc complexes and the ESIPT OFF-ON switch induced by H₂PPI.

Scheme 2.

Structure of **2** with two different zinc-binding sites (**I** and **II**), and its zinc complex formation.

Scheme 3.

Possible formation of PPi adducts from zinc complexes.