Published in final edited form as:

Epilepsy Res. 2015 January ; 0: 183–196. doi:10.1016/j.epilepsyres.2014.11.009.

Spatiotemporal differences in the c-fos pathway between C57BL/6J and DBA/2J mice following flurothyl-induced seizures: a dissociation of hippocampal Fos from seizure activity

Sridhar B. Kadiyala^{1,a}, Dominick Papandrea^{1,2,a}, Karina Tuz¹, Tara M. Anderson², Sachidhanand Jayakumar¹, Bruce J. Herron^{3,4}, and Russell J. Ferland^{1,2,5,*}

¹Center for Neuropharmacology and Neuroscience, Albany Medical College, Albany, NY 12208, USA

²Department of Biology, Center for Biotechnology and Interdisciplinary Studies, Rensselaer Polytechnic Institute, Troy, NY 12180, USA

³Wadsworth Center, Albany, NY 12201, USA

⁴Department of Biomedical Sciences, School of Public Health, SUNY Albany, Albany, NY 12201, USA

⁵Department of Neurology, Albany Medical College, Albany, NY 12208, USA

Abstract

Significant differences in seizure characteristics between inbred mouse strains highlight the importance of genetic predisposition to epilepsy. Here, we examined the genetic differences between the seizure-resistant C57BL/6J (B6) mouse strain and the seizure-susceptible DBA/2J (D2) strain in the phospho-Erk and Fos pathways to examine seizure-induced neuronal activity to uncover potential mechanistic correlates to these disparate seizure responsivities. Expression of neural activity markers was examined following 1, 5, or 8 seizures, or after 8 seizures, a 28 day rest period, and a final flurothyl rechallenge. Two brain regions, the hippocampus and ventromedial nucleus of the hypothalamus (VMH), had significantly different Fos expression profiles following seizures. Fos expression was highly robust in B6 hippocampus following one seizure and remained elevated following multiple seizures. Conversely, there was an absence of Fos (and phospho-Erk) expression in D2 hippocampus following one generalized seizure that increased with multiple seizures. This lack of Fos expression occurred despite intracranial electroencephalographic recordings indicating that the D2 hippocampus propagated ictal discharge during the first flurothyl seizure suggesting a dissociation of seizure discharge from Fos and phospho-Erk expression. Global transcriptional analysis confirmed a dysregulation of the c-fos

© 2014 Elsevier B.V. All rights reserved.

*Correspondence to : Russell J. Ferland, Ph.D., Albany Medical College, Center for Neuropharmacology and Neuroscience, 47 New Scotland Avenue, MC-136, Albany, NY 12208, Phone: (518) 262-0172, Fax: (518) 262-5799, ferlanr@mail.amc.edu.

^aThese authors contributed equally to this work

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

pathway in D2 mice following 1 seizure. Moreover, global analysis of RNA expression differences between B6 and D2 hippocampus revealed a unique pattern of transcripts that were co-regulated with Fos in D2 hippocampus following 1 seizure. These expression differences could, in part, account for D2's seizure susceptibility phenotype. Following 8 seizures, a 28 day rest period, and a final flurothyl rechallenge, ~85% of B6 mice develop a more complex seizure phenotype consisting of a clonic-forebrain seizure that uninterruptedly progresses into a brainstem seizure. This seizure phenotype in B6 mice is highly correlated with bilateral Fos expression in the VMH and was not observed in D2 mice, which always express clonic-forebrain seizures upon flurothyl retest. Overall, these results illustrate specific differences in protein and RNA expression in different inbred strains following seizures that precede the reorganizational events that affect seizure susceptibility and changes in seizure semiology over time.

Keywords

epilepsy; seizure; flurothyl; hippocampus; ventromedial nucleus of the hypothalamus; Fos; Erk; genetics; genomics

Introduction

Epileptogenesis is defined as a change that occurs in the brain over time ultimately resulting in increased seizure severity (Dichter, 2006). While epileptogenesis is difficult to investigate in the human brain, the utilization of rodent epilepsy models provides a means to uncover how the epileptic brain changes over time. Understanding epileptogenesis and the pathways involved in its progression will lead to advancements in the treatment of epilepsy.

The repeated-flurothyl model is a seizure paradigm that uses the chemoconvulsant flurothyl to examine seizure susceptibility and epileptogenesis (Applegate, et al., 1997, Ferland and Applegate, 1999, Samoriski and Applegate, 1997). In this model, mice are given 8 flurothyl-induced seizures, one per day, for 8 days. This exposure results in decreasing seizure thresholds over 8 seizure trials and the expression of generalized clonic seizures in all C57BL/6J (B6) mice. Following a 28 day rest period and final flurothyl rechallenge, 70–90% of B6 mice display a new seizure type characterized by a generalized clonic seizure that secondarily generalizes into a brainstem seizure (denoted as a forebrain→brainstem seizure) (Applegate, et al., 1997, Ferland and Applegate, 1999, Samoriski and Applegate, 1997). The emergence of this new, more complex seizure phenotype with time suggests an underlying evolution of an epileptogenic process (Applegate, et al., 1997, Ferland and Applegate, 1999, Samoriski and Applegate, 1997), and involves the recruitment of new structures into the seizure propagation network necessary for the expression of a more complex seizure phenotype (Ferland and Applegate, 1998). Thus, the repeated-flurothyl model serves as an important paradigm for elucidating brain structures and mechanisms involved in changes in seizure complexity/severity over time and epileptogenesis.

We have previously demonstrated that inbred mouse strains have distinct seizure responsivities following exposure to this repeated-flurothyl model indicating that genetic factors control the observed seizure traits (Papandrea, et al., 2009, Papandrea, et al., 2009). In fact, both chemical and electrical seizure inducing stimuli have uncovered novel genetic

quantitative trait loci that modify seizure responses (Ferraro, et al., 1995, Ferraro, et al., 2002). Therefore, utilization of genetically divergent inbred mouse strains with differential sensitivity to the repeated-flurothyl model can uncover genetic susceptibilities and mechanistic processes involved in epileptogenesis, ultimately aiding in the elucidation genes that mediate the underlying pathophysiology of epilepsy.

Here, we examined an important aspect of seizure responsivity in the repeated-flurothyl model: interstrain differences in seizure-induced changes in neuronal activity and protein expression. To examine neural activation in specific brain regions following seizures, we utilized Fos expression, a well-established marker for neuronal activation (Sagar, et al., 1988) that has been widely utilized following clonic and tonic seizures to map seizure-induced activity (Andre, et al., 1998, Ferland, et al., 1998, Hirsch, et al., 1997, Klein, et al., 2004, Labiner, et al., 1993, Loscher and Ebert, 1996, McCown, et al., 1995, Ribak, et al., 1997, Samoriski, et al., 1997, Shehab, et al., 1992). However, no studies have examined changes in Fos expression in the brain between inbred strains of mice to map differences in neural activity following seizures. Therefore, the goal of our study was to examine Fos expression differences following exposure to the repeated flurothyl-model both between and within two inbred strains with different seizure responsivities as Fos expression variances may indicate differences in reorganizational processes in mice exposed to the repeated-flurothyl model.

Materials and methods

Animals

Adult male DBA/2J (D2) and C57BL/6J (B6) mice were obtained from Jackson Laboratories at 6 weeks of age. All mice were acclimated to the housing facility for one week prior to seizure testing. Mice were kept on a 12h light-dark cycle with lights on at 6:00 am and were given unlimited access to food and water. Flurothyl testing was conducted between 7:00 am and 1:00 pm. All testing was performed under approval of the Institutional Animal Care and Use Committees of the Albany Medical College, Rensselaer Polytechnic Institute and the Wadsworth Center (NY State Department of Health), in accordance with the National Institutes of Health's *Guide for the Care and Use of Laboratory Animals*.

Experimental design

Mice were acclimated to the testing room for at least 30 min before testing commenced. Mice were placed in a closed chamber and a 10% flurothyl solution (bis(2,2,2-trifluoroethyl) ether; Sigma-Aldrich) was infused at a constant rate of 100 μ l/min through a glass syringe on to a gauze pad suspended at the top of the chamber. Since flurothyl is a highly volatile chemoconvulsant, mice are exposed to the flurothyl vapors immediately. Upon the start of the generalized seizure, defined as a loss of postural control, the top of the chamber was removed exposing the mouse to room air. A single mouse was placed in the chamber at a time and exposed to flurothyl. The latency to a generalized seizure onset (used as a surrogate measurement of the generalized seizure threshold (GST) since the infusion rate is held constant between animals), and the behaviorally assigned seizure grades from the scale listed below were recorded. One flurothyl-induced seizure was administered per day for 8

consecutive days (induction phase). Mice were then given a 28-day rest period (incubation phase) in which the mice remained in their home cage. Following the incubation phase, mice were exposed once again to flurothyl.

The seizure behaviors were scored according to a number classification system (Applegate, et al., 1997, Samoriski and Applegate, 1997): Grade 1 – a loss of posture, clonus of hindlimbs and/or forelimbs, and facial clonus including chewing; Grade 2 -grade 1 and low intensity bouncing; Grade 3 - grade 1 or 2 followed by wild running and hopping; Grade 4 - grade 3 followed by hindlimb and/or forelimb treading; Grade 5 - grade 3 or 4 followed by bilateral tonic extension of the forelimbs; Grade 6 - grade 5 followed by bilateral extension of the hindlimbs; and Grade 7 - grade 6 followed by death. Seizure grades 1–2 are classified as forebrain seizures since these seizures are clonic and require forebrain structures for their expression (Engel, et al., 1978, Ferland, et al., 1998, Lothman, et al., 1985, Namba, et al., 1991, Samoriski and Applegate, 1997, Samoriski, et al., 1997). Grades 3–7 denote a seizure type that begins as a clonic-forebrain seizure where the animal loses its postural control, regains its posture, and rapidly progresses into a brainstem seizure (we refer to such seizures as forebrain→brainstem seizures to denote the seizure progression of this type). Brainstem seizures are denoted brainstem since their expression is controlled by a brainstem seizure network (Browning, et al., 1985, Browning, et al., 1981, Browning, et al., 1981, Ferland and Applegate, 1998, Merrill, et al., 2003, Samoriski, et al., 1997, Samoriski, et al., 1998).

B6 and D2 mice were sacrificed at various time points throughout the repeated-flurothyl model, following 1 seizure (trial 1), 5 seizures (trial 5), 8 seizures (trial 8), or 8 seizures, a 28 day incubation period, and flurothyl retest. Control animals, not having flurothyl exposure, were sacrificed at each time point as the experimental groups.

Fos immunohistochemistry

Fos immunohistochemistry was performed as previously described (Ferland, et al., 1998). Briefly, 90 minutes following the last seizure test (or at the matched time point for controls), all mice were given an overdose of sodium pentobarbital or avertin, and perfused transcardially with phosphate buffered saline (PBS) followed by cold 4% paraformaldehyde (PFA)/PBS. Brains were removed from the skull and post-fixed in 4% PFA/PBS for 2–3 d at 4°C. Brains were then cryoprotected in a 30% sucrose/PBS solution, sectioned into 40–50 µm coronal sections using a freezing microtome (Leica Microsystems), and preserved for free-floating section immunohistochemistry.

For 3,3'-diaminobenzidine (DAB) immunostaining, sections were thoroughly washed in PBS/Triton X-100 (0.04%) (PBS-Tx) and were incubated in 3% hydrogen peroxide for 30 min, followed by additional washes in PBS-Tx. The tissue was blocked in 10% normal goat serum (Chemicon) made in PBS-Tx for at least 1 h. The primary rabbit polyclonal IgG antibody directed against Fos (1:1500; sc-52, Santa Cruz Biotechnology, USA) was diluted 1:10,000 in 0.3% normal goat serum in PBS-Tx and incubated with the tissue sections overnight at 4°C on a shaker. Tissues were then washed 3 times with PBS-Tx. Biotinylated goat anti-rabbit secondary antibodies (1:200) (Vectastain kit; Vector Laboratories) were incubated with the tissue sections for 1 h. Following three more washes in PBS-Tx, tissue sections were incubated with an avidin-biotin-horseradish peroxidase solution for 1 h. The

sections again were extensively washed and the antibody complex visualized using 0.03% hydrogen peroxide and DAB with nickel intensification (Vector Laboratories). Tissue sections were mounted onto microscope slides and coverslipped with Cytoseal 60 (Richard-Allan Scientific). Tissue sections were processed in batches, with each experimental group being represented in each group to control for time/day of staining. Sections were imaged using a Zeiss AxioImager Z1 microscope (Carl Zeiss Microimaging) equipped with an AxioCam MRc camera (Carl Zeiss Microimaging) and processed with AxioVision Rel. 4.5 software.

Analysis of Fos expression

Analysis of Fos expression throughout the brain was scored as follows (modified from Samoriski et al. 1998): 0 - no observed Fos, 1 - scattered Fos expression, 2 - low levels of Fos expression, 3 - moderate levels of Fos staining, 4 - high levels of Fos expression, 5 - very high levels of Fos expression. The score for each brain area per group was averaged to give the overall Fos expression per experimental group. Data were assessed blind to both seizure trial and strain.

To determine the changes in Fos expression in the subregions of the hippocampus, we calculated integrated densities using Image J (Schneider, et al., 2012). Linear adjustments were made uniformly across all images. Thresholds for each image were computed and the integrated intensity of the image was calculated. The integrated density measurement in Image J takes into account the area of the selected region. Given small differences in background staining, the background intensity was determined for each section and subtracted for the integrated density of the selected region.

pERK immunohistochemistry

Fifteen minutes following a seizure, mice (including non-seizure controls) were transcardially perfused with 4% PFA/PBS, brains were harvested, and post-fixed for 2 h at RT (Houser, et al., 2008). Brains were cryoprotected in 30% sucrose/PBS and sectioned in the coronal plane at a thickness of 30 μ m. Tissue slices were mounted on Superfrost Plus microscope slides (ThermoFisher Scientific) and were air-dried overnight at RT. Tissue sections were permeabilized with PBS-Tx and blocked for 1 hr at RT with 10% goat serum. pERK antibodies were diluted (1:500, Cell Signaling) in 1% goat serum in PBS-Tx and incubated overnight at 4°C. Tissue sections were rinsed thoroughly and incubated for 1 h at RT with Alexa Fluor 546 goat anti-rabbit antibody (1:500, Life Technologies), washed and mounted with Fluoromount-G antifade solution (Southern Biotechnology). Images were acquired using a Zeiss Z1 microscope equipped with an ORCA-Flash 4.0 digital camera (Hamamatsu) and Neurolucida software (MBF Bioscience).

Preparation of nuclear extracts for western blot analysis

After euthanizing the animals, brains were quickly removed and immersed in homogenization buffer (described below) at 4°C, followed by dissection of the hippocampus. Each hippocampi was homogenized using a Dounce homogenizer with a B-type pestle in 0.5 mL of ice cold homogenization buffer containing 10 mM Tris-HCl buffer (pH 7.5), 0.32 M sucrose, 1 mM EDTA, 1 mM EGTA, 5 mM dithiothreitol (DTT), 10 mM

sodium β -glycerophosphate, 0.5% Nonidet P-40, and protease and phosphatase inhibitors (Roche Applied Science). After centrifugation at 1,000 g for 10 min at 4° C, the pellet obtained was resuspended in 0.2 mL of ice cold extraction buffer containing 50 mM Tris-HCl buffer (pH 7.5), 10% glycerol, 400 mM NaCl, 1 mM EDTA, 1 mM EGTA, 5 mM DTT, 0.5% Nonidet P-40, and protease and phosphatase inhibitors (Roche Applied Science). Suspensions were kept on a nutator for 30 min at 4° C, followed by centrifugation at 20,000 g for 5 min at 4° C to obtain supernatants as nuclear extracts. Final extracts were stored at -80° C until use.

Western blot

Western blotting was performed as previously described (Hsiao, et al., 2009, Tuz, et al., 2013). Briefly, protein concentrations were determined by bicinchoninic acid (BCA) assay based on protein standards (BCA Protein Assay kit, Thermo Scientific). Protein samples (10 μ g) were boiled in Laemmli buffer for 10 min. Proteins were separated on 10% Tris/Glycine SDS polyacrylamide gels at 100 V for 2 h and were transferred to PVDF microporous membrane (Immobilon-FL, Millipore) for 2 h at 100 V. After blocking the membrane with 5% skim milk in TBS-T [100 mM Tris (pH 7.4), 150 mM NaCl and 0.01% Triton-X100] for 1 h at RT, the membrane was incubated overnight with primary antibodies against Fos (1:200, rabbit polyclonal IgG; sc-52, Santa Cruz Biotechnology) or p84 (1:50,000, mouse monoclonal IgG; ab487, Abcam) diluted in blocking solution. After washing with TBS-T, blots were incubated with a secondary antibody (anti-rabbit HRP, 1:2000 (for Fos detection) or anti-mouse heavy chain, 1:1000 (for p84 detection)). Immunodetection was performed using a chemiluminescent substrate (Super Signal West Femto Maximum Sensitivity substrate, Thermo Scientific) and acquired with a G:Box iChemi XT imaging system (Syngene, Synoptics). p84 was used as a nuclear loading control.

Gene expression analysis

Hippocampi from B6 mice and D2 mice were isolated 120 min after 1, 3 or 8 flurothyl-induced seizures and stored in RNAlater as per the manufacturer's recommendation (Qiagen). Since Fos protein levels are upregulated 90 min following a seizure, we harvested hippocampi for RNA expression 120 min after the final seizure to capture RNA changes that are regulated by seizure-induced Fos protein increases. Individual hippocampi were homogenized in 1 ml of Qiazol (Qiagen) and the aqueous phase was transferred to RNeasy mini columns and processed according to the manufacturer's protocol (Qiagen). RNA integrity was confirmed with an Agilent Bioanalyzer, which gave a RIN value of >8 for all samples.

One hundred micrograms of total RNA from sixteen independent samples were submitted to the Wadsworth Center Genomics core facility for microarray analysis. Samples were labeled with the Agilent Low RNA Linear Amplification kit according to the manufacturer's specifications (Agilent). Cy3-labeled cRNA target samples were hybridized to 4X44k mouse development expression arrays (Agilent). For all samples, dye incorporation frequency was >15.

One color-based gene expression normalization and analysis was carried out using Genespring GX Ver 12.6 with the GCRMA algorithm. Data were filtered to focus analysis on samples with a present or marginal flag in at least one sample. Statistical analysis was carried by an unpaired t-test (by strain) or ANOVA (strain by day) and false positives were minimized with a Benjamini-Hochberg multiple testing correction factor (cutoff, $P < 0.05$). To determine if any pathways or classes of genes were coordinately deregulated following repeated flurothyl-induced seizures, we performed functional annotation clustering using the Database for Annotation, Visualization and Integrated Discovery (DAVID) (Huang, et al., 2009).

Electroencephalographic recordings

Bipolar, twisted electrodes were implanted into the hippocampus of B6 and D2 mice ($n = 4$ /strain) at 6 weeks of age. Mice were anesthetized with isoflurane (induction of surgical plane anesthesia with 3% isoflurane and a maintenance concentration of 1.5% isoflurane) and then placed into a Stoelting mouse stereotaxic apparatus. Twisted bipolar wire electrodes with 0.5 mm separation between tips (250 μ m diameter stainless steel wire electrodes with polyimide insulation, Plastics One) were used. Hippocampal electrodes were placed using coordinates from bregma (AP: 1.6; ML: 0.7; DV: 1.85) (Paxinos and Franklin, 2001). Stainless steel screws were placed over the occipital lobe serving as a ground. Mice were allowed to recover from surgery for 1 week. Electrode-implanted mice received one flurothyl-induced seizure (clonic-forebrain seizure), while recording the electrographic activity in the hippocampus. EEG was recorded using an A-M Systems amplifier (signal amplified at 100X) and filtered (1 Hz to 1000 Hz band pass), and then digitalized at a sampling rate of 1024 with a notch filter of 60Hz. Stored data were analyzed offline for electrographic seizure activity. All recordings of EEG activity were performed using SciWorks software (Datawave Technologies). At the conclusion of the experiment, mice were deeply anesthetized and perfused transcardially with PBS followed by ice cold 4% PFA. Brains were kept in 4% PFA for 24 h, and later cryopreserved in 30% sucrose. Brains were sectioned to verify correct electrode placement.

Statistics

One-way analysis of variance (ANOVA) followed by Newman-Keuls post-hoc comparisons was used to assess changes within strains. Repeated measures ANOVA followed by Newman-Keuls post-hoc comparisons were utilized to determine significance across repeated flurothyl seizure inductions (kindling). GSTs on day 8 of the induction phase and on retest were compared using Student's t-test. Fisher exact tests were used to assess differences in the percentage of animals changing their seizure phenotype following flurothyl retest. Student's t-tests were used to determine significance between strains for the Fos quantification. All statistical analyses were performed using Statistica (StatSoft).

Results

Generalized seizure thresholds and seizure phenotypes observed throughout the repeated-flurothyl model in C57BL/6J and DBA/2J mice

Since flurothyl infusion rates were held constant, latencies to generalized seizures were used as a surrogate for generalized seizure thresholds (GST) (Fig. 1) as has been previously described (Ferland and Applegate, 1999, Franck, et al., 1989, Hashimoto, et al., 2006, Oommen, et al., 2007, Papandrea, et al., 2009, Samoriski and Applegate, 1997, Stafstrom, et al., 1992); however, concentrations of flurothyl necessary for generalized seizure expression are also reported (Table 1). All C57BL/6J (B6) mice tested in the repeated-flurothyl model had a high initial latency to generalized seizure expression compared to DBA/2J (D2) mice, which significantly decreased over the course of eight repeated flurothyl exposures, plateauing on seizure trial 5 (Fig. 1; $F_{7,161}=142.65$, $P<0.0001$; Table 1). However, D2 mice had a low initial latency to generalized seizure expression that remained consistent over the course of repeated flurothyl exposures (Fig. 1; Table 1). All B6 and D2 mice tested displayed clonic-forebrain seizures during the 8 induction phase seizure trials. Following a 28-day incubation period and final flurothyl rechallenge, there was a significant difference in the percentage of B6 mice changing their seizure phenotype to a more complex seizure that did not occur in D2 mice (Fig. 1 (bars); Fisher exact test, $P<0.02$). Approximately 85% of B6 mice displayed a forebrain→brainstem seizure (a seizure characterized as a clonic-forebrain seizure that uninterruptedly progresses into a seizure having brainstem-tonic manifestations) with the remaining ~15% of B6 mice having clonic-forebrain seizures (no change in seizure phenotype to a forebrain→brainstem seizure) (Fig. 1). Conversely, none of the D2 mice displayed a change in seizure phenotype to a forebrain→brainstem seizure upon flurothyl retest (Fig. 1).

Differences in Fos expression within and between C57BL/6J and DBA/2J mice following a single or repeated flurothyl seizures (induction phase)

Given these very disparate seizure characteristics between B6 and D2 mice, we next examined neural activity between these two strains of mice, using Fos as a neural activity readout, to further define their divergent seizure phenotypes. Expression of Fos was examined in various anatomical regions of the brain following 1, 5, or 8 flurothyl-induced clonic-forebrain seizures in B6 and D2 mice (Table 2). Fos expression was significantly different in the hippocampi of B6 and D2 mice and was dependent on the number of flurothyl-induced seizures (Figs. 2 and 3; Table 2).

In B6 hippocampi, 1, 5, or 8 flurothyl-induced seizures resulted in a significant increase in Fos expression in the dentate gyrus (DG), and CA3 and CA1 fields (DG: $F_{4,72}=68.9$, $P<0.0001$; CA3: $F_{4,72}=11.4$, $P<0.0001$; CA1: $F_{4,72}=12.9$, $P<0.0001$) (Figs. 2 and 3). The largest observed increase in Fos expression in the B6 hippocampus was the dentate gyrus (Figs. 2 and 3). However, whereas the D2 hippocampus did show a significant increase in Fos expression following 5 and 8 seizures (DG: $F_{4,52}=108.9$, $P<0.0001$; CA3: $F_{4,52}=7.7$, $P<0.0001$; CA1: $F_{4,52}=9.3$, $P<0.0001$), no significant differences were observed in comparing 0 versus 1 seizure (DG: $P=0.69$; CA3: $P=0.32$; CA1: $P=0.13$) (Figs. 2 and 3).

Direct comparisons of these hippocampal regions between B6 and D2 mice demonstrated significant differences in Fos expression. In DG, while there were no Fos differences at baseline (no seizures), there were significant differences following 1 or repeated seizures with D2 mice never achieving B6 levels of Fos expression (Figs. 2 and 3). In CA1, no Fos differences were noted at baseline, but B6 and D2 differed in the amount of Fos expression only following 1 and 5 seizures (Figs. 2 and 3). No differences in CA3 Fos expression were observed between B6 and D2 mice (Figs. 2 and 3).

To support and extend these results, we performed western blot analyses of hippocampi of B6 and D2 mice following 1, 5, or 8 flurothyl-induced seizures (Fig. 4). In agreement with the immunohistological observations of Fos, there is a significant increase in Fos expression in the hippocampus of B6 mice following one seizure that stays elevated following subsequent seizures and retest ($F_{3,15}=4.64$, $P<0.02$; Fig. 4). Conversely, there were no significant differences in Fos expression in the hippocampi of D2 mice following a single seizure ($P>0.75$; Fig. 4). But with repeated seizures, Fos expression in D2 hippocampi can significantly increase ($F_{3,15}=5.43$, $P<0.01$; Fig. 4). These differences in Fos expression were confirmed at the RNA level for *Fos* (Fig. 5). Interestingly, not only did *Fos* RNA show these changes, but the *Fos*-related genes, *FosB*, *FosI1*, and *FosI2* also showed similar changes in mRNA expression (Fig. 5), suggesting that this signaling pathway maybe be differentially reactive to a single seizure between these two strains of mice.

Phosphorylation of Erk increases in the hippocampus of C57BL/6J mice but not in DBA/2J mice following one flurothyl-induced seizure

Erk1 and Erk2 are mitogen-activated protein kinases that mediate proliferation, differentiation, synaptic plasticity and cell death. Erks are phosphorylated upon activation (pErk) and translocate from the cytosol to the nucleus. Erk1/2 are activated following seizures induced chemically or electrically (Baraban, et al., 1993, Garrido, et al., 1998, Houser, et al., 2008, Kim, et al., 1994). pErk 1/2 activation is upstream of *c-fos* gene induction resulting in increases in *c-fos* mRNA expression (Monje, et al., 2003). Since increased Fos expression occurs following 1 flurothyl-induced seizure in B6 mice, but not in D2 mice, we examined whether pErk1/2 levels were increased differentially in these two strains. As expected, pErk1/2 levels increased in B6 hippocampi after 1 seizure. However, no increases in pErk levels were observed in D2 hippocampi following a single seizure (Figure 6; particularly in the molecular layer).

Intrahippocampal EEG recordings reveal electrographic seizure events in both B6 and D2 mice

Since flurothyl-induced seizures are generalized seizures, it was surprising that the hippocampus of D2 mice did not have an upregulated expression of Fos, particularly in comparison to the robust increase in Fos expression that is seen in B6 hippocampus following 1 seizure (Figs. 2 and 3). To determine whether the lack of increased Fos expression in the hippocampus of D2 mice is a result of a lack of propagation of ictal discharge through the hippocampus, we recorded hippocampal EEG in mice having flurothyl-induced seizures. As expected from the observation in B6 mice of increased hippocampal Fos expression following a seizure, we were able to record ictal discharge

within the hippocampus of B6 mice (Fig. 7). Despite the fact that D2 mice fail to express hippocampal Fos following 1 flurothyl-induced seizure, we were able to record electrographic seizure discharges in the hippocampus following flurothyl-induced seizures (Fig. 7). This indicates that the use of Fos neural activity maps as a short-term window into excitability changes in neurons following a behavioral seizure can have mixed results, since Fos expression may not capture all of the neurons propagating ictal discharge through them.

Gene expression differences in the hippocampus of C57BL/6J and DBA/2J mice following repeated flurothyl-induced seizures

We considered the possibility that the delay in Fos activation in the hippocampus of D2 mice after 1 seizure could impact global transcription levels. Analysis of variance tests were conducted to identify transcripts that were significantly deregulated after 1, 3 or 8 seizures compared to baseline levels. Hierarchical clustering of these significant transcripts identified gene relationships between genetic background, number of seizures experienced, and relative gene expression (Fig. 8A). These analyses yielded a complex pattern of expression that was dependent on both conditions and transcript levels resulting in 5 clusters of genes in the second tier clusters. Cluster 1 is principally defined by genes that are increased following 3 or 8 seizures, while cluster 2 is characterized by genes that increased following 3 seizures, but is reduced following 8 seizures. Cluster 3, which includes Fos, has relatively low levels of expression of cluster members at baseline that increase after 1 seizure in B6, but not in D2 hippocampus. Cluster 4 is characterized by decreases in gene expression following 8 seizures; however, this cluster has comparatively higher levels of expression in D2 hippocampus following 3 seizures compared to B6 hippocampus following 3 seizures. Finally, cluster 5 represents genes that are comparatively higher at baseline and following 1 seizure, than after multiple seizures, but there is a subset of these transcripts that are differentially regulated between D2 and B6 in the opposite direction of Fos after a single flurothyl seizure. That is, this cluster of genes has increased expression at baseline in both D2 and B6, which are also elevated in D2 hippocampus following 1 seizure, but that is decreased in B6 hippocampus following 1 seizure.

Among these clusters, Fos and its family members, along with a relatively small set of transcripts that includes early growth response genes and *Bdnf* (which are known targets of Fos) were increased significantly in B6 hippocampus following 1 seizure (Fig. 8B; cluster 3). Moreover, *JunD* and regulators of G-protein coupled receptors (*Rgs2* and *Rgs4*) were also significantly elevated in B6, with the latter gene family being previously implicated in electrical kindling models, but not acute seizure models (Liang and Seyfried, 2005). However, these increases were not observed in D2 hippocampus following 1 seizure (transcript levels were equivalent to B6 and D2 control hippocampus), with this gene cluster only increasing their expression following 3 seizures, indicating a global delay in transcription of certain sets of genes after 1 seizure in D2 mice. However, an additional group of genes in cluster 5 showed increased expression in D2 after 1 seizure that did not occur in B6 hippocampus following a single seizure, indicating that if there is a global delay in a transcriptional mechanism, it is not exclusively suppressing transcription. Closer analysis of these two subsets of array data revealed a number of genes that could play important roles in epileptogenesis (Fig. 8B). We are currently investigating the nature of

these phenomena to develop a hypothetical model of how these genes could mediate seizure progression.

Differences in Fos expression between C57BL/6J and DBA/2J mice following 8 flurothyl-induced seizures, a 28-day incubation phase, and a final flurothyl retest

When B6 mice are exposed to the repeated-flurothyl model, which consists of 8 flurothyl-induced seizures (induction phase), a 28-day incubation phase, and a final flurothyl retest, this final flurothyl challenge results in ~85% of B6 mice having a more complex seizure than previously observed during the induction phase. This new seizure type consists of a generalized clonic-forebrain seizure (as seen originally during the induction phase) that rapidly and uninterruptedly transitions into a seizure with brainstem and tonic manifestations called a forebrain→brainstem seizure (Papandrea, et al., 2009). Conversely, whereas D2 mice have the same type of seizures as B6 mice during the 8 seizure induction phase, D2 mice do not evolve into a more complex seizure semiology. In our analysis of Fos neural activity maps between B6 and D2 mice following 8 flurothyl-induced seizures (induction phase), a 28-day incubation phase, and a final flurothyl retest, the ventromedial nucleus of the hypothalamus (VMH) was the structure having the most dramatic differences in Fos expression (Fig. 9, Table 2). B6 and D2 mice that received no flurothyl-induced seizures had little detectable Fos expression in the VMH. Following 1, 5, or 8 flurothyl-induced seizures, Fos expression in the VMH was scattered in both B6 and D2 mice (Fig. 9, Table 2). When retested following the 28-day incubation period, B6 mice having a forebrain→brainstem seizure had high bilateral expression of Fos in the VMH (Fig. 9). Conversely, B6 mice not having a forebrain→brainstem seizure, but solely expressing a clonic-forebrain seizure had scattered Fos expression bilaterally in the VMH. Upon flurothyl retest for D2 mice, all mice had clonic forebrain seizures. Of the D2 mice that were retested with flurothyl, the majority had scattered Fos expression bilaterally in the VMH (Fig. 9) with 2 animals having low unilateral Fos expression in the VMH.

Discussion

In the current study, we utilized Fos to assess neural activity following repeated seizures in divergent inbred mouse strains. Our results indicate that changes in neural activity depend in part on the number of seizures experienced during the repeated-flurothyl model as well as the genetic background of the mice. In the hippocampus, stark differences were observed between B6 and D2 mice with dramatic differences in the dentate gyrus following 1 seizure. Also, the change in seizure phenotype observed in B6 mice to forebrain→brainstem seizures upon flurothyl retest was associated with high bilateral Fos expression in the VMH. This bilateral activation of the VMH was not seen in D2 mice, which do not have a change in seizure phenotype upon flurothyl retest. These data highlight the importance of genetic background modifiers in susceptibility to seizures as well as their progression. Further consideration of mixed genetic backgrounds in studies exploring seizure susceptibility in transgenic and knockout animals is warranted.

Our data show a difference in Fos expression in the hippocampus between B6 and D2 mice after 1 flurothyl-induced seizure. That is, after 1 flurothyl-induced seizure, B6 mice have

significant increases in hippocampal Fos expression that is not observed in D2 hippocampus. A similar result was observed for hippocampal pErk1/2 levels between B6 and D2 mice after 1 seizure. Phosphorylation of Erk1/2 can be induced by a variety of stimuli, including growth factors and extracellular glutamate, through increases in intracellular calcium levels (Dolmetsch, et al., 2001, Ha and Redmond, 2008, Xia, et al., 1996). Increases in intracellular calcium mediated through Erk1/2 phosphorylation activates the cAMP response element binding protein (CREB) leading to transcription of *c-fos* (Monje, et al., 2003). Reduced Erk activation in the hippocampus of D2 mice indicates that B6 hippocampus is potentially more hyperexcitable than D2 hippocampus. In fact, RNA analysis of other Fos-related proteins, including FosB and its transcriptional partner JunD, revealed a global dysregulation of these transcription factor pathways in D2 hippocampus following 1 seizure. Interestingly, a number of other non-Fos related transcripts followed this trend in the hippocampus of D2 mice after 1 seizure with transcript levels comparable to control D2 and B6 hippocampus. Notably, this phenomena was not limited to transcriptional suppression as there were a comparable number of genes with elevated hippocampus transcription levels in D2 mice after 1 seizure as well as in controls, which were not increased in B6 hippocampus following 1 seizure. Together, these differences indicate a global generalized deregulation of mRNA transcription in D2 hippocampus following a seizure, which may contribute to D2's seizure susceptibility.

One interesting conclusion from these studies was that while there were differences in Fos expression between B6 and D2 mice in the dentate gyrus and CA1 fields, there were no measurable differences in seizure-induced Fos expression in the CA3 fields. The dentate gyrus is the first synapse of the classic tri-synaptic hippocampal circuit and is considered a gatekeeper for determining whether discharge propagates into the CA fields (Heinemann, et al., 1992, Lothman, et al., 1992). However, entorhinal cortex can also access the hippocampal network via propagation of discharge directly to the CA3 and CA1 fields without dentate gyrus involvement (Empson and Heinemann, 1995). Thus, given our Fos results in D2 mice, this strain may have both abnormalities in gating discharge through the dentate gyrus as well as through the secondary pathway via the entorhinal cortex to the CA fields. Future electrophysiological studies specifically focused on examining these differences are needed to understand these phenomena.

Although D2 mice do not have increased Fos expression in the hippocampus following 1 flurothyl seizure, our hippocampal EEG recordings revealed that D2 hippocampal neurons depolarize during the initial flurothyl-induced seizure. We observed robust ictal discharge in the D2 hippocampus during this initial seizure. Thus, depolarization in D2 hippocampus during a seizure may not effectively activate downstream signal transduction pathways, such as the Fos pathway. This inability to convert seizure discharge into a transcriptional response indicated that there may be a threshold of stimulus, below which the depolarization event cannot be further propagated to the signal transduction machinery in D2 mice. This model is supported by the lack of pErk1/2 increased phosphorylation in D2 hippocampus following an initial flurothyl-induced seizure. This stimulus threshold is overcome by additional seizures resulting in subsequent activation of cellular signaling cascades. We suggest that the inability of the D2 hippocampus to induce Fos expression following 1

seizure results in the loss of subsequent activation of Fos-mediated downstream signaling, which may have an important role in regulating seizure susceptibility.

The suppression of Fos in D2 hippocampus, following a single generalized seizure in which ictal discharge was found to propagate, demonstrates a potential caveat in using Fos as a neuronal activity marker that is dependent on genetic background. Previously, it had been shown that electrically kindled seizures could induce robust expression of the *c-fos* gene in brain regions only in rats that showed bursts of population spikes in those regions having increased Fos expression (Labiner, et al., 1993). However, similar synchronous burst firing during kindled seizures was unable to induce *c-fos* mRNA expression in the substantia nigra (Labiner, et al., 1993). This finding clearly highlights the ramifications of using Fos as marker of neuronal activation. Since Fos expression does not mark all cells firing in bursts uniformly, its reliability and dependency as a universal marker across the central nervous system for neuronal activation should be carefully considered when used. In our study, we added to this caveat by showing that although the D2 hippocampus has burst firing during a seizure, it cannot activate Fos expression following one seizure. This is compared to the same structure in a different inbred strain, B6, which can express Fos following a single seizure. Interestingly, this delay in D2 Fos expression is overcome with repeated seizures. Lastly, this study further demonstrates that different inbred strains of mice can have differences in their Fos neural activity maps following a similar stimulus, even within the same neuroanatomical structure.

Following flurothyl induction, a 28-day rest period and flurothyl retest, B6 mice have a change in behavioral seizure phenotype from a forebrain seizure to a forebrain→brainstem seizure. The change in seizure phenotype at flurothyl retest in B6 mice correlates with significant bilateral expression of Fos in the VMH. This alteration in seizure phenotype is likely the result of an enhanced propagation of ictal-seizure discharge from the forebrain seizure circuitry to the brainstem seizure system through the VMH (Ferland and Applegate, 1998, Samoriski, et al., 1998). Interestingly, no D2 mice had a change in seizure phenotype upon flurothyl retest or had evidence for a reorganizational change in the VMH as indicated by Fos immunohistochemistry. A potential explanation for these phenomena is that the reorganization that occurs in the brain in the VMH of B6 mice, required for the change in seizure phenotype, does not occur in D2 mice. This indicates that genetically-mediated neuroanatomical reorganization is possibly contributing to these differences. Current work in our laboratory is aimed at analyzing these changes in BXD recombinant inbred strains of mice (Johnson, et al., 1992, Peirce, et al., 2004, Plomin, et al., 1991) to map quantitative trait loci responsible for setting seizure threshold and modulating epileptogenesis.

In conclusion, our results illustrate changes in the neural activity of specific brain regions during flurothyl-induced epileptogenesis, and we suggest that a change or reorganization in the brain is occurring during the repeated-flurothyl model. The fundamental difference in regional neuronal activity following seizures in two distinct inbred strains of mice may help to explain why certain patient populations are at increased risk for the development of epilepsy and for the progression to more complex seizures over time. The deregulation of subsets of gene transcripts between two inbred mouse strains following exposure to the same seizure model also provides potential insight into molecular changes that could account for

strain differences in seizure susceptibility that may possibly translate to a better understanding of human epilepsy. Elucidating the pathophysiological processes underlying these plasticity changes during epileptogenesis could potentially lead to the development of new therapeutics directed against novel targets.

Acknowledgments

This work was supported by an NIH/NINDS R01NS064283 grant to RJF and an NIH/NAIMS R01AR054828 to BJH. We would also like to acknowledge the Advanced Genomics Technology core at the Wadsworth Center for assistance with the RNA array studies.

References

1. Andre V, Pineau N, Motte JE, Marescaux C, Nehlig A. Mapping of neuronal networks underlying generalized seizures induced by increasing doses of pentylenetetrazol in the immature and adult rat: a c-Fos immunohistochemical study. *Eur J Neurosci*. 1998; 10:2094–2106. [PubMed: 9753096]
2. Applegate CD, Samoriski GM, Ozduman K. Effects of valproate, phenytoin, and MK-801 in a novel model of epileptogenesis. *Epilepsia*. 1997; 38:631–636. [PubMed: 9186244]
3. Baraban JM, Fiore RS, Sanghera JS, Paddon HB, Pelech SL. Identification of p42 mitogen-activated protein kinase as a tyrosine kinase substrate activated by maximal electroconvulsive shock in hippocampus. *J Neurochem*. 1993; 60:330–336. [PubMed: 8417154]
4. Browning RA, Nelson DK, Mogharreban N, Jobe PC, Laird HE 2nd. Effect of midbrain and pontine tegmental lesions on audiogenic seizures in genetically epilepsy-prone rats. *Epilepsia*. 1985; 26:175–183. [PubMed: 3987648]
5. Browning RA, Simonton RL, Turner FJ. Antagonism of experimentally induced tonic seizures following a lesion in the midbrain tegmentum. *Epilepsia*. 1981; 22:595–601. [PubMed: 7285885]
6. Browning RA, Turner FJ, Simonton RL, Bundman MC. Effect of midbrain and pontine tegmental lesions on the maximal electroshock seizure pattern in rats. *Epilepsia*. 1981; 22:583–594. [PubMed: 7285884]
7. Dichter MA. Models of epileptogenesis in adult animals available for antiepileptogenesis drug screening. *Epilepsy Res*. 2006; 68:31–35. [PubMed: 16377136]
8. Dolmetsch RE, Pajvani U, Fife K, Spotts JM, Greenberg ME. Signaling to the nucleus by an L-type calcium channel-calmodulin complex through the MAP kinase pathway. *Science*. 2001; 294:333–339. [PubMed: 11598293]
9. Empson RM, Heinemann U. The perforant path projection to hippocampal area CA1 in the rat hippocampal-entorhinal cortex combined slice. *J Physiol*. 1995; 484 (Pt 3):707–720. [PubMed: 7623286]
10. Engel J Jr, Wolfson L, Brown L. Anatomical correlates of electrical and behavioral events related to amygdaloid kindling. *Ann Neurol*. 1978; 3:538–544. [PubMed: 677819]
11. Ferland RJ, Applegate CD. Decreased brainstem seizure thresholds and facilitated seizure propagation in mice exposed to repeated flurothyl-induced generalized forebrain seizures. *Epilepsy Res*. 1998; 30:49–62. [PubMed: 9551844]
12. Ferland RJ, Applegate CD. The role of the ventromedial nucleus of the hypothalamus in epileptogenesis. *Neuroreport*. 1998; 9:3623–3629. [PubMed: 9858370]
13. Ferland RJ, Applegate CD. Bidirectional transfer between electrical and flurothyl kindling in mice: evidence for common processes in epileptogenesis. *Epilepsia*. 1999; 40:144–152. [PubMed: 9952259]
14. Ferland RJ, Nierenberg J, Applegate CD. A role for the bilateral involvement of perirhinal cortex in generalized kindled seizure expression. *Exp Neurol*. 1998; 151:124–137. [PubMed: 9582260]
15. Ferraro TN, Golden GT, Smith GG, Berrettini WH. Differential susceptibility to seizures induced by systemic kainic acid treatment in mature DBA/2J and C57BL/6J mice. *Epilepsia*. 1995; 36:301–307. [PubMed: 7614915]

16. Ferraro TN, Golden GT, Smith GG, DeMuth D, Buono RJ, Berrettini WH. Mouse strain variation in maximal electroshock seizure threshold. *Brain Res.* 2002; 936:82–86. [PubMed: 11988233]
17. Franck JE, Ginter KL, Schwartzkroin PA. Developing genetically epilepsy-prone rats have an abnormal seizure response to flurothyl. *Epilepsia.* 1989; 30:1–6. [PubMed: 2912711]
18. Garrido YC, Sanabria ER, Funke MG, Cavalheiro EA, Naffah-Mazzacoratti MG. Mitogen-activated protein kinase is increased in the limbic structures of the rat brain during the early stages of status epilepticus. *Brain Res Bull.* 1998; 47:223–229. [PubMed: 9865854]
19. Ha S, Redmond L. ERK mediates activity dependent neuronal complexity via sustained activity and CREB-mediated signaling. *Dev Neurobiol.* 2008; 68:1565–1579. [PubMed: 18837011]
20. Hashimoto Y, Araki H, Suemaru K, Gomita Y. Effects of drugs acting on the GABA-benzodiazepine receptor complex on flurothyl-induced seizures in Mongolian gerbils. *Eur J Pharmacol.* 2006; 536:241–247. [PubMed: 16581068]
21. Heinemann U, Beck H, Dreier JP, Ficker E, Stabel J, Zhang CL. The dentate gyrus as a regulated gate for the propagation of epileptiform activity. *Epilepsy Res Suppl.* 1992; 7:273–280. [PubMed: 1334666]
22. Hirsch E, Danover L, Simler S, Pereira de Vasconcelos A, Maton B, Nehlig A, Marescaux C, Vergnes M. The amygdala is critical for seizure propagation from brainstem to forebrain. *Neuroscience.* 1997; 77:975–984. [PubMed: 9130779]
23. Houser CR, Huang CS, Peng Z. Dynamic seizure-related changes in extracellular signal-regulated kinase activation in a mouse model of temporal lobe epilepsy. *Neuroscience.* 2008; 156:222–237. [PubMed: 18675888]
24. Hsiao YC, Tong ZJ, Westfall JE, Ault JG, Page-McCaw PS, Ferland RJ. Ahi1, whose human ortholog is mutated in Joubert syndrome, is required for Rab8a localization, ciliogenesis and vesicle trafficking. *Hum Mol Genet.* 2009; 18:3926–3941. [PubMed: 19625297]
25. Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc.* 2009; 4:44–57. [PubMed: 19131956]
26. Johnson TE, DeFries JC, Markel PD. Mapping quantitative trait loci for behavioral traits in the mouse. *Behav Genet.* 1992; 22:635–653. [PubMed: 1363267]
27. Kim YS, Hong KS, Seong YS, Park JB, Kuroda S, Kishi K, Kaibuchi K, Takai Y. Phosphorylation and activation of mitogen-activated protein kinase by kainic acid-induced seizure in rat hippocampus. *Biochem Biophys Res Commun.* 1994; 202:1163–1168. [PubMed: 7519421]
28. Klein BD, Fu YH, Ptacek LJ, White HS. c-Fos immunohistochemical mapping of the audiogenic seizure network and tonotopic neuronal hyperexcitability in the inferior colliculus of the Frings mouse. *Epilepsy Res.* 2004; 62:13–25. [PubMed: 15519128]
29. Labiner DM, Butler LS, Cao Z, Hosford DA, Shin C, McNamara JO. Induction of c-fos mRNA by kindled seizures: complex relationship with neuronal burst firing. *J Neurosci.* 1993; 13:744–751. [PubMed: 8381172]
30. Liang, D.; Seyfried, TN. Gene expression changes in kindling. In: Corcoran, ME.; Moshe, SL., editors. *Kindling 6*. Springer; New York: 2005. p. 181–188.
31. Loscher W, Ebert U. The role of the piriform cortex in kindling. *Prog Neurobiol.* 1996; 50:427–481. [PubMed: 9015822]
32. Lothman EW, Hatlelid JM, Zorumski CF. Functional mapping of limbic seizures originating in the hippocampus: a combined 2-deoxyglucose and electrophysiologic study. *Brain Res.* 1985; 360:92–100. [PubMed: 4075185]
33. Lothman EW, Stringer JL, Bertram EH. The dentate gyrus as a control point for seizures in the hippocampus and beyond. *Epilepsy Res Suppl.* 1992; 7:301–313. [PubMed: 1334669]
34. McCown TJ, Duncan GE, Johnson KB, Breese GR. Metabolic and functional mapping of the neural network subserving inferior collicular seizure generalization. *Brain Res.* 1995; 701:117–128. [PubMed: 8925273]
35. Merrill MA, Clough RW, Jobe PC, Browning RA. Role of the superior colliculus and the intercollicular nucleus in the brainstem seizure circuitry of the genetically epilepsy-prone rat. *Epilepsia.* 2003; 44:305–314. [PubMed: 12614385]
36. Monje P, Marinissen MJ, Gutkind JS. Phosphorylation of the carboxyl-terminal transactivation domain of c-Fos by extracellular signal-regulated kinase mediates the transcriptional activation of

- AP-1 and cellular transformation induced by platelet-derived growth factor. *Mol Cell Biol.* 2003; 23:7030–7043. [PubMed: 12972619]
37. Namba H, Iwasa H, Kubota M, Hagihara Y, Yamaura A. Changes of hippocampal glucose utilization subsequent to amygdaloid-kindled generalized seizures. *Epilepsia.* 1991; 32:27–32. [PubMed: 1985827]
 38. Oommen J, Kraus AC, Fisher RS. Intraventricular administration of gabapentin in the rat increases flurothyl seizure threshold. *Neurosci Lett.* 2007; 417:308–311. [PubMed: 17363166]
 39. Papandrea D, Anderson TM, Herron BJ, Ferland RJ. Dissociation of seizure traits in inbred strains of mice using the flurothyl kindling model of epileptogenesis. *Exp Neurol.* 2009; 215:60–68. [PubMed: 18950623]
 40. Papandrea D, Kukol WS, Anderson TM, Herron BJ, Ferland RJ. Analysis of flurothyl-induced myoclonus in inbred strains of mice. *Epilepsy Res.* 2009; 87:130–136. [PubMed: 19744831]
 41. Paxinos, G.; Franklin, KBJ. The mouse brain in stereotaxic coordinates. Academic Press; San Diego, CA: 2001.
 42. Peirce JL, Lu L, Gu J, Silver LM, Williams RW. A new set of BXD recombinant inbred lines from advanced intercross populations in mice. *BMC Genet.* 2004; 5:7. [PubMed: 15117419]
 43. Plomin R, McClearn GE, Gora-Maslak G, Neiderhiser JM. Use of recombinant inbred strains to detect quantitative trait loci associated with behavior. *Behav Genet.* 1991; 21:99–116. [PubMed: 2049054]
 44. Ribak CE, Manio AL, Navetta MS, Gall CM. In situ hybridization for c-fos mRNA reveals the involvement of the superior colliculus in the propagation of seizure activity in genetically epilepsy-prone rats. *Epilepsy Res.* 1997; 26:397–406. [PubMed: 9127720]
 45. Sagar SM, Sharp FR, Curran T. Expression of c-fos protein in brain: metabolic mapping at the cellular level. *Science.* 1988; 240:1328–1331. [PubMed: 3131879]
 46. Samoriski GM, Applegate CD. Repeated generalized seizures induce time-dependent changes in the behavioral seizure response independent of continued seizure induction. *J Neurosci.* 1997; 17:5581–5590. [PubMed: 9204939]
 47. Samoriski GM, Piekut DT, Applegate CD. Differential spatial patterns of Fos induction following generalized clonic and generalized tonic seizures. *Exp Neurol.* 1997; 143:255–268. [PubMed: 9056388]
 48. Samoriski GM, Piekut DT, Applegate CD. Regional analysis of the spatial patterns of Fos induction in brain following flurothyl kindling. *Neuroscience.* 1998; 84:1209–1222. [PubMed: 9578407]
 49. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods.* 2012; 9:671–675. [PubMed: 22930834]
 50. Shehab S, Coffey P, Dean P, Redgrave P. Regional expression of fos-like immunoreactivity following seizures induced by pentylenetetrazole and maximal electroshock. *Exp Neurol.* 1992; 118:261–274. [PubMed: 1306485]
 51. Stafstrom CE, Thompson JL, Holmes GL. Kainic acid seizures in the developing brain: status epilepticus and spontaneous recurrent seizures. *Brain Res Dev Brain Res.* 1992; 65:227–236.
 52. Tuz K, Hsiao YC, Juarez O, Shi B, Harmon EY, Phelps IG, Lennartz MR, Glass IA, Doherty D, Ferland RJ. The Joubert syndrome-associated missense mutation (V443D) in the Abelson-helper integration site 1 (AHI1) protein alters its localization and protein-protein interactions. *J Biol Chem.* 2013; 288:13676–13694. [PubMed: 23532844]
 53. Xia Z, Dudek H, Miranti CK, Greenberg ME. Calcium influx via the NMDA receptor induces immediate early gene transcription by a MAP kinase/ERK-dependent mechanism. *J Neurosci.* 1996; 16:5425–5436. [PubMed: 8757255]

Highlights

- Hippocampal Fos is different in C57BL/6J and DBA/2J mice following seizures
- Fos expression dissociates from ictal discharge in the hippocampus of DBA/2J mice
- Subsets of genes are deregulated in hippocampus between C57BL/6J and DBA/2J mice
- Ventromedial n. of the hypothalamus is involved in expression of complex seizures

Figure 1. Comparisons of repeated flurothyl-induced latencies to generalized seizures between C57BL/6J and DBA/2J inbred strains of mice

The latency to loss of postural control (generalized seizure onset) was determined in C57BL/6J (B6) and DBA/2J (D2) mice using a 10% flurothyl solution. B6 mice have high latencies to generalized seizure expression (compared to D2 mice) that decreased significantly with repeated seizures and remained low on retest. In contrast, D2 mice have low initial latencies to generalized seizure expression that do not decrease with repeated seizures and essentially plateaus across the induction period. Upon retest, approximately 85% of B6 mice expressed a more complex seizure comprised of a clonic seizure that rapidly and uninterruptedly transitioned into a brainstem-tonic seizure (forebrain→brainstem seizure), whereas none of the D2 mice expressed this more complex seizure.

Figure 2. Changes in seizure-induced Fos expression in the hippocampus of two diverse inbred strains of mice having repeated flurothyl-induced generalized clonic seizures

C57BL/6J (B6) and DBA/2J (D2) mice were given flurothyl-induced seizures and were sacrificed following 1) one seizure, 2) five seizures, 3) eight seizures, or 4) following 8 seizures, a 28-d incubation phase, and a final flurothyl challenge (retest). Fos immunohistochemistry was examined to determine neural activity map differences between these two strains of mice. Whereas B6 mice show significant increases in the hippocampus following a single seizure that remains elevated with subsequent seizures, D2 mice show minimal changes in Fos expression following a single seizure, and only with repeated seizures, does Fos expression increase and remain elevated. $N = 4/\text{strain/seizure number}$. CA1, cornu ammonis 1; CA3, cornu ammonis 3; DG, dentate gyrus. Scale bar equals 500 μm .

Figure 3. Quantitative differences in hippocampal Fos expression in two diverse inbred strains of mice following repeated flurothyl-induced generalized clonic seizures

C57BL/6J (B6) and DBA/2J (D2) mice were sacrificed following 1) one seizure, 2) five seizures, 3) eight seizures, or 4) following 8 seizures, a 28-d incubation phase, and a final flurothyl challenge (retest). Fos immunohistochemistry was performed and quantified by measuring the integrated density of Fos immunolabeling in the dentate gyrus, CA3 and CA1 regions of the hippocampus. N = 4/strain/seizure number. * denotes a significance in comparing B6 and D2 mice following equivalent seizure numbers ([Dentate gyrus: 1 seizure: $t_{24}=8.5$, $P<0.0001$; 5 seizures: $t_{22}=6.1$, $P<0.0001$; 8 seizures: $t_{25}=4.0$, $P<0.0005$; retest: $t_{18}=4.7$, $P<0.0002$]; [CA1: 1 seizure: $t_{24}=2.8$, $P<0.01$; 5 seizures: $t_{22}=2.9$, $P<0.008$]). CA1, cornu ammonis 1; CA3, cornu ammonis 3.

Figure 4. Western blot analysis of Fos expression in the hippocampus of C57BL/6J and DBA/2J mice following 1, 5, or 8 flurothyl-induced seizures

Top, Bar graphs representing the averaged expression levels of hippocampal Fos normalized to p84 (nuclear loading control) protein expression following 1, 5, or 8 flurothyl-induced seizures in C57BL/6J (B6) and DBA/2J (D2) mice. In B6 mice, there is a significant increase in Fos protein following a single seizure that remains significantly elevated from control levels following repeated seizures. For D2 mice, there is only a significant increase in Fos levels following 5 and 8 flurothyl-induced seizures, but not following a single seizure. N = 4/strain/seizure number. * represents a significance of $P < 0.05$ as compared to the 0 seizures group; # represents a significance of $P < 0.04$ as compared to the 0 seizures group. Bottom, Representative western blot demonstrating the changes in hippocampal Fos protein levels following a single or repeated seizures in B6 and D2 mice.

Figure 5. RNA analysis of *Fos* expression, and *Fos*-related genes, in the hippocampus of C57BL/6J and DBA/2J mice following 1, 3, and 8 fluoroethyl-induced seizures
 Graph representing the averaged normalized intensity values of *Fos*, *Fosl1*, *Fosl2*, and *FosB* mRNA expression. The relative abundance of mRNAs for *Fos*, *Fosl1*, *Fosl2*, and *FosB* all increased significantly in C57BL/6J hippocampus following 1 seizure ($P < 0.02$). However, these mRNA levels did not significantly increase in DBA/2J hippocampus following 1 seizure. * represents a significance of $P < 0.05$.

Figure 6. pERK labeling in the hippocampus of C57BL/6J and DBA/2J mice following 1 flurothyl-induced seizure

C57BL/6J (B6) and DBA/2J (D2) mice were given a single flurothyl-induced seizure and were sacrificed 15 minutes following the end of the seizure. B6 and D2 coronal brain sections were immunostained with pErk antibodies. pErk labeling (red) increased after one seizure in B6 mice, particularly in the hippocampal molecular layer, while no differences in pErk were found in D2 hippocampus. N = 3/strain/seizure number. CA1, cornu ammonis 1; CA3, cornu ammonis 3; DG, dentate gyrus, ml, molecular layer. Scale bar equals 200 μ m.

Figure 7. While Fos expression is not observed in the hippocampus of DBA/2J mice following one flurothyl-induced seizure, there is ictal discharge in the hippocampus

While the hippocampi of DBA/2J mice do not have evidence of seizure-induced increases in Fos expression following an initial flurothyl-induced seizure, there is propagation of ictal discharge through the hippocampus. Representative electroencephalograms (EEG) in the hippocampus of a C57BL/6J mouse (top) and a DBA/2J mouse (bottom) during an initial flurothyl-induced seizure (N=5/strain). Lines with numbers 1, 2 and 3 indicate the region of the EEG time scale, magnified four times to demonstrate the field depolarization shifts.

Figure 8. Global gene expression in C57BL/6J and DBA/2J hippocampus following repeated flurothyl seizures

Hierarchical clustering analysis of 12,034 significantly deregulated mRNAs between C57BL/6J (B6) and DBA/2J (D2) hippocampus, 2 hours after a single (1), three (3) or eight (8) seizures compared to baseline expression (0). (A) Gene clusters were sorted by conditions (strain and seizure number) as well as gene entities to reveal a temporal pattern of expression that clusters in 5 groups of transcripts in the second tier of clustering analysis. (B) Sorting by conditions led to pair wise (i.e., B6(8) and D2(8) co-cluster) co-clusters for 3 and 8 seizure treatment groups, but D2 hippocampus exposed to a single seizure was more similar to B6 and D2 baseline expression patterns than B6 hippocampus stimulated with a single seizure (B, top: shared decreases in gene expression between B6 and D2 controls and D2 having 1 seizure; B, bottom: shared increases in gene expression between B6 and D2 controls and D2 having 1 seizure). These two clusters had several notable genes that have been implicated in seizure progression or epileptogenesis.

Figure 9. The ventromedial nucleus of the hypothalamus shows bilateral Fos expression in mice having an evolution of more complex seizures (forebrain→brainstem seizures)

Immunostaining for Fos in the ventromedial nucleus of the hypothalamus (VMH) of C57BL/6J (B6) and DBA/2J (D2) mice following 1) eight seizures or 2) following 8 seizures, a 28-d incubation phase, and a final flurothyl retest. Following 8 seizures, B6 and D2 mice displayed clonic-forebrain seizures and scattered Fos expression in the VMH. Following a final flurothyl rechallenge, B6 mice have clonic-forebrain seizures that rapidly progress into brainstem-tonic seizures, with such mice displaying bilateral and high levels of Fos expression in the VMH. However, D2 mice, which only have clonic-forebrain seizures upon flurothyl rechallenge, display scattered Fos expression in the VMH. N = 4/strain/seizure number.

Table 1

Flurothyl concentrations resulting in the expression of generalized seizures (generalized seizure thresholds)

Seizure Trial	C57BL/6J [flurothyl] mg $\pm$ SEM	DBA/2J [flurothyl] mg $\pm$ SEM
1	69.5 $\pm$ 1.5	43.5 $\pm$ 2.5
2	55.5 $\pm$ 1.5	42.3 $\pm$ 1.8
3	46.0 $\pm$ 1.3	37.5 $\pm$ 2.0
4	40.8 $\pm$ 1.0	43.2 $\pm$ 3.2
5	36.5 $\pm$ 0.7	44.8 $\pm$ 1.3
6	33.3 $\pm$ 0.7	42.8 $\pm$ 1.3
7	34.0 $\pm$ 0.8	41.7 $\pm$ 1.7
8	36.2 $\pm$ 1.0	35.8 $\pm$ 1.2
retest	40.3 $\pm$ 1.7	35.7 $\pm$ 3.8

Table 2

Median scores per group for Fos expression in various brain areas in B6 and D2 mice following 1 seizure, 5 seizures, 8 seizures, and flurothyl retest following 8 seizures and a 28-day incubation period. Qualitative measurements were made blind to the observer using the following scale: 0 = no observed Fos, 1 = scattered Fos expression, 2 = low Fos expression, 3 = moderate Fos expression, 4 = high Fos expression, 5 = very high Fos expression.

Brain Region	1 seizure		5 seizures		8 seizures		Retest	
	B6	D2	B6	D2	B6	D2	B6	D2
<i>Neocortex</i>	3	3	3.5	3.5	3.5	3	4	4
<i>Cingulate cortex</i>	3	3	4	3.5	3.5	3	4	4
<i>Piriform cortex</i>	4	4	4	4	4	4	4	4
<i>Perirhinal cortex</i>	2	3	3	3	3	3	3	3
<i>Bed nucleus stria terminalis</i>	3	3	3	3	3	3	3.5	3.5
<i>Ventral pallidum</i>	1.5	1	1	1	1	1	1	0.5
<i>Clastrum</i>	3	3	3	3	3.5	3	3	3.5
<i>Hypothalamus</i>								
<i>Paraventricular</i>	3.5	2	3	4	3.5	3.5	3.5	3
<i>Zona incerta</i>	2	2	2	3	3	3	3	3
<i>Ventromedial nucleus (VMH)</i>	0.5	1	1	0	1.5	1	4	0.5
<i>Dorsomedial nucleus (DMH)</i>	2.5	1	2	3	2	3	4	2.5
<i>Hippocampal formation</i>								
<i>Hilus</i>	3	1	3	2.5	3.5	3	3	3
<i>Dentate gyrus</i>	5	1	4	3.5	3.5	4	4	4
<i>CA 1</i>	3	2	4	2.5	4	4	3	3
<i>CA 3</i>	3.5	2	4	3	4	4	3	3
<i>Subiculum</i>	3	1	2	1.5	2	1	1	2
<i>Amygdala nuclei</i>								
<i>Anterior amygdaloid area</i>	1	1	1.5	1	1	1	1.5	2
<i>Medial</i>	2.5	2	4	2.5	3	3	4	4
<i>Basomedial</i>	3	4	3.5	3	4	3.5	4	4
<i>Basolateral</i>	2	3	3	3	3	3	3	3.5
<i>Cortical</i>	3	4	4	3.5	3.5	4	4	4

Brain Region	1 seizure		5 seizures		8 seizures		Retest	
	B6	D2	B6	D2	B6	D2	B6	D2
Lateral	2.5	3	3	2.5	3	2.5	3	3
Intercalated	3	4	4	4	4	4	4	4
Amygdalostriatal transition	3	4	3	1.5	2	1	2.5	2
Amygdalopiriform	1	3	3	3.5	4	4	4	4
Amygdalohippocampal	1	3	3	3	4	3.5	4	4