Published in final edited form as:

Acta Parasitol. 2014 October ; 59(4): 686–689. doi:10.2478/s11686-014-0304-5.

A new species of coccidia (Apicomplexa: Eimeriidae) from Cordillera striped shrew-rat, *Chrotomys whiteheadi* (Rodentia: Muridae), from the Philippines

Chris T. McAllister^{1,*}, R. Scott Seville², Donald W. Duszynski³, and Sarah E. Bush⁴

¹Science and Mathematics Division, Eastern Oklahoma State College, Idabel, Oklahoma, USA 74745

²Department of Zoology and Physiology, University of Wyoming, Casper, Wyoming, USA 82601

³Department of Biology, University of New Mexico, Albuquerque, New Mexico, USA 87131

⁴Department of Biology, University of Utah, Salt Lake City, Utah, USA 84112

Abstract

During July 2011, a single Cordillera striped shrew-rat (*Chrotomys whiteheadi*) was collected from the Philippines and its faeces examined for coccidian parasites. It harboured an eimerian that we describe here as new. Oocysts of *Eimeria macarthuri* sp. n. were spheroidal to subspheroidal with a bi-layered wall and measured (length $\times$ width, L $\times$ W) 18.2 $\times$ 17.0 μ m, with an L/W ratio of 1.1. A micropyle, oocyst residuum and polar granule were absent. Sporocysts were ovoidal, 9.0 $\times$ 6.4 μ m, with an L/W ratio of 1.3. A nipple-like Stieda body was present as well as a substieda body. A granular sporocyst residuum was present. To our knowledge, *E. macarthuri* represents the only coccidian ever described from a rodent of the Philippines.

Keywords

Apicomplexa; Coccidia; taxonomy; oocysts; Rodentia; Muridae; *Eimeria macarthuri*; Philippines

Introduction

The Cordillera striped shrew-rat, *Chrotomys whiteheadi* Thomas is one of five members of the genus (Carlton and Musser 2005) that is endemic to the Philippines. This species is restricted to northern Luzon (Benguet, Kalinga and Mountain provinces), where it is common in the northern and central Cordillera occurring from 925–2,700 m (Heaney et al. 2008). The species occurs in primary, secondary, and disturbed lower montane forest to mossy forest in the central Cordillera (Sanborn, 1952; Rabor, 1955; Largen, 1985) and agricultural areas adjacent to forest. Although this rat can exist in pristine habitats, it is the most habitat-tolerant native species and can also be found in highly disturbed croplands, vegetable gardens, farms, and rice fields or essentially anywhere where there are

earthworms (Ricart et al. 2007). However, the species is not found in grasslands or in open agricultural areas.

Nothing is known about the coccidia of *C. whiteheadi* nor, for that matter, any other rodent in the Philippines. Herein, we provide a description of a new species of *Eimeria* from *C. whiteheadi*.

Materials and Methods

During July 2011, a single adult (242 g) male *C. whiteheadi* was collected at 680 m elevation at Mt. Cagua, (Cagayan Province, Luzon Island), Philippines, and examined for coccidia. A fresh faecal sample was placed in an individual vial containing 2.5% (w/v) aqueous potassium dichromate ($K_2Cr_2O_7$). The sample was shipped to the USA and examined for coccidia by light microscopy after flotation in Sheather's sugar solution (specific gravity = 1.30). Measurements were taken on 17 sporulated oocysts using a calibrated ocular micrometer and reported in micrometres (μm) with means followed by the ranges in parentheses; photographs were taken using Nomarski interference-contrast optics. Oocysts were ~515 days old when measured and photographed. Descriptions of oocysts and sporocysts follow guidelines of Wilber et al. (1998) as follows: oocyst length (L) and width (W), their ranges and ratios (L/W), micropyle (M), oocyst residuum (OR), polar granule(s) (PG), sporocyst length (L) and width (W), their ranges and ratio (L/W), sporocyst (SP), Stieda body (SB), substieda body (SSB), parastieda body (PSB), sporocyst residuum (SR), sporozoites (SZ) anterior (ARB) and posterior (PRB) refractile bodies, and nucleus (N). A host voucher was accessioned into Kansas Museum of Natural History (KUMNH). Photosyntypes of sporulated oocysts were accessioned into the United States National Parasite Collection (USNPC), Beltsville, Maryland, USA.

Results

The single *R. whiteheadi* was found to be passing oocysts of an eimerian which we describe here as new.

Eimeria macarthuri sp. nov. (Figs. 1–4)

Description of sporulated oocyst: Oocyst with 4 sporocysts; shape spheroidal—subspheroidal; bi-layered wall, colourless, ~1.2 thick, textured outer layer ~0.8 thick, smooth inner layer ~0.4 thick; L $\times$ W: 18.2 $\times$ 17.0 (16–20 $\times$ 15–19); L/W: 1.1 (1.0–1.1); M, OR, PG: all absent.

Description of sporocyst and sporozoites: SP ovoidal, with a smooth uni-layered wall, ~0.4 thick; L $\times$ W: 9.0 $\times$ 6.4 (8–10 $\times$ 6–7); L/W: 1.3 (1.2–1.6); SB nipple-like, SSB body present, PSB body absent; SR: granular, composed of moderately-sized granules in a compact mass or dispersed between SZ; SZ: (not measured) sausage-shaped with spheroidal ARB and PRB; single N slightly posterior to midpoint.

Taxonomic summary

Type host: Cordillera striped shrew-rat, *Chrotomys whiteheadi* Thomas, 1895 (Rodentia: Muridae). Collected 22 July 2011.

Type specimens: Symbiotype host in the Kansas Museum of Natural History (KUMNH # 168307, Parasite # P-4698). Photosyntype of sporulated oocyst deposited in the USNPC as No. 107957.

Type locality: Mt. Cagua, Cagayan Province, Luzon Island, Philippines (18.236°N, 122.104°E, elev. 680 m).

Prevalence: In one of one (100%) of the type host.

Sporulation: Unknown. Specimens were collected in the field, stored in K₂Cr₂O₇ solution, and not examined until 515 days later.

Site of infection: Unknown, oocysts recovered from faeces.

Etymology: The specific epithet is given in honour of General Douglas MacArthur (1880–1964), for his prominent role in the Pacific Theater of WW II, particularly in the Philippines, and the only man ever to become a field marshal in the Philippine Army.

Remarks: As there are no previous coccidians described from any rodent in the Philippines, we restrict our comparisons to eimerians described from rodents found in the general region (i.e., Malaysia) (Colley and Mullin, 1971; Mullin et al. 1972). The only species similar in size to the new species is *Eimeria miyairii* Ohira, 1912, originally described from common brown rat, *Rattus norvegicus* from Japan (Ohira 1912), but also reported from “wild rats” in Japan by Matubayasi (1938), Whitehead’s spiny rat, *Maxomys* (= *Rattus*) *whiteheadi* from Seangor, West Malaysia, and Malaysian field rat, *Rattus tiomanicus* from Pekan, West Malaysia (Ohira 1912; Mullin et al. 1972).

The first reasonable description of any coccidium from *Rattus* was of *E. miyairii* by Ohira (1912), but it was written in Japanese and was not read by Western workers. The result was that for many years the species now known as *Eimeria nieschulzi* Dieben, 1924, was thought to be *E. miyairii* (Pellérdy 1974, p. 584) because their oocysts were somewhat similar. Roudabush (1937) translated Ohira’s 1912 paper and, using isolates of oocysts from different colleagues to infect lab rats (presumably *R. norvegicus*), was able to definitively distinguish, in great detail, between the endogenous developmental stages of the three eimerians known from rats at that time, *E. miyairii*, *E. nieschulzi*, and *Eimeria separata* Becker and Hall, 1931. He did not, however, mention the differences between their sporulated oocysts. Matubayasi (1938), unaware of Roudabush’s (1937) paper, added further confusion when he studied sporogony and endogenous development and provided line drawings and photomicrographs of the sporulated oocysts of three eimerians he found in wild rats in Japan: *E. carinii* (which was actually *E. miyairii*), *E. miyairii* (which was actually *E. nieschulzi*) and *E. separata*. Levine and Husar (1979) found *E. miyairii* oocysts in two wild *R. norvegicus* trapped in Illinois, USA; using a translation of Ohira’s (1912) paper, they cleared up the nomenclatural confusion between *E. carinii*, *E. miyairii*, and *E.*

nieschulzi, and redescribed the sporulated oocyst of *E. miyairii* in detail. Our new species differ from *E. miyairii* by having sporocysts that are smaller (9.0×6.4 [$8-10 \times 6-7$] vs. 12.0×8.0 [$11-14 \times 7-9$]) resulting in a smaller L/W ratio (1.3 vs. 1.6), by lacking a polar granule, and by possessing refractile bodies in the sporozoites. Our sporulated oocysts also differs from those of *E. nieschulzi* in shape (nearly spheroidal vs. ovoidal), size (18.2×17.0 [$16-20 \times 15-19$] vs. 23×17 [$16-26 \times 13-21$]), and L/W ratio (1.1 vs. 1.3), and the latter has a distinct polar granule which our new species lacks. Their sporocysts also are distinctly different in size and our species has a substieda body, which the sporocysts of *E. nieschulzi* lack.

Finally, *Eimeria garumarana* Bandyopadhyay and Dasgupta, 1984, described from the common field rat, *Rattus rattus brunneusculus* (Hodgson) in India, has oocysts that are similar in size and other features to the form we describe here. Its sporulated oocysts are subspheroidal and measure 18×17.9 ($17-20 \times 16.5-19$) and lack an oocyst residuum and polar granule. Its ovoidal sporocysts are 9.8×7.5 ($9-11 \times 6-8$) and have both a nipple-like Stieda body and a granular sporocyst residuum. There are two differences that distinguish these oocysts from ours, both of which could be questioned: 1) Bandyopadhyay and Dasgupta (1984) stated throughout their paper that there is a distinct micropyle in their oocyst, ~ 2.0 wide, but their line drawing does not have a micropyle and they did not provide a photomicrograph; and 2) their sporocysts were not reported to have a substieda body (which the authors could have missed), which our form does have. Given these discrepancies, the distance of the Philippines from India (> 4400 km), being separated by part of the Asian continent, and the difference in host genera we are confident that our form is a new species.

Discussion

The mammalian fauna of the Philippine Islands is species-rich and remarkably diverse, and comprises what could be the largest concentration of endemic mammals of any country. In addition, the following murid rodents have been introduced into the Philippines (widespread unless otherwise noted): house mouse, *Mus musculus*, rice-field rat, *Rattus argentiventer* (Mindanao only), Polynesian rat, *Rattus exulans*, Himalayan field rat, *Rattus nitidus* (northern Luzon only), *R. norvegicus*, Tanezumi rat, *Rattus tanezumi* (formerly included in *R. rattus*), and *R. tiomanicus* (Palawan only) (Heaney et al. 2010). At least two of these rodents in other locales have been reported to harbor coccidia (see Levine and Ivens 1990). We hope that our initial report of a coccidian from an endemic rodent on this Pacific island will stimulate others to survey additional mammals of the Philippines for these apicomplexan parasites in order to increase our knowledge of their biodiversity.

Acknowledgements

We thank P. A. Pilitt for expert curatorial assistance. We also thank N. Antoque, J. A. Cantil, S. Knutie, and S. Villa for their assistance in the field. This study was supported, in part, by the National Science Foundation BS&I grant DEB 0743491 to R. M. Brown, S. E. Bush, D. H. Clayton, and R. G. Moyle and grants from the National Center for Research Resources (5P20RR016474-12) and the National Institute of General Medical Sciences (8P20GM103432-12) from the National Institutes of Health to R. S. Seville.

References

- Bandyopadhyay S, Dasgupta B. *Eimeria garumarana*, nsp. from *Rattus rattus brunneusculus* (Hodgson). Journal of the Bengal Natural History Society. 1984; 3:28–35.
- Carlton, MD.; Musser, GG. Order Rodentia. In: Wilson, DE.; Reeder, DM., editors. Mammal species of the world, a taxonomic and geographic reference. Vol. 2. 2005. p. 745-1599.
- Colley FC, Mullin SW. New species of *Eimeria* (Protozoa: Eimeriidae) from Malaysian rats. Journal of Protozoology. 1971; 18:601–604. [PubMed: 5167309]
- Heaney, L.; Balette, D.; Alviola, P.; Duya, MR. *Chrotomys whiteheadi*. IUCN 2013. IUCN Red List of Threatened Species. Version 2013.2. 2008. www.iucnredlist.org. Downloaded on 30 January 2014
- Heaney, L.; Dolar, ML.; Balette, DS.; Esselstyne, JA.; Rickart, EA.; Sedlock, JL. Synopsis of Philippine mammals. Field Museum of Natural History. 2010. [www.http://archive.fieldmuseum.org/philippine_mammals/](http://archive.fieldmuseum.org/philippine_mammals/). Downloaded on 22 February 2014
- Largen MJ. Taxonomically and historically significant specimens of mammals in the Merseyside County Museums, Liverpool. Journal of Mammalogy. 1985; 66:412–418.
- Levine ND, Husar TJ. Rediscovery and redescription of *Eimeria miyairii* Ohira, 1912 from the Norway rat. Proceedings of the Helminthological Society of Washington. 1979; 46:135–137.
- Levine, ND.; Ivens, V. The coccidian parasites of rodents. Boca Raton, Florida: CRC Press; 1990. p. 228
- Matubayasi H. Studies on the parasitic Protozoa in Japan. IV. Coccidia parasitic in wild rats (*Epimys rattus alexandrinus* and *E. norvegicus*). Annotationes Zoologicae Japonenses. 1938; 17:144–163.
- Mullin SW, Colley FC, Stevens GC. Coccidia of Malaysian mammals: New host records and the description of three new species of *Eimeria*. Journal of Protozoology. 1972; 19:260–263. [PubMed: 5032224]
- Ohira T. Ein Beitrag zur Coccidiumforschung. Ueber ein bei Ratten gefundenes Coccidium (*Eimeria miyairii*). Mitt Medicinische Gesellschaft zu Tokio. 1912; 26:1045–1055.
- Pellérdy, LP. Coccidia and coccidiosis (2nd ed.). Joint Publication of Verlag Paul Parey, Berlin and Hamburg and Akadémiai Kiadó, Budapest. 1974. p. 959
- Rabor DS. Notes on the mammals and birds of the central northern Luzon highlands, Philippines. Part I: Notes on mammals. Silliman Journal. 1955; 2:193–218.
- Ricart EA, Balette DS, Heaney LR. Habitat disturbance and the ecology of small mammals in the Philippines. Journal of Environmental Science and Management. 2007; 10:34–41.
- Roudabush RL. The endogenous phases in the life cycle of *Eimeria nieschulzi*, *Eimeria separata*, and *Eimeria miyairii* coccidian parasites of the rat. Iowa State College Journal of Science. 1937; 11:135–163.
- Sanborn CC. Philippine zoological expedition 1946–1947. Mammals. Fieldiana Zoology. 1952; 33:89–158.
- Wilber PG, Duszynski DW, Upton SJ, Seville RS, Corliss JO. A revision of the taxonomy and nomenclature of the *Eimeria* spp. (Apicomplexa: Eimeriidae) from rodents in the Tribe Marmotini (Sciuridae). Systematic Parasitology. 1998; 39:113–135.

Figs. 1–3.

Nomarski interference-contrast photomicrographs of oocysts of *E. macarthuri* n. sp.

Abbreviations: oocyst wall (OW), Stieda body (SB), substieda body (SSB). Scale bars = 10 μm for all figures.

Fig. 4.
Composite line drawing of oocyst of *Eimeria macarthuri* n. sp. Scale bar = 10 μ m.