Published in final edited form as:

Am J Prev Med. 2015 January ; 48(1): 104–107. doi:10.1016/j.amepre.2014.08.023.

Home and Workplace Built Environment Supports for Physical Activity

Deepti Adlakha, MUD, Aaron J. Hipp, PhD, Christine Marx, MA, Lin Yang, PhD, Rachel Tabak, PhD, Elizabeth A. Dodson, PhD, and Ross C. Brownson, PhD

Brown School (Adlakha, Hipp, Dodson, Brownson), Prevention Research Center (Hipp, Tabak, Dodson, Brownson), Division of Public Health Sciences, Department of Surgery, School of Medicine, (Marx, Yang), and the Alvin J. Siteman Cancer Center, School of Medicine, (Brownson), Washington University in St. Louis, St. Louis, Missouri

Abstract

Background—Physical inactivity has been associated with obesity and related chronic diseases. Understanding built environment (BE) influences on specific domains of physical activity (PA) around homes and workplaces is important for public health interventions to increase population PA.

Purpose—To examine the association of home and workplace BE features with PA occurring across specific life domains (work, leisure, and travel).

Methods—Between 2012 and 2013, telephone interviews were conducted with participants in four Missouri metropolitan areas. Questions included sociodemographic characteristics, home and workplace supports for PA, and dietary behaviors. Data analysis was conducted in 2013; logistic regression was used to examine associations between BE features and domain-specific PA.

Results—In home neighborhoods, seven of 12 BE features (availability of fruits and vegetables, presence of shops and stores, bike facilities, recreation facilities, crime rate, seeing others active, and interesting things) were associated with leisure PA. The global average score of home neighborhood BE features was associated with greater odds of travel PA (AOR=1.99, 95% CI=1.46, 2.72), leisure PA (AOR=1.84, 95% CI=1.44, 2.34), and total PA (AOR=1.41, 95% CI=1.04, 1.92). Associations between workplace neighborhoods' BE features and workplace PA were small, but in the expected direction.

Conclusions—This study offers empirical evidence on BE supports for domain-specific PA. Findings suggest that diverse, attractive, and walkable neighborhoods around workplaces support walking, bicycling, and use of public transit. Public health practitioners, researchers, and worksite

© 2014 Elsevier Inc. on behalf of American Journal of Preventive Medicine. All rights reserved.

Address correspondence to: Deepti Adlakha, MUD, One Brookings Drive, Washington University in St. Louis, Campus Box 1196, Saint Louis MO 63130. deeptia@wustl.edu.

No financial disclosures were reported by the authors of this paper.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

leaders could benefit by utilizing worksite domains and measures from this study for future BE assessments.

Introduction

Physical inactivity is a primary modifiable risk factor for obesity and related chronic diseases.¹ Evidence shows a steep rise in rates of physical inactivity among urban populations.² Fewer than 50% of adults and 40% of youth meet U.S. physical activity (PA) guidelines, with the built environment (BE) identified as a reason for limited PA.^{3–6}

Elements of home neighborhood BE (e.g., streets, sidewalks) have been widely investigated.⁷ However, their impact on specific domains of everyday life in which PA occurs (e.g., at work, while travelling) needs exploration. Because adults spend about half of their waking hours at workplaces,⁸ the influence of workplace neighborhoods is potentially important. This study examines relationships between home and workplace BE features, domain-specific PA (work, travel, leisure), and total PA.

Methods

In 2012–2013, telephone interviews (N=2,015) were conducted with employed adults residing in four Missouri metropolitan areas: St. Louis, Kansas City, Springfield, and Columbia. Eligibility criteria included: (1) ability to speak English or Spanish; (2) age 21–65 years; (3) employed in at least one occupation (20 hours/week) at a single worksite; and (4) worksite has five employees. Exclusion criteria included: (1) presence of medical condition that interferes with ability to walk; (2) pregnancy; and (3) self-employment. Given lack of consensus on what defines a unique neighborhood,⁹ census tracts were used as the primary definition. Two samples were drawn: a (1) representative sample, with oversampling of census tracts in small metropolitan areas; and (2) high-density/minority sample drawn from census tracts with high population density and high minority population (African American and Hispanic). Detailed study design, participant recruitment, and sociodemographic characteristics (Appendix Table 1) have been described in supplementary text and related studies.^{10, 11}

Twelve items from the Physical Activity Neighborhood Environment Survey (PANES)¹² (Table 1) were used to measure BE features around homes and workplaces. Items were identical across both neighborhoods. Reliability of items has been previously established, with reliability coefficients above 0.60.¹⁰ Four-point Likert scale response options ranging from 1 (strongly agree) to 4 (strongly disagree) were combined as “agree” (*strongly agree, agree*) and “disagree” (*disagree, strongly disagree*). A global average score for home and workplace neighborhood BE features was constructed using Likert scale responses from all 12 items.

In 2013, a series of multiple logistic regressions were used to identify BE predictors of meeting CDC-recommended levels (≥150 minutes/week) for work, travel, leisure, and total PA (dichotomous outcomes) using SPSS, version 21 (IBM Corp., Armonk NY). Pairwise interactions between PANES items and interactions between home and workplace global average scores were tested. Physical activity was measured with the International Physical

Activity Questionnaire (IPAQ).¹³ Models were adjusted for age, sex, race, education, and income. Descriptive statistics (Appendix Tables 2–4) and details on data analysis are provided in the supplementary text. The IRBs of Washington University in St. Louis and University of Missouri-Columbia approved study procedures.

Results

In home neighborhoods, seven of 12 BE features (availability of fruits and vegetables, presence of shops and stores, bike facilities, recreation facilities, crime rate, seeing others active, and interesting things to look at) were associated with leisure PA (Table 1). Six home neighborhood BE features significantly predicted travel PA. Except for crime, no clear associations of individual home neighborhood BE features with workplace PA or total PA were observed. The global average score of home neighborhood BE features predicted significant increases in adjusted odds of engagement in travel PA (AOR=1.99, 95% CI=1.46, 2.72), leisure PA (AOR=1.84, 95% CI=1.44, 2.34), and total PA (AOR=1.41, 95% CI=1.04, 1.92).

A smaller number of associations between workplace neighborhood BE features and workplace PA were significant. Three of 12 workplace neighborhood BE features (bike facilities, interesting things to look at, and crime rate) significantly predicted workplace PA (Table 2). Although other workplace BE features were not significant, overall adjusted odds of engagement in workplace PA were in the expected direction. Certain BE features in workplace neighborhoods (e.g., healthy restaurants, bicycle and recreation facilities) significantly supported travel and leisure PA. Bike facilities and presence of interesting things in workplace neighborhoods were significantly associated with PA across all domains. Sidewalks and walkable access to transit were significantly associated with total PA around workplaces.

Similar to home neighborhoods, global average score of workplace BE features was associated with travel PA (AOR=2.13, 95% CI=1.62, 2.80), leisure PA (AOR=1.34, 95% CI=1.09, 1.64), and total PA (AOR=1.30, 95% CI=1.00, 1.68). However, the association between global average score of workplace neighborhood supports with workplace PA was insignificant (AOR=1.01, 95% CI=0.82, 1.25). No significant interactions (pairwise, global average scores) were found between home and workplace BE features.

Discussion

This study is among the first to examine associations between home and workplace BE features (individual, global average) and PA domains around homes and workplaces. Previous studies have demonstrated significant associations between individual BE features and total PA,^{14–16} predominantly in home neighborhoods.

The presence of physically active people and interesting things around homes improved odds of leisure and travel PA. Three BE features around workplaces (presence of shops and stores, bicycle facilities, and recreation facilities) were associated with travel PA, implying higher engagement in active travel. Bicycling facilities around workplaces increased odds of engagement in travel PA. Interesting things around workplaces resulted in a greater

likelihood of engagement in PA across all domains. Around workplaces, public transit access and presence of sidewalks were significantly associated with total PA, while healthy restaurants promoted leisure and travel PA. These associations imply diverse, attractive, and walkable workplace neighborhoods can serve as incentives for walking, bicycling, and active commuting. Cumulatively, BE features predicted increased odds of travel, leisure, and total PA around homes and workplaces, providing additional evidence of BE supports for specific PA domains.

Crime is a frequently cited barrier to PA.¹⁷ However, its association with PA was inconsistent in this analysis. Previous studies have yielded similar inconclusive results acknowledging that the impact of perceived safety from crime on walking in residential neighborhoods needs careful examination.^{14, 17} This may relate to the complexity of measuring crime (e.g., time of occurrence, people's perceptions, and coping mechanisms influence PA differently).¹⁴

The cross-sectional study design limits causal inference.¹⁸ Certain census tracts were oversampled to increase variability by race/ethnicity and walkability, but this was not adjusted for in statistical models. Self-reported PA and neighborhood measures are subject to bias (e.g., social desirability of PA; physically active people may notice more neighborhood destinations). Another limitation of this study and PA literature in general is a lack of consensus on measuring workplace PA (e.g., lack of tested items, inadequate details on types of workplace PA).^{19, 20} The IPAQ measure of workplace PA is meant to capture only work-related PA.^{21, 22} However, without a clear description of workplace PA, study participants could be reporting time spent in PA during work hours (e.g., walking or bicycling to a restaurant during lunch) as workplace, leisure, or travel PA.

Data on BE characteristics to support PA across specific domains (work, travel, and leisure) provide information to guide design decisions for healthy living around homes and workplaces.^{23, 24} Addition of key worksite domains can guide the development of objective measures of worksite PA to explore combined effects of BE around homes and workplaces. Additional research is needed to advance measurement and evaluation of workplace PA.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the NCI Centers for Transdisciplinary Research on Energetics and Cancer (TREC) (Washington University in St. Louis, U54 CA155496). The opinions or assertions contained herein are the private ones of the authors and are not considered as official or reflecting the views of the National Institutes of Health.

References

1. Brownson RC, Hoehner CM, Day K, Forsyth A, Sallis JF. Measuring the built environment for physical activity: state of the science. *Am J Prev Med.* 2009; 36(4S):S99–123. e12. [PubMed: 19285216]
2. Guthold R, Ono T, Strong KL, Chatterji S, Morabia A. Worldwide variability in physical inactivity a 51-country survey. *Am J Prev Med.* 2008; 34(6):486–94. [PubMed: 18471584]

3. CDC. Division of Nutrition, Physical Activity and Obesity. 2009. www.cdc.gov/nccdphp/dnpa/index.htm
4. Saelens BE, Sallis JF, Frank LD. Environmental correlates of walking and cycling: Findings from the transportation, urban design and planning literatures. *Ann Behav Med*. 2003; 25(2):80–91. [PubMed: 12704009]
5. Saelens BE, Handy SL. Built environment correlates of walking: a review. *Med Sci Sports Exerc*. 2008; 40(7S):S550–66. [PubMed: 18562973]
6. Heath GW, Brownson RC, Kruger J, Miles R, Powell KE, Ramsey LT. The effectiveness of urban design and land use and transport policies and practices to increase physical activity: A systematic review. *J Phys Act Health*. 2006; 3 (Suppl 1):S55–76.
7. Kaczynski AT, Henderson KA. Environmental correlates of physical activity: a review of evidence about parks and recreation. *Leis Sci*. 2007; 29(4):315–54.
8. U.S. Bureau of Labor Statistics. American Time Use Survey. 2013. <http://www.bls.gov/tus/>
9. Foster KA, Hipp JA. Defining Neighborhood Boundaries for Social Measurement: Advancing Social Work Research. *Soc Work Res*. 2011; 35(1):25–35.
10. Hoehner CM, Budd EL, Marx CM, Dodson EA, Brownson RC. Development and reliability testing of the Worksite and Energy Balance Survey. *J Public Health Manag Pract*. 2013; 19(3 Suppl 1):S105–13. [PubMed: 23529049]
11. Patterson RE, Colditz GA, Hu FB, et al. The 2011–2016 Transdisciplinary Research on Energetics and Cancer (TREC) initiative: rationale and design. *Cancer Causes Control*. 2013; 24(4):695–704. [PubMed: 23378138]
12. Sallis JF, Kerr J, Carlson JA, et al. Evaluating a brief self-report measure of neighborhood environments for physical activity research and surveillance: Physical activity neighborhood environment scale (PANES). *J Phys Act Health*. 2010; 7(4):533–40. [PubMed: 20683096]
13. IPAQ. International physical activity questionnaires. IPAQ: short last 7 days telephone format. http://www.ipaq.ki.se/questionnaires/IPAQ_SHORT_LAST_7_TELEPHONE-revised_8-23-02.doc
14. Ding D, Adams M, Sallis J, et al. Perceived neighborhood environment and physical activity in 11 countries: Do associations differ by country? *Int J Behav Nutr Phys Act*. 2013; 10:57.10.1186/1479-5868-10-57. [PubMed: 23672435]
15. Adams M, Ding D, Sallis J, et al. Patterns of neighborhood environment attributes related to physical activity across 11 countries: a latent class analysis. *Int J Behav Nutr Phys Act*. 2013; 10:34.10.1186/1479-5868-10-34. [PubMed: 23497187]
16. Moore K, Diez Roux AV, Auchincloss A, et al. Home and work neighbourhood environments in relation to body mass index: the Multi-Ethnic Study of Atherosclerosis (MESA). *J Epidemiol Community Health*. 2013; 67(10):846–53. [PubMed: 23868527]
17. Foster S, Giles-Corti B. The built environment, neighborhood crime and constrained physical activity: an exploration of inconsistent findings. *Prev Med*. 2008; 47(3):241–51. [PubMed: 18499242]
18. Glass TA, Goodman SN, Hernan MA, Samet JM. Causal inference in public health. *Annu Rev Public Health*. 2013; 34:61–75. [PubMed: 23297653]
19. Steele R, Mummery K. Occupational physical activity across occupational categories. *J Sci Med Sport*. 2003; 6(4):398–407. [PubMed: 14723390]
20. Wood TM. Issues and Future Directions in Assessing Physical Activity: An Introduction to the Conference Proceedings. *Research Q for Exerc Sport*. 2000; 71(2):ii–iv.
21. Hagströmer M, Oja P, Sjöström M. The International Physical Activity Questionnaire (IPAQ): a study of concurrent and construct validity. *Public Health Nutr*. 2006; 9(06):755–62. [PubMed: 16925881]
22. Alexander A, Bergman P, Hagstromer M, Sjoström M. The IPAQ environmental module (Reliability testing). *J Publ Health*. 2006; 14(2):76–80.
23. Brownson RC, Boehmer TK, Luke DA. Declining rates of physical activity in the United States: what are the contributors? *Annu Rev Publ Health*. 2005; 26:421–43.

24. Frank L, Sallis J, Conway T, Chapman J, Saelens B, Bachman W. Many pathways from land use to health: Associations between neighbourhood walkability and active transportation, body mass index, and air quality. *J Am Plann Assoc.* 2006; 72(1):75–87.

Table 1

Home neighborhood predictors of meeting CDC recommended levels of work, travel, leisure, and total PA

Variable	Work PA (aOR ^a , 95% CI)	Travel PA (aOR, 95% CI)	Leisure PA (aOR, 95% CI)	Total PA (aOR, 95% CI)
Large selection of fresh fruits and vegetables	1.06 (0.84–1.35)	1.27 (0.94–1.72)	1.33 (1.05–1.68)	1.23 (0.92–1.63)
Opportunities to purchase fast food	0.77 (0.58–1.02)	1.20 (0.85–1.70)	0.92 (0.70–1.21)	0.77 (0.55–1.09)
Presence of healthy restaurants	0.99 (0.80–1.22)	1.28 (0.98–1.68)	1.12 (0.91–1.39)	1.10 (0.84–1.43)
10–15 minute walk to a transit stop	0.89 (0.70–1.13)	1.06 (0.78–1.45)	1.01 (0.80–1.28)	0.90 (0.66–1.22)
Sidewalks on most streets	0.94 (0.72–1.23)	1.61 (1.11–2.34)	1.23 (0.95–1.59)	1.01 (0.73–1.40)
Shops, stores, or markets	1.08 (0.89–1.32)	1.33 (1.03–1.71)	1.22 (1.01–1.48)	1.16 (0.91–1.49)
Facilities to bicycle	0.90 (0.71–1.09)	1.31 (1.02–1.67)	1.27 (1.05–1.53)	1.09 (0.86–1.39)
Recreation facilities	1.06 (0.84–1.34)	1.46 (1.08–1.97)	1.27 (1.02–1.59)	1.27 (0.96–1.67)
Crime rate makes it unsafe to walk	0.80 (0.64–0.98)	0.99 (0.76–1.27)	1.25 (1.02–1.54)	1.05 (0.81–1.36)
Traffic makes it difficult to walk	0.80 (0.63–1.03)	0.94 (0.70–1.25)	1.23 (0.97–1.56)	1.04 (0.73–1.41)
See people being physically active	1.03 (0.82–1.30)	1.51 (1.13–2.03)	1.59 (1.27–2.00)	1.24 (0.94–1.64)
Interesting things to look at	0.95 (0.77–1.17)	1.31 (1.01–1.70)	1.68 (1.37–2.07)	1.18 (0.92–1.53)
Global Average Home Neighborhood Score	0.91 (0.71–1.16)	1.99 (1.46–2.72)	1.84 (1.44–2.34)	1.41 (1.04–1.92)

Boldface indicates statistical significance (p<0.05). n range: 1784–1862

PA, Physical Activity

^a Adjusted for age (continuous), and sex, race, education, income (all categorical).

Table 2

Workplace neighborhood predictors of meeting CDC recommended levels of work, travel, leisure, and total PA

Variable	Work PA (aOR ^a , 95% CI)	Travel PA (aOR, 95% CI)	Leisure PA (aOR, 95% CI)	Total PA (aOR, 95% CI)
Large selection of fresh fruits and vegetables	1.16 (0.95–1.43)	1.22 (0.95–1.57)	1.23 (1.01–1.50)	1.08 (0.84–1.39)
Opportunities to purchase fast food	0.87 (0.69–1.10)	0.85 (0.63–1.15)	1.01 (0.81–1.27)	1.17 (0.86–1.58)
Presence of healthy restaurants	0.98 (0.79–1.22)	1.53 (1.16–2.02)	1.26 (1.02–1.55)	1.05 (0.83–1.38)
10–15 minute walk to a transit stop	1.21 (0.94–1.56)	1.34 (0.96–1.87)	1.27 (0.99–1.62)	1.36 (1.01–1.84)
Sidewalks on most streets	1.10 (0.88–1.38)	1.30 (0.97–1.74)	1.21 (0.97–1.51)	1.38 (1.05–1.81)
Shops, stores, or markets	1.05 (0.86–1.27)	1.51 (1.19–1.93)	1.14 (0.94–1.38)	1.16 (0.92–1.48)
Facilities to bicycle	1.29 (1.06–1.57)	1.63 (1.28–2.07)	1.33 (1.10–1.61)	1.54 (1.20–1.98)
Recreation facilities	1.15 (0.94–1.40)	1.60* (1.25–2.05)	1.25 (1.03–1.51)	1.20 (0.94–1.53)
Crime rate makes it unsafe to walk	0.76 (0.60–0.96)	1.06 (0.80–1.40)	1.19 (0.96–1.49)	0.92 (0.69–1.22)
Traffic makes it difficult to walk	0.90 (0.74–1.10)	1.15 (0.91–1.47)	1.16 (0.96–1.41)	1.02 (0.80–1.31)
See people being physically active	1.14 (0.94–1.40)	1.44 (1.21–1.86)	1.22 (1.01–1.49)	1.18 (0.92–1.51)
Interesting things to look at	1.22 (1.00–1.59)	1.90 (1.47–2.45)	1.21 (1.00–1.46)	1.33 (1.05–1.70)
Global Average Workplace Neighborhood Score	1.01 (0.82–1.25)	2.13 (1.62–2.80)	1.34 (1.09–1.64)	1.30 (1.00–1.68)

Boldface indicates statistical significance ($p < 0.05$). n range: 1708–1864

PA, Physical Activity

^a Adjusted for age (continuous), and sex, race, education, income (all categorical).