Published in final edited form as:

Circ Res. 2014 October 10; 115(9): 770–780. doi:10.1161/CIRCRESAHA.115.304666.

CD36-Dependent 7-Ketocholesterol Accumulation in Macrophages Mediates Progression of Atherosclerosis in Response to Chronic Air Pollution Exposure

Xiaoquan Rao^{#1}, Jixin Zhong^{#1}, Andrei Maiseyeu¹, Bhavani Gopalakrishnan², Frederick A. Villamena², Lung-Chi Chen³, Jack R Harkema⁴, Qinghua Sun^{2,5}, and Sanjay Rajagopalan¹

¹Division of Cardiovascular Medicine, University of Maryland Baltimore, Maryland, USA

²Department of Pharmacology, Davis Heart & Lung Research Institute, The Ohio State University, Columbus, Ohio, USA

³The Department of Environmental Medicine, New York University School of Medicine, Tuxedo, New York, USA

⁴Center for Integrative Toxicology and Department of Pathobiology and Diagnostic Investigation, Michigan State University, East Lansing, Michigan, USA

⁵College of Public Health, The Ohio State University, Columbus, Ohio, USA

[#] These authors contributed equally to this work.

Abstract

Rationale—Air pollution exposure has been shown to potentiate plaque progression in humans and animals. Our previous studies have suggested a role for oxidized lipids in mediating adverse vascular effect of air pollution. However, the types of oxidized lipids formed in response to air pollutants and how this occurs and their relevance to atherosclerosis is not fully understood.

Objective—To investigate the mechanisms by which particulate matter < 2.5 μ m (PM_{2.5}) induces progression of atherosclerosis.

Methods and Results—Atherosclerosis-prone ApoE^{-/-} or LDLR^{-/-} mice were exposed to filtered air or concentrated ambient PM_{2.5} using a versatile aerosol concentrator enrichment system for 6 months. PM_{2.5} increased 7-ketocholesterol (7-KCh), an oxidatively modified form of cholesterol, in plasma IDL/LDL fraction and in aortic plaque concomitant with progression of atherosclerosis and increased CD36 expression in plaque-macrophages from PM_{2.5}-exposed mice. Macrophages isolated from PM_{2.5}-exposed mice displayed increased uptake of oxidized lipids without alterations in their efflux capacity. Consistent with these finding, CD36-positive macrophages displayed a heightened capacity for oxidized lipid uptake. Deficiency of CD36 on

Address correspondence to: Dr. Sanjay Rajagopalan University of Maryland, Baltimore 20 Penn St Health Science Facility II Baltimore, MD 21201 Tel: 410-328-4716 Fax: 410-328-8527 srajagopalan@medicine.umaryland.edu; Dr. Qinghua Sun The Ohio State University 460 W 12th Avenue Floor 3, Room 388 Biomedical Research Tower Columbus, OH 43210-1252 Tel: 614-247-1506 Fax: 614-688-4233 sun.224@osu.edu.

DISCLOSURES The authors declare no conflict of interests relevant to this paper.

hematopoietic cells diminished the effect of air pollution on 7-KCh accumulation, foam cell formation, and atherosclerosis.

Conclusions—Our results suggest a potential role for CD36-mediated abnormal accumulations of oxidized lipids such as 7-KCh in air pollution induced atherosclerosis progression.

Keywords

Air pollution; 7-ketocholesterol; particulate matter; atherosclerosis; CD36; lipid metabolites; foam cells; macrophage

INTRODUCTION

Both epidemiologic and empirical evidence suggest an important effect of inhaled particulate matter ($<2.5\mu\text{m}$, $\text{PM}_{2.5}$) in mediating cardiovascular events.^{1, 2} The impact of air pollution on global cardiovascular disease and daily mortality is significant and substantial,^{1, 2} given its pervasive presence and ubiquitous exposure, with recent data suggesting that $>17\%$ of global adult mortality is directly attributable to anthropogenic components of $\text{PM}_{2.5}$, primarily through ischemic heart disease events.³ In 2005, 89% of the world's population lived in areas where the WHO Air Quality Guideline of $10\mu\text{g}/\text{m}^3$ $\text{PM}_{2.5}$ (annual average) was exceeded.⁴ We and others have previously demonstrated that controlled exposure to air pollution results in potentiation of atherosclerosis in genetically susceptible models of atherosclerosis, which have been corroborated in human studies using surrogate measures such as coronary artery calcium and carotid intima media thickness.^{2, 5, 6} We have also previously shown that chronic exposure to $\text{PM}_{2.5}$ increases generation of derivatives of oxidized phospholipids with resultant mobilization of mononuclear cells from bone marrow to the circulation, lungs, and the vascular wall via a toll-like receptor 4 (TLR4) mechanism, providing a direct pathway by which air pollution may potentiate atherosclerosis.⁷ However, important questions remain, including the role of other oxidation such as oxidatively modified forms of cholesterol which are fundamental to atherosclerosis lesion progression.^{8, 9} In the present study, we found an increase of 7-ketocholesterol (7-KCh), a major component of oxysterols, in the intermediate density lipoproteins (IDL) and low density lipoprotein (LDL) fraction of plasma lipoproteins and within aortic plaque. In addition, air pollution enhanced CD36-mediated uptake of 7-KCh by macrophages. Disruption of CD36 in hematopoietic cells abolished air pollution-induced 7-KCh accumulation in aortic wall and foam cell formation. Our study thus provides new insights into adverse cardiovascular effects of air pollution exposure by linking air pollution mediated lipoprotein oxidation with CD36 mediated abnormal deposition of oxidized lipids in atherosclerotic plaque and vascular dysfunction.

METHODS

For detailed methods please refer to supplementary section.

Animal studies and exposure protocol

Eight-week-old Apolipoprotein E^{-/-} (ApoE^{-/-}) or LDLR^{-/-} male mice were purchased from Jackson Laboratory (Bar Harbor, ME) and housed at constant temperature ($22 \pm 2^\circ\text{C}$) on a

12-h light/dark cycle. The mice were maintained on regular chow (catalog #862, Harlan, Haslett, MI). Assignments to PM_{2.5} group vs. filtered air (FA) group were randomized. The animal exposure and the monitoring of the exposure environment and ambient aerosol were performed. Exposure to concentrated ambient PM_{2.5} was performed at the Ohio Air Pollution Exposure System for Interrogation of Systemic Effects (OASIS-1) using a versatile aerosol concentrator and exposure system (VACES) as previously described.^{6, 10, 11} In this system, ultrafine (PM < 0.1 µm) particles are not excluded. We have previously demonstrated that atherosclerosis progression enhanced in ApoE^{-/-} mice fed high-fat chow and concomitantly exposed to PM_{2.5} in Tuxedo, NY.⁶ However, since the biological effects of PM_{2.5} may differ based on concomitant diet and composition and size of particulate matter (which in turn is dependent on emission sources), we designed our experiments with a different exposure environment (Columbus, OH regional) and in the context of normal diet conditions¹². Throughout this manuscript, concentrated ambient PM_{2.5} exposure is referred to as PM_{2.5} unless specified otherwise.

PM_{2.5} concentration and composition analysis in the exposure chamber

The overall mean ambient PM_{2.5} level during the exposure periods was $9.1 \pm 7.3 \mu\text{g}/\text{m}^3$. Mean exposure enrichment over ambient levels in the chamber was ≈ 10 -fold higher. Particles from the exposures were collected on Teflon filters (Gelman Teflo, 37 mm, 0.2-mm pore, Gelman Sciences, Ann Arbor, MI). Filters were weighed before and after collecting samples. Analysis of PM_{2.5} composition was performed as previously described.¹¹

7-KCh and cholesterol levels

Liquid chromatography/mass spectrometry was used to determine the level of 7-KCh and cholesterol in the serum and tissues as described previously.^{13, 14} Lipoproteins were isolated by fast protein liquid chromatography. 7-KCh and cholesterol level in lipoprotein fraction were determined by Liquid chromatography/mass spectrometry.

Bone marrow transplantation

Bone marrow transplantation was performed as previously described.¹⁵ Briefly, LDLR^{-/-} mice were irradiated at the dose of 950 Rad. Mice were i.v. injected with 15×10^6 bone marrow cells from C57BL/6 or CD36^{-/-} mice 24 h after irradiation.

Data analyses

All the data in the manuscript are expressed as mean $\pm$ SEM unless otherwise specified (Supplemental Table I). Difference between two groups was tested by student's t test. Differences among groups were tested by two-way ANOVA and Bonferroni's post hoc test using Graphpad Prism 5.01 software. P values of < 0.05 were considered statistically significant.

RESULTS

PM_{2.5} concentrations during the study period

ApoE^{-/-} mice were exposed to filtered air or concentrated PM_{2.5} for a total duration of 6 months. The daily ambient air PM_{2.5} concentration (mean ± SD) during exposure hours was 9.1 ± 7.3 µg/m³. The concentration of PM_{2.5} in the exposure chamber during exposure hours was approximately 11 fold higher than the corresponding ambient level (101.6 ± 30.7 µg/m³). The mean elemental composition of exposure, as measured by energy-dispersive X-ray fluorescence (ED-XRF) analysis, is presented in Supplemental Table I.

Atherosclerosis burden with PM_{2.5}.

After exposure with filtered air (FA) or PM_{2.5} for 6 months, ApoE^{-/-} mice were euthanized and atherosclerotic plaque burden was quantified at the level of the aortic sinus. Starting at the level of the aortic valve leaflet, serial 5µm sections, covering a total length of 600 µm were stained with hematoxylin and eosin (H&E) to evaluate the severity of atherosclerosis. PM_{2.5} exposure significantly increased lesion area, lipid and collagen content of atherosclerotic plaque (Fig. 1A-D). The plaque distribution and lipid content in the aorta is depicted from representative FA and PM_{2.5} exposed animals stained with Oil-Red-O (Fig. 1E).

7-KCh levels in mice exposed to PM_{2.5}.

A preponderance of lipids present in plaques is composed of oxidized cholesterol derivatives (oxysterols).¹⁶ 7-Ketocholesterol (7-KCh), a major component of oxysterols in advanced plaque, has been postulated to play an important role in endothelial dysfunction and atherogenesis.¹⁷⁻²⁰ We therefore evaluated levels of 7-KCh in the tissues of exposed mice. No significant difference of 7-KCh was detected in liver and plasma between PM_{2.5}-exposed and FA-exposed group (Supplemental Fig. I). The level of 7-KCh was ≈7-fold higher in the aorta, while levels in the lung did not differ in response to PM_{2.5} exposure (8.58 ± 2.08 vs. 1.25 ± 0.45 µg/g tissue, p<0.05, n=5 in aorta; 4.55 ± 0.28 vs. 4.98 ± 0.35 µg/g lung tissue in PM_{2.5} vs. FA, p>0.05, n=5 in lung, Fig. 2A & B). We also assessed the amount of cholesterol (CHL) in the same tissues. We failed to detect differences in CHL content between PM_{2.5} and FA exposure group in either the aorta or lung (Fig. 2C & D). When adjusted for levels of cholesterol, the ratio of 7-KCh/CHL increased ≈10- fold in the aorta but not in the lung of PM_{2.5} group (Fig. 2E & F). As 7-KCh has been demonstrated to accumulate in the plasma lipoproteins, with this representing a facile pathway of delivery to the arterial vessel wall, we quantified 7-KCh level in the lipoprotein fraction. As depicted in Fig. 2G, plasma very low density lipoprotein (VLDL) and IDL/LDL fractions contained the bulk of 7-KCh, with these levels being confined to the IDL/LDL fraction once the cholesterol content of the lipoprotein fractions were adjusted for. The 7-KCh/ CHL in plasma LDL was ≈2-fold higher in PM_{2.5}-exposed mice (Fig. 2G & H).

Effect of PM_{2.5} on 7-KCh and cholesterol uptake and efflux

The transport and uptake of oxysterols into macrophages and endothelial cells results in progressive accumulation of lipids in the arterial wall in atherosclerosis.²¹ To test the impact

of PM_{2.5} exposure on macrophage mediated sterol accumulation, we used peritoneal macrophages derived from PM_{2.5}/FA exposed mice. Prior studies have demonstrated that these cells are a reasonable surrogate for plaque macrophages and can be isolated in large enough quantities for our assays.^{22, 23} As shown in Fig. 3A, the uptake of 7-KCh by peritoneal exudate macrophages (PEM) derived from PM_{2.5}-exposed mice was higher than that of FA-exposed mice ($9.4 \pm 1.19\%$ vs. $8.18 \pm 0.76\%$, $p < 0.05$; $n = 6$, Fig. 3A). To additionally validate these results, we also used bone marrow derived macrophages (BMDM) to investigate the effect of collected PM_{2.5} particles on cholesterol/oxysterol transport. After overnight treatment with PM_{2.5}, the ability of 7-KCh uptake by BMDM increased (Supplemental Fig. II). In contrast, no significant difference in 7-KCh uptake by cultured endothelial cells (HUVECs) was detected between PM_{2.5}-treated and vehicle-treated group (Supplemental Fig. III-A). These results suggest that PM_{2.5} exposure promotes uptake of 7-KCh selectively in macrophages but not in endothelial cells. Unmodified cholesterol uptake by PEM was not affected by PM_{2.5} treatment (Fig. 4B). Pretreatment with 7-KCh did not increase the cholesterol uptake by PEM (Fig. 3B).

In addition to uptake, efflux is another process that affects the accumulation of oxysterols in the cell. PM_{2.5} did not impact the efflux of 7-KCh in PEM derived from mice exposed to PM_{2.5} (Fig. 3C). Efflux of cholesterol in PEM was also not influenced by the treatment of PM_{2.5} (Fig. 3D). However, pretreatment of 7-KCh reduced the cholesterol efflux in PEM (Fig. 3D).

PM_{2.5} promotes CD36-mediated 7-KCh uptake by macrophages

Scavenger receptors such as CD36 have been reported to mediate oxLDL internalization in macrophages, and have been implicated in the pathogenesis of atherosclerosis.^{24, 25} To verify the involvement of CD36 in air pollution-induced 7-KCh accumulation, we evaluated the total expression of CD36 on circulating monocytes after PM_{2.5} exposure for 12 weeks. PM_{2.5} exposure doubled CD36 expression on CD11b⁺ monocytes (Figure 4A). To assess the expression of CD36 on plaque-infiltrating macrophage, aortic sinus sections of FA- and PM-exposed ApoE^{-/-} mice were stained with F4/80 and CD36. Most F4/80⁺ macrophages infiltrating the plaque expressed CD36, with an increased number of CD36-expressing macrophages observed in the plaque of PM_{2.5}-exposed mice (Figure 4B & C).

Since CD36 expression was increased in the monocytes and macrophages of PM_{2.5}-exposed mice, we investigated whether PM_{2.5} facilitates 7-KCh uptake through a CD36 dependent mechanism. First, we tested the association between CD36 expression and 7-KCh uptake. Bone marrow derived macrophages (BMDMs) of WT mice were incubated with 7-KCh-loaded LDL (labeled with fluorescent dye, BODIPY) for 8 h and subjected to flow cytometric analysis of CD36 expression and 7-KCh-LDL uptake after extensive washes. CD36-positive BMDMs were more efficient in taking up 7-KCh-LDL compared to CD36-negative BMDMs (Fig. 4D). Next, we investigated whether PM_{2.5} could promote intracellular translocation of CD36. We detected cell surface and total CD36 expression on BMDMs after *in-vitro* treatment with PM_{2.5}. Surface expression of CD36 on macrophages decreased after *in-vitro* treatment with PM_{2.5} (50 µg/ml) for 24 hours (Supplemental Fig. IV-A). Further, we detected total expression of CD36 by staining for CD36 after

permeabilization. There was a small but statistically significant increase in total CD36 (Supplemental Fig. IV-B), suggesting that the decrease of surface CD36 is caused by internalization rather than suppression of synthesis of CD36. To exclude the non-specific effect of lipopolysaccharide (LPS) that is usually found in PM particles, cells were incubated with LPS inhibitor polymyxin (25 µg/ml) and PM_{2.5} particles (50 µg/ml) together. Blockade of LPS with polymyxin did not reverse the effect of PM particles on CD36 internalization (Supplemental Fig. V). In contrast, no difference in the expression of SR-A (scavenger receptor class A, also known as CD204) was observed between PM_{2.5}- and vehicle-treated cells (Supplemental Fig. IV-C & -D). In line with this, non-specific phagocytosis induced by microsphere (1 µm latex beads) did not affect 7-KCh uptake (Supplemental Fig. VI).

Loss of CD36 attenuates air pollution effects on atherosclerosis

To demonstrate the role of CD36 in mediating the effects of PM_{2.5} *in vivo*, LDLR^{-/-} mice were transplanted with bone marrow cells isolated from WT or CD36^{-/-} mice followed by 3-month exposure to PM_{2.5}. PM_{2.5} exposure promoted atherosclerotic plaque development in LDLR^{-/-} mice transplanted with WT bone marrows while deficiency of CD36 on hematopoietic cells reduced plaque volume in response to PM_{2.5} (Fig. 5A & B). Lipid deposition in response to PM_{2.5} exposure was reduced in mice transplanted with CD36^{-/-} bone marrows in comparison with mice transplanted with WT marrow (Fig. 5C & D).

Confocal analysis of aortic sinus sections suggested PM_{2.5} exposure elevated 7-KCh level in the plaque in LDLR^{-/-} mice transplanted with WT bone marrows compared with FA, while 7-KCh level in the plaque of LDLR^{-/-} mice transplanted with CD36^{-/-} bone marrows was comparable to mice transplanted with WT marrow and exposed to FA (Fig. 6A & B). Most macrophages were co-localized with 7-KCh and there were also a lot of 7-KCh single positive areas surrounding macrophages (Fig. 6A). To test whether the increased 7-KCh level was caused by increased plaque size, 7-KCh positive area was normalized by plaque area. As shown in Figure 6C, percentage of 7-KCh positive area increased in the plaque of LDLR^{-/-} mice transplanted with WT bone marrows and exposed to PM_{2.5} when compared to FA, but not in LDLR^{-/-} mice with CD36^{-/-} bone marrows, suggesting that PM_{2.5} increased 7-KCh level independent of plaque lesion size and CD36 deficiency can blunt the effect of PM_{2.5}. Similar results were obtained when normalized with macrophage area (F4/80⁺), indicating increase of 7-KCh in aortic plaque is not due to the increase of macrophage (Fig. 6D). To further confirm the role of CD36 in oxysterol uptake, PEMs isolated from FA- or PM_{2.5}-exposed LDLR^{-/-} mice with WT or CD36^{-/-} bone marrows were incubated with DiI-labeled LDL, oxLDL, or 7-KCh-loaded LDL for 8 h. Fluorescence intensity of the cells was measured by flow cytometry after extensive washes. As depicted in Fig. 6E, similar levels of LDL uptake were observed in all 4 groups. However, uptake of oxLDL or 7-KCh-LDL was enhanced by PM_{2.5} exposure in WT macrophages and loss of CD36 reduced the uptake of oxLDL or 7-KCh-LDL by macrophages. The effect of air pollution on oxLDL or 7-KCh-LDL uptake was markedly diminished by CD36 deficiency (Fig. 6E).

PM_{2.5} increases foam cell formation, which is blunted by CD36 deficiency

Foam cell formation from macrophages with subsequent fatty streaks plays an important role in the pathogenesis of atherosclerosis. Modified LDL including oxidized LDL (oxLDL) or acetylated LDL promotes foam cell formation while LDL in its native form is not atherogenic.^{26, 27} Likewise, there was a marked increase in foam cell formation when macrophages were incubated with 7-KCh-LDL compared with LDL (Fig. 7A & 7D). Dil-labeled 7-KCh-LDL was prepared by loading 7-KCh (2mM) to Dil-labeled LDL. Fluorescence intensity of macrophages incubated with Dil-labeled 7-KCh-LDL was significantly higher compared with that incubated with Dil-labeled LDL, suggesting 7-KCh-LDL rather than native LDL accumulates in macrophages and promotes foam cell formation (Fig. 7A & 7D). To determine the effect of PM_{2.5} and CD36 in the foam cell formation, PEMs isolated from FA- or PM_{2.5}-exposed LDLR^{-/-} mice with WT or CD36^{-/-} bone marrows were incubated with 7-KCh-LDL for 36 h. PM_{2.5} exposure promoted foam cell formation as evidenced by Oil Red O staining and confocal visualization of Dil-labeled 7-KCh-LDL. There was no difference in foam cell formation between FA and PM_{2.5} in CD36 deficient macrophages suggesting that this effect of PM_{2.5} in increasing 7-KCh uptake occurs via CD36 pathways (Fig. 7B, C, E & F).

DISCUSSION

In this study we provide evidence for a novel mechanism by which inhaled particulate matter may promote atherosclerosis. 7-KCh, an oxysterol extensively implicated in atherosclerosis, increased with chronic exposure to PM_{2.5} and preferentially accumulated in the aortic plaque in two different hypercholesterolemic rodent models. 7-KCh was almost exclusively present in the IDL/LDL fraction of plasma lipoproteins and selectively accumulated in macrophages through a CD36 dependent process. Consistent with a role for CD36 in air pollution mediated effects, hematopoietic deficiency of CD36 exerted a protective effect on atherosclerosis, inflammation and lipid accumulation in response to air pollution. CD36⁺ macrophages derived from PM_{2.5} exposed animals preferentially accumulated 7-KCh without demonstrating abnormalities in cholesterol efflux pathways. Our results provide new insights into mechanisms underlying air pollution-induced atherosclerosis progression by linking air pollution with vascular lipid dysregulation.

Multiple studies have confirmed an important pro-atherogenic effect of diesel exhaust particles, PM_{2.5}, and ultrafine components of air pollution.^{5, 6, 10} Limited human cross-sectional studies have demonstrated higher carotid intima-media thickness (CIMT) and coronary artery calcium scores in patients exposed to higher levels of air pollution.^{28, 29} Abnormalities in endothelial function have also been described in both acute and chronic exposure studies (primarily in animal models).^{30, 31} We and others have demonstrated increased lipid accumulation within plaques of animals exposed to air pollution together with macrophage accumulation.^{6, 32-34} Increased expression of oxidative stress markers such as iNOS, nitrotyrosine and CD36 have been previously demonstrated in plaques after both PM_{2.5} and diesel exhaust exposure.^{34, 35} While the pro-atherogenic effects of PM_{2.5} in animal models are consistent with the epidemiologic evidence linking PM_{2.5} with atherosclerotic events such as acute myocardial infarction and stroke, the precise pathways

that sustain progression of plaque with exposure resulting in unstable coronary syndromes are poorly understood.² In atherosclerosis, a critical step is recognition and internalization of oxidized LDL (oxLDL) by specific macrophage scavenger receptors (with CD36 accounting for 60-70% of oxLDL uptake).³⁶ CD36 has been shown to mediate oxLDL internalization in macrophages, with its expression correlating with exposure measures such as alveolar macrophages positive for inhaled particles³⁴. We therefore investigated this pathway and the mechanisms that may lead to CD36 mediated foam cell formation in response to PM_{2.5} exposure. CD36 expression was increased in plaque macrophages from PM_{2.5} exposed animals. In our studies, we found a marked increase in 7-KCh in the aorta of exposed hyperlipidemic animals with co-localization primarily in plaque macrophages. Interestingly, nonspecific phagocytosis mediated by in-vitro exposure to microspheres did not enhance 7-KCh uptake. PM_{2.5} exposure did enhance the uptake of 7-KCh. These results suggest that PM_{2.5} promotes the accumulation of 7-KCh by enhancing CD36-mediated internalization rather than non-specific phagocytosis. Consistent with our findings, other air pollutants have also been shown to increase CD36 expression. Bai et al. reported a 1.5 to 2-fold increase of CD36 in plaque, accompanying with 1.5 to 3-fold increase of lipid content after exposure to diesel exhaust.³⁴ CD36 expression was correlated positively with increased entry of monocytes into atherosclerotic plaques.³⁴ Robertson et al have suggested that CD36 mediates the vasoconstrictor effect of circulating vasoactive factors released in response to ozone exposure.³⁷ However, our study does not exclude other scavenger receptors such as LOX-1, CD68, and SR-BI that may also be involved in enhanced atherogenesis. Exposure to engine emissions increased lectin-like oxLDL receptor (LOX-1), plasma oxLDL, and vascular infiltration of monocyte/macrophage in ApoE^{-/-} mice, which were attenuated by the blockade of LOX-1 using neutralizing anti-LOX-1 Ab.³⁸ These findings raise the possibility that air pollutants such as PM_{2.5} as well as gaseous components of vehicle emission may also influence atherogenesis through similar mechanisms (scavenger receptors-mediated dysregulation of oxidized lipids).

We have previously demonstrated that chronic exposure to PM_{2.5} increased the generation of oxidized phospholipid derivatives of 1-palmitoyl-2-arachidonoyl-sn-glycero-3-phosphorylcholine (oxPAPC) such as 1-palmitoyl-2-(5-oxovaleroyl)-sn-glycero-3-phosphocholine (POVPC) and 1-palmitoyl-2-glutaroyl-sn-glycero-3-phosphocholine (PGPC) in the lungs.⁷ The generation of oxPAPC resulted in activation of NF B and pro-inflammatory genes including chemokines such as chemokine (CC motif) ligand 2 (CCL2) through a TLR4/NADPH oxidase dependent pathway.⁷ Mice deficient in NADPH oxidase 2 (NOX2) and TLR4 demonstrated reduced vascular inflammation and improved vascular dysfunction seen with PM_{2.5} exposure.⁷ TLR4 deficiency also abolished circulating increase of Ly6C^{hi} monocytes in response to long term PM_{2.5} exposure, providing a cellular mechanism through which exposure may facilitate inflammation in atherosclerosis.⁷ We have postulated previously that since PAPC is abundantly present in tissues such as the lung (dominant phospholipid in the surfactant fluid of lung), the generation of oxidized forms of PAPC could easily occur with air pollution exposure resulting in TLR4 activation and transcription of pro-inflammatory genes and chemokines. The role of oxidation products other than oxPAPC that may arise with air pollution exposure such as oxidatively modified forms of cholesterol has not been investigated thus far. Since air pollution exposure serves

as a potent redox trigger, facilitating the generation of a range of oxidatively modified products, it is entirely possible that other oxidation products of cholesterol are also generated. In atherosclerosis, lesion formation and progression is fundamentally dependent on the generation of oxidized forms of cholesterol, which are rapidly taken up by macrophages in lesions via scavenger receptor dependent and independent pathways, resulting in lipid accumulation within the vessel wall. 7-KCh is a major oxidation product present in oxLDL and within atherosclerotic plaque.^{16, 17} In current study, we detected elevated levels of 7-KCh in plasma lipoproteins (IDL/LDL) after chronic exposure to air pollution. 7-KCh but not CHL increased in the LDL/IDL fraction of plasma lipoproteins of PM_{2.5}-exposed mice, raising the possibility that elevated 7-KCh in the aorta may result from increased accumulation, through delivery of ApoB lipoproteins enriched in 7-KCh. Consistent with these findings, deficiency of CD36 markedly decreased the internalization of oxidized LDL and 7-KCh-loaded LDL but had no effect on the uptake of native LDL. Macrophages with low expression of CD36 had a significantly lower ability to uptake oxidized and 7-KCh-loaded LDL compared with macrophages with high expression of CD36. Furthermore, *ex vivo* macrophage uptake of 7-KCh-loaded LDL or oxLDL as well as foam cell formation were enhanced by PM_{2.5} exposure, results that were partially abolished by CD36 deficiency. We further demonstrated in carefully performed in-vivo experiments that bone marrow deficiency of CD36 attenuates PM_{2.5} mediated effects on atherosclerotic plaque and lipid accumulation. F4/80+ macrophages in plaque co-localized with 7-KCh within the plaque with abundant 7-KCh surrounding macrophage, suggesting release of 7-KCh presumably from apoptotic macrophages containing the oxysterol. CD36 deficiency in bone marrow derived cells also reduced atherosclerotic lesion in FA-exposed mice. This result is in accordance with previous reports.^{36, 39, 40} Although Moore et al reported that CD36 deficiency in ApoE^{-/-} background did not affect atherosclerotic progression in their work⁴¹, subsequent studies including those from Sheedy et al suggest CD36 deletion did reduce atherosclerotic plaque in ApoE^{-/-} mice.^{40, 42, 43} The differences noted in these studies may be attributable to the use of two different mouse strains by the groups as pointed out previously.⁴⁴ Our result that LDLR^{-/-} mice with CD36-null bone marrow were protected from atherosclerosis is indeed consistent with another previous report that transplantation of CD36-null bone marrow reduces atherosclerotic lesion formation.³⁹

The finding in the current investigation that CD36 mediates air pollution induced 7-KCh accumulation and atherosclerosis progression provides complementary mechanisms to our prior studies linking TLR4 to abnormal vascular effects and cellular inflammation associated with air pollution exposure.⁷ Oxidized phospholipids such as ox-PAPC may result in pro-inflammatory effects through NF B mechanisms that may synergistically interact with accumulation of lipids such as 7-KCh in macrophages to accelerate atherogenesis. Based on our findings of increased 7-KCh in lipoproteins such as VLDL and LDL/IDL and lack of increase in lung derived 7-KCh, it is likely that 7-KCh is continually formed in the vascular compartment during air pollution exposure particularly on the surface of cholesterol loaded lipoproteins such as LDL and IDL. We posit that the continual exposure of LDL to air pollution particularly at the alveolar-capillary membrane, likely plays a role in this process. LDL/IDL containing oxidized lipids such as 7-KCh and oxPAPC in response to air pollution

may be delivered to the atherosclerotic aorta expressing scavenger receptors and CD36. A recent study suggested CD36 may also activate TLR4 and induces sterile inflammation by forming a TLR4/TLR6 heterodimer, thus providing additional pathways by which TLR4 may interact with pathways involving CD36.⁴⁵ In the current study we did not measure oxPAPC in LDL fraction although prior studies have demonstrated increased oxPAPC in circulating lipoproteins including LDL and HDL.^{46, 47} Although oxPAPC may not be the major component of oxLDL or oxidized lipids within plaque, it certainly may represent an additional oxidation product that is delivered via scavenger receptors and may accumulate within atherosclerosis. Additionally the presence of oxPAPC may result in pro-inflammatory effects within the vascular wall.⁴⁸ The relative contribution of 7-KCh versus other oxidized lipids in air pollution mediated atherosclerosis progression will require additional studies.

Mass transfer of 7-KCh can also occur in the opposite direction from within the cell to a wide range of phospholipid-containing acceptors, including high density lipoprotein (HDL).⁴⁹ Recently, ATP-binding cassette sub-family G member 1 (ABCG1) has been suggested to promote the export of 7-KCh and 7 β -hydroxycholesterol (7 β -HC) to HDL, and to exert cytoprotective effects.^{17, 50} In our study, efflux assays performed with both 7-KCh and CHL demonstrated that pre-treatment with PM_{2.5} does not influence 7-KCh and CHL efflux in either macrophages or endothelial cells. Interestingly, we observed increased CHL uptake in endothelial cells and a decrease of CHL efflux in macrophages when cells were pre-treated with 7-KCh. These findings suggest a feed-forward effect on lipid accumulation following 7-KCh exposure/accumulation. These results are consistent with those of Gelissen and co-workers who demonstrated that the presence of 7-KCh suppresses the export of CHL from macrophage to Apolipoprotein A-I acceptor.⁵¹ Additionally, studies have suggested that 7-KCh suppresses incorporation of CHL to plasma membrane and may thus reduce CHL efflux to cholesterol acceptors.⁵²⁻⁵⁴

Our study has a number of important limitations that must be acknowledged. Peritoneal macrophages from mice exposed to PM_{2.5} were used to assess 7-KCh uptake in plaque macrophage in light of the large amount of cells to perform the assays. Prior studies however have demonstrated that these cells are however reasonable surrogates for plaque macrophages.^{22, 23} We believe that despite this limitation, these studies help provide a framework of understanding of how exposure to PM_{2.5} may modulate uptake of 7-KCh. Accumulation of 7-KCh in plaque depends importantly on circulating levels of 7-KCh in lipoproteins, 7-KCh uptake and retention of 7-KCh in the sites of lesion formation. In the present study, we have not provided insights on to how 7-KCh accumulates selectively in lipoproteins, nor have we provided information on the precise locus of 7-KCh formation in LDL. We postulate that plasma lipoproteins may undergo oxidation at the alveolar-capillary membrane interface in response to repetitive PM_{2.5} exposure. This may potentially explain lack of parenchymal lung accumulation of 7-KCh as was noted in the study.

In summary our findings provide a novel mechanism by which air pollution may enhance the development of atherosclerosis via linking CD36-mediated abnormal accumulation of oxidized lipid in vasculature with air pollution induced lipid oxidation. Further studies are required to determine which components of PM_{2.5} are required for lipid oxidation, other

receptors involved in the deposition of oxidized lipid in vascular wall, and the precise compartment where 7-KCh is generated.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGEMENTS

We would like to thank Aixia Wang and Yuntao Bai from Davis Heart and Lung Research Institute, The Ohio State University for their assistance on the exposure of PM_{2.5}.

SOURCES OF FUNDING This work was supported by ES018900 to Dr. Sun; RO-1ES015146, and RO-1ES017290 RO-1ES019616 to Dr. Rajagopalan and an Environmental Protection Agency grant (RD83479701) to Dr. Harkema and Dr. Rajagopalan. Dr. Zhong was supported by American Heart Association Grant 13POST17210033 and National Natural Science Foundation of China grant 81101553/H1604. Dr. Maisseyeu was supported by an American Heart Association NCRP Scientist Development Grant 13SDG14500015.

Nonstandard Abbreviations and Acronyms

7-KCh	7-ketocholesterol
7β-HC	7β-hydroxycholesterol
ABCG1	ATP-binding cassette sub-family G member 1
BMDM	bone marrow derived macrophage
CHL	cholesterol
CIMT	carotid intima-media thickness
ED-XRF	energy-dispersive X-ray fluorescence
FA	filtered air
H&E	hematoxylin and eosin
HDL	high density lipoprotein
IDL	intermediate density lipoprotein
LDL	low density lipoprotein
LPS	lipopolysaccharide
OASIS	Ohio Aerosol Exposure System for Interrogation of Systemic Effects
oxLDL	oxidized LDL
PAPC	1-palmitoyl-2-arachidonyl-sn-glycero-3-phosphorylcholine
PEM	peritoneal exudate macrophage
PM	particulate matter
SR-A	scavenger receptor class A
TLR4	Toll-like receptor 4
VACES	Versatile Aerosol Concentrator Enrichment System

VLDL very low density lipoproteins

REFERENCES

1. Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, Amann M, Anderson HR, Andrews KG, Aryee M, Atkinson C, Bacchus LJ, Bahalim AN, Balakrishnan K, Balmes J, Barker-Collo S, Baxter A, Bell ML, Blore JD, Blyth F, Bonner C, Borges G, Bourne R, Boussinesq M, Brauer M, Brooks P, Bruce NG, Brunekreef B, Bryan-Hancock C, Bucello C, Buchbinder R, Bull F, Burnett RT, Byers TE, Calabria B, Carapetis J, Carnahan E, Chafe Z, Charlson F, Chen H, Chen JS, Cheng AT, Child JC, Cohen A, Colson KE, Cowie BC, Darby S, Darling S, Davis A, Degenhardt L, Dentener F, Des Jarlais DC, Devries K, Dherani M, Ding EL, Dorsey ER, Driscoll T, Edmond K, Ali SE, Engell RE, Erwin PJ, Fahimi S, Falder G, Farzadfar F, Ferrari A, Finucane MM, Flaxman S, Fowkes FG, Freedman G, Freeman MK, Gakidou E, Ghosh S, Giovannucci E, Gmel G, Graham K, Grainger R, Grant B, Gunnell D, Gutierrez HR, Hall W, Hoek HW, Hogan A, Hosgood HD 3rd, Hoy D, Hu H, Hubbell BJ, Hutchings SJ, Ibeanusi SE, Jacklyn GL, Jasrasaria R, Jonas JB, Kan H, Kanis JA, Kassebaum N, Kawakami N, Khang YH, Khatibzadeh S, Khoo JP, Kok C, Laden F, Lalloo R, Lan Q, Lathlean T, Leasher JL, Leigh J, Li Y, Lin JK, Lipshultz SE, London S, Lozano R, Lu Y, Mak J, Malekzadeh R, Mallinger L, Marcenes W, March L, Marks R, Martin R, McGale P, McGrath J, Mehta S, Mensah GA, Merriman TR, Micha R, Michaud C, Mishra V, Mohd Hanafiah K, Mokdad AA, Morawska L, Mozaffarian D, Murphy T, Naghavi M, Neal B, Nelson PK, Nolla JM, Norman R, Olives C, Omer SB, Orchard J, Osborne R, Ostro B, Page A, Pandey KD, Parry CD, Passmore E, Patra J, Pearce N, Pelizzari PM, Petzold M, Phillips MR, Pope D, Pope CA 3rd, Powles J, Rao M, Razavi H, Rehfuess EA, Rehm JT, Ritz B, Rivara FP, Roberts T, Robinson C, Rodriguez-Portales JA, Romieu I, Room R, Rosenfeld LC, Roy A, Rushton L, Salomon JA, Sampson U, Sanchez-Riera L, Sanman E, Sapkota A, Seedat S, Shi P, Shield K, Shivakoti R, Singh GM, Sleet DA, Smith E, Smith KR, Stapelberg NJ, Steenland K, Stockl H, Stovner LJ, Straif K, Straney L, Thurston GD, Tran JH, Van Dingenen R, van Donkelaar A, Veerman JL, Vijayakumar L, Weintraub R, Weissman MM, White RA, Whiteford H, Wiersma ST, Wilkinson JD, Williams HC, Williams W, Wilson N, Woolf AD, Yip P, Zielinski JM, Lopez AD, Murray CJ, Ezzati M, AlMazroa MA, Memish ZA. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: A systematic analysis for the global burden of disease study 2010. *Lancet*. 2012; 380:2224–2260. [PubMed: 23245609]
2. Brook RD, Rajagopalan S, Pope CA 3rd, Brook JR, Bhatnagar A, Diez-Roux AV, Holguin F, Hong Y, Luepker RV, Mittleman MA, Peters A, Siscovick D, Smith SC Jr, Whitsel L, Kaufman JD. Particulate matter air pollution and cardiovascular disease: An update to the scientific statement from the American Heart Association. *Circulation*. 2010; 121:2331–2378. [PubMed: 20458016]
3. Evans J, van Donkelaar A, Martin RV, Burnett R, Rainham DG, Birkett NJ, Krewski D. Estimates of global mortality attributable to particulate air pollution using satellite imagery. *Environmental research*. 2012
4. Brauer M, Amann M, Burnett RT, Cohen A, Dentener F, Ezzati M, Henderson SB, Krzyzanowski M, Martin RV, Van Dingenen R, van Donkelaar A, Thurston GD. Exposure assessment for estimation of the global burden of disease attributable to outdoor air pollution. *Environmental science & technology*. 2012; 46:652–660. [PubMed: 22148428]
5. Araujo JA, Barajas B, Kleinman M, Wang X, Bennett BJ, Gong KW, Navab M, Harkema J, Sioutas C, Lusk AJ, Nel AE. Ambient particulate pollutants in the ultrafine range promote early atherosclerosis and systemic oxidative stress. *Circulation research*. 2008; 102:589–596. [PubMed: 18202315]
6. Sun Q, Wang A, Jin X, Natanzon A, Duquaine D, Brook RD, Aguinaldo JG, Fayad ZA, Fuster V, Lippmann M, Chen LC, Rajagopalan S. Long-term air pollution exposure and acceleration of atherosclerosis and vascular inflammation in an animal model. *JAMA : the journal of the American Medical Association*. 2005; 294:3003–3010.
7. Kampfrath T, Maisyey A, Ying Z, Shah Z, Deiluiis JA, Xu X, Kherada N, Brook RD, Reddy KM, Padture NP, Parthasarathy S, Chen LC, Moffatt-Bruce S, Sun Q, Morawietz H, Rajagopalan S.

- Chronic fine particulate matter exposure induces systemic vascular dysfunction via nadph oxidase and tlr4 pathways. *Circulation research*. 2011; 108:716–726. [PubMed: 21273555]
8. Stocker R, Keaney JF Jr. Role of oxidative modifications in atherosclerosis. *Physiological reviews*. 2004; 84:1381–1478. [PubMed: 15383655]
 9. Libby P, Ridker PM, Maseri A. Inflammation and atherosclerosis. *Circulation*. 2002; 105:1135–1143. [PubMed: 11877368]
 10. Sun Q, Yue P, Kirk RI, Wang A, Moatti D, Jin X, Lu B, Schecter AD, Lippmann M, Gordon T, Chen LC, Rajagopalan S. Ambient air particulate matter exposure and tissue factor expression in atherosclerosis. *Inhalation toxicology*. 2008; 20:127–137. [PubMed: 18236227]
 11. Xu Z, Xu X, Zhong M, Hotchkiss IP, Lewandowski RP, Wagner JG, Bramble LA, Yang Y, Wang A, Harkema JR, Lippmann M, Rajagopalan S, Chen LC, Sun Q. Ambient particulate air pollution induces oxidative stress and alterations of mitochondria and gene expression in brown and white adipose tissues. *Particle and fibre toxicology*. 2011; 8:20. [PubMed: 21745393]
 12. Brook RD, Franklin B, Cascio W, Hong Y, Howard G, Lipsett M, Luepker R, Mittleman M, Samet J, Smith SC Jr, Tager I. Air pollution and cardiovascular disease: A statement for healthcare professionals from the expert panel on population and prevention science of the american heart association. *Circulation*. 2004; 109:2655–2671. [PubMed: 15173049]
 13. McDonald JG, Thompson BM, McCrum EC, Russell DW. Extraction and analysis of sterols in biological matrices by high performance liquid chromatography electrospray ionization mass spectrometry. *Methods in enzymology*. 2007; 432:145–170. [PubMed: 17954216]
 14. Sevanian A, Seraglia R, Traldi P, Rossato P, Ursini F, Hodis H. Analysis of plasma cholesterol oxidation products using gas- and high-performance liquid chromatography/mass spectrometry. *Free radical biology & medicine*. 1994; 17:397–409. [PubMed: 7835746]
 15. Babaev VR, Ding L, Reese J, Morrow JD, Breyer MD, Dey SK, Fazio S, Linton MF. Cyclooxygenase-1 deficiency in bone marrow cells increases early atherosclerosis in apolipoprotein e- and low-density lipoprotein receptor-null mice. *Circulation*. 2006; 113:108–117. [PubMed: 16380543]
 16. Brown AJ, Jessup W. Oxysterols and atherosclerosis. *Atherosclerosis*. 1999; 142:1–28. [PubMed: 9920502]
 17. Terasaka N, Yu S, Yvan-Charvet L, Wang N, Mzhavia N, Langlois R, Pagler T, Li R, Welch CL, Goldberg IJ, Tall AR. Abcg1 and hdl protect against endothelial dysfunction in mice fed a high-cholesterol diet. *The Journal of clinical investigation*. 2008; 118:3701–3713. [PubMed: 18924609]
 18. Deckert V, Duverneuil L, Poupon S, Monier S, Le Guern N, Lizard G, Masson D, Lagrost L. The impairment of endothelium-dependent arterial relaxation by 7-ketocholesterol is associated with an early activation of protein kinase c. *British journal of pharmacology*. 2002; 137:655–662. [PubMed: 12381679]
 19. Myoishi M, Hao H, Minamino T, Watanabe K, Nishihira K, Hatakeyama K, Asada Y, Okada K, Ishibashi-Ueda H, Gabbiani G, Bochaton-Piallat ML, Mochizuki N, Kitakaze M. Increased endoplasmic reticulum stress in atherosclerotic plaques associated with acute coronary syndrome. *Circulation*. 2007; 116:1226–1233. [PubMed: 17709641]
 20. Pedruzzi E, Guichard C, Ollivier V, Driss F, Fay M, Prunet C, Marie JC, Pouzet C, Samadi M, Elbim C, O'Dowd Y, Bens M, Vandewalle A, Gougerot-Pocidalo MA, Lizard G, Ogier-Denis E. Nad(p)h oxidase nox-4 mediates 7-ketocholesterol-induced endoplasmic reticulum stress and apoptosis in human aortic smooth muscle cells. *Molecular and cellular biology*. 2004; 24:10703–10717. [PubMed: 15572675]
 21. Ross R. Atherosclerosis—an inflammatory disease. *The New England journal of medicine*. 1999; 340:115–126. [PubMed: 9887164]
 22. Yokode M, Kita T, Kikawa Y, Ogorochi T, Narumiya S, Kawai C. Stimulated arachidonate metabolism during foam cell transformation of mouse peritoneal macrophages with oxidized low density lipoprotein. *The Journal of clinical investigation*. 1988; 81:720–729. [PubMed: 3125226]
 23. Winzerling JJ, Jouni ZE, McNamara DJ. Human peritoneal monocytic cells: Lipoprotein uptake and foam cell formation. *Life sciences*. 1998; 62:501–513. [PubMed: 9464462]
 24. Kunjathoor VV, Febbraio M, Podrez EA, Moore KJ, Andersson L, Koehn S, Rhee JS, Silverstein R, Hoff HF, Freeman MW. Scavenger receptors class a-i/ii and cd36 are the principal receptors

- responsible for the uptake of modified low density lipoprotein leading to lipid loading in macrophages. *The Journal of biological chemistry*. 2002; 277:49982–49988. [PubMed: 12376530]
25. Zeng Y, Tao N, Chung KN, Heuser JE, Lublin DM. Endocytosis of oxidized low density lipoprotein through scavenger receptor cd36 utilizes a lipid raft pathway that does not require caveolin-1. *The Journal of biological chemistry*. 2003; 278:45931–45936. [PubMed: 12947091]
 26. Goldstein JL, Ho YK, Basu SK, Brown MS. Binding site on macrophages that mediates uptake and degradation of acetylated low density lipoprotein, producing massive cholesterol deposition. *Proceedings of the National Academy of Sciences of the United States of America*. 1979; 76:333–337. [PubMed: 218198]
 27. Jones NL, Reagan JW, Willingham MC. The pathogenesis of foam cell formation: Modified ldl stimulates uptake of co-incubated ldl via macropinocytosis. *Arteriosclerosis, thrombosis, and vascular biology*. 2000; 20:773–781.
 28. Hoffmann B, Moebus S, Mohlenkamp S, Stang A, Lehmann N, Dragano N, Schmermund A, Memmesheimer M, Mann K, Erbel R, Jockel KH. Residential exposure to traffic is associated with coronary atherosclerosis. *Circulation*. 2007; 116:489–496. [PubMed: 17638927]
 29. Kunzli N, Jerrett M, Garcia-Esteban R, Basagana X, Beckermann B, Gilliland F, Medina M, Peters J, Hodis HN, Mack WJ. Ambient air pollution and the progression of atherosclerosis in adults. *PloS one*. 2010; 5:e9096. [PubMed: 20161713]
 30. Miller MR, Borthwick SJ, Shaw CA, McLean SG, McClure D, Mills NL, Duffin R, Donaldson K, Megson IL, Hadoke PW, Newby DE. Direct impairment of vascular function by diesel exhaust particulate through reduced bioavailability of endothelium-derived nitric oxide induced by superoxide free radicals. *Environmental health perspectives*. 2009; 117:611–616. [PubMed: 19440501]
 31. Mills NL, Tornqvist H, Gonzalez MC, Vink E, Robinson SD, Soderberg S, Boon NA, Donaldson K, Sandstrom T, Blomberg A, Newby DE. Ischemic and thrombotic effects of dilute diesel-exhaust inhalation in men with coronary heart disease. *The New England journal of medicine*. 2007; 357:1075–1082. [PubMed: 17855668]
 32. Chen LC, Quan C, Hwang JS, Jin X, Li Q, Zhong M, Rajagopalan S, Sun Q. Atherosclerosis lesion progression during inhalation exposure to environmental tobacco smoke: A comparison to concentrated ambient air fine particles exposure. *Inhalation toxicology*. 2010; 22:449–459. [PubMed: 20235771]
 33. Quan C, Sun Q, Lippmann M, Chen LC. Comparative effects of inhaled diesel exhaust and ambient fine particles on inflammation, atherosclerosis, and vascular dysfunction. *Inhalation toxicology*. 2010; 22:738–753. [PubMed: 20462391]
 34. Bai N, Kido T, Suzuki H, Yang G, Kavanagh TJ, Kaufman JD, Rosenfeld ME, van Breemen C, Eeden SF. Changes in atherosclerotic plaques induced by inhalation of diesel exhaust. *Atherosclerosis*. 2011; 216:299–306. [PubMed: 21435644]
 35. Ying Z, Kampfrath T, Thurston G, Farrar B, Lippmann M, Wang A, Sun Q, Chen LC, Rajagopalan S. Ambient particulates alter vascular function through induction of reactive oxygen and nitrogen species. *Toxicological sciences : an official journal of the Society of Toxicology*. 2009; 111:80–88. [PubMed: 19182107]
 36. Febbraio M, Podrez EA, Smith JD, Hajjar DP, Hazen SL, Hoff HF, Sharma K, Silverstein RL. Targeted disruption of the class b scavenger receptor cd36 protects against atherosclerotic lesion development in mice. *The Journal of clinical investigation*. 2000; 105:1049–1056. [PubMed: 10772649]
 37. Robertson S, Colombo ES, Lucas SN, Hall PR, Febbraio M, Paffett ML, Campen MJ. Cd36 mediates endothelial dysfunction downstream of circulating factors induced by o3 exposure. *Toxicological sciences : an official journal of the Society of Toxicology*. 2013; 134:304–311. [PubMed: 23650127]
 38. Lund AK, Lucero J, Harman M, Madden MC, McDonald JD, Seagrave JC, Campen MJ. The oxidized low-density lipoprotein receptor mediates vascular effects of inhaled vehicle emissions. *American journal of respiratory and critical care medicine*. 2011; 184:82–91. [PubMed: 21493736]
 39. Febbraio M, Guy E, Silverstein RL. Stem cell transplantation reveals that absence of macrophage cd36 is protective against atherosclerosis. *Arteriosclerosis, thrombosis, and vascular biology*. 2004; 24:2333–2338.

40. Sheedy FJ, Grebe A, Rayner KJ, Kalantari P, Ramkhalawon B, Carpenter SB, Becker CE, Ediriweera HN, Mullick AE, Golenbock DT, Stuart LM, Latz E, Fitzgerald KA, Moore KJ. Cd36 coordinates nlrp3 inflammasome activation by facilitating intracellular nucleation of soluble ligands into particulate ligands in sterile inflammation. *Nature immunology*. 2013; 14:812–820. [PubMed: 23812099]
41. Moore KJ, Kunjathoor VV, Koehn SL, Manning JJ, Tseng AA, Silver JM, McKee M, Freeman MW. Loss of receptor-mediated lipid uptake via scavenger receptor a or cd36 pathways does not ameliorate atherosclerosis in hyperlipidemic mice. *The Journal of clinical investigation*. 2005; 115:2192–2201. [PubMed: 16075060]
42. Kuchibhotla S, Vanegas D, Kennedy DJ, Guy E, Nimako G, Morton RE, Febbraio M. Absence of cd36 protects against atherosclerosis in apoe knock-out mice with no additional protection provided by absence of scavenger receptor a i/ii. *Cardiovascular research*. 2008; 78:185–196. [PubMed: 18065445]
43. Park YM. Cd36, a scavenger receptor implicated in atherosclerosis. *Experimental & molecular medicine*. 2014; 46:e99. [PubMed: 24903227]
44. Witztum JL. You are right too! *The Journal of clinical investigation*. 2005; 115:2072–2075. [PubMed: 16075051]
45. Stewart CR, Stuart LM, Wilkinson K, van Gils JM, Deng J, Halle A, Rayner KJ, Boyer L, Zhong R, Frazier WA, Lacy-Hulbert A, El Khoury J, Golenbock DT, Moore KJ. Cd36 ligands promote sterile inflammation through assembly of a toll-like receptor 4 and 6 heterodimer. *Nature immunology*. 2010; 11:155–161. [PubMed: 20037584]
46. Navab M, Anantharamaiah GM, Reddy ST, Van Lenten BJ, Ansell BJ, Fonarow GC, Vahabzadeh K, Hama S, Hough G, Kamranpour N, Berliner JA, Lusis AJ, Fogelman AM. The oxidation hypothesis of atherogenesis: The role of oxidized phospholipids and hdl. *Journal of lipid research*. 2004; 45:993–1007. [PubMed: 15060092]
47. Groeneweg M, Vergouwe MN, Scheffer PG, Vermue HP, Sollewijn Gelpke MD, Sijbers AM, Leitinger N, Hofker MH, de Winther MP. Modification of ldl with oxidized 1-palmitoyl-2-arachidonoyl-sn-glycero-3-phosphorylcholine (oxpapc) results in a novel form of minimally modified ldl that modulates gene expression in macrophages. *Biochimica et biophysica acta*. 2008; 1781:336–343. [PubMed: 18534202]
48. Furnkranz A, Schober A, Bochkov VN, Bashtrykov P, Kronke G, Kadl A, Binder BR, Weber C, Leitinger N. Oxidized phospholipids trigger atherogenic inflammation in murine arteries. *Arteriosclerosis, thrombosis, and vascular biology*. 2005; 25:633–638.
49. Gelissen IC, Rye KA, Brown AJ, Dean RT, Jessup W. Oxysterol efflux from macrophage foam cells: The essential role of acceptor phospholipid. *Journal of lipid research*. 1999; 40:1636–1646. [PubMed: 10484610]
50. Engel T, Kannenberg F, Fobker M, Nofer JR, Bode G, Lueken A, Assmann G, Seedorf U. Expression of atp binding cassette-transporter abcg1 prevents cell death by transporting cytotoxic 7beta-hydroxycholesterol. *FEBS letters*. 2007; 581:1673–1680. [PubMed: 17408620]
51. Gelissen IC, Brown AJ, Mander EL, Kritharides L, Dean RT, Jessup W. Sterol efflux is impaired from macrophage foam cells selectively enriched with 7-ketocholesterol. *The Journal of biological chemistry*. 1996; 271:17852–17860. [PubMed: 8663356]
52. Gaus K, Dean RT, Kritharides L, Jessup W. Inhibition of cholesterol efflux by 7-ketocholesterol: Comparison between cells, plasma membrane vesicles, and liposomes as cholesterol donors. *Biochemistry*. 2001; 40:13002–13014. [PubMed: 11669638]
53. Gaus K, Kritharides L, Schmitz G, Boettcher A, Drobnik W, Langmann T, Quinn CM, Death A, Dean RT, Jessup W. Apolipoprotein a-1 interaction with plasma membrane lipid rafts controls cholesterol export from macrophages. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*. 2004; 18:574–576. [PubMed: 14734645]
54. Maor I, Mandel H, Aviram M. Macrophage uptake of oxidized ldl inhibits lysosomal sphingomyelinase, thus causing the accumulation of unesterified cholesterol-sphingomyelin-rich particles in the lysosomes. A possible role for 7-ketocholesterol. *Arteriosclerosis, thrombosis, and vascular biology*. 1995; 15:1378–1387.

Novelty and Significance

What Is Known?

- Exposure to air pollution contributes to cardiovascular morbidity and mortality.
- Oxidized lipids promote atherosclerosis.
- Formation of oxidized lipid in the lung is increased in animals exposed to air pollution.

What New Information Does This Article Contribute?

- In mice, exposure to air pollution increases accumulation of oxidized lipids such as 7-ketocholesterol in macrophages and the aortic wall.
- CD36 links air pollution and abnormal accumulation of oxidized lipid in macrophages and atherosclerotic lesion.
- Deficiency of CD36 reduces ability of macrophage to take up oxidized lipids and it diminishes the effect of air pollution on atherogenesis.

Both epidemiologic and empirical evidence suggest an adverse effect of air pollution on cardiovascular disease. However, the underlying mechanisms remain poorly understood. Previous studies have provided evidence of acceleration of atherosclerosis by air pollution in experimental models. Limited human studies using surrogate markers of subclinical cardiovascular disease such as coronary artery calcium and carotid intima media thickness support these findings. We asked whether air pollution contributes to the formation oxidized lipids in plasma lipoproteins and whether these are deposited in to the arterial wall. We found that the levels of 7-ketocholesterol were increased in plasma lipoproteins in response to air pollution exposure in atherosclerosis-prone mice. These results suggest that air pollution increases CD36-mediated accumulation of oxidized lipids in macrophages and the arterial wall and thereby, promotes atherogenesis. This study provides new insights into mechanisms underlying air pollution-induced atherosclerosis progression by linking air pollution with the formation of oxidized lipids.

Figure 1. Plaque burden increased upon PM exposure

ApoE^{-/-} 2.5 mice were exposed to FA or PM_{2.5} for 6 months and plaque burden was evaluated. A, Representative images of H&E, Masson's trichrome and Oil-Red-O staining indicate plaque area, collagen and lipid content in the aortic sinus of FA- or PM_{2.5}-exposed ApoE^{-/-} mice. B, Statistical analysis indicates the plaque area in the aortic root (600 μm sections starting from aortic valve leaflets) of FA- or PM_{2.5}-exposed mice. C, Bar graph shows collagen content in the aortic sinus of PM_{2.5}- or FA-exposed mice. D, Statistical analysis of Oil-Red-O staining indicates level of lipid content in the sinus of FA- or PM_{2.5}-exposed mice. E, Representative images show Oil-Red-O staining of plaque at the surface of the entire aorta. n=6. Results are shown as mean ± SEM. *, *P*<0.05.

Figure 2. PM_{2.5} exposure increased the accumulation of 7-ketocholesterol (7-KCh) but not cholesterol level in aortic wall

Aorta, lung and plasma were collected from mice exposed to PM_{2.5} or FA.

VLDL/LDL/IDL/HDL fractions were isolated from plasma by fast protein liquid chromatography (FPLC). 7-KCh and cholesterol were detected by liquid chromatography/mass spectrometry (LC/MS). A, 7-KCh level in the aorta of PM_{2.5}-exposed mice. B, 7-KCh level in the lung of PM_{2.5}-exposed mice. C, Cholesterol (CHL) content in aorta in FA- and PM-exposed mice. D, Lung CHL level in FA and PM_{2.5}-exposed mice. E, Ratio of 7-KCh/CHL in the aorta of FA/ PM_{2.5}-exposed mice. F, Ratio of 7-KCh/CHL in the lung of FA/ PM_{2.5}-exposed mice. G, 7-KCh level increased in the VLDL/IDL/LDL fraction of PM_{2.5}-exposed mice. H, The ratio of 7-KCh/CHL in the LDL fraction of FA and PM_{2.5}-exposed mice. n=5/group. *, P<0.05.

Figure 3. Effects of PM exposure on 7-KCh/CHL uptake and efflux

A, 7-KCh uptake in peritoneal exudate macrophages (PEMs): PEMs isolated from mice exposed to FA or PM_{2.5} for 3 months were incubated with ³H-labeled 7-ketocholesterol (1 μCi/ml) for 3 hours. Cells and medium were then harvested for the detection of radioactivity. B, Cholesterol uptake in PEMs: PEMs isolated from mice exposed to FA or PM for 3 months were incubated with H³-labeled cholesterol (1 μCi/ml) for 3 hours with or without the presence of 7-KCh (20 ug/ml). Cells and medium were then harvested for the detection of radioactivity. C, 7-KCh efflux in PEMs: PEMs isolated from mice exposed to

FA or PM_{2.5} were incubated with ³H-labeled 7-ketocholesterol (1 µCi/ml) for 6 hours. After washing out the remaining 7-ketocholesterol, cells were incubated with HDL (50 ug/ml) overnight. Cells and medium were then collected to detect 7-ketocholesterol. D, Cholesterol efflux in PEMs: PEMs isolated from mice exposed to FA or PM_{2.5} were incubated with H³-labeled cholesterol (1 µCi/ml) for 6 hours, followed by 1 hour incubation with 7-KCh or vehicle. After washing out the remaining H³-labeled cholesterol and 7-KCh, cells were incubated with 50 ug/ml HDL overnight. Cells and medium were then collected to detect radioactivity. n=6. *, *P*<0.05.

Figure 4. PM exposure enhances CD36-mediated uptake in macrophages

A, CD36 expression was analyzed on circulating monocytes which were gated based on expression of CD11b and F4/80. B & C, CD36-expressing macrophages in the plaque of PM_{2.5}-exposed mice: sinus sections of FA- or PM_{2.5}-exposed mice were used for the confocal detection of CD36 (Red) and F4/80 (Green). Representative images (B) and statistical analysis (C) were shown. D, BMDMs were incubated with bodipy-labeled 7-ketocholesterol LDL (7-KCh-LDL, 10 µg/ml) for 8 hours. Cells were then harvested and stained with PE-labeled anti-CD36, PE-Cy5-labeled CD11b, and APC-labeled F4/80, followed by analyzing on LSRII flow cytometer. CD36-positive and -negative macrophages were gated for the analysis of bodipy intensity.

Figure 5. Loss of CD36 diminished promotive effect of air pollution on atherosclerosis

A, LDLR^{-/-} mice transplanted with WT or CD36^{-/-} bone marrows were exposed to FA or PM_{2.5} for 3 months. Aortic sinus sections were used for the detection of plaque burden by H&E staining and lipid content detection by Oil-Red-O staining. A, representative image of H&E staining. B, statistical analysis of plaque volume. C, representative images of Oil-Red-O staining. D, statistical analysis of Oil-Red-O positive area. *, $P < 0.05$ when compared with FA; #, $P < 0.05$ when compared with WT.

Figure 6. Loss of CD36 reduces PM_{2.5}-mediated uptake of oxysterol

A - D, Aortic sinus sections from FA- or PM_{2.5}-exposed LDLR^{-/-} mice with WT or CD36^{-/-} bone marrows were used for the detection of 7-KCh (Green) and macrophage (F4/80, Red). Representative images (A) and statistical analysis of 7-KCh area (B), 7-KCh/Plaque ratio (C), and 7-KCh/F4/80 ratio (D) were shown. E, PEMs isolated from FA- or PM_{2.5}-exposed LDLR^{-/-} mice with WT or CD36^{-/-} bone marrows were incubated with DiI-labeled LDL (10 µg/ml), oxLDL (10 µg/ml), or 7-KCh-LDL (10 µg/ml) for 8 hours. Cells were then harvested for the analysis of DiI intensity on LSRII flow cytometer. Flow histograms are shown. n=6. *, P<0.05 when compared with FA; #, P<0.05 when compared with WT.

Figure 7. PM_{2.5} increases foam cell formation in response to 7-ketocholesterol LDL (7-KCh-LDL) and CD36 siRNA transfection can blunt this effect

A, PEMs isolated from WT mice were incubated with Dil-labeled LDL (5 μ g/ml), Dil-labeled oxLDL (5 μ g/ml), or Dil-labeled 7-KCh-LDL (5 μ g/ml) for 36 hours. Cells were then fixed with 4% paraformaldehyde and stained with Oil-red-O. Representative images of Oil-red-O staining and Dil fluorescence imaging were shown. B, PEMs isolated from FA- or PM_{2.5}-exposed LDLR^{-/-} mice with WT or CD36^{-/-} bone marrows were incubated with 7-KCh-LDL (5 μ g/ml) for 36 hours. Cells were then fixed with 4% paraformaldehyde and stained with Oil-red-O. Representative images were shown. C, PEMs isolated from FA- or

PM_{2.5}-exposed LDLR^{-/-} mice with WT or CD36^{-/-} bone marrows were incubated with DiI-labeled 7-KCh-LDL (5 µg/ml) for 36 hours. Cells were then visualized under confocal microscope and representative images of DiI fluorescence were shown. D. Statistical bar graph of % Oil-red-O positive area/cell area for figure A. E, Statistical bar graph of % Oilred-O positive area/cell area for figure B. F, Statistical bar graph of % fluorescence positive area/cell area for figure C.