Published in final edited form as:

Biochemistry. 2005 August 2; 44(30): 10181–10188. doi:10.1021/bi050599l.

The biphasic kinetics of Colchicine-Tubulin interaction : Role of amino acids surrounding the A-ring of bound colchicine molecule

Suvroma Gupta^{1,‡}, Mithu Banerjee^{1,‡}, Asim Poddar¹, Asok Banerjee⁴, Gautam Basu³, Debjani Roy², and Bhabatarak Bhattacharya^{1,*}

¹Department of Biochemistry, Bose Institute, Centenary Campus, Calcutta 700 054, India

²Distributed informatics center, Bose Institute, Centenary Campus, Calcutta 700 054, India

³Department of Biophysics, Bose Institute, Centenary Campus, Calcutta 700 054, India

⁴Department of Biochemistry, The University of Texas Health Science Center, San Antonio, Texas 78229-3900, USA

Abstract

Isotypes of vertebrate tubulin have variable amino acid sequences which are clustered at their C-terminal ends. Isotypes bind colchicine at different on-rates and affinity constants. The kinetics of colchicine binding to purified (unfractionated) brain tubulin have been reported to be biphasic under pseudo-first order conditions. Experiments with individual isotypes established that the presence of β_{III} in the purified tubulin is responsible for the biphasic kinetics. Since the isotypes mainly differ at the C-termini, the colchicine binding kinetics of unfractionated tubulin and the β_{III} isotype, cleaved at the C-termini, have been tested under pseudo-first order conditions. Removal of the C-termini made no difference to the nature of the kinetics. Sequence alignment of different β isotypes of tubulin showed that besides the C-terminal region, there are differences in the main body as well. In order to establish whether these differences lie at the colchicine binding site or not, homology modeling of all β tubulin isotypes was done. We found that the isotypes differed from each other in the amino acids located near the A-ring of colchicine at the colchicine-binding site on β -tubulin. While the β_{III} isotype has two hydrophilic residues (Serine²⁴² and Threonine³¹⁷) both β_{II} and β_{IV} have two hydrophobic residues (Leucine²⁴² and Alanine³¹⁷). β_{II} has Isoleucine at position 318, while β_{III} and β_{IV} have Valine at that position. Thus these alterations in the nature of the amino acids surrounding the colchicine site could be responsible for the different colchicine binding kinetics of the different isotypes of tubulin.

The heterodimeric protein tubulin consists of two similar but distinct polypeptide chains called α and β (1,2). In many organisms, both α and β tubulin exist as numerous isotypes, distinguished by their unique C-terminal sequences (3-9). Thus, tubulin prepared from mammalian brain contains four β -tubulin classes designated as β_I , β_{II} , β_{III} and β_{IV} accounting

*Corresponding author, Communicating Address: Prof. B. Bhattacharyya, Department of Biochemistry, Bose Institute, Centenary Campus, P1/12, CIT Scheme VII M, Calcutta - 700 054, India. FAX: 91-33-2334-3886; TEL: 91-33-2355-0256, Email : bablu@boseinst.ernet.in.

[‡]Both share equal credits

respectively for 3, 58, 25 and 13% of brain tubulin (10). The kinetics of colchicine binding to purified (unfractionated) brain tubulin have been reported to be biphasic under pseudo-first order conditions (11). The origin of the two phases in the binding kinetics was not clear until it was shown that the bovine kidney tubulin, which lacks the β_{III} tubulin isotype (which accounts for 25% of the total brain tubulin) binds colchicine in a monophasic manner (12). In order to know the functional significance of the isotypes, the individual dimer $\alpha\beta_{II}$, $\alpha\beta_{III}$ and $\alpha\beta_{IV}$, have been separated and purified using immunoaffinity columns (10,13). Detailed studies on the colchicine binding properties of these purified isotypes have shed light on the observed biphasic kinetics of the colchicine-tubulin interaction (14). The results of colchicine binding to purified β_{III} and β_{III} depleted isotype fraction clearly demonstrated that the on-rates and the affinity constants for the binding of colchicine to the different tubulin isotypes differ considerably and hence accounts for the origin of the biphasic kinetics observed in unfractionated tubulin (14,15,16).

The β -tubulin isotypes differ from each other apparently at the C-termini ($\beta^{430-444}$). This C-terminal part of the β -subunit, containing a large number of glutamic acid residues, has been found to be flexible, solvent exposed and sensitive to proteolysis (17,18). The carboxy termini of tubulin attract attention for their role in tubulin polymerization. The binding of microtubule associated proteins as well as Ca^{+2} takes place at the C-termini and regulates the self assembly of tubulin (19,20). Colchicine does not bind to the C-termini of tubulin, however, several properties of colchicine-tubulin interactions such as pH-sensitivity, off-rate, on-rate and the stability of the colchicine binding site are regulated by the C-terminal region of α -tubulin (21). Recently, Pal et al. suggested that the negatively charged C-terminus of α -tubulin is involved in ion pair formation with the positively charged residues of the main body of tubulin (22). They termed this interaction as the “Tail-body interaction” which was shown to control many properties of tubulin. Since the β -isotypes also differ mainly at their C-termini, we wanted to find out whether the C-terminus of β -tubulin is involved in a similar type of interaction as is the C-terminus of α -tubulin and thereby affects the colchicine-binding kinetics. Although the C-termini are not a part of the colchicine binding site, it is possible that through this “Tail-body interaction” they can influence the kinetics of the colchicine-tubulin interaction. Thus, it would be interesting to see whether the kinetics of colchicine binding become monophasic with C-termini depleted tubulin.

In the present study, we therefore measured the kinetics of colchicine binding to unfractionated tubulin lacking the C-terminal region of either only β ($\alpha\beta_s$), or both α and β tubulin ($\alpha_s\beta_s$). In this paper, we will refer to these two forms of tubulin as hybrid tubulin and tubulin S, respectively. Surprisingly, we found that like $\alpha\beta$ -tubulin, the colchicine binding to $\alpha\beta_s$ and $\alpha_s\beta_s$ tubulin remained biphasic under pseudo-first order conditions. The C-termini plays no role in controlling the colchicine-tubulin binding kinetics. So our next focus was to compare amino acid sequences at the colchicine binding site on the tubulin main body among different isotypes. Amino acid residues defining the binding site of colchicine on tubulin have been recently identified from the crystal structure of tubulin-colchicine-stathmin-domain complex, solved at 3.5 Å resolution (23). From the crystal structure it has been observed that the colchicine binding site is mostly buried in the intermediate domain of the β subunit while the B-ring side chain of colchicine interacts with the α subunit. Analysis of all the β -tubulin isotype sequences along with the renal β tubulin sequence was performed

using the β subunit of the crystal structure as a template. It has been reported that the colchicine (Figure 1B) where the C-10 methoxy group of ring C has been replaced by a hydrogen atom and MD [2-methoxy-5-(2',4'-dimethoxy phenyl)-2,4,6-cycloheptatriene-1-one], (Figure 1E), an A-ring analogue of AC (Figure 1D) which has a hydrogen in place of 3'-methoxy group of ring A, do not recognize different isotypes (24). So it was also important to analyze the amino acids lying at the binding site of different colchicine analogues having different substituents in the A and C ring of colchicine. To unravel and decipher this mystery, the structure of the colchicine molecule and also the amino acids at the colchicine binding site of tubulin have been scrutinized in detail.

Material and Methods

PIPES, GTP, EGTA, PMSF, colchicine and subtilisin were purchased from Sigma Chemical Co (St. Louis, MO). All other reagents were of analytical grade.

Tubulin preparation and estimation

Tubulin was isolated from goat brain by two cycles of temperature-dependent assembly and disassembly in PEM buffer (50 mM PIPES, 1 mM EGTA, 0.5 mM MgCl_2 , pH 6.9), in the presence of 1 mM GTP, followed by two more cycles in 1M glutamate buffer (25). The purified tubulin, free of MAPs, was stored in aliquots at -70°C . Protein concentration was estimated by the method of Lowry et al (26) using bovine serum albumin as the standard.

Purification of $\alpha\beta_{III}$

$\alpha\beta_{III}$ isotype was purified from bovine brain tubulin according to Banerjee et al (13). Briefly, PC-tubulin was passed successively through two immunoaffinity columns, first through an anti- β_{II} column that removes $\alpha\beta_I$ and $\alpha\beta_{II}$ and followed by a second anti- β_{IV} column that removes $\alpha\beta_{IV}$. Purified $\alpha\beta_{III}$ was concentrated using an Amicon XM 50 ultrafiltration membrane, dialyzed in MES buffer containing 8M glycerol and was frozen in aliquots at -80°C . Before use, the purified isotype was passed through a Sephadex G-25 column equilibrated in PEM buffer to remove the glycerol.

Proteolysis and the preparation of hybrid tubulin ($\alpha\beta_s$) and tubulin S ($\alpha_s\beta_s$)

Tubulin was digested with subtilisin in PEM buffer containing 1 mM GTP (27). Digestion of tubulin with subtilisin was done using a tubulin to enzyme ratio of 100:1 (w/w). The reaction was terminated by the addition of 1% by volume of 1% (w/v) PMSF in DMSO. Digestion at 4°C resulted in the cleavage of the C-terminus of the β subunit only and the product $\alpha\beta_s$ is termed hybrid tubulin, digestion at 30°C cleaved C-terminus of both the subunits resulting in tubulin S ($\alpha_s\beta_s$). The products were subjected to SDS – polyacrylamide slab gel containing 9% acrylamide. The picture of the gel is shown in Figure 2.

Colchicine binding kinetics

The kinetics of colchicine binding to tubulin samples were studied at 37°C under pseudo first order conditions where the drug was present in large excess over tubulin. The samples were excited at 380 nm to reduce the inner filter effect due to the high drug concentration,

and the emission was monitored at 437 nm. All fluorescence values were corrected for the inner filter effect using the equation (28)

$$F_{\text{corr}} = F_{\text{obs}} \cdot \text{antilog} (A_{380} + A_{437}) / 2$$

where F_{obs} and F_{corr} were the observed and the corrected fluorescence values, and A_{380} and A_{437} were the absorbances at the excitation and the emission wavelength, respectively. The biphasic plot was analyzed (29) in terms of two parallel reactions as,

$$F_{\text{max}} - F_t = A \cdot e^{-\alpha t} + B \cdot e^{-\beta t}$$

where A and B were the amplitudes, α and β were the rate constants for the fast and slow phases, respectively. The analysis was performed using non-linear curve-fitting softwares Origin 5.0 and MINSQ (from MicroMath Scientific Software, Salt Lake City, Utah). The apparent on-rate constants were calculated as, $k_{\text{on,app}} = \alpha/c$ where c was the concentration of the drug.

Multiple sequence alignment

Sequences corresponding to the β chain of the crystal structure (PDB ID 1SA0) and kidney β tubulin (GenBank accession no. **P05217**) were extracted from www.rcsb.org/pdb and www.ncbi.nlm.nih.gov respectively. Sequences corresponding to all β tubulin isoforms were collected from literature (7). All sequences were aligned using the multiple sequence alignment programme CLUSTALW (30).

Three dimensional model building

Homology modeling of different isoforms were performed using the β -subunit of the crystal structure as the template using the program MODELLER (31). The stereo chemical quality of the models were analyzed by the program PROCHECK (32) and the corresponding Ramachandran plots (33). The homology modeled isoforms were then subjected to energy minimization using the conjugate gradient method until the final energy derivative reached 0.001 Kcal/mole. Each of the modeled β isoforms was then superimposed separately on the β -subunit of the crystal structure. The RMS deviation of superimposition was in the range of 0.2-0.3 Å for the isoforms. The colchicine binding sites of the superimposed β isoforms were determined from the position of colchicine at the β subunit of the crystal structure.

Tubulin residues lying within 5 Å and 8 Å from the center of mass of the colchicine molecule were studied.

Results and Discussion

Biphasic kinetics of colchicine binding to tubulin and the role of its C-termini

The C-termini of tubulin are not only responsible for distinguishing the isoforms of tubulin but also profoundly influence other properties like polymerization (19,34,35), colchicine

binding (21), chaperone activity (36), bis-ANS binding etc (37). There are reports that the presence of the β_{III} isotype gives rise to the biphasic binding kinetics of colchicine-tubulin interaction since the removal of this isotype produces monophasic binding curve. Different combinations of tubulin isotypes namely, ($\alpha\beta_{II} + \alpha\beta_{IV}$) exhibit monophasic pattern whereas biphasic kinetics were observed when mixture of isotypes, i.e. ($\alpha\beta_{III} + \alpha\beta_{II}$) or ($\alpha\beta_{III} + \alpha\beta_{IV}$), were tested (14). The kinetics of colchicine binding with $\alpha\beta$, $\alpha\beta_s$ and $\alpha_s\beta_s$ tubulin was measured at 37°C under pseudo first order conditions using colchicine fluorescence at 430 nm. The tubulin concentration was 3 μ M and the colchicine concentration was 50 μ M in all cases. The kinetics of colchicine binding remained biphasic with $\alpha\beta_s$ tubulin (Figure 3). The apparent on-rate constants were 98.1 $M^{-1}s^{-1}$ for the fast phase and 26.3 $M^{-1}s^{-1}$ for the slower phase (Table 1). This biphasic nature of colchicine binding persists even when the C-terminus of both α and β subunits were cleaved ($\alpha_s\beta_s$). The on-rate constants were 118 $M^{-1}s^{-1}$ and 26.5 $M^{-1}s^{-1}$ for the fast and slow phases respectively. Since it has been previously demonstrated that the slow phase in the binding kinetics is due to the presence of the $\alpha\beta_{III}$ isotype (14), we studied the effect of subtilisin cleavage on the binding kinetics for purified $\alpha\beta_{III}$ isotype. As shown in Figure 4, the kinetics of colchicine binding for $\alpha\beta_{III}$ isotype does not change significantly upon removal of the C-terminus from β_{III} . The on-rate constants were 9.8 $M^{-1}s^{-1}$ and 9.3 $M^{-1}s^{-1}$ for native and subtilisin cleaved $\alpha\beta_{III}$ respectively. These results clearly indicate that the C-terminus of β -tubulin doesn't play any role in the interaction of colchicine with tubulin and that different tubulin isotypes must differ in other part (s) of the tubulin molecule. From the above observations it can be concluded that kinetics of the tubulin-colchicine interaction is not a “tail” specific phenomenon of the β -tubulin isotype.

Mapping the Colchicine binding site

Our experiments with C-termini cleaved tubulin showed biphasic kinetics similar to native tubulin (uncleaved) so it became clear that the C-termini, which lies far away from the colchicine binding site plays no role in regulating colchicine tubulin binding kinetics. From a sequence alignment of the β -subunit of the crystal structure and the β isotypes of tubulin (Figure 5), it has been found that the isotypes differ in many other regions of the sequence besides the C-terminal end. The amino acids differing in the case of the β isotypes and the β subunit of crystal structure have been marked bold. Amino acids lying within 5 Å from the bound colchicine molecule, and which differs among different isotypes and the β subunit of the crystal structure, have been marked bold as well as underlined in Figure 5.

In order to study whether these changes occur at the colchicine binding site or not, the three dimensional models of individual β isotypes (β_I , β_{II} , β_{III} and β_{IV}) were constructed using homology modeling. The result is presented in Table 2.

Distinguishing isotypes on the basis of residues surrounding the bound colchicine molecule in tubulin

It can be noted from Table 2 that amino acid residues lying within 5 Å and 8 Å from the colchicine molecule, in the crystal structure, are identical for β_I and β_{IV} . Isotypes β_{II} and β_{IV} differ at only position 318 : Isoleucine³¹⁸ for β_{II} while Valine³¹⁸ for β_{IV} . However, isotype β_{III} has three changes: Serine²⁴², Threonine³¹⁷, Valine³⁵³ in place of Leucine²⁴²,

Alanine³¹⁷, Threonine³⁵³ for β_{IV} . In spite of having a single difference in amino acid residue a combination of ($\alpha\beta_{II} + \alpha\beta_{IV}$) tubulin exhibits monophasic kinetics for colchicine tubulin interaction. Addition of $\alpha\beta_{III}$ isotype to either $\alpha\beta_{II}$ or $\alpha\beta_{IV}$ to form ($\alpha\beta_{III} + \alpha\beta_{II}$) or ($\alpha\beta_{III} + \alpha\beta_{IV}$) changes the binding kinetics from monophasic to biphasic (14) (with appearance of the slow phase of binding). Thus it seems that $\alpha\beta_{III}$ is responsible for the biphasic kinetic pattern. From Table 2 we see that $\alpha\beta_{III}$ has a much wider variation in the nature of amino acid residues (two within 5 Å and in addition one within 8 Å) at the colchicine binding site compared to that of $\alpha\beta_{II}$ and $\alpha\beta_{IV}$. These differences in nature of amino acids may explain the difference in the association rates as well as the affinity constant values for the $\alpha\beta_{III}$ -colchicine interaction.

Previously, an analysis was done to define a part of the colchicine binding site by correlating the experimentally determined association constants with some available primary sequences (38). Primary β -tubulin sequences of some organisms with known binding affinity constants were compared and it was suggested that the relative affinities of different tubulins for colchicine depend upon residues in the immediate vicinity of β^{316} . It has been reported from a comparison of residues with the experimental association constants that tubulin with Isoleucine³¹⁶ in β_I (38) must bind colchicine significantly more weakly than with Valine³¹⁶ in β_{II} (38), but more strongly than Methionine³¹⁶ in *Caenorhabditis elegans* or Phenylalanine³¹⁶ in *Saccharomyces cerevisiae* system (38). An examination of the residues lying at the colchicine binding site for $\alpha\beta_{III}$, $\alpha\beta_{II}$ and $\alpha\beta_{IV}$ revealed that there are differences around β^{316} . Besides this there are also differences in other regions surrounding the colchicine molecule. β_{II} has an Isoleucine residue at position 318 whereas β_{IV} has a Valine residue at the same position. The colchicine binding affinity constant for $\alpha\beta_{IV}$ is about fourteen fold greater than that of $\alpha\beta_{II}$ whereas the corresponding rate constant for $\alpha\beta_{IV}$ is twice that of $\alpha\beta_{II}$. Indeed the side chains of Isoleucine³¹⁸ and Valine³¹⁸ differ only in the presence or absence of an ethyl group. It is clear from Figure 6A, that the distance between the carbon of 3-methoxy group of ring A and the side chain methyl carbon of Valine³¹⁸ in β_{IV} is 4.21 Å. On the other hand the distance between the hydrogen of 3-methoxy group of ring A and the side chain ethyl carbon of Isoleucine³¹⁸ in β_{II} is 3.62 Å (Figure 6B). The ethyl group, being a bulkier group than a methyl group, can enter into van der Waals repulsive interaction with the 3-methoxy group of ring A of colchicine which results in the lowering of affinity constant and rate constant of β_{II} relative to β_{IV} . The affinity constant for colchicine binding for $\alpha\beta_{IV}$ is about 28 fold higher than that of $\alpha\beta_{III}$ and the rate constant is about eight times greater. When β_{III} is compared to β_{IV} , there are two changes within 5 Å of the colchicine molecule. Leucine²⁴² and Alanine³¹⁷ of $\alpha\beta_{IV}$ have been replaced by Serine²⁴² and Threonine³¹⁷ respectively in $\alpha\beta_{III}$ (Figure 6C). Thus we find that there is a shift from a hydrophobic to a hydrophilic environment. Since there are evidences that colchicine binding involves hydrophobic interaction we can conclude that these two alterations lowers the K_a (affinity constant) of colchicine binding as well as the on-rate constant of $\alpha\beta_{III}$ relative to $\alpha\beta_{IV}$. In case of kidney β tubulin, which lacks the β_{III} isotype (12), the apparent on-rate constant of binding is very close to that of the faster binding component ($\alpha\beta_{IV}$) of brain tubulin.

When we compare colchicine binding kinetics of $\alpha\beta_{II}$ and $\alpha\beta_{III}$ we find that the rate constant of $\alpha\beta_{II}$ is about four times that of $\alpha\beta_{III}$, whereas the corresponding affinity constant for colchicine binding is about two times greater. Isoleucine³¹⁸, Leucine²⁴², Alanine³¹⁷ in $\alpha\beta_{II}$ have been replaced by Valine³¹⁸, Serine²⁴², Threonine³¹⁷ in $\alpha\beta_{III}$. Thus the hydrophobic residues (Leucine and Alanine) in the case of isotype $\alpha\beta_{II}$ have been replaced by hydrophilic residues (Serine and Threonine) in case of $\alpha\beta_{III}$. This difference in the nature of residues might be responsible for the differential binding kinetics as well as the difference in affinity constant between β_{II} and β_{III} . Therefore, the variation in the nature of residues encompassing the colchicine molecule influences the kinetics and affinity constant of the colchicine-tubulin interaction. Interestingly we found that the variation in nature of residues (in a proximal region of 5 Å) lies around the A-ring of colchicine only, as discussed later.

Effect of colchicine structure on kinetics of colchicine-tubulin interaction

The kinetics of colchicine-tubulin interaction is mainly controlled by the B-ring and the C-7 side chain of colchicine. Thus, AC (Figure 1D; having only the A and C-ring of colchicine) binding to tubulin is instantaneous and reversible whereas colchicine binding is slow and poorly reversible (39, 40). However, the biphasic kinetics of colchicine binding to unfractionated tubulin indicates that the rate of the colchicine-tubulin interaction is also controlled at the protein level. The binding of desacetamidocolchicine (DAAC) (Figure 1C), to tubulin follows biphasic kinetics similar to those of colchicine. Moreover like colchicine the affinity constant of DAAC for $\alpha\beta_{III}$ is much less than that for $\alpha\beta_{II}$ and $\alpha\beta_{IV}$ (15). Therefore, it is obvious that the B-ring substituent does not distinguish different β isotypes. Again there is a report that AC [2-methoxy-5-(2', 3', 4'-trimethoxyphenyl) tropone] (Figure 1D), which is devoid of the B-ring, also exhibits biphasic kinetics in its binding to tubulin (16). So at this point it is clear that the kinetics of binding of colchicine to tubulin may be a feature of rings A and C. Indeed, there are reports that colchicine analogues with modified C-ring such as colchicide (Figure 1B) and MD [2-methoxy-5-(2',4'-dimethoxy phenyl)-2,4,6-cycloheptatriene-1-one] (Figure 1E) with a modified A-ring do not recognize tubulin isotypes (24,41). Colchicide has a modified C-ring, where the C-10 methoxy group of the ring C has been replaced by a hydrogen atom. This analogue does not show any fluorescence upon tubulin binding nor can it distinguish kinetically among different tubulin isotypes (24). MD, a fast binding colchicine analogue like AC (without B-ring), possesses a modified A-ring, where 3' methoxy group in A-ring is replaced by a hydrogen (41). Like colchicide, MD is unable to differentiate among tubulin isotypes. One thing is common to both analogues that the replacement of a bulky methoxy group by a hydrogen atom abolished completely their ability to recognize different tubulin isotypes. So, may be the presence of a methoxy group is the crucial determinant for this type of recognition. Since a larger methoxy group is replaced by a smaller hydrogen atom, methoxy group of those analogues may play some role in generating biphasic kinetics of the colchicine-tubulin interaction. From Figure 6B it appears that the 3-methoxy group of ring A of colchicine suffers steric repulsion with residue Isoleucine³¹⁸ in β_{II} . Such interaction is absent in the case of β_{IV} (Figure 6A) as well as in the case of β_{III} (Figure 6C) since both of them have a Valine residue at the same position and Isoleucine happens to be much bulkier than Valine.

Acknowledgments

We are grateful to Prof. Richard F. Ludueña of the Department of Biochemistry, University of Texas Health Science Center, USA and Dr. Pradeep Parrack of the Biochemistry Department of Bose Institute, Calcutta, India for their critical reading of the manuscript.

The financial support from CSIR, Government of India is thankfully acknowledged. Additional support was provided by grants to A. Banerjee (National Institutes of Health CA59711 and US Army Breast Cancer Research Program DAMD17-98-1-8244) and to R. F. Ludueña (Welch Foundation AQ-0726 and US Army Breast Cancer Research Program DAMD17-01-1-0411).

References

1. Olmsted JB, Borisy GG. Microtubules. *Annu Rev Biochem.* 1973; 42:507–540. [PubMed: 4581232]
2. Dustin P. Microtubules. *Sci Am.* 1980; 243:66–76. [PubMed: 7423178]
3. Villasante A, Wang D, Dobner P, Dolph P, Lewis SA, Cowan NJ. Six mouse alpha-tubulin mRNAs encode five distinct isotypes: testis-specific expression of two sister genes. *Mol Cell Biol.* 1986; 6:2409–2419. [PubMed: 3785200]
4. Sullivan KF, Cleveland DW. Identification of conserved isotype-defining variable region sequences for four vertebrate beta tubulin polypeptide classes. *Proc Natl Acad Sci U S A.* 1986; 83:4327–4331. [PubMed: 3459176]
5. Wang D, Villasante A, Lewis SA, Cowan NJ. The mammalian beta-tubulin repertoire: hematopoietic expression of a novel, heterologous beta-tubulin isotype. *J Cell Biol.* 1986; 103:1903–1910. [PubMed: 3782288]
6. Monteiro MJ, Cleveland DW. Sequence of chicken c beta 7 tubulin. Analysis of a complete set of vertebrate beta-tubulin isotypes. *J Mol Biol.* 1988; 199:439–446. [PubMed: 3351937]
7. Sullivan KF. Structure and utilization of tubulin isotypes. *Annu Rev Cell Biol.* 1988; 4:687–716. [PubMed: 3058169]
8. Luduena RF. Are tubulin isotypes functionally significant. *Mol Biol Cell.* 1993;445–457. [PubMed: 8334301]
9. Lewis SA, Wang DH, Cowan NJ. Microtubule-associated protein MAP2 shares a microtubule binding motif with tau protein. *Science.* 1988; 242:936–939. [PubMed: 3142041]
10. Banerjee A, Roach MC, Wall KA, Lopata MA, Cleveland DW, Luduena RF. A monoclonal antibody against the type II isotype of beta-tubulin. Preparation of isotypically altered tubulin. *J Biol Chem.* 1988; 263:3029–3034. [PubMed: 3277964]
11. Garland DL. Kinetics and mechanism of colchicine binding to tubulin: evidence for ligand-induced conformational change. *Biochemistry.* 1978; 17:4266–4272. [PubMed: 708711]
12. Banerjee A, Luduena RF. Kinetics of association and dissociation of colchicine-tubulin complex from brain and renal tubulin. Evidence for the existence of multiple isotypes of tubulin in brain with differential affinity to colchicine. *FEBS Lett.* 1987; 219:103–107. [PubMed: 3595871]
13. Banerjee A, Roach MC, Trcka P, Luduena RF. Preparation of a monoclonal antibody specific for the class IV isotype of beta-tubulin. Purification and assembly of alpha beta II, alpha beta III, and alpha beta IV tubulin dimers from bovine brain. *J Biol Chem.* 1992; 267:5625–5630. [PubMed: 1544937]
14. Banerjee A, Luduena RF. Kinetics of Colchicine Binding to Purified β -Tubulin Isotypes from Bovine Brain. *J Biol Chem.* 1992; 267:13335–13339. [PubMed: 1618835]
15. Banerjee A, D'Hoore A, Engelborghs Y. Interaction of desacetamidocolchicine, a fast binding analogue of colchicine with isotypically pure tubulin dimers alpha beta II, alpha beta III, and alpha beta IV. *J Biol Chem.* 1994; 269:10324–10329. [PubMed: 8144613]
16. Banerjee A, Engelborghs Y, D'Hoore A, Fitzgerald TJ. Interactions of a bicyclic analog of colchicine with beta-tubulin isoforms alphabeta (II), alphabeta (III) and alphabeta (IV). *Eur J Biochem.* 1997; 246:420–424. [PubMed: 9208933]
17. Ponstingl H, Little M, Krauhs E, Kempf T. Carboxy-terminal amino acid sequence of alpha-tubulin from porcine brain. *Nature.* 1979; 282:423–425. [PubMed: 574193]

18. Sackett DL, Bhattacharyya B, Wolff J. Tubulin subunit carboxyl termini determine polymerization efficiency. *J Biol Chem.* 1985; 260:43–45. [PubMed: 3965457]
19. Serrano L, de la Torre J, Maccioni RB, Avila J. Involvement of the carboxyl-terminal domain of tubulin in the regulation of its assembly. *Proc Natl Acad Sci U S A.* 1984; 81:5989–5993. [PubMed: 6385005]
20. Littauer UZ, Givon D, Thierauf M, Ginzburg I, Ponstingl H. Common and distinct tubulin binding sites for microtubule-associated proteins. *Proc Natl Acad Sci U S A.* 1986; 83:7162–7166. [PubMed: 3463956]
21. Mukhopadhyay K, Parrack PK, Bhattacharyya B. The carboxy terminus of the alpha subunit of tubulin regulates its interaction with colchicine. *Biochemistry.* 1990; 29:6845–6850. [PubMed: 2204421]
22. Debnath P, Mahapatra P, Manna T, Chakrabarti P, Bhattacharya B, Banerjee A, Basu G, Roy S. Conformational Properties of α -Tubulin Tail Peptide: Implications for Tail-Body Interaction. *Biochemistry.* 2001; 40:15512–15519. [PubMed: 11747426]
23. Ravelli RB, Gigant B, Curmi PA, Jourdain I, Lachkar S, Sobel A, Knossow M. Insight into tubulin regulation from a complex with colchicine and a stathmin-like domain. *Nature.* 2004; 428:198–202. [PubMed: 15014504]
24. Dumortier C, Yan Q, Bane S, Engelborghs Y. Mechanism of tubulin-colchicine recognition: a kinetic study of the binding of the colchicine analogues colchicine and isocolchicine. *Biochem J.* 1997; 327:685–688. [PubMed: 9581543]
25. Hamel E, del Campo AA, Lowe MC, Lin CM. Interactions of taxol, microtubule-associated proteins and guanine nucleotides in tubulin polymerization. *J Biol Chem.* 1981; 256:11887–11894. [PubMed: 6117556]
26. Lowry OH, Rosenbrough NJ, Farr AA, Randall RJ. Protein measurement with the Folin phenol reagent. *J Biol Chem.* 1951; 193:265–275. [PubMed: 14907713]
27. Bhattacharyya B, Sackett DL, Wolff J. Tubulin, hybrid dimers, and tubulin S. Stepwise charge reduction and polymerization. *J Biol Chem.* 1985; 260:10208–10216. [PubMed: 3894367]
28. Lakowicz, JR. Principles of Fluorescence Spectroscopy. Plenum Press; New York and London: 1983. Instrumentation for fluorescence spectroscopy; p. 19-49.
29. Lambeir A, Engelborghs Y. A fluorescence stopped flow study of colchicine binding to tubulin. *J Biol Chem.* 1981; 256:3279–3282. [PubMed: 7204403]
30. Thompson JD, Higgins DG, Gibson TJ. The CLUSTAL _X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* 1994; 22:4673–4680. [PubMed: 7984417]
31. Sali A, Blundell TL. Comparative protein modelling by satisfaction of spatial restraints. *J Mol Biol.* 1993; 234:779–815. [PubMed: 8254673]
32. Laskowski RA, MacArthur MW, Moss DS, Thornton JM. PROCHECK: A program to check the stereochemical quality of protein structure. *J Appl Crystallogr.* 1993; 26:283–291.
33. Sheik SS, Sundarajan P, Hussain AS, Sekar K. Ramachandran plot on the web. *Bioinformatics.* 2002; 18:1548–1549. [PubMed: 12424132]
34. Bhattacharyya B, Sackett DL, Wolff J. Tubulin, hybrid dimers, and tubulin S. Stepwise charge reduction and polymerization. *J Biol Chem.* 1985; 260:10208–10216. [PubMed: 3894367]
35. Rai SS, Wolff J. The C terminus of beta-tubulin regulates vinblastine-induced tubulin polymerization. *Proc Natl Acad Sci U S A.* 1998; 95:4253–4257. [PubMed: 9539723]
36. Guha S, Manna TK, Das KP, Bhattacharyya B. Chaperone-like activity of tubulin. *J Biol Chem.* 1998; 273:30077–30080. [PubMed: 9804759]
37. Horowitz P, Prasad V, Luduena RF. Bis(1,8-anilinonaphthalenesulfonate). A novel and potent inhibitor of microtubule assembly. *J Biol Chem.* 1984; 259:14647–14650. [PubMed: 6548750]
38. Burns GR. Analysis of the colchicine-binding site of β -tubulin. *FEBS Lett.* 1992; 297:205–208. [PubMed: 1544399]
39. Bane S, Puett D, Macdonald TL, Williams RC. Binding to tubulin of the colchicine analog 2-methoxy-5-(2', 3', 4'-trimethoxyphenyl) tropone. Thermodynamic and kinetic aspects. *J Biol Chem.* 1984; 259:7391–7398. [PubMed: 6736010]

40. Ray K, Bhattacharyya B, Biswas BB. Role of B-ring of colchicine in its binding to tubulin. *J Biol Chem.* 1981; 256:6241–6244. [PubMed: 7240202]
41. Dumortier C, Potenziano JL, Bane S, Engelborghs Y. The mechanism of tubulin-colchicine recognition--a kinetic study of the binding of a bicyclic colchicine analogue with a minor modification of the A ring. *Eur J Biochem.* 1997; 249:265–269. [PubMed: 9363778]

Abbreviations used

C-termini	carboxy termini
β_I	beta isotype I
β_{II}	beta isotype II
β_{III}	beta isotype III
β_{IV}	beta isotype IV

FIGURE 1. Structure of drugs

(**A**) Colchicine, (**B**) Colchicide, (**C**) DAAC (desacetamido colchicine), (**D**) 2-methoxy-5-(2', 3', 4'-trimethoxyphenyl) tropone (AC) and (**E**) 2-methoxy-5-(2',4'-dimethoxy phenyl)-2,4,6-cycloheptatriene-1-one (MD).

Lane.	1	2	3
Sample	$\alpha\beta$	$\alpha\beta_s$	$\alpha_s\beta_s$

FIGURE 2. Subtilisin digestion of tubulin

Lane1, undigested tubulin ($\alpha\beta$); lane 2, hybrid tubulin obtained by enzymatic digestion of tubulin at 4°C ($\alpha\beta_s$) ; lane 3, tubulin S obtained by the digestion of tubulin at 30°C ($\alpha_s\beta_s$).

FIGURE 3. Kinetics of colchicine binding to $\alpha\beta$ -tubulin

3 μ M tubulin (mixture of all isotypes) was incubated with 50 μ M colchicine at 37 $^{\circ}$ C and the binding was monitored fluorometrically at 430 nm. The samples were excited at 380 nm to reduce the inner filter effect due to the high drug concentration, and the emission was monitored at 437 nm. Panel **A**, $\alpha\beta$; panel **B**, $\alpha\beta_s$ shows the plot of $(F_m - F_t)$ versus time. Inset in each panel shows the plot of $\ln(F_m - F_t)$ versus time.

FIGURE 4. Kinetics of colchicine binding to purified β_{III} tubulin isotype
 3 μ M purified isotype $\alpha\beta_{III}$ tubulin was incubated with 50 μ M colchicine at 37°C and the binding was monitored fluorometrically at 430 nm. Panel **A**, $\alpha\beta_{III}$ isotype, panel **B**, subtilisin digested $\alpha\beta_{III}$ isotype shows the plot of $(F_m - F_t)$ versus time. Inset in each panel shows the plot of $\ln(F_m - F_t)$ versus time.

1SA0B	MREIVHIQAGQCGNQIGAKFWEVISDEHGIDPTGSYHGSDSLQLERINVYNEA A GNKYV	60
β_1	MREIVHIQAGQCGNQIGAKFWEVISDEHGIDPTGTYHGSDSLQLDRISVYNEA T GGKYV	60
β_{11}	MREIVHIQAGQCGNQIGAKFWEVISDEHGIDPTGSYHGSDSLQLERINVYNEA A GNKYV	60
β_{111}	MREIVHIQAGQCGNQIGAKFWEVISDEHGIDP S GNVVGSDSLQLERINVYNEA S SHKYV	60
β_{1v}	MREIVHIQAGQCGNQIGAKFWEVISDEHGIDPTGTYHGSDSLQLERINVYNEA T GGKYV	60
β_{kidney}	MREIVHLQAGQCGNQIGAKFWEVISDEHGIDPTGTYHGSDSLQLERINVYNEA T GGKYV	60

1SA0B	PRAILVDLEPGTMDSVRS G PFGQIFRPDNFVFGQSGAGNNWAKGHYTEGAELVDSVLDV V	120
β_1	PRAILVDLEPGTMDSVRS G PFGQIFRPDNFVFGQSGAGNNWAKGHYTEGAELVDSVLD - V	119
β_{11}	PRAILVDLEPGTMDSVRS G PFGQIFRPDNFVFGQSGAGNNWAKGHYTEGAELVDSVLD - V	119
β_{111}	PRAILVDLEPGTMDSVRS A F G H L FRPDNFI F GQSGAGNNWAKGHYTEGAELVDSVLD - V	119
β_{1v}	PRAVLVDLEPGTMDSVRS G PFGQIFRPDNFVFGQSGAGNNWAKGHYTEGAELVDSVLD - V	119
β_{kidney}	PRAVLVDLEPGTMDSVRS G PFGQIFRPDNFVFGQSGAGNNWAKGHYTEGAELVDSVLDV V	120

1SA0B	RKE S ESCDCLQGFLTHSLGGGTSGMGTL L LISKIREEYPDRIMNTFSV V SPKVS D TVV	180
β_1	RKE A ESCDCLQGFLTHSLGGGTSGMGTL L LISKIREEYPDRIMNTFSV V SPKVS D TVV	179
β_{11}	RKE A ESCDCLQGFLTHSLGGGTSGMGTL L LISKIREEYPDRIMNTFSV M SPKVS D TVV	179
β_{111}	RKE C ENCDCLQGFLTHSLGGGTSGMGTL L LISKVREEYPDRIMNTFSV V SPKVS D TVV	179
β_{1v}	RKE A ESCDCLQGFLTHSLGGGTSGMGTL L LISKIREEYPDRIMNTFSV V SPKVS D TVV	179
β_{kidney}	RKE A ESCDCLQGFLTHSLGGGTSGMGTL L LISKIREEYPDRIMNTFSV V SPKVS D TVV	180

1SA0B	EPYNATLS V HQLVENTDETYCIDNEALYDICFRTLKL T PTYGDLNHLVS A TMSGV T T C L	240
β_1	EPYNATLS V HQLVENTDETYCIDNEALYDICFRTLKL T PTYGDLNHLVS G TMSGV T T C L	239
β_{11}	EPYNATLS V HQLVENTDETY S IDNEALYDICFRTLKL T PTYGDLNHLVS A TMSGV T T C L	239
β_{111}	EPYNATLS I HQLVENTDETYCIDNEALYDICFRTLKL A TPTYGDLNHLVS A TMSGV T T C S	239
β_{1v}	EPYNATLS V HQLVENTDETYCIDNEALYDICFRTLKL T PTYGDLNHLVS G TMSGV T T C L	239
β_{kidney}	EPYNATLS V HQLVENTDETYCIDNEALYDICFRTLKL T PTYGDLNHLVS A TMSGV T T C L	240

1SA0B	RFPGQLNADLRKLAVNMVFPRLHFFMPGFAPLTSRGSQQYRALTVPELTQQMF D AK N MM	300
β_1	RFPGQLNADLRKLAVNMVFPRLHFFMPGFAPLTSRGSQQYRALTV P DLTQQV F DA K NMM	299
β_{11}	RFPGQLNADLRKLAVNMVFPRLHFFMPGFAPLTSRGSQQYRALTVPELTQQMF D SK N MM	299
β_{111}	RFPGQLNADLRKLAVNMVFPRLHFFMPGFAPL T ARGSQYRALTVPELTQQMF D AK N MM	299
β_{1v}	RFPGQLNADLRKLAVNMVFPRLHFFMPGFAPLTSRGSQQYRALTVPELTQQMF D AK N MM	299
β_{kidney}	RFPGQLNADLRKLAVNMVFPRLHFFMPGFAPLTSRGSQQYRALTVPELTQQMF D AK N MM	300

1SA0B	AACDPRHGRYLTV A A VFRGRMSMKEVDEQ M L N VQ N KNSSYFVEWIPNNVKT A VCDIPPRG	360
β_1	AACDPRHGRYLTV A A VFRGRMSMKEVDEQ M L N VQ N KNSSYFVEWIPNNVKT A VCDIPPRG	359
β_{11}	AACDPRHGRYLTV A A I F RGRMSMKEVDEQ M L N VQ N KNSSYFVEWIPNNVKT A VCDIPPRG	359
β_{111}	AACDPRHGRYLTV A T VFRGRMSMKEVDEQ M L A I Q SKNSSYFVEWIPNNV K VAVCDIPPRG	359
β_{1v}	AACDPRHGRYLTV A A VFRGRMSMKEVDEQ M L N VQ N KNSSYFVEWIPNNVKT A VCDIPPRG	359
β_{kidney}	AACDPRHGRYLTV A A VFRGRMSMKEVDEQ M L N VQ N KNSSYFVEWIPNNVKT A VCDIPPRG	360

1SA0B	LKM- S ATFIGNSTAIQELFKRISEQFTAMFRKAF L HWTGEGMDEMEFTEAESNMNDLV	419
β_I	LKM- A VTFIGNSTAIQELFKRISEQFTAMFRKAF L HWTGEGMDEMEFTEAESNMNDLV	418
β_{II}	LKM- S ATFIGNSTAIQELFKRISEQFTAMFRKAF L HWTGEGMDEMEFTEAESNMNDLV	418
β_{III}	LKM- S STFIGNSTAIQELFKRISEQFTAMFRKAF L HWTGEGMDEMEFTEAESNMNDLV	418
β_{IV}	LKMS A VTFIGNSTAIQELFKRISEQFTAMFRKAF L HWTGEGMDEMEFTEAESNMNDLV	419
β_{kidney}	LKM- S ATFIGNSTAIQELFKRISEQFTAMFRKAF L HWTGEGMDEMEFTEAESNMNDLV	419

1SA0B	SEYQYQDATADE QGEFEEGEDEA -----	445
β_I	SEYQYQDATA EEEDFGEAE EA-----	443
β_{II}	SEYQYQDATADE QGEFEEGEDEA -----	444
β_{III}	SEYQYQDATA EEEGEMYEDDEESESQGPK	449
β_{IV}	SEYQYQDATA EEEGEFEEAE EEVA-----	445
β_{kidney}	SEYQYQDATA EEEGEFEEAE EEVA-----	445

FIGURE 5. Sequence alignment of β isotypes of tubulin

1SA0B: Tubulin β subunit of the tubulin-colchicine-stathmin-domain complex (crystal structure); β_I : vertebrate beta isotype I; β_{II} : vertebrate beta isotype II; β_{III} : vertebrate beta isotype III; β_{IV} : vertebrate isotype IV; β_{kidney} : kidney beta tubulin sequence (Gen Bank accession no. P05217). The amino acids differing in case of β isotypes and the β subunit of the crystal structure have been marked bold. The amino acids within 5 Å distance from the bound colchicine molecule which are different among different isotypes and the β subunit of the crystal structure have been marked bold as well as underlined.

FIGURE 6. Residues of different β -tubulin isotypes lying in close proximity of colchicine molecule

The colchicine molecule occupies the upper right hand side in each panel. The numbers over the dotted line indicate the corresponding distance in Å between the atoms of the colchicine molecule and the nearby amino acid residues. Panel (A) shows residues of β_{IV} denoted by B353, B317, B318 and B242, panel (B) shows the residues of β_{II} denoted by B353, B317, B318 and B242, and panel (C) shows the residues of β_{III} denoted by B353, B317, B318 and B242.

Table 1
Comparison of the colchicine-binding parameters for tubulin, digested tubulin and purified $\alpha\beta_{III}$

Different types of tubulin	Kinetic Pattern	k_{on} (fast) $M^{-1}s^{-1}$	k_{on} (slow) $M^{-1}s^{-1}$
$\alpha\beta$ tubulin	biphasic	78.1	16.03
$\alpha\beta_s$	biphasic	98.12	26.3
$\alpha_s\beta_s$	biphasic	118	26.5
$\alpha\beta_{III}$	monophasic	9.8	-
Subtilisin digested $^*\alpha\beta_{(s)III}$	monophasic	9.3	-

* $\alpha\beta_{(s)III}$ indicates β -C-terminus digested isotype $\alpha\beta_{III}$

Table 2
Residues of different β -Tubulin isotypes lying in close proximity to colchicine molecule

Crystal structure ^I	Isotype 1 ² β_I	Isotype 2 ² β_{II}	Isotype 3 ² β_{III}	Isotype 4 ² β_{IV}
GLN (133)				
ASN (167)				
GLU (200)				
TYR (202)				
GLY (237)				
VAL (238)				
THR (239)				
THR (240)				
CYS (241)				
LEU (242)			SER	
ARG (243)				
GLN (247)				
LEU (248)				
ASN (249)				
ALA (250)				
ASP (251)				
LEU (252)				
ARG (253)				
LYS (254)				
LEU (255)				
ALA (256)				
VAL (257)				
ASN (258)				
MET (259)				
VAL (260)				
PHE (268)				
LEU (313)				
THR (314)				
VAL (315)				
ALA (316)				
ALA (317)			THR	
VAL (318)		ILE		
ILE (347)				
PRO (348)				
ASN (349)				
ASN (350)				

Crystal structure ¹	Isotype 1 ² β_I	Isotype 2 ² β_{II}	Isotype 3 ² β_{III}	Isotype 4 ² β_{IV}
VAL (351)				
LYS (352)				
THR (353)			VAL	
ALA (354)				
THR (376)				
PHE (377)				
ILE (378)				
GLY (379)				
ASN (380)				

¹Residues in the crystal structure, lying within 8 Å (normal front face) and 5 Å (bold front face) of colchicine are shown. The residue numbering follows the convention used in the crystal structure (1SA0). The residue numbering in the crystal structure differs from the sequence numbering (see Figure 5) since the crystal structure omits residue numbers 45-46 and 361- 368 to suit proper superimposition of the α and β chain in 1SA0.

²For the isotypes, a residue is shown only if it differs from the equivalent residue in the crystal structure.