CASE REPORT

Pulmonary arteriovascular malformation: a rare cause of unexplained hypoxia and acute dyspnoea in young patients

Nousheen Iqbal,¹ Karim Abdur Rehman,² Javaid Ahmed Khan,³ Tanveer Ul Haq⁴

¹Department of Medicine, Aga Khan Hospital, Karachi, Sindh, Pakistan

²Aga Khan University Hospital, Karachi, Sindh, Pakistan

³Department of Pulmonary Medicine, Aga Khan University Hospital, Karachi, Sindh, Pakistan

⁴Department of Radiology, Aga Khan University, Karachi, Sindh, Pakistan

Correspondence to

Dr Nousheen Iqbal,
nousheen.iqbal@aku.edu

Accepted 6 December 2014

SUMMARY

Pulmonary arteriovenous malformations (PAVMs) are anomalous vascular connections between arteries and veins in the lung and comprise of two types, simple and complex. PAVMs are associated with congenital conditions such as hereditary haemorrhagic telangiectasia along with acquired causes. We present a case of a 26-year-old man who presented with dyspnoea, palpitations and decreased oxygen saturation as an initial presentation of PAVM, which was treated successively with embolisation.

BACKGROUND

Pulmonary arteriovenous malformations (PAVMs) are anomalous connections between arteries and veins in the lungs supplied by the pulmonary artery in 95% of cases.¹ PAVMs are mostly unilateral and arise from the lower lung lobes. Common clinical presentations range from cyanosis, brain abscess, stroke and fatal haemoptysis.² A PAVM presenting as dyspnoea is quite rare. In our case, the patient reported to ER with predominant symptoms of dyspnoea and palpitations; workup showed polycythaemia and disproportionate hypoxia. Our literature search revealed a paucity of data consistent with the clinical presentation of our case, making it imperative to report it. Our patient did not have a diagnosis of haemorrhagic telangiectasia (HHT).

CASE PRESENTATION

A 26-year-old man presented to the Emergency Department of the Aga Khan University Hospital, Karachi with symptoms of dyspnoea on mild exertion and episodic palpitations for 2 days, lasting for 10–15 min each, and recurring and resolving spontaneously. No precipitating factors or associated symptoms could be identified. The patient was a regular smoker. There was no family history of cardiac disease and no history of nasal bleed, haemoptysis, or medical or surgical history.

Physical examination revealed oxygen saturation of 85% on room air while sitting and 88% while lying down, on pulse oximetry. No clubbing was noted. A chest and precordium examination was unremarkable.

INVESTIGATIONS

A complete blood count showed polycythaemia with haemoglobin of 19.2 g/dL and haematocrit of 55%. ECG showed pre-mature ventricular contractions. A chest X-ray showed a small well-defined

nodular opacity in right anterior fourth intercostal space, in the posteroanterior view (figure 1). Arterial-blood gases revealed hypoxia pH=7.44, PCO₂=22 mm Hg, PO₂=64 mm Hg with oxygen saturations=93% on 0.28% FiO₂ with an A-a gradient of 108. Echocardiography was normal. Troponins were negative. A CT of the chest with contrast was ordered to rule out pulmonary embolism or PAVM. To our surprise, the scan revealed multiple arteriovenous malformations (AVMs) in the lower lobes of the lungs bilaterally: three on the left (figure 2) and two on the right (figure 3). A nidus of AVM in right lower lobe measured 1.1 cm and the nidus of the largest AVM on left lower lobe measured 1 cm.

DIFFERENTIAL DIAGNOSIS

- ▶ Pulmonary embolism
- ▶ Valvular heart disease
- ▶ Shunts

TREATMENT

Digital subtraction angiogram was performed, which showed one major and one minor fistula on the right, and two major and one minor fistula on the left. The three major fistulas were embolised and successfully occluded with platinum detachable coils.

Figure 1 Chest X-ray posteroanterior view showing rounded opacity in right anterior fourth intercostal space.

To cite: Iqbal N, Rehman KA, Khan JA, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2014-207222

Figure 2 CT of the chest with contrast showing three arteriovenous malformations (AVMs) on the left of the lung.

OUTCOME AND FOLLOW-UP

A follow-up in the clinic showed marked improvement in the patient's presenting symptoms and maintained oxygen saturation (96%) on room air. In addition, no hypoxia or dyspnoea was observed after the patient was made to take a 6 min walk.

After the completion of treatment, follow-up in a case such as this requires evaluation of adequate oxygen delivery and signs of hypoxia. In addition, coils and feeding vessels also need to be evaluated with a repeat CT scan. Initial follow-up in uncomplicated cases is on a 3-month basis, which can subsequently be extended to 6–12 months and more until a visit to the pulmonologist is only required every 3 years unless the patient develops symptoms.³

DISCUSSION

PAVMs are fistulous connections between pulmonary artery and veins with high flow and low resistance. Around 70% of PAVMs are associated with HHT whereas only 20% of patients with HHT develop PAVM. Shunting of microemboli bypasses the lungs and may result in development of transient ischaemic attacks (TIAs), a cerebrovascular accident or a brain abscess.² The exact mechanism of the formation of PAVMs is not known, however, it is hypothesised that it is due to the failure in

resorption of vascular septa.³ A review of the Mayo Clinic experience reported a morbidity of 26–33% and mortality of 8–16% of patients when left untreated.²

Patients with PAVM lesions of >2 mm are usually well compensated and are asymptomatic despite impaired gas exchange. The slow progression of the disease allows the body's compensatory mechanisms to respond adequately: polycythaemia⁴ and increased cardiac output help the body to transport adequate oxygen to the tissues.⁵ Dyspnoea is a rare presentation of this disease. For patients with PAVM who present with dyspnoea, other causes should be considered. Our patient had predominant symptoms of dyspnoea and palpitation, which brought him to the hospital. He had an adequate polycythaemic response, and all other investigations for cardiopulmonary disease were negative.

Why our patient decompensated with palpitations and dyspnoea at an early age despite no comorbidities is not known, but he significantly improved postembolisation. We present this case to demonstrate that some patients with PAVMs do not cope well with the compensation required with right to left shunts, and the mechanism is still unclear.

Learning points

- ▶ Dyspnoea is a rare presentation of pulmonary arteriovenous malformations (PAVMs).
- ▶ Clinical vigilance should be maintained in a patient presenting with dyspnoea in any age group, as in this case.
- ▶ The combination of disproportionate hypoxia, polycythaemia and chest X-ray findings of nodular opacity are important indicators of PAVMs.
- ▶ In cases where haemoptysis, dyspnoea or hypoxia are unexplained, PAVM should be listed as a differential and diagnostic tests for it should be undergone.

Contributors NI wrote the manuscript and review the literature. KAR made the literature search and helped in writing the manuscript. JAK supervised and reviewed the manuscript. TUH performed the intervention on the patient.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 White RI Jr, Mitchell SE, Barth KH, *et al*. Angioarchitecture of pulmonary arteriovenous malformations: an important consideration before embolotherapy. *AJR Am J Roentgenol* 1983;140:681–6.
- 2 Swanson KL, Prakash U, Stanson AW. eds. *Pulmonary arteriovenous fistulas: Mayo Clinic experience, 1982–1997*. Mayo Clinic Proceedings. Elsevier, 1999.
- 3 Meek ME, Meek JC, Beheshti MV. eds. *Management of Pulmonary Arteriovenous Malformations. Seminars in interventional radiology*. Thieme Medical Publishers.
- 4 Santhirapala V, Williams LC, Tighe HC, *et al*. Arterial oxygen content is precisely maintained by graded erythrocytotic responses in settings of high/normal serum iron levels, and predicts exercise capacity: an observational study of hypoxaemic patients with pulmonary arteriovenous malformations. *PLoS ONE* 2014;9:e90777.
- 5 Howard LS, Santhirapala V, Murphy K, *et al*. Cardiopulmonary exercise testing demonstrates maintenance of exercise capacity in patients with hypoxemia and pulmonary arteriovenous malformations. *Chest* 2014;146:709–18.

Figure 3 CT of the chest showing two arteriovenous malformations (AVMs) on the right of the lung.

Copyright 2014 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow