Published in final edited form as:

Biochem Biophys Res Commun. 2015 January 2; 456(1): 213–218. doi:10.1016/j.bbrc.2014.11.061.

Characterization of the minimal length of functional SecA in *Escherichia coli*

Bing Na, Zhipeng You, Hsiuchin Yang, and Phang C. Tai*

Department of Biology, Center of Biotechnology and Drug Design, Georgia State University, Atlanta, Georgia 30303

Abstract

Previous studies showed that certain regions of *E. coli* SecA can be deleted from its N- and/or C-termini to complement a SecA *amber ts* mutant. In this study, we determined and characterized the dispensability of both ends of SecA molecules. With N-terminal intact or 9-aa deleted, 826aa (SecA₁₋₈₂₆ and SecA₁₀₋₈₂₆ respectively) is the minimum for complementation activity, while with N-terminus deleted by 2-21aa, SecA₂₂₋₈₂₉ is the minimum. Further deletion at the C-terminus of SecA₁₋₈₂₆/SecA₁₀₋₈₂₆/SecA₂₂₋₈₂₉ abolished the complementation activity in the cells. A hydrophobic amino acid is required for the 826th residue in the minimal-length SecAs. Chemical crosslinking and gel filtration result showed that both purified SecA₂₂₋₈₂₈ and SecA₂₂₋₈₂₉ could form a dimer. Moreover, the *in vitro* ATPase and protein translocation activities of SecA₂₂₋₈₂₈ and SecA₂₂₋₈₂₉ were similar, though lower than wild-type SecA. The active mutants had more truncated SecA in soluble than membrane-bound form, but was more stably embedded in membranes. In contrast, the inactive mutants tended to have truncated SecA more membrane-bound than soluble form, and were more loosely bound and easily chased out. Thus, the loss of complementation appears to be related to their altered subcellular localization and stability in the membranes. This study defines the substantial regions of N- and C-termini of SecA that may be deleted without losing complementation activity.

Keywords

E. coli SecA; N-/C-terminal deletions; complementation; translocation; membrane-binding

1. Introduction

The majority of proteins destined for the periplasm or outer membrane in *Escherichia coli* are translocated by the general Sec secretion pathway [1-4]. The bacterial Sec translocase has been extensively studied in the Gram-negative bacterium *E. coli*. The core of the Sec

© 2014 Elsevier Inc. All rights reserved.

* Corresponding Author: Phang C. Tai 161 Jesse Hill Jr. Drive, 592 Petit Science Center Department of Biology Georgia State University Atlanta, GA 30303 Telephone: 404-413-5303 Fax: 404-413-5301 biopct@gsu.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

translcon consists of SecA ATPase that acts as a molecular motor, and a heterotrimeric integral membrane protein complex composed of SecYEG [5-7]. SecB is a molecular chaperone that targets preproteins [2,8] to SecA. Other Sec components, SecD, SecE, YajC and YidC, facilitate protein translocation [9,10].

SecA is an essential component of bacterial Sec translocase that interacts with several translocation components through different regions: ATP [11], preprotein [12], SecYEG [13], SecB [14], lipids [15] and RNA [16]. SecA consists of two separable soluble functional domains: the N-terminal 68 k-Da domain that possesses high ATPase activity, and the C-terminal 34 k-Da domain that down-regulates the ATPase activity and is involved in dimerization and SecB/lipid binding [17,18]. The N-68 domain contains a nucleotide binding domain (NBD) [11] and an intramolecular regulator of ATP hydrolysis (IRA2) [19]. SecA recognizes different substrates through the ligands specificity in the N-68 domain [20]. Biochemical and biophysical studies have shown that native SecA forms stable dimer in solution [21-24]. While in solution, SecA undergoes monomer-dimer equilibrium reactions, and the oligomeric state of SecA depends on temperature, ionic strength and protein concentration [8]. The crystal structures of *E. coli* SecA show that SecA formed a dimer in solution. SecA dimerization has been attributed to the C-terminal domain as well as N68-domain [25].

It has been reported that SecA deleted of N-terminal 8-amino acids, though lacking interaction with SecE, is functional in cells [26], and deleted of certain both N-/C-termini retains complementation activity for a SecA *ts* mutant [24,27-29]. We previously identified two specific N-terminal domains that are crucial for initial SecA/membrane interactions [30]. The first small helix, the linker and part of the second helix (2-22) were found to be dispensable for SecA activity in complementing the growth of the SecA *ts* mutant. However, deletions of N-terminal aminoacyl residues #23-25 resulted in severe progressive retardation of growth. Moreover, a decrease of SecA activity caused by N-terminal deletions correlated to the loss of SecA membrane binding, formation of lipid-specific domains and channel activity [30]. Here we further investigated the roles of the C-terminal residues of SecA with or without N-terminal deletions. We identified the regions of N- and/or C-termini that can be deleted without losing complementation activity in *E. coli*.

2. Materials and methods

2.1. Media, bacterial strains and plasmids

Culture medium TAG containing 10 g/L of Tryptone, 5 g/L of NaCl, 40 mM potassium phosphate buffer (pH 7.0), 7.6 mM ammonium sulfate, 1.6 mM sodium citrate with 0.5% glucose was used as liquid and solid (with 1.5% agar) medium. Where appropriated, 100 µg/ml of ampicillin was added.

The bacterial strains and plasmids used in this work are listed in Table 1. *E. coli* strain DH5α was used as a host for plasmid construction and SecA *ts/amber* mutant BL21.19 (secA13(Am) supF(Ts) trp(An) zch::Tn10 recA::CAT claA::KAN) for complementation without over-expression and induced for protein over-expression [31].

Plating efficiency of colony forming units (CFU) at 30 °C and 42 °C was as described [24,28,30].

2.2. Overexpression and purification of mutant SecAs

BL21.19 cultures containing plasmids were grown at 30°C in TAG medium until OD₆₀₀ 0.8 and then were induced by 0.5 mM IPTG at 20°C overnight. The cell culture was collected by centrifugation. SecA was purified according to procedures previously described [32]. The purification of SecA₂₂₋₈₂₈ and SecA₂₂₋₈₂₉ was similar to that of SecA except that they were eluted on SP-Sepharose column by 100 mM NaCl at pH 6.1 which separated well from wild-type SecA which was eluted at higher NaCl concentration.

2.3. Separation of cytosol and membrane SecA

BL21.19/pET5a-SecA₂₂₋₈₂₈ and BL21.19/pET5a-SecA₂₂₋₈₂₉ were grown in TAG/Amp medium at 42°C to deplete wild-type SecA until BL21.19/pET5a-SecA₂₂₋₈₂₈ stopped growing. One ml of cell culture with OD₆₀₀ 1.0 was harvested, and the pellet was re-suspended in 0.5 ml of 30 mM Tris-HCl, pH 8.0 containing 1 mM dithiothreitol (DTT), and 50 µg lysozyme in 10 mM EDTA (pH 8.0). The samples were treated by freezing-thawing 3 times to lyse the cells. MgCl₂ (10 mM) and 6 µg DNase I were added. The mixtures were sonicated in ice-water bath, and centrifuged to remove unbroken cells, followed by centrifugation to pellet membranes. The membranes were dissolved in the same volume as cytosol. SecAs were detected by immunoblotting and quantified by Bio-Rad GS710 densitometer.

2.4 In vitro chase of [³⁵S]-SecA in membrane

[³⁵S]-Met labeled SecA₂₂₋₈₂₈ and SecA₂₂₋₈₂₉ membranes were prepared from strains grown in TAG medium with Amp at 30°C. 0.5 mM of IPTG was added to the cells when OD₆₀₀ reached 0.8. After 90 min, the cells were labeled with 5 mCi [³⁵S]-Met for 10 min, harvested, washed and resuspended in 30 mM Tris-HCl, pH 8.0 containing 1 mM DDT. The cytosol and membrane fractions were separated as described [33]. [³⁵S]-labeled-membranes (0.01A₂₆₀) were incubated at 37°C for 15 min with respective purified nonradioactive SecA₂₂₋₈₂₈ or SecA₂₂₋₈₂₉. The membranes were collected by centrifugation and the SecA was quantified as above.

2.5. Chemical crosslinking of SecA truncates and other in vitro activity assays

Purified SecAs were cross-linked as described [25]. Reaction mixtures (20 µl) contained: 20 µg purified SecAs, 1 mM 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide (EDC) in 0.1 M 2-Morpholinoethanesulfonic acid, monohydrate (MES) pH 6.4. Samples were separated by 5 % SDS-PAGE and detected immunologically. Three levels activities of ATPase (intrinsic, membrane and translocation) were assayed as described [34]. *In vitro* translocation of [³⁵S]-Met-labeled proOmpA was as described [32].

2.6. Reagents and chemicals

All reagents and are reagent grades and obtained from Sigma unless otherwise indicated.

3. Results and discussion

3.1. Maximal N- and C-terminal deletions of SecA to retain complementation activity

Based on our previous work on the deletion of N-termini, we further define the minimal length of N-/C-termini of active SecA for complementation [30]. Wild-type SecA (901 residues) was deleted from its N- and/or C- termini and tested for complementing *ts* strain BL21.19 at 42°C (Table 1).

With SecA N-terminal SecA₂₋₉, 826th amino acid is the minimal length at the C-terminal for complementation activity, identical to intact SecA; the deletion of first 9 SecA-aminoacyl-residues have no effect on complementation activity. With SecA₂₋₂₁, C-terminal 829th residue is the minimum (Table 1). SecA_{1-825/10-825} and SecA₂₂₋₈₂₈ completely lost their complementation activity, as reflected in the difference of the colony forming units (Fig. 1A). Therefore, the minimal length at the SecA C-terminus to complement can vary, depending on the length of the N-terminal region. It is interesting to note that the first 9 aminoacyl residues (SecA₂₋₉) have no effect on SecA function, and that the additional deletion of N-terminal 12 residues SecA₂₋₂₁ can be compensated by C-terminal three residues (from 826 to 829). Previously, N-/C-truncated SecAs have been shown to be active, or inactive, depending on the constructs, and the level of protein expression in the cells [24,28,29,35].

3.2. The importance of 826th aminoacyl residue in the hydrophobic side of α -helix

Since the active and inactive truncated SecA in our constructs differ only in one amino acid (SecA_{1-825/1-826}, SecA_{10-825/10-826} and SecA_{22-828/22-829}), amino acid type might be important for SecA function. To test this possibility, the last amino acid was mutagenized and tested for complementation (Table 1).

For SecA₁₋₈₂₆-SecA₁₀₋₈₂₆, mutating Leu⁸²⁶ to hydrophobic Ile, Phe, Met and Val retained complementation while mutations to hydrophilic Ser, Glu, Gln, Arg, or Lys lost complementation (Table 1). Mutations (L826G/L826P) breaking α -helix could not complement either. SecA₁₋₈₂₅-SecA₁₀₋₈₂₅ could not complement regardless whether Thr⁸²⁵ was mutagenized into hydrophobic or hydrophilic residues (Table 1), indicating that SecA₁₀₋₈₂₆ is the minimal length of SecA, and the first 9 amino acids were dispensable regardless whether C-terminus is intact.

For SecA₁₋₈₂₆, Val⁸²⁹₂₂₋₈₂₉ could be replaced without losing complementation activity (Table 1). SecA₂₂₋₈₂₈ or mutating Arg⁸²⁸ could not complement (Table 1), indicating that 22-829aa is the minimal length for complementation. Leu⁸²⁶ of SecA₂₂₋₈₂₉ can be replaced with hydrophobic residues (SecA₂₂₋₈₂₉L826M and SecA₂₂₋₈₂₉L826V), but not hydrophilic residues (SecA₂₂₋₈₂₉L826E and SecA₂₂₋₈₂₉L826R) for complementation (Table 1). Hence, hydrophobic 826th residue is required for the minimal length SecAs to complement BL21.19. The reported crystal structure of *E. coli* SecA lacks several residues [36]. Modeling supplemented by the crystal structures of *Bacillus subtilis* [37] and *Mycobacterium tuberculosis* SecA [38-40] using SWISS-MODEL server [41-43] shows overall structure of *E. coli* SecA (SFig. 1A), similar to *B. subtilis* and *M. tuberculosis* SecA. The hydrophobic Leu⁸²⁶ was located in a long α -helix_{810-828aa} (SFig. 1B). In the α -helix,

Phe⁸¹¹, Leu⁸¹⁵, Leu⁸¹⁸, Val⁸²² and Leu⁸²⁶ appear every 3-4 amino acids forming one turn of the α -helix, and compose a hydrophobic side (SFig.1 B). It is conceivable that a hydrophobic residue is required at the 826th position to maintain the hydrophobicity and the α -helix integrity. Lys⁸²⁸ is not located in the hydrophobic side, so it could be changed to hydrophobic or hydrophilic amino acids without impairing complementation activity as long as Val⁸²⁹ is intact. On the other hand, Val⁸²⁹ is located outside of the α -helix_{810-828aa}. Therefore, the change of Val⁸²⁹ will not affect the structure of the α -helix_{810-828aa} and the hydrophobic side, thus having no effect on the complementation activity.

The α -helix_{810-828aa} forms a Helix-Turn-Helix structure with α -helix_{762-782aa} [44] (SFig.1 C), which was proposed to function as an intramolecular regulator of ATP hydrolysis (IRA1) [45]. Like α -helix_{810-828aa}, α -helix_{762-782aa} also forms a hydrophobic side (SFig. 1B), which interacts with the hydrophobic side of α -helix_{799-828aa} (SFig.1 B, C). These two α -helices further interact with the hydrophobic C-terminal portion (651-667aa) of the α -helix_{621-667aa} (Fig.1C). This long α -helix_{621-667aa} stretches throughout SecA (SFig.1A) with one end interacting with NBD and the other end with IRA1 regulatory domain, enabling the communication between the functional and regulatory domains [19].

3.3. ATPase and translocation activity do not account for complementation

To better understand the mechanism underlying the complementation ability of the truncated SecA, the complementation active and non-active SecA proteins were overexpressed and purified. Since SecA_{1-825/826} failed to be overexpressed in our hands, we purified SecA₂₂₋₈₂₈ and SecA₂₂₋₈₂₉ and used them as representatives of the complementation non-active and active minimal length SecA proteins, respectively. The intrinsic-, membrane- and translocation-ATPase levels of the two complementation-active and non-active truncated SecAs were similar, though greatly reduced comparing to wild-type (data not shown), indicating that high ATPase activity is not obligatory for cell viability, as shown previously [30].

We next compared the protein translocation activity of SecA₂₂₋₈₂₈ and SecA₂₂₋₈₂₉ (Fig. 1B). The two mutant proteins, although one supported cell viability and the other did not (Fig. 1A), showed similar impaired protein translocation activity, much lower than wild-type SecA (Fig. 1B). Thus, high protein translocation activity seems not essential for cell viability either [30].

Taken together, the data suggest that *in vitro* ATPase and protein translocation activities, although important, are not directly correlated to cell viability, at least in the complementation ability under the conditions used. As suggested earlier [30], SecA needs not run at the maximal capacity *in vivo* to support the cell viability. The results suggest that the *in vitro* ATPase and protein translocation activities, albeit routinely used as criteria in evaluating SecA activity, do not necessarily represent the reflect SecA functions of the protein *in vivo*,

3.4. Dimer formation of truncated SecAs

SecA was found in equilibrium of monomer and dimer in solution [8] and exists as a dimer [25]. The C-terminal domain of SecA is involved in dimerization [22,25]. It has been reported that SecA₁₁₋₈₃₁ functions as a monomer [29] but has been disputed [24,28,46]. Deletion of the SecA C-terminus may affect its dimerization ability, thus impairing its complementation ability. We therefore examined the oligomeric state of the truncated SecAs by gel filtration and by chemical crosslinking. Both SecA₂₂₋₈₂₈ and SecA₂₂₋₈₂₉ were eluted in the size exclusion chromatography at around 160 kDa which is the size of dimers (Fig. 2 A). Similar to previous data [46], when wild-type SecA was treated with EDC [22], several high molecular weight bands were observed at around 200 kDa and above (Fig. 2 B, lanes 2-4), representing different crosslinking species of SecA. The band intensity increased with longer incubation time with EDC. A similar pattern of the high molecular weight bands were observed when SecA₂₂₋₈₂₈ and SecA₂₂₋₈₂₉ were treated with EDC (Fig. 2 B, lanes 6-8 and 10-12), suggesting different crosslinking sites in the molecules. The crosslinking products of SecA₂₂₋₈₂₈ and SecA₂₂₋₈₂₉ were observed even when their concentration was low (data not shown). The gel filtration and crosslinking results indicate that substantial N-/C-terminal deletions did not affect SecA dimerization, and dimerization alone did not affect complementation activity of truncated SecA.

Previous study showed that the region of SecA from 675-831aa was involved in dimer formation [25], and that the four cysteine residues in SecA were not involved in dimerization [47]. SecA may form dimer via non-covalent bonds, like salt bridge, hydrophobic interaction and van der Waals' interaction, but probably not by one disulfide bond. One amino acid loss is less likely to result in significant change in these non-covalent interactions. Since both SecA₂₂₋₈₂₈ and SecA₂₂₋₈₂₉ form dimers, the loss of complementation activity is unlikely due to loss of dimerization.

3.5. The difference of subcellular distribution and stability of SecA truncates in the membranes

Since SecA is involved in lipid binding, and functions in the membrane [15], we examined their subcellular localization. Wild-type SecA distributes equally between cytosol and membrane fractions, while SecA51(Ts) mutant protein localizes in favor of the membrane-bound form [31]. When SecA was deleted from N- and/or C-termini, the complementation active derivatives of SecA, SecA₂₂₋₉₀₁, SecA₂₂₋₈₃₁ and SecA₂₂₋₈₂₉, had increased ratio of cytosol form, which was 60-70% of their total proteins (Fig. 3A, data not shown). On the contrary, the complementation inactive SecA₂₂₋₈₂₈, had significantly reduced cytosol form, which is about 30% of its total proteins. Similar observation was found in other active and inactive C-terminal truncated SecA mutants, SecA₁₋₈₂₈, SecA₁₋₈₂₆ and SecA₁₋₈₂₅ (Fig. 3A, data not shown). Since SecA functions in the membrane, and inactive SecA (SecA₁₋₈₂₅ and SecA₂₂₋₈₂₈) had more membrane-bound form, we determined whether these forms have normal conformations that are embedded in the membrane and resistant to urea treatment [32]. The membranes containing SecA₁₋₈₂₅ or SecA₁₋₈₂₆ were treated with 6 M urea: Urea-washed SecA₁₋₈₂₅ membrane contained less than 40% of total SecA₁₋₈₂₅ (Fig. 3B lanes 1 and 2), while more than 90% of SecA₁₋₈₂₆ remained in the membrane (lanes 3 and 4) under same conditions. These results indicated that most SecA₁₋₈₂₆ integrated properly into the

membrane [32]. In contrast, SecA₁₋₈₂₅ only loosely bound to membranes. To substantiate the result, we determined the stability of SecA₂₂₋₈₂₈ and SecA₂₂₋₈₂₉ on membrane. The membranes of BL21.19/pET5a-SecA₂₂₋₈₂₈ and BL21.19/pET5a-SecA₂₂₋₈₂₉ labeled with [³⁵S]-Met were chased by increasing amount of respective purified unlabeled SecA₂₂₋₈₂₈ and SecA₂₂₋₈₂₉. The amount of labeled SecA₂₂₋₈₂₈ and SecA₂₂₋₈₂₉ on the membranes after chase was quantified (Fig. 3C). Only about 10% of labeled SecA₂₂₋₈₂₉ was chased off by 10 µg of unlabeled SecA₂₂₋₈₂₉, indicating SecA₂₂₋₈₂₉ was stably embedded in the membrane. In contrast, about 70% of SecA₂₂₋₈₂₈ was chased out by 10 µg of unlabeled SecA₂₂₋₈₂₈, indicating its instability in membranes. These data suggested that the nature of the binding of inactive SecA₂₂₋₈₂₈ to the membranes differ significantly to that of its active counterpart SecA₂₂₋₈₂₉.

The localization and integration of SecA in the membrane is critical for SecA functions. Membrane is the site where SecA functions. On the other hand, membrane is the site of the limiting biochemical defect in SecA51 (Ts) mutant protein [31]. The inactive mutant (SecA₂₂₋₈₂₈) with changed conformation is enriched in membranes in cells. The loss of the C-terminal lipid binding site may weaken its interaction with membranes. On the contrary, SecA₂₂₋₈₂₉ is active *in vivo* even with lowered ATPase and translocation activity *in vitro*. SecA₂₂₋₈₂₉ has slightly reduced membrane-bound form (Fig. 3B) but stably integrated into the membrane (Fig. 3C).

In summary, we define the dispensable regions of N- and/or C- termini SecA that could be deleted without losing complementation activity in the cells. The minimum lengths of functional *E. coli* SecA are 10-826aa and 22-829aa for active complementation. The conformation and stability of SecA in the membrane appear to contribute to the *in vivo* activity.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work was supported in part by a NIH grant GM34766. NB was a fellow of the Molecular Basis of Diseases Program at Georgia State University. The Georgia State University Biology Core Facilities were partly supported by Georgia Research Alliance, and Center of Biotechnology and Drug Design.

References

1. Manting EH, Driessen AJ. Escherichia coli translocase: the unravelling of a molecular machine. *Mol Microbiol.* 2000; 37:226–238. [PubMed: 10931320]
2. Does, C.v.d.; Keyzer, J.d.; Laan, M.v.d.; Driessen, AJM. *Methods in Enzymology.* 87-98: 2003. Reconstitution of purified bacterial preprotein translocase in liposomes.
3. Mori H, Ito K. The Sec protein-translocation pathway. *Trends Microbiol.* 2001; 9:494–500. [PubMed: 11597451]
4. Driessen AJ, Manting EH, van der Does C. The structural basis of protein targeting and translocation in bacteria. *Nat Struct Biol.* 2001; 8:492–498. [PubMed: 11373615]
5. Collinson I. The structure of the bacterial protein translocation complex SecYEG. *Biochem Soc Trans.* 2005; 33:1225–1230. [PubMed: 16246086]

6. Breyton C, Haase W, Rapoport TA, Kuhlbrandt W, Collinson I. Three-dimensional structure of the bacterial protein-translocation complex SecYEG. *Nature*. 2002; 418:662–665. [PubMed: 12167867]
7. Veenendaal AK, van der Does C, Driessen AJ. The protein-conducting channel SecYEG. *Biochim Biophys Acta*. 2004; 1694:81–95. [PubMed: 15546659]
8. Woodbury RL, Hardy SJ, Randall LL. Complex behavior in solution of homodimeric SecA. *Protein Sci*. 2002; 11:875–882. [PubMed: 11910030]
9. Nishiyama K. Function of YidC in the protein insertion into the cytoplasmic membrane of *E. coli*. *Tanpakushitsu Kakusan Koso*. 2004; 49:959–960. [PubMed: 15168502]
10. Duong F, Wickner W. The SecDFyajC domain of preprotein translocase controls preprotein movement by regulating SecA membrane cycling. *Embo J*. 1997; 16:4871–4879. [PubMed: 9305629]
11. Mitchell C, Oliver D. Two distinct ATP-binding domains are needed to promote protein export by *Escherichia coli* SecA ATPase. *Mol Microbiol*. 1993; 10:483–497. [PubMed: 7968527]
12. Chou YT, Gierasch LM. The conformation of a signal peptide bound by *Escherichia coli* preprotein translocase SecA. *J Biol Chem*. 2005; 280:32753–32760. [PubMed: 16046390]
13. Duong F. Binding, activation and dissociation of the dimeric SecA ATPase at the dimeric SecYEG translocase. *Embo J*. 2003; 22:4375–4384. [PubMed: 12941690]
14. Fekkes P, de Wit JG, Boorsma A, Friesen RH, Driessen AJ. Zinc stabilizes the SecB binding site of SecA. *Biochemistry*. 1999; 38:5111–5116. [PubMed: 10213615]
15. Breukink E, Nouwen N, van Raalte A, Mizushima S, Tommassen J, de Kruijff B. The C terminus of SecA is involved in both lipid binding and SecB binding. *J Biol Chem*. 1995; 270:7902–7907. [PubMed: 7713885]
16. Park SK, Kim DW, Choe J, Kim H. RNA helicase activity of *Escherichia coli* SecA protein. *Biochem Biophys Res Commun*. 1997; 235:593–597. [PubMed: 9207202]
17. Dapic V, Oliver D. Distinct membrane binding properties of N- and C-terminal domains of *Escherichia coli* SecA ATPase. *J Biol Chem*. 2000; 275:25000–25007. [PubMed: 10835419]
18. Nakatogawa H, Mori H, Ito K. Two independent mechanisms down-regulate the intrinsic SecA ATPase activity. *J Biol Chem*. 2000; 275:33209–33212. [PubMed: 10982786]
19. Sianidis G, Karamanou S, Vrontou E, Boulias K, Repanas K, Kyripides N, Politou AS, Economou A. Cross-talk between catalytic and regulatory elements in a DEAD motor domain is essential for SecA function. *Embo J*. 2001; 20:961–970. [PubMed: 11230120]
20. Papanikou E, Karamanou S, Baud C, Frank M, Sianidis G, Keramissanou D, Kalodimos CG, Kuhn A, Economou A. Identification of the preprotein binding domain of SecA. *J Biol Chem*. 2005
21. Driessen AJ. SecA, the peripheral subunit of the *Escherichia coli* precursor protein translocase, is functional as a dimer. *Biochemistry*. 1993; 32:13190–13197. [PubMed: 8241173]
22. Akita M, Shinkai A, Matsuyama S, Mizushima S. SecA, an essential component of the secretory machinery of *Escherichia coli*. exists as homodimer, *Biochem Biophys Res Commun*. 1991; 174:211–216.
23. Shinkai A, Akita M, Matsuyama S, Mizushima S. Quantitative renaturation from a guanidinedenatured state of the SecA dimer, a 200 KDa protein involved in protein secretion in *Escherichia coli*. *Biochem Biophys Res Commun*. 1990; 172:1217–1223. [PubMed: 2147102]
24. Wang H, Na B, Yang H, Tai PC. Additional in vitro and in vivo evidence for SecA functioning as dimers in the membrane: dissociation into monomers is not essential for protein translocation in *Escherichia coli*. *J Bacteriol*. 2008; 190:1413–1418. [PubMed: 18065528]
25. Hirano M, Matsuyama S, Tokuda H. The carboxyl-terminal region is essential for Sec-A dimerization. *Biochem Biophys Res Commun*. 1996; 229:90–95. [PubMed: 8954088]
26. Mori H, Sugiyama H, Yamanaka M, Sato K, Tagaya M, Mizushima S. Amino-terminal region of SecA is involved in the function of SecE for protein translocation into *Escherichia coli* membrane vesicles. *J Biochem (Tokyo)*. 1998; 124:122–129. [PubMed: 9644254]
27. Mori H, Sugiyama H, Yamanaka M, Sato K, Tagaya M, Mizushima S. Amino-terminal region of SecA is involved in the function of SecG for protein translocation into *Escherichia coli* membrane vesicles. *J Biochem*. 1998; 124:122–129. [PubMed: 9644254]

28. Das S, Stivison E, Folta-Stogniew E, Oliver D. Reexamination of the role of the amino terminus of SecA in promoting its dimerization and functional state. *J Bacteriol.* 2008; 190:7302–7307. [PubMed: 18723626]
29. Or E, Boyd D, Gon S, Beckwith J, Rapoport T. The bacterial ATPase SecA functions as a monomer in protein translocation. *The Journal of biological chemistry.* 2005; 280:9097–9105. [PubMed: 15618215]
30. Floyd JH, You Z, Hsieh YH, Ma Y, Yang H, Tai PC. The dispensability and requirement of SecA N-terminal aminoacyl residues for complementation, membrane binding, lipid-specific domains and channel activities. *Biochem Biophys Res Commun.* 2014
31. Cabelli RJ, Dolan KM, Qian LP, Oliver DB. Characterization of membrane-associated and soluble states of SecA protein from wild-type and SecA51(TS) mutant strains of *Escherichia coli*. *J Biol Chem.* 1991; 266:24420–24427. [PubMed: 1837021]
32. Chen X, Xu H, Tai PC. A significant fraction of functional SecA is permanently embedded in the membrane. SecA cycling on and off the membrane is not essential during protein translocation. *J Biol Chem.* 1996; 271:29698–29706. [PubMed: 8939903]
33. Tai PC, Tian G, Xu H, Lian JP, Yu JN. In vitro protein translocation into *Escherichia coli* inverted membrane vesicles. *Methods Cell Biol.* 1991; 34:167–187. [PubMed: 1943799]
34. Lill R, Dowhan W, Wickner W. The ATPase activity of SecA is regulated by acidic phospholipids, SecY, and the leader and mature domains of precursor proteins. *Cell.* 1990; 60:271–280. [PubMed: 2153463]
35. Karamanou S, Sianidis G, Gouridis G, Pozidis C, Papanikolaou Y, Papanikou E, Economou A. *Escherichia coli* SecA truncated at its termini is functional and dimeric. *FEBS Lett.* 2005; 579:1267–1271. [PubMed: 15710424]
36. Papanikolaou Y, Papadovasilaki M, Ravelli RB, McCarthy AA, Cusack S, Economou A, Petratos K. Structure of dimeric SecA, the *Escherichia coli* preprotein translocase motor. *J Mol Biol.* 2007; 366:1545–1557. [PubMed: 17229438]
37. Hunt JF, Weinkauff S, Henry L, Fak JJ, McNicholas P, Oliver DB, Deisenhofer J. Nucleotide control of interdomain interactions in the conformational reaction cycle of SecA. *Science.* 2002; 297:2018–2026. [PubMed: 12242434]
38. Owens MU, Schmidt MG, King CH, Quinn FD. Identification of export proteins from *Mycobacterium tuberculosis* that interact with SecA. *Ann N Y Acad Sci.* 1996; 797:240–241. [PubMed: 8993366]
39. Owens MU, Swords WE, Schmidt MG, King CH, Quinn FD. Cloning, expression, and functional characterization of the *Mycobacterium tuberculosis* secA gene. *FEMS Microbiol Lett.* 2002; 211:133–141. [PubMed: 12076803]
40. Sharma V, Arockiasamy A, Ronning DR, Savva CG, Holzenburg A, Braunstein M, Jacobs WR Jr. Sacchettini JC. Crystal structure of *Mycobacterium tuberculosis* SecA, a preprotein translocating ATPase. *Proc Natl Acad Sci U S A.* 2003; 100:2243–2248. [PubMed: 12606717]
41. Guex N, Peitsch MC. SWISS-MODEL and the Swiss-PdbViewer: an environment for comparative protein modeling. *Electrophoresis.* 1997; 18:2714–2723. [PubMed: 9504803]
42. Peitsch MC. Protein modeling by E-mail. *Bio/Technology.* 1995; 13:658–660.
43. Schwede T, Kopp J, Guex N, Peitsch MC. SWISS-MODEL: An automated protein homology-modeling server. *Nucleic Acids Res.* 2003; 31:3381–3385. [PubMed: 12824332]
44. Karamanou S, Vrontou E, Sianidis G, Baud C, Roos T, Kuhn A, Politou AS, Economou A. A molecular switch in SecA protein couples ATP hydrolysis to protein translocation. *Mol Microbiol.* 1999; 34:1133–1145. [PubMed: 10594836]
45. Vrontou E, Karamanou S, Baud C, Sianidis G, Economou A. Global co-ordination of protein translocation by the SecA IRA1 switch. *J Biol Chem.* 2004; 279:22490–22497. [PubMed: 15007058]
46. Wang H, Ma Y, Hsieh YH, Yang H, Li M, Wang B, Tai PC. SecAAA trimer is fully functional as SecAA dimer in the membrane: existence of higher oligomers? *Biochem Biophys Res Commun.* 2014; 447:250–254. [PubMed: 24704204]

47. Rajapandi T, Oliver D. Carboxy-terminal region of Escherichia coli SecA ATPase is important to promote its protein translocation activity in vivo. *Biochem Biophys Res Commun.* 1994; 200:1477–1483. [PubMed: 8185602]

Highlights

1. SecA retains complementation activity lacking substantial regions of N/C-termini.
2. SecA₂₋₉ is not essential; the minimal functional SecA is SecA₁₀₋₈₂₆.
3. SecA₂₂₋₈₂₉ is the minimal length for functional SecA.
4. In minimal length of functional SecA, hydrophobic 826th is required.
5. Functionality of truncated SecAs relates to the stability in membranes.

Figure 1.

Activities of truncated SecA mutants. (A) Plating efficiency of SecA and its truncated mutants as percent at 42°C/30°C. (B) *In vitro* protein translocation. Newly synthesized proOmpA was incubated with the same amount of SecA, SecA₂₂₋₈₂₉ or SecA₂₂₋₈₂₉ with SecA-depleted membranes.

Figure 2.

(A) Size chromatography. Proteins (1 mg/ml) were chromatographed on a Superose 12-10/30 column. The arrow indicates the elution peak SecA₂₂₋₈₂₈ and SecA₂₂₋₈₂₉. (B) EDC crosslinking of SecA, SecA₂₂₋₈₂₈ and SecA₂₂₋₈₂₉. Molecular weight markers were as indicated.

Figure 3.

The subcellular distribution and stability of SecA truncates. (A). SecA distributions in cytosol and membrane were detected by immunoblotting. (B). SecA₁₋₈₂₅ was loosely bound to membrane. Membranes were washed by 6 M urea (wm) or 30 mM Tris-HCl pH 7.6 (m), and SecA were analyzed. (C). Retention of [³⁵S]-SecA on membranes. [³⁵S-Met]-Membranes of SecA₂₂₋₈₂₈ and SecA₂₂₋₈₂₉ were incubated at 37°C for 15 minutes with 0-5 μg of cold purified SecA₂₂₋₈₂₈ or SecA₂₂₋₈₂₉ respectively. The membranes were collected and [³⁵S]-labeled-SecA were analyzed. [³⁵S]-labeled-SecA without cold protein was set as 100%.

TABLE 1

Summary of complementation activity of N-/C- terminal truncated SecA mutants.

(2-9) N-termi deletion	42°C	(2-21) N-term deletions	42°C
SecA ₁₀₋₈₂₉ - SecA ₁₀₋₉₀₁	+++	SecA ₂₂₋₈₂₉ - SecA ₂₂₋₉₀₁	+++
SecA ₁₀₋₈₂₈	+++	SecA ₂₂₋₈₂₈	-
SecA ₁₀₋₈₂₆	+++	SecA ₂₂₋₈₂₈ K828D,E,I,F,M,V,L	-
SecA ₁₀₋₈₂₆ L826G,S,E,R,K	-	SecA ₂₂₋₈₂₉ V829G, D,E,R,K,I,F,M,L	+++
SecA ₁₀₋₈₂₆ L826I,F,M,V	+++	SecA ₂₂₋₈₂₉ L826E,R	-
SecA ₁₀₋₈₂₈ T825M,V,L,E	-	SecA ₂₂₋₈₂₉ L826M,V	+++
SecA ₁₀₋₈₂₁	-	SecA ₂₂₋₈₂₉ K828R,H,F,V,L	+++

¶ The numbers indicate the beginning and ending amino acid of SecA variants. The (2-9) N-terminal deletions were identical in complementation efficiency as intact SecA in all tests including mutations (for simplification, data not shown).